

Estrogen-Related Receptor α (ERR α) and G Protein-Coupled Estrogen Receptor (GPER) Synergistically Indicate Poor Prognosis in Patients with Triple-Negative Breast Cancer

This article was published in the following Dove Press journal:
OncoTargets and Therapy

Shuang Ye^{1,*}
Yuanyuan Xu^{1,*}
Ling Wang¹
Kewen Zhou¹
Jiehua He²
Jiabin Lu²
Qitao Huang²
Peng Sun²
Tinghuai Wang¹

¹Department of Physiology, Zhongshan School of Medicine, Sun Yat-Sen University, Guangzhou 510080, People's Republic of China; ²State Key Laboratory of Oncology in South China, Collaborative Innovation Center for Cancer Medicine, Department of Pathology, Sun Yat-Sen University Cancer Center, Guangzhou 510060, People's Republic of China

*These authors contributed equally to this work.

Purpose: The present study aims to demonstrate the correlation between estrogen-related receptor α (ERR α) and G protein-coupled estrogen receptor (GPER) expression and its predictive role in the prognosis of patients with triple-negative breast cancer (TNBC).

Methods: A retrospective review of 199 cases of TNBC was conducted to assess the GPER and ERR α expression, and its clinicopathologic and prognostic implications. Subsequently, the effects of ERR α and GPER on cell viability, migration, and invasion induced by estrogen were also investigated in vitro.

Results: Compared to TNBCs with ERR α low expression, ERR α -high patients exhibited higher nuclear grade, more frequent lymph nodal metastasis, a higher rate of local recurrence, and distant metastasis. Survival analyses revealed that ERR α -high patients had decreased overall survival (OS), local recurrence-free survival (LRFS), and distant disease-free survival (DDFS) than ERR α -low patients. The GPER expression level positively correlated with ERR α ($R=0.167$, $P=0.18$), and TNBCs with ERR α -low/GPER-low demonstrated the best survival outcomes among groups. In vitro, E2 significantly enhanced cell viability, migration, and invasion in BT-549 and MDA-MB-231 cell lines, which was associated with the increased expression of ERR α . Moreover, the overexpression of ERR α induced by estrogen and G1 (GPER agonist) was reversed by knocking down of GPER and blocking the MAPK signaling with PD98059 in both cell lines.

Conclusion: Our findings suggest that ERR α and GPER synergistically predict unfavorable prognosis in TNBCs. Mechanically, GPER mediates the upregulation expression of ERR α induced by estrogen and promotes cell viability, migration, and invasion.

Keywords: triple-negative breast cancer, GPER, ERR α , estrogen, prognosis

Introduction

Identification of molecular subtypes of breast cancer (BC) has led to a better understanding of its biologic behavior, which provides valuable information for the individualized treatment for BCs. Triple-negative breast cancer (TNBC) is an aggressive subtype of breast cancer (BC), which is negative for estrogen receptor (ER), progesterone receptor (PR), and human epidermal growth factor receptor 2 (HER2). TNBC accounts for 15–20% of all BC, which is always poorly differentiated and predominantly occurs in young women.^{1–4} Despite the development of novel clinical therapies in TNBCs, such as Poly-ADP-ribose polymerase inhibitors

Correspondence: Peng Sun; Tinghuai Wang
Email sunpeng1@sysucc.org.cn;
wangth@mail.sysu.edu.cn

and immunotherapy, however, TNBC still demonstrates worse prognosis than non-TNBC, with a median OS of 28 months, approximately.^{5,6} Recurrence and distant metastasis are the leading causes of mortality in patients with TNBC,^{4,7,8} and recent studies suggest that younger patients with TNBC may have a higher incidence of recurrence or metastasis when compared with their older counterparts. However, the underlying mechanism is still unclear.^{9–11} One of the hypotheses is that female patients with TNBC at a younger age exhibit higher estrogen levels, suggesting that estrogen may still affect the tumor biological behavior. Given the lack of activated ER signaling in TNBC, it is generally believed that estrogen may indirectly be involved in the regulation of malignant phenotype in TNBC by activating effectors related to ER signaling or binding other receptors.

Tumorous aggressive behaviors are closely related to high-energy metabolism.^{12–15} Estrogen-related receptor α (ERR α) is an orphan nuclear receptor, which is also highly homologous with classic estrogen receptor α (ER α). ERR α expression is found in tissues requiring high energy metabolism as well as various tumorous tissues,^{16,17} which has been reported to act as a metabolic regulator *in vitro*.¹⁸ Clinicopathological studies have revealed that high expression of ERR α is associated with increased TNM stage of breast cancer, more frequent recurrence, and related poor survival outcome.^{19,20} Growing evidences indicate that the ERR family promotes cell migration and invasion by regulating the Warburg effect and metabolism-related genes^{17,21–23} Although estrogen is not a natural ligand for the ERR family, ERRs do share target genes, coregulatory proteins, ligands, and sites of action with the ERs.²⁴ Li et al²⁵ have found that physiological levels of 17 β -estradiol (E2) regulate the expression of ERR α in SKBR3 breast cancer cells which suggest that ERRs can actively influence the estrogenic response, and the pharmacological modulation of ERR family activity may be clinically valuable in estrogen-related cancer, especially in breast cancer.

On the other hand, G protein-coupled estrogen receptor (GPER) has been identified as a novel estrogen membrane receptor.²⁶ Estrogen binds to and activates GPER in the plasma membrane, and ligand-bound GPER induces activations of adenylyl cyclase, cAMP, calcium mobilization, and mitogen-activated protein kinase (MAPK).^{26,27} Previous studies by other investigators and us have revealed that GPER is commonly expressed in TNBC, and estrogen promotes cell migration and invasion in TNBC via GPER signaling.^{28–30} Clinically, TNBCs with high expression of GPER demonstrate worse OS, LRFS, and DDFS than in the GPER-low

group, especially in the premenopausal setting.^{31,32} Interestingly, a study by Li et al²⁵ also demonstrates that GPER agonists, including G1 and tamoxifen, as well as overexpression of GPER, could lead to an increase of ERR α mRNA level, suggesting that GPER/ERR α -mediated signaling may be relevant to breast cancer cells escaping inhibitory control by tamoxifen. A recent study by Kotula-Balak et al³³ also implies the existence of an interaction between GPER and ERR α in tumorous Leydig cells.

Therefore, the current study aimed to demonstrate the correlation between ERR α and GPER expression and its clinicopathologic and prognostic implications in patients with TNBC. Moreover, we also determine the role of ERR α /GPER in breast cancer cell proliferation, migration, and invasion and explore the preliminary mechanism of the interactions between GPER and ERR α *in vitro*.

Materials and Methods

Clinicopathologic Features Analysis

Clinicopathologic data of patients diagnosed as invasive breast carcinoma at the Sun Yat-sen University Cancer Center (SYSUCC) during the year 2000–2010 were retrieved. All patients were operated at SYSUCC and did not receive neoadjuvant therapy. Formalin-fixed paraffin-embedded (FFPE) tissue specimens were stained routinely with hematoxylin and eosin (H&E). ER, PR, and HER2 status were determined on immunohistochemical (IHC) staining. ER, and PR status were classified as negative using a cut-off of 1%, according to the American Society of Clinical Oncology/College of American Pathologists (ASCO/CAP) guidelines.³⁴ HER2 status was defined as negative with 0, 1+ as well as 2+ on IHC without HER2 gene amplification on fluorescence *in situ* hybridization (FISH).³⁵ The H&E and IHC slides from a total of 199 cases were retrospectively reviewed by two pathologists (PS, JH) to reconfirm the diagnosis of TNBC. Clinicopathological features, including age at diagnosis, menopausal status, laterality of tumor, nuclear grade, tumor size, lymph node metastasis, TNM stage, were analyzed. The tumor staging was based on the TNM stage was assessed according to the criteria established by the 8th edition American Joint Committee on Cancer (AJCC 8th) staging manual for breast cancer.

Immunohistochemistry Staining

FFPE tumor tissues were deparaffinized by xylene and then rehydrated by a series of graded ethanol. Antigen retrieval was conducted using EDTA buffer (GPER, pH=9.0; ERR α , pH=8.0). Subsequently, tissues were

incubated with anti-GPER primary antibody (1:100 dilution, Abcam, UK) and anti-ERR α antibody (1:100 dilution, Novus Biologicals, USA) for 50 min at 37°C. Secondary antibodies (Zhongshan Golden Bridge Biotechnology, China) were incubated for 30 min at 37°C, followed by the staining of DAB (Zhongshan Golden Bridge Biotechnology, China) and then hematoxylin. Endometrial carcinoma tissue was used as positive staining control, which has been demonstrated to express GPER and ERR α .^{36,37} Finally, the slides were mounted and semi-quantified by two experienced pathologists (PS, JH) blind to the outcome by consensus. The expression of GPER and ERR α was calculated as the product of dyeing intensity (grade as 0, negative; 1, weak; 2, moderate; and 3, strong staining) and positive-staining scope (0, <5%; 5%–24%; 25%–49%; 50%–74%; 75%–100%). For the division of high and low expression groups of GPER and ERR α , the receiver-operating characteristic (ROC) curves were plotted to generate the optimal cut-off values. The immunoreactive score of IRS=0.7 and 0.8 was used throughout the manuscript for GPER and ERR α , respectively.

Cell Culture

Human mammary epithelial cell MCF-10A and breast cancer cell lines BT-549, MDA-MB-231 were obtained from the Institute of Biochemistry and Cell Biology, Shanghai Institute of Biological Sciences (Shanghai, China). Cells were cultivated at 37°C in a humidified atmosphere with 5% CO₂. MCF-10A and BT-549 were cultured in RPMI 1640 medium (Gibco, USA), and MDA-MB-231 was cultured in Leibovitz's L-15 medium (Hyclone, USA). Both medium were supplemented with 10% FBS (Gibco, USA). Cells passaged more than five times were abandoned in our experiments. The information about associated drugs in this experiment were as follows: 17 β -estradiol (E2, MilliporeSigma, Germany), G1 (MilliporeSigma, Germany), G15 (MilliporeSigma, Germany), the MEK inhibitor PD98059 (#167869-21-8, MedChemExpress, USA). The E2 concentration used in the present study was 100nM, which had been reported to show the most significant effects according to our previous time-concentration experiments.²⁹

Protein Extraction and Western Blotting

Cells were lysed using Triton-X lysis buffer, and then the protein concentration was measured by the BCA kit (Thermo Fisher Scientific, USA). 30 μ g of protein samples

per lane was separated by polyacrylamide gel electrophoresis and then transferred onto PVDF membrane (Thermo Fisher Scientific, USA). 5% BSA (Thermo Fisher Scientific, USA) dissolved in TBST (Beijing Solarbio Science and Technology, China) was used to block the membrane. The band was incubated with the primary antibody overnight at 4°C. Primary antibodies were applied as follows: anti-GPER (1:5000 dilution, Abcam, UK), anti-ERR α (1:5000 dilution, Cell Signaling Technology, USA), anti-GAPDH (1:5000 dilution, Cell Signaling Technology, USA), and anti- β -actin (1:5000 dilution, Cell Signaling Technology). Secondary antibody (1:5000 dilution, Santa Cruz Biotechnology) incubation was carried out at room temperature for 2h. Protein bands were then detected using chemiluminescence reagent (Millipore, UK) on the ChemiDocTM imaging system (Biorad, USA) and analyzed semi-quantitatively using the Image lab software (Biorad, USA). β -actin and GAPDH were served as internal control.

RNA Extraction and Quantitative Real-Time PCR (qRT-PCR)

RNA was extracted using Trizol[®] (Thermo Fisher Scientific, USA) following the manufacturer's construction and then quantified on NanodropTM spectrophotometer (Thermo Fisher Scientific, USA). For reverse transcription, 1 μ g total RNA was added according to the PrimeScript RT reagent Kit (Takara Japan). Quantitative real-time PCR reaction was performed using the SYBR Premix ExTaqTM Kit (Takara, Japan) on the BIO-RAD CFX96 system. Cycling conditions were as follows: denaturation at 95°C for 5 min, followed by 40 cycles of 95°C for 15 s, 60°C for 30 s, 72°C for 30 s. Relative expression levels of the targeted gene were determined by the 2 $^{-\Delta\Delta C_t}$ method and normalized to GAPDH or β -actin. All the primers involved in this study were designed and synthesized by Jierui (China), and sequence information is listed in [Supplementary Table 1](#).

Wound-Healing Assay

To examine cell migration ability under different treatment conditions wound-healing assay was conducted. Briefly, 5 \times 10⁵ cells were seeded into a 6-well plate overnight to adhere. When cells reached 90–100% confluence, 10 μ L sterile micropipette tip was used to scratch one gap vertically at the bottom of the culture plate. PBS was used to wash away cellular debris, and then serum-free and phenol red-free medium was added to eliminate the interference

of E2. An inverted microscope captured the closure of the wound area after the scratch under $200\times$ magnification at pointed time. Each experiment was triplicated. The wound-healing distance of cells was analyzed using ImageJ software.

Cell Viability Assay

Cell Counting Kit-8 (CCK-8) was carried out to measure the cell viability. 1×10^3 cells were plated into a transparent 96-well plate and cultured at 37°C . After consequent drug treatment, 10 μL CCK-8 reagent (Dojindo, Japan) was added into each well and incubated for 1–4 h. The OD values at 450 nm were detected in a microplate spectrometer. Per group was set more than three replicate wells, and experiments were triplicated.

Transwell Invasion Assay

1×10^4 cells suspended in 200 μL serum-free medium were seeded in 8 μm pore sized transwell upper chamber (Corning, USA) coated with 200 $\mu\text{g}/\text{mL}$ Matrigel (BD Science, CA). The lower chamber was added 900 μL complete medium containing 10% FBS. After the incubation of 12 h, the inserts were washed with PBS and then fixed with 4% paraformaldehyde (Beijing Solarbio Science and Technology, China) for 10 min, stained by 0.1% of crystal violet solution (Beijing Solarbio Science and Technology, China) for 30 min. Cells in the upper surface were wiped out softly by cotton swabs. Random six fields per well were photographed under $200\times$ magnification and counted by ImageJ software.

Transfection of siRNA and Plasmid

Commercial siRNA of GPER and ERR α were purchased from RiboBio (Guangzhou, China), while GPER expression plasmid was synthesized by Sino Biological (Beijing, China). Transient transfection using Lipofectamine™ 3000 transfection reagent (Invitrogen, USA) was conducted under the manufacturer's construction. For siRNA, a total of 125 μL Opti-MEM (Invitrogen, USA) was used to incubate 3.75 μL Lipofectamine 3000 reagent and 50 nM siRNA for 10 min and then added into FBS-free culture medium. 6–8 h later, the culture medium was replaced with complete medium containing 10% FBS. To overexpressing GPER, the P3000 reagent was added additionally into the transfection system according to the construction. Subsequent experiments were conducted after 48 h transfection.

Statistical Analyses

Data in this study were analyzed using the SPSS version 26.0 (IBM Inc., USA) and GraphPad Prism 7.0 (GraphPad Software Inc., USA). Eligible patients were considered as the unmatched cohort. To reduce bias, we also developed a 1:1 (ERR α -low: ERR α -high) matched cohort using propensity score matching (PSM) for age, nuclear grade, tumor size, LNM, and TNM stage with a caliper of 0.03. The clinicopathological variables were compared between groups using the Chi-square test. The Pearson correlation coefficient was used to analyze the correlation between the expression of ERR α and GPER. OS, LRFS, and DDFS curves were drawn using the Kaplan–Meier methods and were compared using Log rank tests. Univariate and multivariate analyses of the patients' survival were performed using the Cox proportional hazards regression model and the logistic regression model. For experiments in vitro, the difference between groups was compared using the Student's *t*-test or one-way ANOVA. Two-tailed *p*-value <0.05 was considered statistically significant.

Results

Association Between ERR α Expression and Clinicopathologic Features in TNBCs

The baseline clinicopathological features of 199 TNBCs are summarized in [Table 1](#), of which 54.8% (109/199) was classified into ERR α -low group, while 45.2% (90/199) was classified into ERR α -high group. Examples of IHC staining for ERR α with different intensities have been shown in [Figure 1](#). Compared to ERR α -low group, ERR α -high TNBCs exhibited higher nuclear grade (grade II, 12.2% vs 23.9%; grade III, 87.8% vs 76.1%; $p=0.036$), more frequent LNM (52.2% vs 40.4%; $p=0.014$), and higher TNM stage ($p=0.007$). No significant difference was observed in age, menopausal status, tumor size, and laterality between groups. Besides, 37.7% (75/199) of TNBCs had a high expression of GPER, and these cases were predominantly in the ERR α -high group than in the ERR α -low group (50.0% vs 27.5%; $p=0.001$). The expression of GPER was positively correlated to that of ERR α with Pearson correlation coefficient of 0.167 ($p=0.018$; [Supplementary Figure 1](#)).

Prognostic Implications of ERR α in TNBCs

The patients were followed-up for 2–205 months, with a median of 113 months. Compared to ERR α -low group,

Table I Association Between ERR α Expression and Clinicopathologic Features in TNBCs

Variables	Unmatched Cohort				Matched Cohort [#]			
	All (n=199)	ERR α -Low (n=109)	ERR α -High (n=90)	p*	All (n=163)	ERR α -Low (n=73)	ERR α -High (n=90)	p*
Age at diagnosis (y)								
<45	72 (36.2)	36 (33.0)	36 (40.0)	0.353	71 (43.6)	35 (47.9)	36 (40.0)	0.370
45–55	84 (42.2)	51 (46.8)	33 (36.7)		60 (36.8)	27 (37.0)	33 (36.7)	
>55	43 (21.6)	22 (20.2)	21 (23.3)		32 (19.6)	11 (15.1)	21 (23.3)	
Menopause status								
Pre-menopause	156 (78.4)	87 (79.8)	69 (76.7)	0.591	131 (80.4)	62 (84.9)	69 (76.7)	0.262
Post-menopause	43 (21.6)	22 (20.2)	21 (23.3)		32 (19.6)	11 (15.1)	21 (23.3)	
Laterality of tumor								
Left	105 (52.8)	61 (56.0)	44 (48.9)	0.320	84 (51.5)	40 (54.8)	44 (48.9)	0.553
Right	94 (47.2)	48 (44.0)	46 (51.1)		79 (48.5)	33 (45.2)	46 (51.1)	
Tumor size (pT)								
pT1	53 (26.6)	32 (29.4)	21 (23.3)	0.083	39 (23.9)	18 (24.7)	21 (23.3)	0.469
pT2	123 (61.8)	70 (64.2)	53 (58.9)		101 (62.0)	48 (65.8)	53 (58.9)	
pT3	18 (9.0)	6 (5.5)	12 (13.3)		18 (11.0)	6 (8.2)	12 (13.3)	
pT4	5 (2.5)	1 (0.9)	4 (4.4)		5 (3.1)	1 (1.4)	4 (4.4)	
NG								
Grade II	37 (18.6)	26 (23.9)	11 (12.2)	0.036	21 (12.9)	10 (13.7)	11 (12.2)	0.964
Grade III	162 (81.4)	83 (76.1)	79 (87.8)		142 (87.1)	63 (86.3)	79 (87.8)	
LNM (pN)								
pN0	108 (54.3)	65 (59.6)	43 (47.8)	0.014	83 (50.9)	40 (54.8)	43 (47.8)	0.313
pN1	45 (22.6)	28 (25.7)	17 (18.9)		34 (20.9)	17 (23.3)	17 (18.9)	
pN2	27 (13.6)	11 (10.1)	16 (17.8)		27 (16.6)	11 (15.1)	16 (17.8)	
pN3	19 (9.5)	5 (4.6)	14 (15.6)		19 (11.7)	5 (6.8)	14 (15.6)	
TNM staging								
I	28 (14.1)	20 (18.3)	8 (8.9)	0.007	19 (11.7)	11 (15.1)	8 (8.9)	0.126
IIa	88 (44.2)	49 (45.0)	39 (43.3)		67 (41.1)	28 (28.4)	39 (43.3)	
IIb	33 (16.6)	23 (21.1)	10 (11.1)		27 (16.6)	17 (23.3)	10 (11.1)	
IIIa	27 (13.6)	11 (10.1)	16 (17.8)		27 (16.6)	11 (15.1)	16 (17.8)	
IIIb	4 (2.0)	1 (0.9)	3 (3.3)		4 (2.5)	1 (1.4)	3 (3.3)	
IIIc	19 (9.5)	5 (4.6)	14 (15.6)		19 (11.7)	5 (6.8)	14 (15.6)	
Local recurrence								
Yes	28 (14.1)	9 (8.3)	19 (21.1)	0.009	27 (16.6)	8 (11.0)	19 (21.1)	0.128
No	171 (85.9)	100 (91.7)	71 (78.9)		136 (83.4)	65 (89.0)	71 (78.9)	
Distant metastasis								
Yes	48 (24.1)	13 (11.9)	35 (38.9)	<0.001	47 (28.8)	12 (16.4)	35 (38.9)	0.003
No	151 (75.9)	96 (88.1)	55 (61.1)		116 (71.2)	61 (83.6)	55 (61.1)	
Death of tumor								
Yes	57 (28.6)	12 (11.0)	45 (50.0)	<0.001	57 (35.0)	12 (16.4)	45(50.0)	<0.001
No	142 (71.4)	97 (89.0)	45 (50.0)		106 (65.0)	61 (83.6)	45(50.0)	
GPER expression								
Low	124 (62.3)	79 (72.5)	45 (50.0)	0.001	99 (60.7)	54 (74.0)	45(50.0)	0.003
High	75 (37.7)	30 (27.5)	45 (50.0)		64 (39.3)	19 (26.0)	45(50.0)	

Notes: * χ^2 test comparing proportions among ERR α -low and ERR α -high group; [#]Propensity score matching (PSM) was performed using 1:1 nearest neighbor matching with a caliper of 0.03 to accept a matched pair.

Abbreviations: TNBC, triple-negative breast cancer; LNM, lymph node metastasis; NG, nuclear grade.




Figure 1 Immunohistochemical staining for ER α with different intensity. The staining is graded as negative (A), weak (B), moderate (C), and strong (D). Magnification, $\times 10$.

ER α -high TNBCs demonstrated increased incidences of local recurrence (21.1% vs 8.3%; $p=0.009$), distant metastasis (38.9% vs 11.9%; $p<0.001$), and more patients with high ER α died of tumor (50.0% vs 11.0%; $p<0.001$) in the unmatched cohort. Similarly, in the matched cohort, more ER α -high TNBCs died of tumor (50.0% vs 16.4%, $p<0.001$) and exhibited distant metastasis (38.9% vs 16.4%, $p=0.003$) than TNBCs with low ER α expression (Table 1). The survival curves are shown in Figure 2, ER α -high TNBCs were associated with a decrease in OS, LRFS, and DDFS when comparing to ER α -low group in both cohorts. Similar survival outcomes were observed between GPER-high and GPER-low groups (Supplementary Figure 2). Further survival analysis in the subgroups according to the co-expression status of ER α and GPER revealed that in the unmatched cohort, ER α -high/GPER-high subgroup was significantly associated with the worst survival outcomes of both OS, LRFS, and DDFS, while patients in ER α -low/GPER-

low exhibited a relative excellent survival ($p<0.001$). A decreased DDFS was observed in ER α -low/GPER-high subgroup when comparing to ER α -high/GPER-low subgroup, while the survival curve of OS or LRFS in these two groups was overlapping. In the matched cohort, ER α -low/GPER-low TNBCs had the best OS, LRFS, and DDFS, followed by ER α -high/GPER-low subgroup. ER α -low/GPER-high subgroup displayed an increased OS than patients with ER α -low/GPER-low. However, no significant difference was found between these two subgroups in LRFS and DDFS (Figure 3). A Cox proportional hazards regression model was then used to identify the biomarkers and clinicopathological factors affecting the prognosis of patients with TNBC. The results are summarized in Table 2. In the univariate model, LNM, as well as the expression of ER α and GPER, were identified as prognostic indicators for OS, LRFS, and DDFS, while nuclear grade and tumor size were also associated with OS and DDFS, but not LRFS.




Figure 2 Survival curves of OS, LRFS, and DDFS in ERR α -low and ERR α -high groups. Kaplan-Meier curves of OS (A), LRFS (B), and DDFS (C) between groups in patients with TNBC in the unmatched cohort. Kaplan-Meier curves of OS (D), LRFS (E), and DDFS (F) between groups in patients with TNBC in a matched cohort using propensity score matching for age, nuclear grade, tumor size, LNM and TNM stage with a caliper of 0.03.




Figure 3 Survival curves of OS, LRFS, and DDFS based on the co-expression status of ERR α and GPER. Kaplan-Meier curves of OS (A), LRFS (B), and DDFS (C) among groups in patients with TNBC in the unmatched cohort. Kaplan-Meier curves of OS (D), LRFS (E), and DDFS (F) among groups in patients with TNBC in a matched cohort using propensity score matching for age, nuclear grade, tumor size, LNM and TNM stage with a caliper of 0.03.

Multivariate analysis proved that LNM and the expression of ERR α and GPER were independent factors for OS, LRFS, and DDFS. Besides, tumor size was also suggested as an independent factor OS.

ERR α Mediates Malignant Phenotype of TNBC Induced by Estrogen

Our previous study suggests the overexpression of GPER is associated with poor prognosis, especially in pre-

Table 2 Univariate and Multivariate Survival Analyses for OS, LRFS and DDFS

	Univariable Model				Multivariable Model			
	p	HR	95% CI Lower	95% CI Upper	p	HR	95% CI Lower	95% CI Upper
OS								
NG	0.012	4.48	1.40	14.32	0.085	2.86	0.87	9.44
Tumor size	<0.001	2.47	1.64	3.74	0.003	1.72	1.20	2.47
LNM (negative vs positive)	<0.001	4.05	2.25	7.31	<0.001	3.41	1.87	6.24
ERR α (low vs high)	<0.001	5.66	2.99	10.70	<0.001	3.32	1.70	6.48
GPER (low vs high)	<0.001	4.42	2.53	7.73	0.001	2.80	1.56	5.02
LRFS								
LNM (negative vs positive)	0.001	3.86	1.70	8.79	0.003	3.43	1.50	7.84
ERR α (low vs high)	0.003	3.35	1.51	7.42	0.029	3.58	1.60	8.04
GPER (low vs high)	<0.001	4.45	2.01	9.86	0.002	2.46	1.09	5.52
DDFS								
NG	0.019	4.06	1.26	13.07	0.063	3.19	0.94	10.80
Tumor size	0.003	2.05	1.27	3.30	0.104	1.44	0.93	2.23
LNM (negative vs positive)	<0.001	3.39	1.84	6.25	<0.001	3.22	1.71	6.06
ERR α (low vs high)	<0.001	4.24	2.24	8.03	0.036	2.08	1.05	4.11
GPER (low vs high)	<0.001	11.42	5.34	24.44	<0.001	9.08	4.16	19.81

Abbreviations: OS, overall survival; LRFS, local recurrence-free survival; DDFS, distant disease-free survival; NG, nuclear grade; LNM, lymph node metastasis.

menopausal patients with TNBC.³¹ Thus, a χ^2 statistic of the Log rank test was used in the present study to calculate the survival discrimination based on the expression of ERR α and GPER in pre- and post-menopause patients (Table 3). Compared to post-menopausal patients, χ^2 statistic of pre-menopausal patients showed a larger and statistically significant difference between ERR α -high and ERR α -low group on OS (24.28, $p < 0.001$ vs 10.54, $p = 0.012$), LRFS (6.38, $p = 0.012$ vs 3.88, $p = 0.049$), and DDFS (18.07, $p < 0.001$ vs 4.43, $p = 0.035$). A larger χ^2 statistic between GPER-high and GPER-low group was also found on OS (27.72, $p < 0.001$ vs 4.47, $p = 0.035$), LRFS (19.17, $p < 0.001$ vs 0.16, $p = 0.685$), and DDFS (53.35, $p < 0.001$ vs 10.15, $p = 0.001$) in pre-menopausal TNBCs. These may imply the role of estrogen in the tumor progression of TNBC, affecting ERR α /GPER signaling. In vitro, treatment with 100nM E2 significantly enhanced the cell viability, migration, and invasion in both BT-549 and MDA-MB-231 cell lines as compared to the control ($p < 0.05$; Figure 4). Compared to normal breast epithelial cell lines MCF-10A, the expression of ERR α was relatively higher in BT-549 and MDA-MB-231 cells (Figure 5A). We further verify the function of ERR α in TNBC by knocking down ERR α in BT-549 and MDA-MB-231 using RNA interference. Inhibition of ERR α significantly reduced cell viability, and suppressed the wound-healing and invasion in both BT-549 and MDA-

Table 3 The χ^2 Statistic on OS, LRFS, and DDFS for Pre- and Post-Menopause TNBCs Based on ERR α and GPER Expressions (n=199)

Survival	Pre-Menopause		Post-Menopause	
	χ^2	p*	χ^2	p*
ERRα-high vs ERRα-low				
OS	24.28	<0.001	10.54	0.012
LRFS	6.38	0.012	3.88	0.049
DDFS	18.07	<0.001	4.43	0.035
GPER-high vs GPER-low				
OS	27.72	<0.001	4.47	0.035
LRFS	19.17	<0.001	0.16	0.685
DDFS	53.35	<0.001	10.15	0.001

Note: * χ^2 statistic of the Log rank test calculating the discrimination between groups in pre- and post-menopause patients.

Abbreviations: OS, overall survival; LRFS, local recurrence-free survival; DDFS, distant disease-free survival.

MB-231 cells (Figure 5B–D). Moreover, despite the fact that ERR α is not a natural receptor for estrogen, Western blot and qRT-PCR exhibited that treatment with 100nM E2 upregulated ERR α at mRNA and protein level in both cell lines (Figure 5E). We also observed that enhanced cell viability, migration, and invasion induced by E2 were inhibited after transfecting with siEER α (Figure 5B–D and F). These results indicated that ERR α 's effects on trigger the malignant phenotype of TNBC might be regulated by E2.




Figure 4 E2 enhanced cell viability, migration, and invasion in BT-549 and MDA-MB-231 cells. **(A)** Cell viability was detected using CCK-8 assay with the treatment of 100nM E2 or control. **(B)** Cell invasion was evaluated by the Transwell invasion assay with the treatment of 100nM E2 or control. The invasive cells were measured after 12h. **(C)** Cell migration was evaluated by a wound-healing assay with the treatment of 100nM E2 or control. The healing percentage was measured after 24h. ***p* < 0.01.




Figure 5 ERRα mediates the malignant phenotype of TNBC induced by estrogen. **(A)** Relative ERRα expression in normal mammary epithelial cells (MCF-10A) and TNBC cells (BT-549 and MDA-MB-231). **(B)** Cell viability was detected using CCK-8 assay with the siERRα and/or treatment of 100nM E2 in BT-549 and MDA-MB-231 cells. **(C)** and **(D)** Cell migration was evaluated by a wound-healing assay with the siERRα and/or treatment of 100nM E2 in BT-549 and MDA-MB-231 cells. The healing percentage was measured after 24h. **(E)** Relative ERRα expression in BT-549 and MDA-MB-231 cells incubated with 100nM E2 or control. **(F)** Cell invasion was evaluated by the Transwell invasion assay with the siERRα and/or treatment of 100nM E2 in BT-549 and MDA-MB-231 cells. The invasive cells were measured after 12h. **p* < 0.05, ***p* < 0.01.

GPER Intermediates the Regulation of $ERR\alpha$ by Estrogen

GPER is reported to commonly expressed in ER-negative breast cancer.^{32,38} Western blotting showed that the expression of GPER was low in MDA-MB-231 cells and relatively high in BT-549 cells (Figure 6A). Thus, we knocked down GPER in BT-549 cells using siGPER, and overexpressed GPER in MDA-MB-231 cells by transfecting plasmid, respectively, to investigate whether GPER intermediate the interaction between E2 and $ERR\alpha$. Our result revealed that both E2 and G1 (GPER agonist) significantly increased $ERR\alpha$ mRNA level as compared to control in both cell lines, especially in BT-549. This effect was suppressed in BT-549 cells by siGPER (Figure 6B). Consistently, overexpression of GPER notably increased $ERR\alpha$ mRNA level in MDA-MB-231 cells, while no change was observed with pretreatment of G15, which is regarded as a GPER antagonist (Figure 6B). Activation of the ERK1/2 MAPK signaling has been reported to be required for the activation of GPER ligands, such as E2 and G1.^{27,39} We also found that increased $ERR\alpha$ mRNA levels induced by E2 and G1 were markedly suppressed after the administration of ERK1/2 MAPK inhibitor PD98059 in both cell lines (Figure 6C).

Discussion

Estrogen receptor signaling is essential for the prognosis and treatment of ER-positive breast cancer. Application of endocrinotherapy, such as tamoxifen or toremifene, significantly improves the clinical outcome of patients with breast cancer. TNBC, which demonstrates a higher degree of malignancy and rapid progression,^{40,41} is proved to be lack of ER signaling activation. However, the clinical relevance of estrogen and TNBC remains controversial. A study by Contreras-Zárate et al found that estradiol induces BDNF/TrkB signaling in TNBC to promote brain metastases.⁴² Girgert et al suggested that 17 β -estradiol significantly increased proliferation in TNBC in vitro, along with the activation of Src, EGFR, Cyclin D1, and CREB signaling.⁴³ However, McNamara et al demonstrated that enzymes known to modulate levels of estrogens were detected in TNBC and were associated with longer disease-free survival, suggesting the presence of, and a potential protective effect of estrogens in TNBC.⁴⁴ Current findings speculate the effect of estrogen in TNBC may be mediated by other activating effectors related to ER signaling or other receptors.

Consistent with previous studies,^{19,20} we observed that high expression of $ERR\alpha$ was associated with aggressive clinicopathologic features in TNBC, including high NG, frequent LNM, and advanced TNM stage, as well as poor survival outcomes. Age and menopausal status were




Figure 6 GPER intermediates the regulation of $ERR\alpha$ by estrogen. (A) Relative GPER expression in BT-549 and MDA-MB-231 cells. (B) Relative $ERR\alpha$ expression in BT-549 and MDA-MB-231 cells with treatment of 100nM E2, G1, or G15. (C) Relative $ERR\alpha$ expression in BT-549 and MDA-MB-231 cells overexpressing or knocking down of GPER with treatment of 100nM E2, G1, or G15. (D) Effects of blocking of ERK1/2 MAPKs with PD98059 on $ERR\alpha$ expression with treatment of 100nM E2, G1, or G15. * $p < 0.05$, ** $p < 0.01$.

neither associated with $ERR\alpha$ expression nor affecting the survival of TNBCs in this cohort. However, larger discrimination in survival curves between $ERR\alpha$ -high (or GPER-high) and $ERR\alpha$ -low (or GPER-low) group was observed in premenopausal patients than postmenopausal patients, which consistent with our previous study.³¹ These findings implied that $ERR\alpha$ and GPER might mediate the tumor-promoting effect of estrogen in TNBC. Notably, we also reveal a positive correlation between $ERR\alpha$ and GPER expressions, and the co-expression status of $ERR\alpha$ /GPER could further discriminate the survival tendency, which may provide a novel estrogen-related model for predicting the prognosis of TNBCs.

$ERR\alpha$ has been reported as tumorigenic by reprogramming the energy metabolism of tumor cells, leading to the production of various biosynthetic precursors, and creating a micromovement for tumor proliferation.^{12,13} Kallen et al⁴⁵ have studied the structure of the ligand-binding domain (LBD) of $ERR\alpha$ by X-ray diffraction and found that the LBD of $ERR\alpha$ is almost entirely occupied by amino acid side chain, which is only possible to bind a ligand with less than 4 carbon atoms. Thus, estrogen is clearly not a natural ligand for $ERR\alpha$. However, our data showed that the malignant phenotypes induced by E2 in TNBC cells were significantly suppressed by $ERR\alpha$. Moreover, E2 treatment increased the protein and mRNA level of $ERR\alpha$ in vitro, suggesting the indirect interactions between E2 and $ERR\alpha$. Interestingly, studies by Li et al²⁵ and Kotula-Balak et al³³ showed that the activation of GPER lead to an increased $ERR\alpha$ level in HER2-positive breast cancer cells and tumorous Leydig cells, respectively. Similarly, we demonstrated that GPER agonists and overexpression of GPER also resulted in an increased $ERR\alpha$ level in TNBC, while E2 induced increased expression of $ERR\alpha$ was inhibited by knocking down of GPER as well as ERK1/2 MAPK signaling inhibitor. These results indicate that GPER may intermediate the regulation of $ERR\alpha$ by E2, and resulted in promoting cell proliferation, migration, and invasion in TNBC, which is consistent with the findings in clinicopathological and prognostic analyses.

The effect of estrogen in breast cancer is no longer rely on the expression of ER (predominantly $ER\alpha$). The role of other subtypes of estrogen receptors, such as $ER\beta$ ^{29,46,47} and $ER\alpha36$,^{48,49} as well as other receptors or effectors related to ER signaling, such as GPER and ERR family, are raising concern in TNBC. However, the cross-talk among different ER subtypes or effectors related to ER

signaling could be extremely complicated. The present study suggests the prognostic role of $ERR\alpha$ in TNBCs and provides a new idea for the understanding of estrogenic effects on the malignant phenotype of TNBC by $ERR\alpha$ -GPER interactions. The underlying mechanism interpreting the interaction between GPER and $ERR\alpha$ should be involved in further investigations. Moreover, given the lack of effective conventional therapies for the treatment of TNBC, Wisinski's team has attempted to conduct a Phase 2 study evaluating single-agent high-dose estradiol targeting $ER\beta$ in patients with advanced TNBC,⁵⁰ while limited efficacy was found. Promising therapeutic options targeting estrogen-related agents, such as GPER, ERR family, $ER\beta$, and $ER\alpha36$, may be warranted in further clinical trials.

Abbreviations

$ERR\alpha$, estrogen-related receptor α ; GPER, G protein-coupled estrogen receptor; TNBC, triple-negative breast cancer; E2, 17 β -estradiol; OS, overall survival; LNM, lymph nodal metastasis; NG, nuclear grade; ER, estrogen receptor; PR, progesterone receptor; HER2, human epidermal growth factor receptor 2; LRFS, local recurrence-free survival; DDFS, distant disease-free survival; FFPE, formalin fixed paraffin-embedded; H&E, haematoxylin and eosin; FISH, fluorescence in situ hybridization; IHC, immunohistochemical staining; qRT-PCR, quantitative real-time PCR; PSM, propensity score matching; ROC, receiver operating characteristic curve; MAPK, mitogen-activated protein kinase; ERK1/2, extracellular regulated kinase 1/2.

Data Sharing Statement

The datasets generated and analyzed during the current study are also available from the corresponding author (Tinghuai Wang) on reasonable request.

Funding

This study is funded by the Guangdong Medical Research Foundation (A2018241) and the National Natural Science Foundation of China (81372818, 81572585, 81902679) in the design of the study and collection, analysis, and interpretation of data and in writing the manuscript.

Ethics Approval and Consent to Participate

All procedures performed in studies involving human participants were approved by the Ethics Committee and

Research Committee of the Sun Yat-sen University Cancer Center (SYSUCC) and followed the ethical standards of the 1964 Helsinki declaration and its later amendments or comparable ethical standards. The formal written informed consent was obtained from all individual participants included in the study.

Funding

This study is funded by the Guangdong Medical Research Foundation (A2018241) and the National Natural Science Foundation of China (81372818, 81572585, 81902679) in the design of the study and collection, analysis, and interpretation of data and in writing the manuscript.

Disclosure

The authors declare that they have no competing interests for this work.

References

- Garcia Fernandez A, Gimenez N, Fraile M, et al. Survival and clinicopathological characteristics of breast cancer patient according to different tumour subtypes as determined by hormone receptor and Her2 immunohistochemistry. A single institution survey spanning 1998 to 2010. *Breast*. 2012;21:366–373. doi:10.1016/j.breast.2012.03.004
- Guney Eskiler G, Cecener G, Egeli U, Tunca B. Triple negative breast cancer: new therapeutic approaches and BRCA status. *APMIS*. 2018;126:371–379. doi:10.1111/apm.12836
- Rastelli F, Biancanelli S, Falzetta A, et al. Triple-negative breast cancer: current state of the art. *Tumori*. 2010;96:875–888. doi:10.1177/548.6505
- Sharma P. Update on the treatment of early-stage triple-negative breast cancer. *Curr Treat Options Oncol*. 2018;19:22. doi:10.1007/s11864-018-0539-8
- He J, Tsang JY, Xu X, et al. AJCC 8th edition prognostic staging provides no better discriminatory ability in prognosis than anatomical staging in triple negative breast cancer. *BMC Cancer*. 2020;20:18. doi:10.1186/s12885-019-6494-3
- Li X, Yang J, Peng L, et al. Triple-negative breast cancer has worse overall survival and cause-specific survival than non-triple-negative breast cancer. *Breast Cancer Res Treat*. 2017;161:279–287. doi:10.1007/s10549-016-4059-6
- Sihto H, Lundin J, Lundin M, et al. Breast cancer biological subtypes and protein expression predict for the preferential distant metastasis sites: a nationwide cohort study. *Breast Cancer Res*. 2011;13:R87. doi:10.1186/bcr2944
- Lin NU, Claus E, Sohl J, Razzak AR, Arnaout A, Winer EP. Sites of distant recurrence and clinical outcomes in patients with metastatic triple-negative breast cancer: high incidence of central nervous system metastases. *Cancer*. 2008;113:2638–2645. doi:10.1002/ncr.23930
- Aapro M, Wildiers H. Triple-negative breast cancer in the older population. *Ann Oncol*. 2012;23(Suppl 6):52–55. doi:10.1093/annonc/mds189
- Anderson WF, Jatoi I, Devesa SS. Distinct breast cancer incidence and prognostic patterns in the NCI's SEER program: suggesting a possible link between etiology and outcome. *Breast Cancer Res Treat*. 2005;90(2):127–137. doi:10.1007/s10549-004-3777-3
- Millikan RC, Newman B, Tse CK, et al. Epidemiology of basal-like breast cancer. *Breast Cancer Res Treat*. 2008;109:123–139. doi:10.1007/s10549-007-9632-6
- De Santis MC, Porporato PE, Martini M, Morandi A. Signaling pathways regulating redox balance in cancer metabolism. *Front Oncol*. 2018;8:126. doi:10.3389/fonc.2018.00126
- Heiden MG, Cantley LC, Thompson CB. Understanding the Warburg effect: the metabolic requirements of cell proliferation. *Science*. 2009;324:1029–1033. doi:10.1126/science.1160809
- Ward PS, Thompson CB. Metabolic reprogramming: a cancer hallmark even warburg did not anticipate. *Cancer Cell*. 2012;21:297–308. doi:10.1016/j.ccr.2012.02.014
- Lane AN, Tan J, Wang Y, Yan J, Higashi RM, Fan TW. Probing the metabolic phenotype of breast cancer cells by multiple tracer stable isotope resolved metabolomics. *Metab Eng*. 2017;43:125–136. doi:10.1016/j.ymben.2017.01.010
- Giguere V, Yang N, Segui P, Evans RM. Identification of a new class of steroid hormone receptors. *Nature*. 1988;331:91–94. doi:10.1038/331091a0
- Xu Z, Liu J, Gu L, Ma X, Huang B, Pan X. Research progress on the reproductive and non-reproductive endocrine tumors by estrogen-related receptors. *J Steroid Biochem Mol Biol*. 2016;158:22–30. doi:10.1016/j.jsbmb.2016.01.008
- Deblois G, Giguere V. Oestrogen-related receptors in breast cancer: control of cellular metabolism and beyond. *Nat Rev Cancer*. 2013;13:27–36. doi:10.1038/nrc3396
- Chang CY, Kazmin D, Jasper JS, Kunder R, Zuercher WJ, McDonnell DP. The metabolic regulator ERR α , a downstream target of HER2/IGF-1R, as a therapeutic target in breast cancer. *Cancer Cell*. 2011;20:500–510. doi:10.1016/j.ccr.2011.08.023
- Cavallini A, Notarnicola M, Giannini R, et al. Oestrogen receptor-related receptor alpha (ERR α) and oestrogen receptors (ER α and ER β) exhibit different gene expression in human colorectal tumour progression. *Eur J Cancer*. 2005;41:1487–1494. doi:10.1016/j.ejca.2005.04.008
- Tam IS, Giguere V. There and back again: the journey of the estrogen-related receptors in the cancer realm. *J Steroid Biochem Mol Biol*. 2016;157:13–19. doi:10.1016/j.jsbmb.2015.06.009
- Zhang S, Liu X, Liu J, Guo H, Xu H, Zhang G. PGC-1 α interacts with microRNA-217 to functionally regulate breast cancer cell proliferation. *Biomed Pharmacother*. 2017;85:541–548. doi:10.1016/j.biopha.2016.11.062
- Audet-Walsh E, Papadopoli DJ, Gravel SP, et al. The PGC-1 α /ERR α axis represses one-carbon metabolism and promotes sensitivity to anti-folate therapy in breast cancer. *Cell Rep*. 2016;14:920–931. doi:10.1016/j.celrep.2015.12.086
- Giguère V. To ERR in the estrogen pathway. *Trends Endocrinol Metab*. 2002;13:220–225. doi:10.1016/S1043-2760(02)00592-1
- Li Y, Birnbaumer L, Teng CT. Regulation of ERR α gene expression by estrogen receptor agonists and antagonists in SKBR3 breast cancer cells: differential molecular mechanisms mediated by G protein-coupled receptor GPR30/GPER-1. *Mol Endocrinol*. 2010;24:969–980. doi:10.1210/me.2009-0148
- Thomas P, Pang Y, Filardo EJ, Dong J. Identity of an estrogen membrane receptor coupled to a G protein in human breast cancer cells. *Endocrinology*. 2005;146:624–632. doi:10.1210/en.2004-1064
- Filardo EJ, Quinn JA, Frackelton AR, Bland KI. Estrogen action via the G protein-coupled receptor, GPR30: stimulation of adenylyl cyclase and cAMP-mediated attenuation of the epidermal growth factor receptor-to-MAPK signaling axis. *Mol Endocrinol*. 2002;16:70–84. doi:10.1210/mend.16.1.0758
- Zhou X, Wang S, Wang Z, et al. Estrogen regulates Hippo signaling via GPER in breast cancer. *J Clin Invest*. 2015;125:2123–2135. doi:10.1172/JCI79573

29. Zhou K, Sun P, Zhang Y, You X, Li P, Wang T. Estrogen stimulated migration and invasion of estrogen receptor-negative breast cancer cells involves an ezrin-dependent crosstalk between G protein-coupled receptor 30 and estrogen receptor beta signaling. *Steroids*. 2016;111:113–120. doi:10.1016/j.steroids.2016.01.021
30. Yu T, Liu M, Luo H, et al. GPER mediates enhanced cell viability and motility via non-genomic signaling induced by 17 β -estradiol in triple-negative breast cancer cells. *J Steroid Biochem Mol Biol*. 2014;143:392–403. doi:10.1016/j.jsbmb.2014.05.003
31. Ye S, Xu Y, Li J, Zheng S, Sun P, Wang T. Prognostic role of GPER/Ezrin in triple-negative breast cancer is associated with menopausal status. *Endocr Connect*. 2019;8:661–671. doi:10.1530/EC-19-0164
32. Steiman J, Peralta EA, Louis S, Kamel O. Biology of the estrogen receptor, GPR30, in triple negative breast cancer. *Am J Surg*. 2013;206:698–703. doi:10.1016/j.amjsurg.2013.07.014
33. Kotula-Balak M, Milon A, Pawlicki P, et al. Insights into the role of estrogen-related receptors α , β and γ in tumor Leydig cells. *Tissue Cell*. 2018;52:78–91. doi:10.1016/j.tice.2018.04.003
34. Hammond MEH, Hayes DF, Dowsett M, et al. American Society of Clinical Oncology/College of American Pathologists guideline recommendations for immunohistochemical testing of estrogen and progesterone receptors in breast cancer. *J Clin Oncol*. 2010;28:2784–2795. doi:10.1200/JCO.2009.25.6529
35. Wolff AC, Hammond MEH, Allison KH, et al. Human epidermal growth factor receptor 2 testing in breast cancer: American Society of Clinical Oncology/College of American Pathologists clinical practice guideline focused update. *J Clin Oncol*. 2018;36:2105–2122. doi:10.1200/JCO.2018.77.8738
36. Skrzypczak M, Schuler S, Lattrich C, Ignatov A, Ortmann O, Treeck O. G protein-coupled estrogen receptor (GPER) expression in endometrial adenocarcinoma and effect of agonist G-1 on growth of endometrial adenocarcinoma cell lines. *Steroids*. 2013;78:1087–1091. doi:10.1016/j.steroids.2013.07.007
37. Watanabe A, Kinoshita Y, Hosokawa K, Mori T, Yamaguchi T, Honjo H. Function of estrogen-related receptor alpha in human endometrial cancer. *J Clin Endocrinol Metab*. 2006;91:1573–1577. doi:10.1210/jc.2005-1990
38. Luo HJ, Luo P, Yang GL, Peng QL, Liu MR, Tu G. G-protein coupled estrogen receptor 1 expression in primary breast cancers and its correlation with clinicopathological variables. *J Breast Cancer*. 2011;14:185–190. doi:10.4048/jbc.2011.14.3.185
39. Gaudet HM, Cheng SB, Christensen EM, Filardo EJ. The G-protein coupled estrogen receptor, GPER: the inside and inside-out story. *Mol Cell Endocrinol*. 2015;418(3):207–219. doi:10.1016/j.mce.2015.07.016
40. Dent R, Trudeau M, Pritchard KI, et al. Triple-negative breast cancer: clinical features and patterns of recurrence. *Clin Cancer Res*. 2007;13:4429–4434. doi:10.1158/1078-0432.CCR-06-3045
41. Gluz O, Liedtke C, Gottschalk N, Pusztai L, Nitz U, Harbeck N. Triple-negative breast cancer—current status and future directions. *Ann Oncol*. 2009;20:1913–1927. doi:10.1093/annonc/mdp492
42. Contreras-Zárate MJ, Day NL, Ormond DR, et al. Estradiol induces BDNF/TrkB signaling in triple-negative breast cancer to promote brain metastases. *Oncogene*. 2019;38:4685–4699. doi:10.1038/s41388-019-0756-z
43. Girgert R, Emons G, Gründker C. 17 β -estradiol-induced growth of triple-negative breast cancer cells is prevented by the reduction of GPER expression after treatment with gefitinib. *Oncol Rep*. 2017;37:1212–1218. doi:10.3892/or.2016.5306
44. McNamara KM, Oguro S, Omata F, et al. The presence and impact of estrogen metabolism on the biology of triple-negative breast cancer. *Breast Cancer Res Treat*. 2017;161:213–227. doi:10.1007/s10549-016-4050-2
45. Kallen J, Lattmann R, Beerli R, et al. Crystal structure of human estrogen-related receptor alpha in complex with a synthetic inverse agonist reveals its novel molecular mechanism. *J Biol Chem*. 2007;282:23231–23239. doi:10.1074/jbc.M703337200
46. Reese JM, Bruinsma ES, Nelson AW, et al. ER β -mediated induction of cystatins results in suppression of TGF β signaling and inhibition of triple-negative breast cancer metastasis. *Proc Natl Acad Sci U S A*. 2018;115:E9580–E9589. doi:10.1073/pnas.1807751115
47. Zhao L, Huang S, Mei S, et al. Pharmacological activation of estrogen receptor beta augments innate immunity to suppress cancer metastasis. *Proc Natl Acad Sci U S A*. 2018;115:E3673–E3681. doi:10.1073/pnas.1803291115
48. Wang X, Zheng N, Dong J, Wang X, Liu L, Huang J. Estrogen receptor- α 36 is involved in icaritin induced growth inhibition of triple-negative breast cancer cells. *J Steroid Biochem Mol Biol*. 2017;171:318–327. doi:10.1016/j.jsbmb.2017.05.009
49. Hu B, Yan W, Wang M, et al. Huaier polysaccharide inhibits the stem-like characteristics of ER α -36(high) triple negative breast cancer cells via inactivation of the ER α -36 signaling pathway. *Int J Biol Sci*. 2019;15:1358–1367. doi:10.7150/ijbs.27360
50. Wisinski KB, Xu W, Tevaarwerk AJ, et al. Targeting estrogen receptor beta in a phase 2 study of high-dose estradiol in metastatic triple-negative breast cancer: a wisconsin oncology network study. *Clin Breast Cancer*. 2016;16:256–261. doi:10.1016/j.clbc.2016.03.005

OncoTargets and Therapy

Publish your work in this journal

OncoTargets and Therapy is an international, peer-reviewed, open access journal focusing on the pathological basis of all cancers, potential targets for therapy and treatment protocols employed to improve the management of cancer patients. The journal also focuses on the impact of management programs and new therapeutic

agents and protocols on patient perspectives such as quality of life, adherence and satisfaction. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/oncotargets-and-therapy-journal>

Dovepress