

Skeletal muscle features in myotonic dystrophy and sarcopenia: do similar nuclear mechanisms lead to skeletal muscle wasting?

M. Malatesta

Dipartimento di Scienze Neurologiche, Neuropsicologiche, Morfologiche e Motorie, Sezione di Anatomia e Istologia, Università di Verona, Italy

Abstract

In the cell nucleus, the gene primary transcripts undergo molecular processing to generate mature RNAs, which are finally exported to the cytoplasm. These mRNA maturation events are chronologically and spatially ordered, and mostly occur on distinct ribonucleoprotein (RNP)-containing structures. Defects in the mRNA maturation pathways have been demonstrated in myotonic dystrophy type 1 (DM1) and type 2 (DM2) whose characteristic multisystemic features are caused by the expansion of two distinct nucleotide sequences: (CTG)_n in the DMPK gene on chromosome 19q13 in DM1, and (CCTG)_n in the ZNF9 gene on chromosome 3q21 in DM2. By combining biomolecular and cytochemical techniques, it has been shown that the basic mechanisms of DMs reside in the accumulation of CUG- or CCUG-containing transcripts in intranuclear *foci* where several RNA-binding proteins necessary for the physiological processing of pre-mRNA are sequestered. Moreover, a nucleoplasmic accumulation of splicing and cleavage factors has been found in DMs. This suggests that the dystrophic phenotype could depend on a general alteration of the pre-mRNA post-transcriptional pathway. Interestingly, the accumulation of pre-mRNA processing factors in the myonuclei of DM1 and DM2 patients is reminiscent of the nuclear alterations typical of sarcopenia, *i.e.*, the loss of muscle mass and function which physiologically occurs during ageing. Consistently, in an *in vitro* study, we observed that satellite-cell-derived DM2 myoblasts show cell senescence alterations and impairment of the pre-mRNA maturation pathways earlier than the myoblasts from healthy patient. These results suggest possible common cellular mechanisms responsible for skeletal muscle wasting in sarcopenia and in myotonic dystrophy.

Skeletal muscle features in myotonic dystrophy

In recent years, the role of mutations in non-protein-coding regions has come to light in the pathogenesis of different neuromuscular diseases,^{1,2} so that the terms *toxic RNA* and *spliceopathy* have been increasingly referred to pathological conditions in which accumulation of mutant RNAs results in a deleterious gain-of-function deregulating transcript processing and protein synthesis in multiple metabolic pathways.

One of the best studied examples of RNA-dominant disease is myotonic dystrophy (DM), an autosomal dominant disorder characterised by a variety of multisystemic features including muscular dystrophy with increased number of centrally located or clumped nuclei in muscle fibres,³ myotonia (muscle hyperexcitability), dilated cardiomyopathy, cardiac conduction defects,⁴ insulin-resistance, cataracts,⁵ and disease-specific serological abnormalities.^{6,7} Two forms of DM are presently known: the more severe DM1-Steinert's disease (OMIM 160900), caused by an expanded (CTG)_n nucleotide sequence in the 3' untranslated region of the Dystrophia Myotonic Protein Kinase (DMPK) gene (OMIM 605377) on chromosome 19q13,⁸⁻¹⁰ and the milder form DM2 (OMIM 602688), caused by the expansion of the tetranucleotidic repeat (CCTG)_n in the first intron of the Zinc Finger Protein (ZNF)-9 gene (OMIM 116955) on chromosome 3q21.¹¹

In the cell nucleus, the gene primary transcripts undergo molecular processing to generate mature RNAs, which are finally exported to the cytoplasm: these mRNA maturation events are chronologically and spatially ordered, and mostly occur on distinct RNP-containing structures.¹² By combining biomolecular techniques with the analysis *in situ* of the nuclear organization and molecular composition, it has been demonstrated that the expanded-CUG- or CCUG-containing transcripts, in DM1 and DM2 cells respectively, are retained in the cell nucleus, and accumulate in the form of RNP-containing focal aggregates.¹³ These nuclear *foci* specifically sequester the alternative splicing regulators CUG-binding protein 1 (CUGBP1) and muscle-blind-like 1 (MBLN1) protein,¹⁴⁻¹⁶ which are necessary for the physiological processing of pre-mRNA, especially for contractile protein synthesis.¹⁷ These focal aggregates are considered as a biomolecular feature of DMs, and have been detected in several adult tissues as well as in cultured cells from DM patients.^{13,18-23} These *foci* also sequester hnRNPs and snRNPs, *i.e.* splicing factors involved in the early phases of the pre-mRNA processing,²⁴ thus strengthening the hypothesis that the multifactorial phenotype of dystrophic patients may result from a more gen-

Correspondence: Manuela Malatesta, Dipartimento di Scienze Neurologiche, Neuropsicologiche, Morfologiche e Motorie, Sezione di Anatomia e Istologia, Università degli Studi di Verona, Strada Le Grazie 8, 37134 Verona, Italy. Tel. +39.045.8027157 - Fax: +39.045.8027163. E-mail: manuela.malatesta@univr.it

Key words: cell nucleus, myotonic dystrophy, sarcopenia.

Received for publication: 28 July 2012.

Accepted for publication: 3 August 2012.

This work is licensed under a Creative Commons Attribution NonCommercial 3.0 License (CC BY-NC 3.0).

©Copyright M. Malatesta, 2012

Licensee PAGEpress, Italy

European Journal of Histochemistry 2012; 56:e36

doi:10.4081/ejh.2012.e36

eral alteration of the pre-mRNA post-transcriptional pathway.

Recently, it has been demonstrated that MBNL1 accumulate in the nuclear *foci* during interphase but, at mitosis, the *foci* relocate to the cytoplasm where they undergo degradation, while newly-formed *foci* develop in the nucleus of the daughter cells as a consequence of *de novo* accumulation of expanded RNAs.²⁵ Therefore, in proliferating cells, the cyclic release from the nucleus of the *foci* and their cytoplasmic degradation would prevent the massive intranuclear sequestration of nuclear factors; on the contrary, in non cycling cells, the nuclear *foci* do not undergo intracellular relocation/degradation and progressively increase in number and size. This dynamic behaviour of nuclear *foci* is compatible with the evidence that in DM patients the most affected organs or tissues are those where non-renewing cells are mainly present, such as the skeletal muscle, heart and the central nervous system, whereas cells from self-renewing tissues (such as skin fibroblasts or layering epithelial cells) are much less affected. Accordingly, measuring intranuclear *foci* in skeletal muscle biopsies taken from patients at different times it has been demonstrated that the MBNL1-containing *foci* actually become larger with increasing patient's age.²⁵

In addition to the formation of intranuclear *foci*, DM1 and DM2 cells show an altered distribution of nuclear ribonucleoprotein (RNP)-containing structures and molecular factors responsible for pre-mRNA transcription and maturation. In particular, by means of ultrastructural immunocytochemistry on skeletal muscle biopsies from DM1 and DM2 patients, it has been shown^{26,27} that splicing and cleavage factors accumulate in the intranuclear functional sites where they usually localize, sometimes ectopi-

cally relocating also into nuclear RNP domains where they generally do not occur (*unpublished results*). This accumulation could hamper the functionality of the splicing machinery and slow down the intranuclear molecular trafficking thus reducing the metabolic activity of myonuclei, consistent with recent findings demonstrating a reduced protein synthesis in DM1 and DM2 myoblasts.^{28,29}

Skeletal muscle features in sarcopenia

During ageing, the skeletal muscle undergoes a progressive loss of mass, strength and function, in the process known as sarcopenia.^{30,31} Sarcopenia affects healthy, physically active subjects: the rate of muscle loss in humans has been estimated to range 1 to 2% per year after age of fifty. Therefore, sarcopenia represents a great risk factor for frailty, loss of independence and physical disability in elderly, since it is associated with decreased functional performance, higher risk of falls and motor function impairment. The mechanisms underlying age-related skeletal muscle wasting and weakness are probably manifold and still remain to be fully elucidated;³⁰ however, although no specific therapy is presently available to counteract its onset or progress, studies performed on humans and other mammals have stressed the importance of physical exercise as an effective, although still debated, approach to prevent or limit the age-related muscle mass loss.³²⁻³⁶

Interestingly, the sarcopenic process is characterised by structural and functional alterations of the skeletal muscle that are reminiscent of myotonic dystrophy. In fact, the aged muscle shows grouped atrophy, fibre size variability and centrally located nuclei.³⁷ In addition, factors involved in the post-transcriptional processing of pre-mRNA have been found to accumulate not only in the nucleoplasmic RNP-containing structures where they usually locate but also in ectopic nuclear domains.³⁸⁻⁴⁰ This intranuclear accumulation/delocalization of RNP structures containing splicing and cleavage factors has been found not only in the skeletal muscle but also in other tissues (*e.g.*, liver, brain) of aged mammals.⁴¹⁻⁴⁴ Moreover, aged cells undergo malfunctions of the degradation systems both in the cytoplasm⁴⁵ and in the nucleus^{46,47} with accumulation of crosslinked insoluble molecules (including non-coding RNAs) which hampers the intracellular transport mechanisms. This suggests that in ageing cells the entire production chain of mRNA, from its synthesis to the cytoplasmic export, becomes less efficient, likely contributing to the reduced capability of cells to positively react to metabolic stimuli, which typically occurs in elderly. This loss of responsive-

ness would have particularly severe effects in skeletal muscles, where a misregulated protein turnover would result in a structural imbalance between muscle protein degradation and the restoring protein synthesis.⁴⁸

Concluding remarks

A recent *in vitro* study⁴⁹ reported that satellite-cell-derived myoblasts from DM2 patients show cell-senescence alterations (*e.g.*, cytoplasmic vacuolisation, reduction of the proteosynthetic apparatus, accumulation of heterochromatin and impairment of the pre-mRNA maturation pathways) earlier than the myoblasts from healthy patients; moreover, when grown in a differentiation medium DM2 myoblasts fuse into multinucleated myotubes exhibiting structural defects similar to those observed in senescent myotubes from healthy patients.⁵⁰ The early occurrence of senescence-related features in satellite cell-derived myoblasts suggests that satellite cells from DM2 patients have a reduced regeneration capability, which would contribute to the muscular dystrophic phenotype.

The cytochemical and ultrastructural evidence demonstrates that the skeletal muscle of DM patients shares intriguing similarities with the muscle from aged individuals in several nuclear features, especially in the altered nuclear RNP-containing structures involved in pre-mRNA transcription and splicing. This opens interesting perspectives on the role of the RNP nuclear components in the onset of muscle cell dysfunctions and encourages comparative studies aimed at detecting common cellular mechanisms at the basis of skeletal muscle wasting.

Finally, it is worth noting that the analysis *in situ* of the organization and molecular composition of nuclear domains is a powerful tool not only for getting information about the DNA/RNA pathways which govern cellular metabolism, but also for detecting the occurrence of cell dysfunctions related to pathological phenotypes.⁵¹⁻⁵³

References

- Osborne RJ, Thornton CA. RNA-dominant diseases. *Hum Mol Genet* 2006;15:R162-9.
- Wheeler TM, Thornton CA. Myotonic dystrophy: RNA-mediated muscle disease. *Curr Opin Neurol* 2007;20:572-6.
- Vihola A, Bassez G, Meola G, Zhang S, Haapasalo H, Paetau A, et al. Histopathological differences of myotonic dystrophy type 1 (DM1) and PROMM/DM2. *Neurology* 2003;60:1854-7.
- Bachinski LL, Udd B, Meola G, Sansone V,

- Bassez G, Eymard B, et al. Confirmation of the type 2 myotonic dystrophy (CCTG)n expansion mutation in patients with proximal myotonic myopathy/proximal myotonic dystrophy of different European origins: a single shared haplotype indicates an ancestral founder effect. *Am J Hum Gen* 2003;73:835-48.
- Meola G, Moxley RT 3rd. Myotonic dystrophy type 2 and related myotonic disorders. *J Neurol* 2004;251:1173-82.
- Day JW, Ricker K, Jacobsen JF, Rasmussen LJ, Dick KA, Kress W, et al. Myotonic dystrophy type 2: molecular, diagnostic and clinical spectrum. *Neurology* 2003;60:657-64.
- Savkur RS, Philips AV, Cooper TA, Dalton JC, Moseley ML, Ranum LP, et al. Insulin receptor splicing alteration in myotonic dystrophy type 2. *Am J Hum Genet* 2004;74:1309-13.
- Brook JD, McCurrach ME, Harley HG, Buckler AJ, Church D, Aburatani H, et al. Molecular basis of myotonic dystrophy: expansion of a trinucleotide (CTG) repeat at the 3' end of a transcript encoding a protein kinase family member. *Cell* 1992;69:385-7.
- Fu YH, Pizzuti A, Fenwick RG Jr, King J, Rajnarayan S, Dunne PW, et al. An unstable triplet repeat in a gene related to myotonic muscular dystrophy. *Science* 1992;255:1256-8.
- Mahadevan M, Tsilfidis C, Sabourin L, Shutter G, Amemiya C, Jansen G, et al. Myotonic dystrophy mutation: an unstable CTG repeat in the 3' untranslated region of the gene. *Science* 1992;255:1253-5.
- Ranum LP, Rasmussen PF, Benzow KA, Koob MD, Day JW. Genetic mapping of a second myotonic dystrophy locus. *Nat Genet* 1998;19:196-8.
- Fakan S. Ultrastructural cytochemical analyses of nuclear functional architecture. *Eur J Histochem* 2004;48:5-14.
- Liquori CL, Ricker K, Moseley ML, Jacobsen JF, Kress W, Naylor SL, et al. Myotonic dystrophy type 2 caused by a CCTG expansion in intron 1 of ZNF9. *Science* 2001;293:864-7.
- Miller JW, Urbinati CR, Teng-Ummuay P, Stenberg MG, Byrne BJ, Thornton CA, et al. Recruitment of human muscleblind proteins to (CUG)(n) expansions associated with myotonic dystrophy. *EMBO J* 2000;19:4439-48.
- Fardaei M, Rogers MT, Thorpe HM, Larkin K, Hamshere MG, Harper PS, et al. Three proteins, MBNL, MBLL and MBXL, co-localize in vivo with nuclear foci of expanded-repeat transcripts in DM1 and DM2 cells. *Hum Mol Genet* 2002;11:805-14.
- Jones K, Jin B, Iakova P, Huichalaf C, Sarkar P, Schneider-Gold C, et al. RNA foci, CUGBP1, and ZNF9 are the primary targets of the mutant CUG and CCUG repeats expanded in myotonic dystrophies type 1

- and type 2. *Am J Pathol* 2011;179:2475-89.
17. Llorian M, Smith CWJ. Decoding muscle alternative splicing. *Curr Opin Genet Develop* 2011;21:380-7.
 18. Taneja KL, McCurrach M, Schalling M, Housman D, Singer RH. Foci of trinucleotide repeat transcripts in nuclei of myotonic dystrophy cells and tissues. *J Cell Biol* 1995;128:995-1002.
 19. Mankodi A, Teng-Umuay P, Krym M, Henderson D, Swanson M, Thornton CA. Ribonuclear inclusions in skeletal muscle in myotonic dystrophy types 1 and 2. *Ann Neurol* 2003;54:760-8.
 20. Mankodi A, Lin XL, Blaxall BC, Swanson MS, Thornton CA. Nuclear RNA foci in the heart in myotonic dystrophy. *Circ Res* 2005;97:1152-5.
 21. Cardani R, Mancinelli E, Sansone V, Rotondo G, Meola G. Biomolecular identification of (CCTG)_n mutation in myotonic dystrophy type 2 (DM2) by FISH on muscle biopsy. *Eur J Histochem* 2004;48:437-42.
 22. Cardani R, Mancinelli E, Rotondo G, Sansone V, Meola G. Muscleblind-like protein 1 nuclear sequestration is a molecular pathology marker of DM1 and DM2. *Eur J Histochem* 2006;50:177-82.
 23. Cardani R, Mancinelli E, Giagnacovo M, Sansone V, Meola G. Ribonuclear inclusions as biomarker of myotonic dystrophy type 2, even in improperly frozen or defrozed skeletal muscle biopsies. *Eur J Histochem* 2009;53:107-11.
 24. Perdoni F, Malatesta M, Cardani R, Giagnacovo M, Mancinelli E, Meola G, et al. RNA/MBNL1-containing foci in myoblast nuclei from patients affected by myotonic dystrophy type 2: an immunocytochemical study. *Eur J Histochem* 2009;53:151-8.
 25. Giagnacovo M, Malatesta M, Cardani R, Meola G, Pellicciari C. Nuclear ribonucleo-protein-containing foci increase in size in non-dividing cells from patients with myotonic dystrophy type 2. *Histochem Cell Biol* 2012; DOI: 10.1007/s00418-012-0984-6
 26. Malatesta M, Giagnacovo M, Cardani R, Meola G, Pellicciari C. RNA processing is altered in skeletal muscle nuclei of patients affected by myotonic dystrophy. *Histochem Cell Biol* 2011;135:419-25.
 27. Malatesta M., Giagnacovo M., Cardani R., Meola G., Pellicciari C. Alteration of RNA processing in skeletal muscle nuclei of patients affected by myotonic dystrophy. *Eur J Histochem* 2011;55/suppl1:14.
 28. Salisbury E, Schoser B, Schneider-Gold C, Wang GL, Huichalaf C, Jin B, et al. Expression of RNA CCUG repeats dysregulates translation and degradation of proteins in myotonic dystrophy 2 patients. *Am J Pathol* 2009;175:748-62.
 29. Huichalaf C, Sakai K, Jin B, Jones K, Wang GL, Schoser B, et al. Expansion of CUG RNA repeats causes stress and inhibition of translation in myotonic dystrophy 1 (DM1) cells. *FASEB J* 2010;24:3706-19.
 30. Thompson LD. Age-related muscle dysfunction. *Exp Gerontol* 2009;44:106-11.
 31. Cruz-Jentoft AJ, Baeyens JP, Bauer JM, Boirie Y, Cederholm T, Landi F, et al. European Working Group on Sarcopenia in Older People. Sarcopenia: European consensus on definition and diagnosis: Report of the European Working Group on Sarcopenia in Older People. *Age Ageing* 2010;39:412-23.
 32. Marcell TJ. Sarcopenia: causes, consequences, and preventions. *J Gerontol A Biol Sci Med Sci* 2003;58:M911-6.
 33. Bautmans I, Van Puyvelde K, Mets T. Sarcopenia and functional decline: pathophysiology, prevention and therapy. *Acta Clin Belg* 2009;64:303-16.
 34. Zancanaro C, Mariotti R, Perdoni F, Nicolato E, Malatesta M. Physical training is associated with changes in NMR and morphometrical parameters of the skeletal muscle in senescent mice. *Eur J Histochem* 2007;51:305-10.
 35. Malatesta M., Fattoretti P., Giagnacovo M., Pellicciari C., Zancanaro C. Structural and functional features of skeletal muscle cell nuclei are modulated by physical exercise in old mice. *Eur J Histochem* 2011;55(Suppl. 1):14.
 36. Curzi D, Salucci S, Marini M, Esposito F, Agnello L, Veicsteinas A, et al. How physical exercise changes rat myotendinous junctions: an ultrastructural study. *Eur J Histochem* 2012;56:e19.
 37. Edstrom E, Altun M, Bergman E, Johnson H, Kullberg S, Ramirez-León V, et al. Factors contributing to neuromuscular impairment and sarcopenia during aging. *Physiol Behav* 2007;92:129-35.
 38. Malatesta M, Perdoni F, Muller S, Zancanaro C, Pellicciari C. Nuclei of aged myofibres undergo structural and functional changes suggesting impairment in RNA processing. *Eur J Histochem* 2009;53: 97-106.
 39. Malatesta M, Perdoni F, Muller S, Pellicciari C, Zancanaro C. Pre-mRNA processing is partially impaired in satellite cell nuclei from aged muscles. *J Biomed Biotechnol* 2010;2010:410405.
 40. Malatesta M, Fattoretti P, Giagnacovo M, Pellicciari C, Zancanaro C. Physical training modulates structural and functional features of cell nuclei in type II myofibers of old mice. *Rejuvenation Res* 2011;14:543-52.
 41. Malatesta M, Bertoni-Freddari C, Fattoretti P, Caporaloni C, Fakan S, Gazzanelli G. Altered RNA structural constituents in aging and vitamin E deficiency. *Mech Ageing Dev* 2003;124:175-81.
 42. Malatesta M, Baldelli B, Battistelli S, Fattoretti P, Bertoni-Freddari C. Aging affects the distribution of the circadian CLOCK protein in rat hepatocytes. *Microsc Res Tech* 2005;68:45-50.
 43. Malatesta M, Fattoretti P, Baldelli B, Battistelli S, Ballelli M, Bertoni-Freddari C. Effects of ageing on the fine distribution of the circadian CLOCK protein in reticular formation neurons. *Histochem Cell Biol* 2007;127:641-7.
 44. Malatesta M, Biggiogera M, Cisterna B, Ballelli M, Bertoni-Freddari C, Fattoretti P. Perichromatin fibrils accumulation in hepatocyte nuclei reveals alterations of pre-mRNA processing during ageing. *DNA Cell Biol* 2010;29:49-57.
 45. Jameson CW. Towards a unified and interdisciplinary model of ageing. *Med Hypotheses* 2004;63:83-6.
 46. Costanzo M, Cisterna B, Zharskaya OO, Zatschina OV, Biggiogera M. Discrete foci containing RNase A are found in nucleoli of HeLa cells after aging in culture. *Eur J Histochem* 2011;55:e15.
 47. Costanzo M, Cisterna B, Zharskaya OO, Zatschina OV, Biggiogera M. Synthesis, movement and destruction of RNAs in the nucleus. *Eur J Histochem* 2011;55(Suppl. 1):2.
 48. Koopman R, van Loon LJC. Aging, exercise and muscle protein metabolism. *J Appl Physiol* 2009;106:2040-8.
 49. Malatesta M, Giagnacovo M, Renna LV, Cardani R, Meola G, Pellicciari C. Cultured myoblasts from patients affected by myotonic dystrophy type 2 exhibit senescence-related features: ultrastructural evidence. *Eur J Histochem* 2011;55:e26.
 50. Giagnacovo M, Costanzo M, Cardani R, Veneroni P, Pellicciari C, Meola G. Ultrastructural features of myotubes derived from myoblasts of patients affected by myotonic dystrophy type 2, after senescence in vitro. *Eur J Histochem* 2011;55(Suppl. 1):15.
 51. Giagnacovo M, Cardani R, Meola G, Pellicciari C, Malatesta M. Routinely frozen biopsies of human skeletal muscle are suitable for morphological and immunocytochemical analyses at transmission electron microscopy. *Eur J Histochem* 2010;54:137-42.
 52. Malatesta M, Meola G. Structural and functional alterations of the cell nucleus in skeletal muscle wasting: the evidence in situ. *Eur J Histochem* 2010;54:e44.
 53. Cenni V, Capanni C, Columbaro M, Ortolani M, D'Apice MR, Novelli G, et al. Autophagic degradation of farnesylated prelamin A as a therapeutic approach to lamin-linked progeria. *Eur J Histochem* 2011;55:e36.