

REVIEW

Mutations in the phosphatidylinositol 3-kinase pathway: role in tumor progression and therapeutic implications in breast cancer

Todd W Miller^{1,2}, Brent N Rexer^{2,3}, Joan T Garrett³ and Carlos L Arteaga^{1,2,3,4*}

Abstract

Mutations in genes that constitute the phosphatidylinositol 3-kinase (PI3K) pathway occur in >70% of breast cancers. Clinical and experimental evidence suggest that PI3K pathway activation promotes resistance to some of the current breast cancer therapies. PI3K is a major signaling hub downstream of human epidermal growth factor receptor (HER)2 and other receptor tyrosine kinases. PI3K activates AKT, serum/glucocorticoid regulated kinase (SGK), phosphoinositide-dependent kinase 1 (PDK1), mammalian target of rapamycin (mTOR), and several other molecules involved in cell cycle progression and survival. In estrogen receptor (ER)+ breast cancer cells, PI3K activation promotes estrogen-dependent and -independent ER transcriptional activity, which, in turn, may contribute to anti-estrogen resistance. Activation of this pathway also confers resistance to HER2-targeted therapies. In experimental models of resistance to anti-estrogens and HER2 inhibitors, pharmacological inhibition of PI3K/AKT/mTOR has been shown to overcome drug resistance. Early clinical data suggest that combined inhibition of either HER2 or ER plus inhibition of the PI3K pathway might be an effective strategy for treatment of respective HER2+ and ER+ breast cancers resistant to standard therapies. Here, we review alterations in the PI3K pathway in breast cancer, their association with therapeutic resistance, and the state of clinical development of PI3K pathway inhibitors.

Introduction

The phosphatidylinositol 3-kinase (PI3K) pathway is the most frequently mutated pathway in breast cancer, with mutation and/or amplification of the genes encoding the PI3K catalytic subunits p110 α (*PIK3CA*) and p110 β (*PIK3CB*), the PI3K regulatory subunit p85 α (*PIK3R1*), receptor tyrosine kinases (RTKs) such as human epidermal growth factor receptor (HER)2 (*ERBB2*) and fibroblast growth factor receptor (FGFR)1, the PI3K activator K-Ras, the PI3K effectors AKT1, AKT2, and phosphoinositide-dependent kinase 1 (PDK1), and loss of the lipid phosphatases PTEN (phosphatase and tensin homolog) and INPP4B (inositol polyphosphate-4-phosphatase, type II) (Table 1). PI3K is activated by growth factor RTKs and G-protein-coupled receptors (Figure 1). PI3K phosphorylates phosphatidylinositol 4,5-bisphosphate (PIP₂) to produce phosphatidylinositol 3,4,5-trisphosphate (PIP₃). In turn, PIP₃ recruits to the plasma membrane several pleckstrin homology (PH) domain-containing proteins, such as PDK1 and AKT, which, upon activation, drive cell cycle progression and survival. Negative regulation of this pathway is conferred by PTEN and INPP4B, which dephosphorylate PIP₃ and PIP₂, respectively. Akt phosphorylates and inactivates Tuberin (TSC2), a GTPase-activating protein of the Ras homologue Rheb. Inactivation of Tuberin allows GTP bound-Rheb to accumulate and activate the mammalian target of rapamycin (mTOR)/Raptor (TORC1) complex, which ultimately regulates protein synthesis and cell growth [1]. mTOR also couples with Rictor to form the TORC2 complex, which phosphorylates and activates AKT at Ser473.

Class IA PI3K isoforms are heterodimeric lipid kinases that contain a p110 catalytic subunit and a p85 regulatory subunit. The three genes *PIK3CA*, *PIK3CB*, and *PIK3CD* encode the homologous p110 α , p110 β , and p110 δ isozymes, respectively. Expression of p110 δ is largely restricted to immune and hematopoietic cells, whereas p110 α and p110 β are ubiquitously expressed. *PIK3CA* mutations are the most common genetic alterations of this pathway in breast cancer, where $\geq 80\%$ occur within the helical (E542K and E545K) and kinase (H1047R)

*Correspondence: carlos.arteaga@vanderbilt.edu

⁴Division of Hematology-Oncology, VUMC, 2220 Pierce Avenue, 777 PRB, Nashville, TN 37232-6307, USA

Full list of author information is available at the end of the article

Table 1. Phosphatidylinositol 3-kinase pathway alterations in human breast cancers by molecular subtype

Gene (protein)	Alteration	Effect on signaling	Frequency			Reference
			Luminal (ER+)	HER2+	Basal (TN)	
<i>ErbB2</i> (HER2)	Gene amplification or overexpression	Hyperactivation of ErbB2 signaling (PI3K, MEK)	10%	~100%	0%	[30-32]
<i>PTEN</i>	Loss-of-function mutation or reduced expression	Hyperactivation of PI3K signaling	29-44%	22%	67%	[6,8,104,105]
<i>PIK3CA</i> (p110 α /PI3K)	Activating mutation	Hyperactivation of PI3K signaling	28-47%	23-33%	8-25%	[6,52,66-68, 105-107]
<i>PIK3CB</i> (p110 β /PI3K)	Amplification	Unknown		5% of all cases		[62]
<i>IGF1R</i> and <i>INSR</i> (IGF-1R, InsR)	Receptor activation, <i>IGF1R</i> amplification	Activates IGF-1R/InsR signaling (PI3K, MEK)	41-48%	18-64%	42%	[108,109]
<i>FGFR1</i>	Amplification, activating mutation	Hyperactivation of FGFR signaling (PI3K, MEK)	8.6-11.6%	5.4%	5.6%	[63,110]
<i>RPS6K1</i> (p70S6K)	Amplification	Unknown		3.8-12.5% of all cases		[111]
<i>INPP4B</i>	Reduced expression or genomic loss	Hyperactivation of PI3K signaling	10-33%	54%	53%	[64,112]
<i>PIK3R1</i> (p85 α /PI3K)	Inactivating mutation	Derepression of catalytic activity of p110 α		2% of all cases		[113]
<i>AKT1</i>	Activating mutation	Hyperactivation of AKT	2.6-3.8%	0%	0%	[65,66, 106,114]
<i>AKT2</i>	Amplification	Hyperactivation of AKT		2.8% of all cases		[115]
<i>EGFR</i>	Amplification	Hyperactivation of EGFR signaling (PI3K, MEK)		0.8% of all cases		[116]
<i>PDK1</i>	Amplification or overexpression	Hyperactivation of PDK1 (AKT, TORC1)	22%	22%	38%	[117]
<i>KRAS</i>	Activating mutation	Hyperactivation of PI3K and MEK		4-6% of all cases		[118,119]

EGFR, epidermal growth factor receptor; ER, estrogen receptor; FGFR, fibroblast growth factor receptor; HER, human epidermal growth factor receptor; IGF-1R, insulin-like growth factor-1 receptor; INPP4B, inositol polyphosphate-4-phosphatase, type II; InsR, insulin receptor; MEK, mitogen-activated protein kinase kinase; PDK1, phosphoinositide-dependent kinase 1; PI3K, phosphatidylinositol 3-kinase; TN, triple negative.

domains of p110 α . Such mutations confer increased catalytic activity through different mechanisms [2], but both induce characteristics of cellular transformation, including growth factor- and anchorage-independent growth, and resistance to anoikis [3]. Temporally regulated expression of the H1047R mutant in the mammary gland of transgenic mice induces mammary tumor formation [4]. Genetic or pharmacological inactivation of *PIK3CA*^{H1047R} expression results in disappearance of mammary tumors. However, some of these recur and become insensitive to PI3K inhibition via c-myc overexpression [5].

PI3K pathway alterations frequently co-occur in breast cancer, suggesting that they confer advantages to cancer cells by different mechanisms. For example, *PIK3CA* mutations sometimes occur in breast tumors harboring PTEN loss or HER2 overexpression [6-8]. p110 α is essential for signaling and growth of tumors driven by *PIK3CA* mutations, RTKs, and/or mutant Ras, whereas p110 β lies downstream of G-protein-coupled receptors and has been shown to mediate tumorigenesis in PTEN-deficient

cells [9]. HER2 overexpression and *PIK3CA* mutations are commonly found in both ductal carcinoma *in situ* and invasive breast cancers. However, *PIK3CA* mutations are found at a lower frequency in intraepithelial neoplastic lesions. This suggests that *PIK3CA* mutations can further augment PI3K pathway activation mediated by other oncogenes such as *ERBB2* (HER2) [10-14].

Molecular analyses have shown that breast cancer is a collection of diseases that generally fit into three subtypes that respond to different therapeutics and exhibit a different natural history. Breast cancers that express estrogen receptor α (ER) and/or progesterone receptor (PR) are hormone-dependent and, as such, respond to therapies that inhibit ER signaling by multiple mechanisms. HER2-positive cancers exhibit amplification or overexpression of the *ERBB2* (HER2) proto-oncogene and respond clinically when treated with HER2-directed therapies. Triple-negative breast cancers (TNBCs), which lack detectable expression of ER, PR, and HER2, have no approved targeted therapy and are treated with traditional chemotherapy. Therefore, we will separately




Figure 1. Diagram of the phosphatidylinositol 3-kinase signaling pathway. Tumor promoters and suppressors are labeled in pink and blue, respectively. Nodes targeted by drugs in clinical development are shown in red. AMPK, AMP-activated protein kinase; GPCR, G-protein-coupled receptor; GSK3, glycogen synthase kinase 3; INPP4B, inositol polyphosphate-4-phosphatase, type II; LKB1, liver kinase B1; PDK1, phosphoinositide-dependent kinase 1; PI3K, phosphatidylinositol 3-kinase; PIP1, phosphatidylinositol monophosphate; PIP2, phosphatidylinositol 4,5-bisphosphate; PIP3, phosphatidylinositol 3,4,5-trisphosphate; PTEN, phosphatase and tensin homolog; RTK, receptor tyrosine kinase.

review the roles of molecular alterations in the PI3K pathway in each breast cancer subtype and their clinical implications.

PI3K pathway inhibitors in clinical development

Several drugs targeting multiple levels of the PI3K network (that is, PI3K, AKT, mTOR) are in clinical development in breast cancer. The first group encompasses ATP mimetics that bind competitively and reversibly to the ATP-binding pocket of p110; some of these compounds also bind and inhibit mTOR (Table 2). Notably, the pan-PI3K and p110 α -specific inhibitors are equally potent against oncogenic mutants of p110 α [15,16]. A second group includes allosteric and ATP-competitive inhibitors of the three isoforms of AKT;

these have also shown antitumor activity in preclinical models [17-19] and recently entered human trials. Allosteric inhibitors such as MK-2206 bind to the PH domain and/or hinge region in AKT to promote an inactive conformation and thus prevent localization (and activation) of AKT to the plasma membrane [19]. The macro-lide rapamycin and its analogs (rapalogs) complex with FK506-binding protein (FKBP12), which then binds to mTOR and inhibits the kinase activity of TORC1 but not TORC2 [20]. Formulation problems with rapamycin and its inability to effectively inhibit phosphorylation of 4E-BP proteins (which are TORC1 substrates that promote translation) prompted the development of analogs that have shown cytostatic activity in preclinical models and clinical trials [20]. Compounds that target

Table 2. Phosphatidylinositol 3-kinase pathway inhibitors in clinical development

Kinase target	Drug	Manufacturer	Mechanism of action
Pan-PI3K	BKM120	Novartis	ATP-competitive
	XL-147	Exelixis	ATP-competitive
	PX-866	Oncothyreon	ATP-competitive
	GDC-0941	Genentech/Roche	ATP-competitive
	CH5132799	Chugai Pharma	ATP-competitive
p110 δ -specific	CAL-101	Calistoga	ATP-competitive
p110 α -specific	BYL719	Novartis	ATP-competitive
	GDC-0032	Genentech/Roche	ATP-competitive
	INK-1117	Intellikine	ATP-competitive
PI3K/mTOR	PKI-587	Pfizer	ATP-competitive
	BEZ235	Novartis	ATP-competitive
	BGT226	Novartis	ATP-competitive
	PF-4691502	Pfizer	ATP-competitive
	GDC-0980	Genentech/Roche	ATP-competitive
	XL-765	Exelixis	ATP-competitive
	SF1126	Semafor	ATP-competitive/peptide-conjugate
	GSK1059615	GSK	ATP-competitive
TORC1	Everolimus (RAD001)	Novartis	Indirect, FKBP12-mediated
	Temsirolimus (CCI-779)	Wyeth/Pfizer	Indirect, FKBP12-mediated
	Ridaferolimus (AP-23573)	Merck/Ariad	Indirect, FKBP12-mediated
TORC1/TORC2	INK-128	Intellikine	ATP-competitive
	OSI-027	OSI Pharm.	ATP-competitive
	AZD-8055	Astrazeneca	ATP-competitive
AKT	AZD5363	Astrazeneca	ATP-competitive
	GDC-0068	Genentech/Roche	ATP-competitive
	GSK690693	GSK	ATP-competitive
	MK-2206	Merck	Allosteric
	VQD002	Vioquest	ATP-competitive

mTOR, mammalian target of rapamycin; PI3K, phosphatidylinositol 3-kinase.

the ATP-binding cleft of mTOR, and are thus active against both TORC1 and TORC2 [21], are also in phase I trials.

Inhibition of TORC1 relieves negative feedback on activators of PI3K (for example, insulin-like growth factor-1 receptor (IGF-1R), insulin receptor substrate 1 (IRS-1), HER3) [22-24], suggesting that direct inhibitors of PI3K may be more effective. However, inhibition of PI3K or AKT also results in feedback upregulation/activation of several RTKs, which, by providing an input to PI3K, may counteract drug action and/or activate other oncogenic pathways such as the mitogen-activated protein kinase kinase (MEK) pathway [17,25-27]. These data suggest that PI3K/AKT/TORC1 inhibitors could be combined with RTK inhibitors to induce an optimal antitumor effect. Consistent with this notion, studies in

human cancer xenografts have shown that combinations of inhibitors targeting HER2 and PI3K, HER2 and AKT, HER2 and TORC1, or epidermal growth factor receptor (EGFR) and AKT are superior to single-agent treatments [17,22,25,26].

PI3K pathway alterations in ER+ breast cancer

Approximately 75% of primary breast cancers express ER and/or PR. Such hormone receptor expression typically indicates a degree of estrogen dependence for cancer cell growth. Treatments for these patients inhibit ER function either by antagonizing ligand binding to ER (tamoxifen and other selective estrogen receptor modulators (SERMs)), downregulating ER (fulvestrant), or blocking estrogen biosynthesis (aromatase inhibitors (AIs)). Although endocrine therapies have changed the natural

history of hormone-dependent breast cancer, >30% of patients with early ER+ breast cancer relapse within 15 years following adjuvant therapy with tamoxifen, and approximately 20% of patients treated with an AI relapse within 9 years [28,29]. A mechanism of resistance to endocrine therapy involves overexpression of HER2 [30-32]. However, <10% of ER+ breast cancers express high HER2 levels, suggesting that for the majority of ER+ breast cancers, mechanisms of escape from endocrine therapy remain to be elucidated.

In addition to its pro-survival and growth-promoting roles, the PI3K pathway interacts with ER directly [33] and indirectly. ER phosphorylation at Ser167 by AKT or p70S6K increases estrogen-induced, tamoxifen-induced, and ligand-independent ER transcriptional activity [33,34]. Additionally, PI3K and Ras contribute to the modulation of ER and transcription cofactors [35-45]. The activation of ER by growth factor RTK signaling is reciprocated in a feed-forward fashion, whereby ER promotes the transcription of genes encoding receptor ligands, RTKs, and signaling adaptors [45-50]. Clinical evidence further suggests that ER may activate the PI3K pathway. For example, neoadjuvant treatment of patients bearing ER+ breast cancer with the AI letrozole reduces P-AKT_{S473}, P-mTOR_{S2448}, and P-S6 tumor levels; these reductions have been shown to correlate with clinical response [51,52]. Emerging evidence also implicates estrogens in the rapid, non-genomic activation of PI3K via IGF-1R/insulin receptor (InsR), EGFR, Src, PI3K, and MEK [53-56].


PI3K pathway activation has been shown to confer anti-estrogen resistance in various experimental models, including in *PTEN*-deficient cells, and in cells over-expressing HER2, IGF-1R, or an activated mutant of AKT1 [33,44,54]. Tumor cells with acquired endocrine resistance have shown upregulation of IGF-1R, InsR, HER2, and EGFR levels as well as PI3K/AKT/mTOR activation [57-61]. Inhibition of the PI3K pathway reverses such anti-estrogen resistance. However, PI3K or AKT inhibition relieves feedback inhibition of the expression and activation of RTKs, which can contribute to drug resistance [17,25,26]. Interestingly, a recent study showed that in ER+ breast cancer cells treated with the PI3K/mTOR inhibitor BEZ235 or with PI3K siRNA, exogenous estradiol prevented drug- and siRNA-induced apoptosis [62]. Since most breast cancers that adapt to anti-estrogen therapy retain ER, these data imply that unopposed estrogen ligands may protect ER+ tumors from the therapeutic effects of PI3K inhibitors used as single agents.

Clinical evidence suggests that activation of PI3K via overexpression of HER2 or FGFR1, or loss of INPP4B also confers anti-estrogen resistance to patients with ER+ breast cancer [30-32,63,64]. Whether other mutations in

the PI3K pathway correlate with anti-estrogen resistance remains to be determined. *PIK3CA* mutations occur in 28 to 47% of ER+ breast cancers. Interestingly, such mutations correlate with good long-term outcome and lower PI3K and TORC1 activation as assessed by gene expression profiling and immunohistochemistry (IHC) in patients bearing ER+ tumors [6,52,65-68]. Despite these findings, preclinical evidence indicates that combined targeting of PI3K and ER is synergistic [69], suggesting that combinations of anti-estrogens and PI3K pathway inhibitors will be clinically more effective than anti-estrogens alone.

The correlations between *PIK3CA* mutations, good patient outcome, and low PI3K pathway activation (measured by gene expression profiling and IHC) beg the need for alternative methods indicative of PI3K pathway activation to identify ER+ tumors at risk of recurrence. For example, a primary breast tumor gene expression signature of *PTEN* loss, derived from a comparison of *PTEN*-expressing versus *PTEN*-negative tumors by IHC, was predictive of poor relapse-free survival following tamoxifen, while *PTEN* status by IHC was not [8]. Breast cancers of the luminal A and luminal B molecular subtypes (based on gene expression profiling) are typically ER+. However, luminal B tumors benefit less from adjuvant anti-estrogen therapy [70]. Of note, a gene expression signature of PI3K activation, based on tumor levels of a panel of phosphoproteins (for example, P-AKT, P-p70S6K) in ER+ tumors, was enriched in luminal B breast cancers [71]. This suggests that luminal B tumors have higher PI3K activity, which may contribute to their lower response to anti-estrogens compared to luminal A tumors [72]. Similarly, we identified a tumor protein signature of PI3K pathway activation that predicts poor outcome following adjuvant endocrine therapy [58]. Therefore, signatures of PI3K activation may complement mutational analyses for the identification of high-risk, PI3K-driven, ER+ tumors.

Further rationale for combined inhibition of PI3K and ER comes from studies using inhibitors of TORC1 or HER2. In patients with ER+ tumors randomized to neoadjuvant letrozole with or without the TORC1 inhibitor everolimus for 4 months before surgery, the addition of everolimus increased clinical response and suppression of tumor cell proliferation [52]. In the TAMRAD study in patients with metastatic ER+ breast cancer who had progressed on an AI, the addition of everolimus to tamoxifen improved the rate of clinical benefit, time-to-progression, and disease-free survival compared to women receiving tamoxifen alone [73]. Most recently, results from the phase III trial BOLERO-2 showed that treatment with everolimus plus the AI exemestane provided a time-to-progression of 10.6 months compared to 4.1 months with the AI alone in post-menopausal



women with advanced breast cancer who had recurred or progressed on prior endocrine therapy [74].

The ER and PI3K pathways appear to exist in an equilibrium, where tumors with low PI3K activation have

high ER levels, and vice versa [58,71]. The interdependence of these pathways is supported by studies showing that inhibition of HER2 with the antibody trastuzumab or the tyrosine kinase inhibitor lapatinib restores or upregulates ER levels or transcriptional activity in breast cancer cells and patient tumors. Furthermore, treatment with AIs or fulvestrant inhibits the growth of HER2+ tumors that had progressed on trastuzumab or lapatinib [75,76]. These data suggest that combined inhibition of ER and HER2, an RTK that potently activates PI3K, may provide more effective control of ER+/HER2+ tumors. Indeed, two clinical trials showed that the addition of trastuzumab or lapatinib to therapy with an AI increased progression-free survival and clinical benefit compared to the AI alone [77,78].

PI3K alterations in HER2+ breast cancer

Most patients bearing breast cancers with amplification or overexpression of HER2 benefit from anti-HER2 therapy. However, most patients with HER2+ metastatic disease eventually acquire resistance to trastuzumab, lapatinib, and the combination [79-81]. HER2 potently activates PI3K via heterodimerization with HER3, and other PI3K pathway activating mutations often coexist in HER2+ cancers (Table 1). Experimental and clinical evidence suggest that mutational activation of the PI3K pathway confers resistance to HER2-directed therapies, perhaps by providing an additional input to this pathway independent of HER2/HER3 dimers. HER2+ breast cancer cell lines are highly sensitive to PI3K and mTOR inhibitors before and after acquiring resistance to trastuzumab or lapatinib [22,82,83] (Figure 2). These data suggest that these drug-resistant cells remain PI3K-dependent, and that patients with trastuzumab- and/or lapatinib-resistant disease would benefit from PI3K pathway inhibitors.

Retrospective analyses of cohorts of patients with HER2+ metastatic breast cancer have shown that tumors harboring *PIK3CA* mutations and/or decreased levels of PTEN have a poor outcome following treatment with trastuzumab compared to HER2+ tumors with a 'wild-type' PI3K pathway [84-87]. In addition, a neoadjuvant study in patients with HER2+ breast cancer showed that both alterations (*PIK3CA* mutations and PTEN loss) were associated with a statistically lower pathological complete response rate to trastuzumab with chemotherapy. However, tumors with decreased PTEN responded to neoadjuvant therapy with lapatinib followed by trastuzumab and chemotherapy [88]. Pending confirmation of this report, these data suggest that PTEN-deficient HER2+ cancer cells still rely heavily on upstream input from HER2 and, therefore, dual blockade of HER2 with trastuzumab and lapatinib is effective against HER2+/PTEN-deficient breast cancers. A few

studies suggest that combined targeting of HER2 and the PI3K pathway is superior to HER2-directed therapy alone. In patients who had progressed on trastuzumab and chemotherapy, the addition of the TORC1 inhibitor everolimus to trastuzumab and chemotherapy conferred a 19 to 44% objective response rate [89-91]. Preclinical studies also suggest that because of the reactivation of HER3 following inhibition of PI3K/AKT/TORC1 in HER2-overexpressing breast cancer cells, PI3K inhibitors should be given in combination with anti-HER2 therapy in patients with HER2 tumors [22,25,92]. At this time, patients with drug-resistant HER2+ breast cancer are a subgroup of intense focus in exploratory trials with PI3K pathway inhibitors.

PI3K pathway mutations in triple-negative breast cancer

Since ER, PR, and HER2 are established molecular markers associated with response to targeted therapies, ER-/PR-/HER2-negative cancers are loosely grouped as TNBCs. Such cancers occur in 10 to 15% of patients, are associated with earlier age at diagnosis, poor prognosis, and *BRCA1* mutations, and are more prevalent in African-American and Hispanic women [93]. By gene expression profiling, TNBCs cluster separately from ER+ and HER2+ cancers, mainly within the basal-like molecular subtype. A recent analysis revealed that TNBCs can be divided into six subtypes [94]. Interestingly, the 'mesenchymal-like' and 'mesenchymal stem-like' subtypes exhibit enrichment for components of growth factor signaling pathways, including inositol phosphate metabolism. Growth of breast cancer cell lines classified as 'mesenchymal-like,' 'mesenchymal stem-like,' or 'luminal androgen receptor subtype' was inhibited by the PI3K/mTOR inhibitor BEZ235. Cell lines of the 'luminal androgen receptor subtype' exhibit a high frequency of *PIK3CA* mutations. In contrast, PTEN status did not correlate with sensitivity to BEZ235. PTEN has functions outside of the PI3K pathway, including in DNA double-strand break repair. In addition, *BRCA1* mutations impair double-strand break repair and correlate with the presence of *PTEN* mutations [95], and *PTEN* knock-down has been shown to sensitize *BRCA1*-mutant cancer cells to poly(ADP-ribose) polymerase (PARP) inhibition [96,97]. Thus, it is conceivable that PTEN-deficient cells may respond to combined PI3K/PARP-directed therapy.

The standard treatment for patients with TNBC includes mainly DNA-damaging chemotherapy. PI3K pathway mutations have been associated with resistance to such agents, likely by promoting cell survival. Also, DNA damage elicits DNA-dependent protein kinase-mediated phosphorylation of AKT [98]. Preclinical studies in diverse cancer cell types have shown that PI3K

inhibitors enhance the apoptotic effects of DNA-damaging agents [99-102]. Clinical trials are ongoing to test such drug combinations in patients with TNBC.

Conclusions

Somatic mutations in the PI3K pathway identify cancers with aberrant activation of, and potential dependence on, this signaling pathway. These attributes may be useful for the selection of patients for trials with PI3K inhibitors. Indeed, a recent analysis of patients with solid tumors enrolled in phase I trials with PI3K/AKT/mTOR inhibitors showed a higher response rate among patients with *PIK3CA*-mutant versus wild-type *PIK3CA* cancers [103]. This suggests that tumors with gain-of-function mutations in the PI3K pathway depend on PI3K signaling, and this dependence can be exploited in patients with such cancers. There is increasing agreement that initial phase II efficacy studies with PI3K inhibitors in patients with advanced disease should be enriched with, if not limited to, patients harboring mutations and/or activation of this pathway. As with other targeted therapies, only a fraction of patients will likely benefit from single-agent PI3K-directed therapy. PI3K pathway inhibitors are being tested in human trials in combination with inhibitors of HER2, MEK, and ER. Early clinical data suggest that this strategy is feasible and that, as single agents, these drugs are well-tolerated. To determine if inhibition of PI3K confers a benefit compared to standard targeted therapies alone will require randomized clinical trials.

Abbreviations

AI, aromatase inhibitor; EGFR, epidermal growth factor receptor; ER, estrogen receptor; FGFR, fibroblast growth factor receptor; HER, human epidermal growth factor receptor; IGF-1R, insulin-like growth factor-1 receptor; IHC, immunohistochemistry; INPP4B, inositol polyphosphate-4-phosphatase, type II; InsR, insulin receptor; MEK, mitogen-activated protein kinase kinase; mTOR, mammalian target of rapamycin; PARP, poly(ADP-ribose) polymerase; PDK1, phosphoinositide-dependent kinase 1; PH, pleckstrin homology; PI3K, phosphatidylinositol 3-kinase; PIP₂, phosphatidylinositol 4,5-bisphosphate; PIP₃, phosphatidylinositol 3,4,5-trisphosphate; PR, progesterone receptor; PTEN, phosphatase and tensin homolog; RTK, receptor tyrosine kinase; siRNA, small interfering RNA; TNBC, triple-negative breast cancer.

Competing interests

The authors declare that they have no competing interests.

Acknowledgements

This work was supported by the National Institutes of Health K99CA142899 (TWM), K08CA143153 (BNR), Breast Cancer Specialized Program of Research Excellence (SPORE) P50CA98131, Vanderbilt-Ingram Cancer Center Support Grant P30CA68485; a grant from the Breast Cancer Research Foundation (CLA); American Cancer Society Clinical Research Professorship Grant CRP-07-234 (CLA) and Postdoctoral Fellowship 118813-PF-10-070-01-TBG (JTG); the Department of Defense BC093376 (JTG) and BC087465 (BNR); the Lee Jeans Translational Breast Cancer Research Program (CLA); and Stand Up to Cancer/ American Association for Cancer Research Dream Team Translational Cancer Research Grant SU2C-AACR-DT0209 (CLA).

Author details

¹Department of Cancer Biology, Vanderbilt-Ingram Cancer Center, Vanderbilt University, Nashville, TN 37232, USA. ²Breast Cancer Research Program, Vanderbilt-Ingram Cancer Center, Vanderbilt University, Nashville, TN 37232,

USA. ³Department of Medicine, Vanderbilt-Ingram Cancer Center, Vanderbilt University, Nashville, TN 37232, USA. ⁴Division of Hematology-Oncology, VUMC, 2220 Pierce Avenue, 777 PRB, Nashville, TN 37232, USA.

Published: 1 November 2011

References

1. Guertin DA, Sabatini DM: Defining the role of mTOR in cancer. *Cancer Cell* 2007, **12**:9-22.
2. Zhao L, Vogt PK: Helical domain and kinase domain mutations in p110 α of phosphatidylinositol 3-kinase induce gain of function by different mechanisms. *Proc Natl Acad Sci U S A* 2008, **105**:2652-2657.
3. Isakoff SJ, Engelman JA, Irie HY, Luo J, Brachmann SM, Pearlman RV, Cantley LC, Brugge JS: Breast cancer-associated PIK3CA mutations are oncogenic in mammary epithelial cells. *Cancer Res* 2005, **65**:10992-11000.
4. Meyer DS, Brinkhaus H, Muller U, Muller M, Cardiff RD, Bentires-Alj M: Luminal expression of PIK3CA mutant H1047R in the mammary gland induces heterogeneous tumors. *Cancer Res* 2011, **71**:4344-4351.
5. Liu P, Cheng H, Santiago S, Raeder M, Zhang F, Isabella A, Yang J, Semaan DJ, Chen C, Fox EA, Gray NS, Monahan J, Schlegel R, Beroukhir M, Mills GB, Zhao JJ: Oncogenic PIK3CA-driven mammary tumors frequently recur via PI3K pathway-dependent and PI3K pathway-independent mechanisms. *Nat Med* 2011, **17**:1116-1120.
6. Perez-Tenorio G, Alkhori L, Olsson B, Waltersson MA, Nordenskjold B, Rutqvist LE, Skoog L, Stal O: PIK3CA mutations and PTEN loss correlate with similar prognostic factors and are not mutually exclusive in breast cancer. *Clin Cancer Res* 2007, **13**:3577-3584.
7. Saal LH, Holm K, Maurer M, Memeo L, Su T, Wang X, Yu JS, Malmström PO, Mansukhani M, Enoksson J, Hibshoosh H, Borg A, Parsons R: PIK3CA mutations correlate with hormone receptors, node metastasis, and ERBB2, and are mutually exclusive with PTEN loss in human breast carcinoma. *Cancer Res* 2005, **65**:2554-2559.
8. Saal LH, Johansson P, Holm K, Gruvberger-Saal SK, She QB, Maurer M, Koujak S, Ferrando AA, Malmström P, Memeo L, Isola J, Bendahl PO, Rosen N, Hibshoosh H, Ringnér M, Borg A, Parsons R: Poor prognosis in carcinoma is associated with a gene expression signature of aberrant PTEN tumor suppressor pathway activity. *Proc Natl Acad Sci U S A* 2007, **104**:7564-7569.
9. Jia S, Roberts TM, Zhao JJ: Should individual PI3 kinase isoforms be targeted in cancer? *Curr Opin Cell Biol* 2009, **21**:199-208.
10. Oda K, Okada J, Timmerman L, Rodriguez-Viciana P, Stokoe D, Shoji K, Taketani Y, Kuramoto H, Knight ZA, Shokat KM, McCormick F: PIK3CA cooperates with other phosphatidylinositol 3'-kinase pathway mutations to effect oncogenic transformation. *Cancer Res* 2008, **68**:8127-8136.
11. Miron A, Varadi M, Carrasco D, Li H, Luongo L, Kim HJ, Park SY, Cho EY, Lewis G, Kehoe S, Iglehart JD, Dillon D, Allred DC, Macconail L, Gelman R, Polyak K: PIK3CA mutations in situ and invasive breast carcinomas. *Cancer Res* 2010, **70**:5674-5678.
12. Li H, Zhu R, Wang L, Zhu T, Li Q, Chen Q, Wang H, Zhu H: PIK3CA mutations mostly begin to develop in ductal carcinoma of the breast. *Exp Mol Pathol* 2010, **88**:150-155.
13. Dunlap J, Le C, Shukla A, Patterson J, Presnell A, Heinrich MC, Corless CL, Troxell ML: Phosphatidylinositol-3-kinase and AKT1 mutations occur early in breast carcinoma. *Breast Cancer Res Treat* 2010, **120**:409-418.
14. Kalinsky K, Heguy A, Bhanot UK, Patil S, Moynahan ME: PIK3CA mutations rarely demonstrate genotypic intratumoral heterogeneity and are selected for in breast cancer progression. *Breast Cancer Res Treat* 2011, **129**:635-643.
15. Folkes AJ, Ahmadi K, Alderton WK, Alix S, Baker SJ, Box G, Chuckowree IS, Clarke PA, Depledge P, Eccles SA, Friedman LS, Hayes A, Hancox TC, Kugendradas A, Lensun L, Moore P, Olivero AG, Pang J, Patel S, Pergl-Wilson GH, Raynaud FI, Robson A, Saghir N, Salphati L, Sohal S, Utsch MH, Valenti M, Wallweber HJ, Wan NC, Wiesmann C, et al: The identification of 2-(1H-indazol-4-yl)-6-(4-methanesulfonyl-piperazin-1-ylmethyl)-4-morpholin-4-yl-thieno[3,2-d]pyrimidine (GDC-0941) as a potent, selective, orally bioavailable inhibitor of class I PI3 kinase for the treatment of cancer. *J Med Chem* 2008, **51**:5522-5532.
16. Maira SM, Stauffer F, Brueggen J, Furet P, Schnell C, Fritsch C, Brachmann S, Chène P, De Pover A, Schoemaker K, Fabbro D, Gabriel D, Simonen M, Murphy L, Finan P, Sellers W, García-Echeverría C: Identification and characterization of NVP-BEZ235, a new orally available dual phosphatidylinositol 3-kinase/mammalian target of rapamycin inhibitor with potent *in vivo* antitumor

- activity. *Mol Cancer Ther* 2008, **7**:1851-1863.
17. Chandralapaty S, Sawai A, Scaltriti M, Rodrik-Outmezguine V, Grbovic-Huezo O, Serra V, Majumder PK, Baselga J, Rosen N: **AKT inhibition relieves feedback suppression of receptor tyrosine kinase expression and activity.** *Cancer Cell* 2011, **19**:58-71.
 18. Rhodes N, Heerding DA, Duckett DR, Eberwein DJ, Knick VB, Lansing TJ, McConnell RT, Gilmer TM, Zhang SY, Robell K, Kahana JA, Geske RS, Kleymenova EV, Choudhry AE, Lai Z, Leber JD, Minthorn EA, Strum SL, Wood ER, Huang PS, Copeland RA, Kumar R: **Characterization of an Akt kinase inhibitor with potent pharmacodynamic and antitumor activity.** *Cancer Res* 2008, **68**:2366-2374.
 19. Hirai H, Sootome H, Nakatsuru Y, Miyama K, Taguchi S, Tsujioka K, Ueno Y, Hatch H, Majumder PK, Pan BS, Kotani H: **MK-2206, an allosteric Akt inhibitor, enhances antitumor efficacy by standard chemotherapeutic agents or molecular targeted drugs *in vitro* and *in vivo*.** *Mol Cancer Ther* 2010, **9**:1956-1967.
 20. Dowling RJ, Topisirovic I, Fonseca BD, Sonenberg N: **Dissecting the role of mTOR: lessons from mTOR inhibitors.** *Biochim Biophys Acta* 2010, **1804**:433-439.
 21. Bhagwat SV, Gokhale PC, Crew AP, Cooke A, Yao Y, Mantis C, Kahler J, Workman J, Bittner M, Dudkin L, Epstein DM, Gibson NW, Wild R, Arnold LD, Houghton PJ, Pachter JA: **Preclinical characterization of OSI-027, a potent and selective inhibitor of mTORC1 and mTORC2: distinct from rapamycin.** *Mol Cancer Ther* 2011, **10**:1394-1406.
 22. Miller TW, Forbes JT, Shah C, Wyatt SK, Manning HC, Olivares MG, Sanchez V, Dugger TC, de Matos Granja N, Narasanna A, Cook RS, Kennedy JP, Lindsley CW, Arteaga CL: **Inhibition of mammalian target of rapamycin is required for optimal antitumor effect of HER2 inhibitors against HER2-overexpressing cancer cells.** *Clin Cancer Res* 2009, **15**:7266-7276.
 23. O'Reilly KE, Rojo F, She QB, Solit D, Mills GB, Smith D, Lane H, Hofmann F, Hicklin DJ, Ludwig DL, Baselga J, Rosen N: **mTOR inhibition induces upstream receptor tyrosine kinase signaling and activates Akt.** *Cancer Res* 2006, **66**:1500-1508.
 24. Carracedo A, Ma L, Teruya-Feldstein J, Rojo F, Salmena L, Alimonti A, Egia A, Sasaki AT, Thomas G, Kozma SC, Papa A, Nardella C, Cantley LC, Baselga J, Pandolfi PP: **Inhibition of mTORC1 leads to MAPK pathway activation through a PI3K-dependent feedback loop in human cancer.** *J Clin Invest* 2008, **118**:3065-3074.
 25. Chakrabarty A, Sanchez V, Kuba MG, Rinehart C, Arteaga CL: **Feedback upregulation of HER3 (ErbB3) expression and activity attenuates antitumor effect of PI3K inhibitors.** *Proc Natl Acad Sci U S A* 2011 [Epub ahead of print].
 26. Garrett JT, Olivares MG, Rinehart C, Granja-Ingram ND, Sánchez V, Chakrabarty A, Dave B, Cook RS, Pao W, McKinley E, Manning HC, Chang J, Arteaga CL: **Transcriptional and posttranslational up-regulation of HER3 (ErbB3) compensates for inhibition of the HER2 tyrosine kinase.** *Proc Natl Acad Sci U S A* 2011, **108**:5021-5026.
 27. Amin DN, Sergina N, Ahuja D, McMahon M, Blair JA, Wang D, Hann B, Koch KM, Shokat KM, Moasser MM: **Resiliency and vulnerability in the HER2-HER3 tumorigenic driver.** *Sci Transl Med* 2011, **2**:16ra17.
 28. Early Breast Cancer Trialists' Collaborative Group: **Effects of chemotherapy and hormonal therapy for early breast cancer on recurrence and 15-year survival: an overview of the randomised trials.** *Lancet* 2005, **365**:1687-1717.
 29. Forbes JF, Cuzick J, Buzdar A, Howell A, Tobias JS, Baum M: **Effect of anastrozole and tamoxifen as adjuvant treatment for early-stage breast cancer: 100-month analysis of the ATAC trial.** *Lancet Oncol* 2008, **9**:45-53.
 30. Ellis MJ, Tao Y, Young O, White S, Proia AD, Murray J, Renshaw L, Faratian D, Thomas J, Dowsett M, Krause A, Evans DB, Miller WR, Dixon JM: **Estrogen-independent proliferation is present in estrogen-receptor HER2-positive primary breast cancer after neoadjuvant letrozole.** *J Clin Oncol* 2006, **24**:3019-3025.
 31. Arpino G, Green SJ, Allred DC, Lew D, Martino S, Osborne CK, Elledge RM: **HER-2 amplification, HER-1 expression, and tamoxifen response in estrogen receptor-positive metastatic breast cancer: a southwest oncology group study.** *Clin Cancer Res* 2004, **10**:5670-5676.
 32. De Laurentiis M, Arpino G, Massarelli E, Ruggiero A, Carlomagno C, Ciardiello F, Tortora G, D'Agostino D, Caputo F, Cancellio G, Montagna E, Malorni L, Zinno L, Lauria R, Bianco AR, De Placido S: **A meta-analysis on the interaction between HER-2 expression and response to endocrine treatment in advanced breast cancer.** *Clin Cancer Res* 2005, **11**:4741-4748.
 33. Campbell RA, Bhat-Nakshatri P, Patel NM, Constantinidou D, Ali S, Nakshatri H: **Phosphatidylinositol 3-kinase/AKT-mediated activation of estrogen receptor alpha: a new model for anti-estrogen resistance.** *J Biol Chem* 2001, **276**:9817-9824.
 34. Yamnik RL, Digilova A, Davis DC, Brodt ZN, Murphy CJ, Holz MK: **S6 kinase 1 regulates estrogen receptor alpha in control of breast cancer cell proliferation.** *J Biol Chem* 2009, **284**:6361-6369.
 35. Coso OA, Chiariello M, Yu JC, Teramoto H, Crespo P, Xu N, Miki T, Gutkind JS: **The small GTP-binding proteins Rac1 and Cdc42 regulate the activity of the JNK/SAPK signaling pathway.** *Cell* 1995, **81**:1137-1146.
 36. Minden A, Lin A, Claret FX, Abo A, Karin M: **Selective activation of the JNK signaling cascade and c-Jun transcriptional activity by the small GTPases Rac and Cdc42Hs.** *Cell* 1995, **81**:1147-1157.
 37. Logan SK, Falasca M, Hu P, Schlessinger J: **Phosphatidylinositol 3-kinase mediates epidermal growth factor-induced activation of the c-Jun N-terminal kinase signaling pathway.** *Mol Cell Biol* 1997, **17**:5784-5790.
 38. Hibi M, Lin A, Smeal T, Minden A, Karin M: **Identification of an oncoprotein- and UV-responsive protein kinase that binds and potentiates the c-Jun activation domain.** *Genes Dev* 1993, **7**:2135-2148.
 39. Derijard B, Hibi M, Wu IH, Barrett T, Su B, Deng T, Karin M, Davis RJ: **JNK1: a protein kinase stimulated by UV light and Ha-Ras that binds and phosphorylates the c-Jun activation domain.** *Cell* 1994, **76**:1025-1037.
 40. Rauscher FJ, 3rd, Cohen DR, Curran T, Bos TJ, Vogt PK, Bohmann D, Tjian R, Franza BR, Jr.: **Fos-associated protein p39 is the product of the jun proto-oncogene.** *Science* 1988, **240**:1010-1016.
 41. DeNardo DG, Cuba VL, Kim H, Wu K, Lee AV, Brown PH: **Estrogen receptor DNA binding is not required for estrogen-induced breast cell growth.** *Mol Cell Endocrinol* 2007, **277**:13-25.
 42. Petz LN, Ziegler YS, Loven MA, Nardulli AM: **Estrogen receptor alpha and activating protein-1 mediate estrogen responsiveness of the progesterone receptor gene in MCF-7 breast cancer cells.** *Endocrinology* 2002, **143**:4583-4591.
 43. Massarweh S, Schiff R: **Unraveling the mechanisms of endocrine resistance in breast cancer: new therapeutic opportunities.** *Clin Cancer Res* 2007, **13**:1950-1954.
 44. Shou J, Massarweh S, Osborne CK, Wakeling AE, Ali S, Weiss H, Schiff R: **Mechanisms of tamoxifen resistance: increased estrogen receptor-HER2/neu cross-talk in ER/HER2-positive breast cancer.** *J Natl Cancer Inst* 2004, **96**:926-935.
 45. Hurtado A, Holmes KA, Geistlinger TR, Hutcherson IR, Nicholson RI, Brown M, Jiang J, Howat WJ, Ali S, Carroll JS: **Regulation of ERBB2 by oestrogen receptor-PAX2 determines response to tamoxifen.** *Nature* 2008, **456**:663-666.
 46. Mukku VR, Stancel GM: **Regulation of epidermal growth factor receptor by estrogen.** *J Biol Chem* 1985, **260**:9820-9824.
 47. Krywicki RF, Figueroa JA, Jackson JG, Kozelsky TW, Shimasaki S, Von Hoff DD, Yee D: **Regulation of insulin-like growth factor binding proteins in ovarian cancer cells by oestrogen.** *Eur J Cancer* 1993, **29A**:2015-2019.
 48. Martinez-Lacaci I, Saceda M, Plowman GD, Johnson GR, Normanno N, Salomon DS, Dickson RB: **Estrogen and phorbol esters regulate amphiregulin expression by two separate mechanisms in human breast cancer cell lines.** *Endocrinology* 1995, **136**:3983-3992.
 49. Guvakova MA, Surmacz E: **Tamoxifen interferes with the insulin-like growth factor I receptor (IGF-IR) signaling pathway in breast cancer cells.** *Cancer Res* 1997, **57**:2606-2610.
 50. Lee AV, Jackson JG, Gooch JL, Hilsenbeck SG, Coronado-Heinsohn E, Osborne CK, Yee D: **Enhancement of insulin-like growth factor signaling in human breast cancer: estrogen regulation of insulin receptor substrate-1 expression *in vitro* and *in vivo*.** *Mol Endocrinol* 1999, **13**:787-796.
 51. Generali D, Fox SB, Brizzi MP, Allevi G, Bonardi S, Aguggini S, Milani M, Bersiga A, Campo L, Dionisio R, Vergoni F, Giardini R, Dogliotti L, Bottini A, Harris AL, Berruti A: **Down-regulation of phosphatidylinositol 3'-kinase/AKT/molecular target of rapamycin metabolic pathway by primary letrozole-based therapy in human breast cancer.** *Clin Cancer Res* 2008, **14**:2673-2680.
 52. Baselga J, Semiglazov V, van Dam P, Manikhas A, Bellet M, Mayordomo J, Campone M, Kubista E, Greil R, Bianchi G, Steinseifer J, Molloy B, Tokaji E, Gardner H, Phillips P, Stumm M, Lane HA, Dixon JM, Jonat W, Rugo HS: **Phase II randomized study of neoadjuvant everolimus plus letrozole compared with placebo plus letrozole in patients with estrogen receptor-positive breast cancer.** *J Clin Oncol* 2009, **27**:2630-2637.
 53. Simoncini T, Hafezi-Moghadam A, Brazil DP, Ley K, Chin WW, Liao JK: **Interaction of oestrogen receptor with the regulatory subunit of**

- phosphatidylinositol-3-OH kinase. *Nature* 2000, **407**:538-541.
54. Miller TW, Pérez-Torres M, Narasanna A, Guix M, Stål O, Pérez-Tenorio G, Gonzalez-Angulo AM, Hennessy BT, Mills GB, Kennedy JP, Lindsley CW, Arteaga CL: **Loss of Phosphatase and Tensin homologue deleted on chromosome 10 engages ErbB3 and insulin-like growth factor-I receptor signaling to promote antiestrogen resistance in breast cancer.** *Cancer Res* 2009, **69**:4192-4201.
 55. Song RX, McPherson RA, Adam L, Bao Y, Shupnik M, Kumar R, Santen RJ: **Linkage of rapid estrogen action to MAPK activation by ERalpha-Shc association and Shc pathway activation.** *Mol Endocrinol* 2002, **16**:116-127.
 56. Song RX, Fan P, Yue W, Chen Y, Santen RJ: **Role of receptor complexes in the extranuclear actions of estrogen receptor alpha in breast cancer.** *Endocr Relat Cancer* 2006, **13** Suppl 1:S3-13.
 57. Massarweh S, Osborne CK, Creighton CJ, Qin L, Tsimelzon A, Huang S, Weiss H, Rimawi M, Schiff R: **Tamoxifen resistance in breast tumors is driven by growth factor receptor signaling with repression of classic estrogen receptor genomic function.** *Cancer Res* 2008, **68**:826-833.
 58. Miller TW, Hennessy BT, Gonzalez-Angulo AM, Fox EM, Mills GB, Chen H, Higham C, Garcia-Echeverria C, Shyr Y, Arteaga CL: **Hyperactivation of phosphatidylinositol-3 kinase promotes escape from hormone dependence in estrogen receptor-positive human breast cancer.** *J Clin Invest* 2010, **120**:2406-2413.
 59. Fan M, Yan PS, Hartman-Frey C, Chen L, Paik H, Oyer SL, Salisbury JD, Cheng AS, Li L, Abbosh PH, Huang TH, Nephew KP: **Diverse gene expression and DNA methylation profiles correlate with differential adaptation of breast cancer cells to the antiestrogens tamoxifen and fulvestrant.** *Cancer Res* 2006, **66**:11954-11966.
 60. Frogne T, Benjaminsen RV, Sonne-Hansen K, Sorensen BS, Nexø E, Laenkholm AV, Rasmussen LM, Riese DJ 2nd, de Cremoux P, Stenvang J, Lykkesfeldt AE: **Activation of ErbB3, EGFR and Erk is essential for growth of human breast cancer cell lines with acquired resistance to fulvestrant.** *Breast Cancer Res Treat* 2009, **114**:263-275.
 61. Massarweh S, Osborne CK, Jiang S, Wakeling AE, Rimawi M, Mohsin SK, Hilsenbeck S, Schiff R: **Mechanisms of tumor regression and resistance to estrogen deprivation and fulvestrant in a model of estrogen receptor-positive, HER-2/neu-positive breast cancer.** *Cancer Res* 2006, **66**:8266-8273.
 62. Crowder RJ, Phommaly C, Tao Y, Hoog J, Luo J, Perou CM, Parker JS, Miller MA, Huntsman DG, Lin L, Snider J, Davies SR, Olson JA Jr, Watson MA, Saporita A, Weber JD, Ellis MJ: **PIK3CA and PIK3CB inhibition produce synthetic lethality when combined with estrogen deprivation in estrogen receptor-positive breast cancer.** *Cancer Res* 2009, **69**:3955-3962.
 63. Turner N, Pearson A, Sharpe R, Lambros M, Geyer F, Lopez-Garcia MA, Natrajan R, Marchio C, Iorns E, Mackay A, Gillett C, Grigoriadis A, Tutt A, Reis-Filho JS, Ashworth A: **FGFR1 amplification drives endocrine therapy resistance and is a therapeutic target in breast cancer.** *Cancer Res* 2010, **70**:2085-2094.
 64. Gewinnee C, Wang ZC, Richardson A, Teruya-Feldstein J, Etemadmoghadam D, Bowtell D, Barretina J, Lin WM, Rameh L, Salmena L, Pandolfi PP, Cantley LC: **Evidence that inositol polyphosphate 4-phosphatase type II is a tumor suppressor that inhibits PI3K signaling.** *Cancer Cell* 2009, **16**:115-125.
 65. Loi S, Haibe-Kains B, Majaj S, Lallemand F, Durbecq V, Larsimont D, Gonzalez-Angulo AM, Pusztai L, Symmans WF, Bardelli A, Ellis P, Tutt AN, Gillett CE, Hennessy BT, Mills GB, Phillips WA, Piccart MJ, Speed TP, McArthur GA, Sotiriou C: **PIK3CA mutations associated with gene signature of low mTORC1 signaling and better outcomes in estrogen receptor-positive breast cancer.** *Proc Natl Acad Sci U S A* 2010, **107**:10208-10213.
 66. Stemke-Hale K, Gonzalez-Angulo AM, Lluch A, Neve RM, Kuo WL, Davies M, Carey M, Hu Z, Guan Y, Sahin A, Symmans WF, Pusztai L, Nolden LK, Horlings H, Berns K, Hung MC, van de Vijver MJ, Valero V, Gray JW, Bernards R, Mills GB, Hennessy BT: **An integrative genomic and proteomic analysis of PIK3CA, PTEN, and AKT mutations in breast cancer.** *Cancer Res* 2008, **68**:6084-6091.
 67. Ellis MJ, Lin L, Crowder R, Tao Y, Hoog J, Snider J, Davies S, DeSchryver K, Evans DB, Steinseifer J, Bandaru R, Liu W, Gardner H, Semiglazov V, Watson M, Hunt K, Olson J, Baselga J: **Phosphatidylinositol-3-kinase alpha catalytic subunit mutation and response to neoadjuvant endocrine therapy for estrogen receptor positive breast cancer.** *Breast Cancer Res Treat* 2010, **119**:379-390.
 68. Campbell IG, Russell SE, Choong DY, Montgomery KG, Ciavarella ML, Hooi CS, Cristiano BE, Pearson RB, Phillips WA: **Mutation of the PIK3CA gene in ovarian and breast cancer.** *Cancer Res* 2004, **64**:7678-7681.
 69. Miller TW, Balko JM, Fox EM, Ghazoui Z, Dunbier A, Anderson H, Dowsett M, Jiang A, Smith RA, Maira S-M, Manning HC, Gonzalez-Angulo AM, Mills GB, Higham C, Chanthaphychith S, Kuba MG, Miller WR, Shyr Y, Arteaga CL: **ERα-dependent E2F transcription can mediate resistance to estrogen deprivation in human breast cancer.** *Cancer Discovery* 2011, **1**:338-351.
 70. Sotiriou C, Pusztai L: **Gene-expression signatures in breast cancer.** *N Engl J Med* 2009, **360**:790-800.
 71. Creighton CJ, Fu X, Hennessy BT, Casa AJ, Zhang Y, Gonzalez-Angulo AM, Lluch A, Gray JW, Brown PH, Hilsenbeck SG, Osborne CK, Mills GB, Lee AV, Schiff R: **Proteomic and transcriptomic profiling reveals a link between the PI3K pathway and lower estrogen receptor (ER) levels and activity in ER+ breast cancer.** *Breast Cancer Res* 2010, **12**:R40.
 72. Sørli T, Perou CM, Tibshirani R, Aas T, Geisler S, Johnsen H, Hastie T, Eisen MB, van de Rijn M, Jeffrey SS, Thorsen T, Quist H, Matese JC, Brown PO, Botstein D, Eystein Lønning P, Børresen-Dale AL: **Gene expression patterns of breast carcinomas distinguish tumor subclasses with clinical implications.** *Proc Natl Acad Sci U S A* 2001, **98**:10869-10874.
 73. Bachelot T, Bourgier C, Cropet C, Guastalla J-P, Ferrero J-M, Leger-Falandry C, Soulie P, Eymard J-C, Debled M, Spaeth D, Legouffe E, Delozier T, El Kouri C, Chidiac J: **TAMRAD: A GINECO randomized phase II trial of everolimus in combination with tamoxifen versus tamoxifen alone in patients (pts) with hormone-receptor positive, HER2 negative metastatic breast cancer (MBC) with prior exposure to aromatase inhibitors (AI).** *Cancer Res* 2010, **70**(24S):Abstract S1-6.
 74. Baselga J: **Everolimus in combination with exemestane for postmenopausal women with advanced breast cancer who are refractory to letrozole or anastrozole: results of the BOLERO-2 phase III trial.** *European Multidisciplinary Cancer Congress Presentation of late breaking abstract No 9LBA September 26, 2011* 2011.
 75. Munzone E, Curigliano G, Rocca A, Bonizzi G, Renne G, Goldhirsch A, Nole F: **Reverting estrogen-receptor-negative phenotype in HER-2-overexpressing advanced breast cancer patients exposed to trastuzumab plus chemotherapy.** *Breast Cancer Res* 2006, **8**:R4.
 76. Xia W, Bacus S, Hegde P, Husain I, Strum J, Liu L, Pauluzzo G, Lyass L, Trusk P, Hill J, Harris J, Spector NL: **A model of acquired autoresistance to a potent ErbB2 tyrosine kinase inhibitor and a therapeutic strategy to prevent its onset in breast cancer.** *Proc Natl Acad Sci U S A* 2006, **103**:7795-7800.
 77. Johnston S, Phippen J Jr, Pivrot X, Lichinitser M, Sadeghi S, Dieras V, Gomez HL, Romieu G, Manikhas A, Kennedy MJ, Press MF, Maltzman J, Florance A, O'Rourke L, Oliva C, Stein S, Pegram M: **Lapatinib combined with letrozole versus letrozole and placebo as first-line therapy for postmenopausal hormone receptor-positive metastatic breast cancer.** *J Clin Oncol* 2009, **27**:5538-5546.
 78. Kaufman B, Mackey JR, Clemens MR, Bapsy PP, Vaid A, Wardley A, Tjulandin S, Jahn M, Lehle M, Feyereislova A, Révil C, Jones A: **Trastuzumab plus anastrozole versus anastrozole alone for the treatment of postmenopausal women with human epidermal growth factor receptor 2-positive, hormone receptor-positive metastatic breast cancer: results from the randomized phase III TAnDEM study.** *J Clin Oncol* 2009, **27**:5529-5537.
 79. Cobleigh MA, Vogel CL, Tripathy D, Robert NJ, Scholl S, Fehrenbacher L, Wolter JM, Paton V, Shak S, Lieberman G, Slamon DJ: **Multinational study of the efficacy and safety of humanized anti-HER2 monoclonal antibody in women who have HER2-overexpressing metastatic breast cancer that has progressed after chemotherapy for metastatic disease.** *J Clin Oncol* 1999, **17**:2639-2648.
 80. Blackwell KL, Burstein HJ, Storniolo AM, Rugo H, Sledge G, Koehler M, Ellis C, Casey M, Vukelja S, Bischoff J, Baselga J, O'Shaughnessy J: **Randomized study of Lapatinib alone or in combination with trastuzumab in women with ErbB2-positive, trastuzumab-refractory metastatic breast cancer.** *J Clin Oncol* 2010, **28**:1124-1130.
 81. Cameron D, Casey M, Press M, Lindquist D, Pienkowski T, Romieu CG, Chan S, Jagiello-Gruszfeld A, Kaufman B, Crown J, Chan A, Campone M, Viens P, Davidson N, Gorbounova V, Raats JJ, Skarlos D, Newstat B, Roychowdhury D, Paoletti P, Oliva C, Rubin S, Stein S, Geyer CE: **A phase III randomized comparison of lapatinib plus capecitabine versus capecitabine alone in women with advanced breast cancer that has progressed on trastuzumab: updated efficacy and biomarker analyses.** *Breast Cancer Res Treat* 2008, **112**:533-543.
 82. Eichhorn PJ, Gili M, Scaltriti M, Serra V, Guzman M, Nijkamp W, Beijersbergen RL, Valero V, Seoane J, Bernards R, Baselga J: **Phosphatidylinositol 3-kinase hyperactivation results in lapatinib resistance that is reversed by the mTOR/phosphatidylinositol 3-kinase inhibitor NVP-BE235.** *Cancer Res* 2008, **68**:9221-9230.

83. Rexer BN, Ham AJ, Rinehart C, Hill S, de Matos Granja-Ingram N, González-Angulo AM, Mills GB, Dave B, Chang JC, Liebler DC, Arteaga CL: **Phosphoproteomic mass spectrometry profiling links Src family kinases to escape from HER2 tyrosine kinase inhibition.** *Oncogene* 2011, **30**:4163-4174.
84. Esteva FJ, Guo H, Zhang S, Santa-Maria C, Stone S, Lanchbury JS, Sahin AA, Hortobagyi GN, Yu D: **PTEN, PIK3CA, p-AKT, and p-p70S6K status: association with trastuzumab response and survival in patients with HER2-positive metastatic breast cancer.** *Am J Pathol* 2010, **177**:1647-1656.
85. Razis E, Bobos M, Kotoula V, Eleftheraki AG, Kalofonos HP, Pavlakis K, Papakostas P, Aravantinos G, Rigakos G, Efstratiou I, Petraki K, Bafaloukos D, Kostopoulos I, Pectasides D, Kalogeras KT, Skarlos D, Fountzilias G: **Evaluation of the association of PIK3CA mutations and PTEN loss with efficacy of trastuzumab therapy in metastatic breast cancer.** *Breast Cancer Res Treat* 2011, **128**:447-456.
86. Berns K, Horlings HM, Hennessy BT, Madiredjo M, Hijmans EM, Beelen K, Linn SC, Gonzalez-Angulo AM, Stemke-Hale K, Hauptmann M, Beijersbergen RL, Mills GB, van de Vijver MJ, Bernards R: **A functional genetic approach identifies the PI3K pathway as a major determinant of trastuzumab resistance in breast cancer.** *Cancer Cell* 2007, **12**:395-402.
87. Wang L, Zhang Q, Zhang J, Sun S, Guo H, Jia Z, Wang B, Shao Z, Wang Z, Hu X: **PI3K pathway activation results in low efficacy of both trastuzumab and lapatinib.** *BMC Cancer* 2011, **11**:248.
88. Dave B, Migliaccio I, Gutierrez MC, Wu MF, Chamness GC, Wong H, Narasanna A, Chakraborty A, Hilsenbeck SG, Huang J, Rimawi M, Schiff R, Arteaga C, Osborne CK, Chang JC: **Loss of phosphatase and tensin homolog or phosphoinositol-3 kinase activation and response to trastuzumab or lapatinib in human epidermal growth factor receptor 2-overexpressing locally advanced breast cancers.** *J Clin Oncol* 2011, **29**:166-173.
89. Morrow PH, Wulf GM, Booser DJ, Moore JA, Flores PR, Krop IE, Winer EP, Hortobagyi GN, Yu D, Esteva FJ: **Phase I/II trial of everolimus (RAD001) and trastuzumab in patients with trastuzumab-resistant, HER2-overexpressing breast cancer.** *J Clin Oncol* 2010, **28**(15):abstract 1014.
90. Jerusalem G, Fasolo A, Dieras V, Cardoso F, Bergh J, Vittori L, Zhang Y, Massacesi C, Sahnoud T, Gianni L: **Phase I trial of oral mTOR inhibitor everolimus in combination with trastuzumab and vinorelbine in pre-treated patients with HER2-overexpressing metastatic breast cancer.** *Breast Cancer Res Treat* 2011, **125**:447-455.
91. Andre F, Campone M, O'Regan R, Manlius C, Massacesi C, Sahnoud T, Mukhopadhyay P, Soria JC, Naughton M, Hurvitz SA: **Phase I study of everolimus plus weekly paclitaxel and trastuzumab in patients with metastatic breast cancer pretreated with trastuzumab.** *J Clin Oncol* 2010, **28**:5110-5115.
92. Serra V, Scaltriti M, Prudkin L, Eichhorn PJ, Ibrahim YH, Chandarlapaty S, Markman B, Rodriguez O, Guzman M, Rodriguez S, Gili M, Russillo M, Parra JL, Singh S, Arribas J, Rosen N, Baselga J: **PI3K inhibition results in enhanced HER signaling and acquired ERK dependency in HER2-overexpressing breast cancer.** *Oncogene* 2011, **30**:2547-2557.
93. Reddy KB: **Triple-negative breast cancers: an updated review on treatment options.** *Curr Oncol* 2011, **18**:e173-179.
94. Lehmann BD, Bauer JA, Chen X, Sanders ME, Chakravarthy AB, Shyr Y, Pietenpol JA: **Identification of human triple-negative breast cancer subtypes and preclinical models for selection of targeted therapies.** *J Clin Invest* 2011, **121**:2750-2767.
95. Saal LH, Gruvberger-Saal SK, Persson C, Lövgren K, Jumppanen M, Staaf J, Jönsson G, Pires MM, Maurer M, Holm K, Koujak S, Subramaniam S, Vallon-Christersson J, Olsson H, Su T, Memeo L, Ludwig T, Ethier SP, Krogh M, Szabolcs M, Murty VV, Isola J, Hishoosh H, Parsons R, Borg A: **Recurrent gross mutations of the PTEN tumor suppressor gene in breast cancers with deficient DSB repair.** *Nat Genet* 2008, **40**:102-107.
96. Mendes-Pereira AM, Martin SA, Brough R, McCarthy A, Taylor JR, Kim JS, Waldman T, Lord CJ, Ashworth A: **Synthetic lethal targeting of PTEN mutant cells with PARP inhibitors.** *EMBO Mol Med* 2009, **1**:315-322.
97. Farmer H, McCabe N, Lord CJ, Tutt AN, Johnson DA, Richardson TB, Santarosa M, Dillon KJ, Hickson I, Knights C, Martin NM, Jackson SP, Smith GC, Ashworth A: **Targeting the DNA repair defect in BRCA mutant cells as a therapeutic strategy.** *Nature* 2005, **434**:917-921.
98. Bozulic L, Surucu B, Hynx D, Hemmings BA: **PKBalpha/Akt1 acts downstream of DNA-PK in the DNA double-strand break response and promotes survival.** *Mol Cell* 2008, **30**:203-213.
99. Hu L, Hofmann J, Lu Y, Mills GB, Jaffe RB: **Inhibition of phosphatidylinositol 3'-kinase increases efficacy of paclitaxel in *in vitro* and *in vivo* ovarian cancer models.** *Cancer Res* 2002, **62**:1087-1092.
100. Ng SSW, Tsao MS, Chow S, Hedley DW: **Inhibition of phosphatidylinositol 3-kinase enhances gemcitabine-induced apoptosis in human pancreatic cancer cells.** *Cancer Res* 2000, **60**:5451-5455.
101. Westhoff MA, Kandenwein JA, Karl S, Vellanki SH, Braun V, Eramo A, Antoniadis G, Debatin KM, Fulda S: **The pyridinylfuranopyrimidine inhibitor, PI-103, chemosensitizes glioblastoma cells for apoptosis by inhibiting DNA repair.** *Oncogene* 2009, **28**:3586-3596.
102. Wallin JJ, Guan J, Prior WW, Edgar KA, Kassees R, Sampath D, Belvin M, Friedman LS: **Nuclear phospho-Akt increase predicts synergy of PI3K inhibition and doxorubicin in breast and ovarian cancer.** *Sci Transl Med* 2010, **2**:48ra66.
103. Janku F, Tsimberidou AM, Garrido-Laguna I, Wang X, Luthra R, Hong DS, Naing A, Falchook GS, Moroney JW, Piha-Paul SA, Wheler JJ, Moulder SL, Fu S, Kurzrock R: **PIK3CA mutations in patients with advanced cancers treated with PI3K/AKT/mTOR axis inhibitors.** *Mol Cancer Ther* 2011, **10**:558-565.
104. Shoman N, Klassen S, McFadden A, Bickis MG, Torlakovic E, Chibbar R: **Reduced PTEN expression predicts relapse in patients with breast carcinoma treated by tamoxifen.** *Mod Pathol* 2005, **18**:250-259.
105. Gonzalez-Angulo AM, Ferrer-Lozano J, Stemke-Hale K, Sahin A, Liu S, Barrera JA, Burgues O, Lluch AM, Chen H, Hortobagyi GN, Mills GB, Meric-Bernstam F: **PI3K pathway mutations and PTEN levels in primary and metastatic breast cancer.** *Mol Cancer Ther* 2011, **10**:1093-1101.
106. Hennessy BT, Gonzalez-Angulo AM, Stemke-Hale K, Gilcrease MZ, Krishnamurthy S, Lee JS, Fridlyand J, Sahin A, Agarwal R, Joy C, Liu W, Stivers D, Baggerly K, Carey M, Lluch A, Monteagudo C, He X, Weigman V, Fan C, Palazzo J, Hortobagyi GN, Nolden LK, Wang NJ, Valero V, Gray JW, Perou CM, Mills GB: **Characterization of a naturally occurring breast cancer subset enriched in epithelial-to-mesenchymal transition and stem cell characteristics.** *Cancer Res* 2009, **69**:4116-4124.
107. Samuels Y, Wang Z, Bardelli A, Silliman N, Ptak J, Szabo S, Yan H, Gazdar A, Powell SM, Riggins GJ, Willson JK, Markowitz S, Kinzler KW, Vogelstein B, Velculescu VE: **High frequency of mutations of the PIK3CA gene in human cancers.** *Science* 2004, **304**:554.
108. Law JH, Habibi G, Hu K, Masoudi H, Wang MY, Stratford AL, Park E, Gee JM, Finlay P, Jones HE, Nicholson RI, Carboni J, Gottardis M, Pollak M, Dunn SE: **Phosphorylated insulin-like growth factor-I/insulin receptor is present in all breast cancer subtypes and is related to poor survival.** *Cancer Res* 2008, **68**:10238-10246.
109. Fu P, Ibusuki M, Yamamoto Y, Hayashi M, Murakami K, Zheng S, Iwase H: **Insulin-like growth factor-1 receptor gene expression is associated with survival in breast cancer: a comprehensive analysis of gene copy number, mRNA and protein expression.** *Breast Cancer Res Treat* 2011, **130**:307-317.
110. Elbaoumy Elsheikh S, Green AR, Lambros MB, Turner NC, Grainge MJ, Powe D, Ellis IO, Reis-Filho JS: **FGFR1 amplification in breast carcinomas: a chromogenic *in situ* hybridisation analysis.** *Breast Cancer Res* 2007, **9**:R23.
111. Monni O, Barlund M, Mousse S, Kononen J, Sauter G, Heiskanen M, Paavola P, Avela K, Chen Y, Bittner ML, Kallioniemi A: **Comprehensive copy number and gene expression profiling of the 17q23 amplicon in human breast cancer.** *Proc Natl Acad Sci U S A* 2001, **98**:5711-5716.
112. Fedele CG, Ooms LM, Ho M, Vieusseux J, O'Toole SA, Millar EK, Lopez-Knowles E, Sriravana A, Gurusu R, Baglietto L, Giles GG, Bailey CG, Rasko JE, Shields BJ, Price JT, Majerus PW, Sutherland RL, Tiganis T, McLean CA, Mitchell CA: **Inositol polyphosphate 4-phosphatase II regulates PI3K/Akt signaling and is lost in human basal-like breast cancers.** *Proc Natl Acad Sci U S A* 2010, **107**:22231-22236.
113. Jaiswal BS, Janakiraman V, Kijavini NM, Chaudhuri S, Stern HM, Wang W, Kan Z, Dbouk HA, Peters BA, Waring P, Dela Vega T, Kenski DM, Bowman KK, Lorenzo M, Li H, Wu J, Modrusan Z, Stinson J, Eby M, Yue P, Kaminker JS, de Sauvage FJ, Backer JM, Seshagiri S: **Somatic mutations in p85alpha promote tumorigenesis through class IA PI3K activation.** *Cancer Cell* 2009, **16**:463-474.
114. Carpten JD, Faber AL, Horn C, Donoho GP, Briggs SL, Robbins CM, Hostetter G, Boguslawski S, Moses TY, Savage S, Uhlir M, Lin A, Du J, Qian YW, Zeckner DJ, Tucker-Kellogg G, Touchman J, Patel K, Mousse S, Bittner M, Schevitz R, Lai MH, Blanchard KL, Thomas JE: **A transforming mutation in the pleckstrin homology domain of AKT1 in cancer.** *Nature* 2007, **448**:439-444.
115. Bellacosa A, de Feo D, Godwin AK, Bell DW, Cheng JQ, Altomare DA, Wan M, Dubeau L, Scambia G, Masciullo V, Ferrandina G, Benedetti Panici P, Mancuso S, Neri G, Testa JR: **Molecular alterations of the AKT2 oncogene in ovarian and breast carcinomas.** *Int J Cancer* 1995, **64**:280-285.

116. Al-Kuraya K, Schraml P, Torhorst J, Tapia C, Zaharieva B, Novotny H, Spichtin H, Maurer R, Mirlacher M, Köchli O, Zuber M, Dieterich H, Mross F, Wilber K, Simon R, Sauter G: **Prognostic relevance of gene amplifications and coamplifications in breast cancer.** *Cancer Res* 2004, **64**:8534-8540.
117. Maurer M, Su T, Saal LH, Koujak S, Hopkins BD, Barkley CR, Wu J, Nandula S, Dutta B, Xie Y, Chin YR, Kim DJ, Ferris JS, Gruvberger-Saal SK, Laakso M, Wang X, Memeo L, Rojzman A, Matos T, Yu JS, Cordon-Cardo C, Isola J, Terry MB, Toker A, Mills GB, Zhao JJ, Murty VV, Hibshoosh H, Parsons R: **3-Phosphoinositide-dependent kinase 1 potentiates upstream lesions on the phosphatidylinositol 3-kinase pathway in breast carcinoma.** *Cancer Res* 2009, **69**:6299-6306.
118. Rochlitz CF, Scott GK, Dodson JM, Liu E, Dollbaum C, Smith HS, Benz CC: **Incidence of activating ras oncogene mutations associated with primary and metastatic human breast cancer.** *Cancer Res* 1989, **49**:357-360.
119. Di Nicolantonio F, Arena S, Tabernero J, Grosso S, Molinari F, Macarulla T, Russo M, Cancelliere C, Zecchin D, Mazzucchelli L, Sasazuki T, Shirasawa S, Geuna M, Frattini M, Baselga J, Gallicchio M, Biffo S, Bardelli A: **Deregulation of the PI3K and KRAS signaling pathways in human cancer cells determines their response to everolimus.** *J Clin Invest* 2010, **120**:2858-2866.

doi:10.1186/bcr3039

Cite this article as: Miller TW, et al.: Mutations in the phosphatidylinositol 3-kinase pathway: role in tumor progression and therapeutic implications in breast cancer. *Breast Cancer Research* 2011, **13**:224.