

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

2,5-Dimethoxybenzonitrile

Bernhard Bugenhagen,^a Yosef Al Jasem^b and Thies Thiemann^{c*}

^aFachbereich Chemie, University of Hamburg, Martin-Luther-King-Platz 6, 20146 Hamburg, Germany, ^bDepartment of Chemical Engineering, United Arab Emirates University, AL Ain, Abu Dhabi, United Arab Emirates, and ^cDepartment of Chemistry, United Arab Emirates University, AL Ain, Abu Dhabi, United Arab Emirates

Correspondence e-mail: thies@uaeu.ac.ae


Received 23 October 2013; accepted 15 November 2013

Key indicators: single-crystal X-ray study; $T = 100$ K; mean $\sigma(\text{C}-\text{C}) = 0.002$ Å; R factor = 0.044; wR factor = 0.116; data-to-parameter ratio = 16.1.

In the title molecule, $\text{C}_9\text{H}_9\text{NO}_2$, the non-H atoms are essentially coplanar with a maximum deviation of 0.027 (2) Å for the C atom of one of the methyl groups. In the crystal, the molecules are arranged into centrosymmetric pairs *via* pairs of C—H...O and C—H...N interactions whereas π – π stacking interactions between the benzene rings [centroid–centroid distance 3.91001 (15) Å] organize them into polymeric strands propagating along the a -axis direction. There is a step of 0.644 (2) Å between the two planar parts of the centrosymmetric pair. In neighboring strands related by the n -glide operation, the aromatic rings are tilted by 29.08 (2)°.

Related literature

For the use of the title compound as a key reagent in the synthesis of pharmaceutically active heterocycles, see: Bergeron *et al.* (2006); Delgado *et al.* (1987). For another method of preparation of the title compound, see: Ushijima *et al.* (2012). For the crystal structures of aromatic nitriles, see: Buschmann *et al.* (1995); Zabinski *et al.* (2007); Zanotti *et al.* (1980).



Experimental

Crystal data

$\text{C}_9\text{H}_9\text{NO}_2$
 $M_r = 163.17$
 Monoclinic, $P2_1/n$
 $a = 3.91001$ (15) Å
 $b = 11.3347$ (4) Å
 $c = 17.8432$ (6) Å
 $\beta = 93.400$ (3)°

$V = 789.40$ (5) Å³
 $Z = 4$
 Mo $K\alpha$ radiation
 $\mu = 0.10$ mm⁻¹
 $T = 100$ K
 $0.60 \times 0.25 \times 0.23$ mm

Data collection

Agilent SuperNova (Dual, Cu at zero, Atlas) diffractometer
 Absorption correction: multi-scan (*CrysAlis PRO*; Agilent, 2013)
 $T_{\min} = 0.899$, $T_{\max} = 1.000$

3225 measured reflections
 1785 independent reflections
 1374 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.026$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.044$
 $wR(F^2) = 0.116$
 $S = 1.06$
 1785 reflections

111 parameters
 H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.21$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.24$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{C8}-\text{H8B}\cdots\text{O1}^i$	0.98	2.65	3.428 (2)	136
$\text{C8}-\text{H8B}\cdots\text{N1}^i$	0.98	2.73	3.504 (2)	136
$\text{C9}-\text{H9B}\cdots\text{N1}^{ii}$	0.98	2.71	3.640 (2)	158

Symmetry codes: (i) $-x + 3, -y + 2, -z + 1$; (ii) $-x + \frac{1}{2}, y + \frac{1}{2}, -z + \frac{1}{2}$.

Data collection: *CrysAlis PRO* (Agilent, 2013); cell refinement: *CrysAlis PRO*; data reduction: *CrysAlis PRO*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008) within *OLEX2* (Dolomanov *et al.*, 2009); molecular graphics: *PLATON* (Spek, 2009); *Mercury* (Macrae *et al.*, 2008); software used to prepare material for publication: *SHELXL97* and *PLATON*.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: GK2594).

References

- Agilent (2013). *CrysAlis PRO*. Agilent Technologies, Yarnton, England.
 Bergeron, R. J., Wiegand, J., McManis, J. S. & Bharti, N. (2006). *J. Med. Chem.* **49**, 7032–7043.
 Buschmann, W. E., Arif, A. M. & Miller, J. S. (1995). *J. Chem. Soc. Chem. Commun.* pp. 2343–2344.
 Delgado, A., Mauleon, D., Rosell, G., Salas, M. L. & Najjar, J. (1987). *Anal. Quim. Ser. C*, **83**, 90–95.
 Dolomanov, O. V., Bourhis, L. J., Gildea, R. J., Howard, J. A. K. & Puschmann, H. (2009). *J. Appl. Cryst.* **42**, 339–341.
 Macrae, C. F., Bruno, I. J., Chisholm, J. A., Edgington, P. R., McCabe, P., Pidcock, E., Rodriguez-Monge, L., Taylor, R., van de Streek, J. & Wood, P. A. (2008). *J. Appl. Cryst.* **41**, 466–470.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.
 Ushijima, S., Moriyama, K. & Togo, H. (2012). *Tetrahedron*, **68**, 4588–4595.
 Zabinski, J., Wolska, I. & Maciejewska, D. (2007). *J. Mol. Struct.* **833**, 74–81.
 Zanotti, G., Bardi, R. & Del Pra, A. (1980). *Acta Cryst.* **B36**, 168–171.

supplementary materials

Acta Cryst. (2013). E69, o1808 [doi:10.1107/S1600536813031309]

2,5-Dimethoxybenzotrile

Bernhard Bugenhagen, Yosef Al Jasem and Thies Thiemann

1. Comment

The aromatic ring (C1—C6) of the title compound is almost coplanar with non-H atoms of all substituents, with torsion angles of 3.2 (2)°, 2.5 (2)° and 178.7 (1)° for the methoxy group (C3—C2—O1—C8), for the methoxy group (C4—C5—O2—C9) and for the nitrile group (C5—C6—C1—C7), respectively. With the length of 1.1492 (19) Å, the triple bond of the nitrile group (C≡N) is at the higher end of the acceptable range of cyano bond lengths (Buschmann *et al.*, 1995; Zanotti *et al.*, 1980), but longer than in comparable alkoxy-substituted benzonitriles (Zabinski *et al.*, 2007). The molecules of the title compound arrange themselves in pairs through C8—H8B···O1 and C8—H8B···N1 interactions (Table 1, Fig. 2). In one pair, the average plane of the aromatic ring (C1—C6) of one molecule has an off-set of 0.644 (2) Å to the respective plane in the other molecule. The stacked centrosymmetric dimers form strands propagating along the *a* axis. Each pair in one strand forms four close contacts C9—H9B···N1 (Table 1) with four pairs of the four neighboring strands (Figure 3). The average plane of the aromatic ring (C1—C6) of a molecule in one strand forms an angle of 29.08 (2)° with the respective average plane of another molecule in the neighboring strand, with the molecules linked by C9—H9B···N1 close contact.

2. Experimental

To triphenylphosphine (870 mg, 3.3 mmol) in dry CH₂Cl₂ (10 ml) was added bromotrichloromethane (650 mg, 3.3 mmol), and the resulting mixture was stirred at rt for 20 min, during which the solution turned from yellow to red-brownish in color. Thereafter, 2,5-dimethoxybenzaloxime (552 mg, 3.05 mmol) was added. The reaction mixture was kept under reflux for 25 min. Then, triphenylphosphine (870 mg, 3.3 mmol) was added, and the mixture stirred for 8 h at reflux. The cooled reaction mixture was concentrated *in vacuo* and subjected directly to column chromatography on silica gel (CH₂Cl₂ – hexane 5: 1) to give the title compound (195 mg, 39%) as colorless needles; m.p. 360 - 361 K (Lit. 354 - 358 K; Ushijima *et al.*, 2012); n_{max} (KBr/cm⁻¹) 2224 (CN), 1582, 1508, 1420, 1287, 1237, 1120, 1039, 879, 815, 753, 704, 488; d_{H} (400 MHz, CDCl₃) 3.77 (3H, s, OCH₃), 3.87 (3H, s, OCH₃), 6.89 (1H, d, ³J = 8.8 Hz), 7.04 (1H, d, ⁴J = 2.8 Hz), 7.07 (1H, dd, ³J = 8.8 Hz, ⁴J = 2.8 Hz); d_{C} (100.5 MHz, CDCl₃) 55.9 (OCH₃), 56.4 (OCH₃), 101.7 (C_{quat}), 112.6 (CH), 116.4 (C_{quat}), 117.5 (CH), 120.8 (CH), 153.1 (C_{quat}), 153.7 (C_{quat}); MS (EI, 70 eV) *m/z* (%) 163 (*M*⁺, 100).

3. Refinement

All carbon-bound hydrogen atoms were placed in calculated positions with C—H distances of 0.95 - 0.98 Å and refined as riding with $U_{\text{iso}}(\text{H}) = xU_{\text{eq}}(\text{C})$, where $x = 1.5$ for methyl and $x = 1.2$ for all other H-atoms.

Computing details

Data collection: *CrysAlis PRO* (Agilent, 2013); cell refinement: *CrysAlis PRO* (Agilent, 2013); data reduction: *CrysAlis PRO* (Agilent, 2013); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008) within OLEX2 (Dolomanov *et al.*, 2009); molecular graphics: *PLATON* (Spek,

2009); *Mercury* (Macrae *et al.*, 2008); software used to prepare material for publication: *SHELXL97* (Sheldrick, 2008) and *PLATON* (Spek, 2009).




Figure 1

A view of the title molecule with displacement ellipsoids shown at the 50% probability level.





Figure 2

Intermolecular interactions between molecules of the title compound. [Symmetry codes: i: x,y,z ; ii: $3-x,2-y,1-z$; iii: $2.5-x,-1/2+y,1/2-z$; iv: $-1/2+x,1.5-y,1/2+z$]

**Figure 3**

The crystal packing diagram showing the C—H...O and C—H...N intermolecular interactions between molecules within pairs (green colored) and between molecules in different strands (blue colored).

2,5-Dimethoxybenzonitrile

Crystal data

$C_9H_9NO_2$

$M_r = 163.17$

Monoclinic, $P2_1/n$

$a = 3.91001$ (15) Å

$b = 11.3347$ (4) Å

$c = 17.8432$ (6) Å

$\beta = 93.400$ (3)°

$V = 789.40$ (5) Å³

$Z = 4$

$F(000) = 344$

$D_x = 1.373$ Mg m⁻³

Melting point = 360–361 K

Mo $K\alpha$ radiation, $\lambda = 0.7107$ Å

Cell parameters from 1290 reflections

$\theta = 3.6$ – 32.0 °

$\mu = 0.10$ mm⁻¹

$T = 100$ K

Block, colourless

$0.60 \times 0.25 \times 0.23$ mm

Data collection

Agilent SuperNova (Dual, Cu at zero, Atlas) diffractometer

Radiation source: SuperNova (Mo) X-ray

Source

Mirror monochromator

Detector resolution: 10.4127 pixels mm⁻¹

ω scans

Absorption correction: multi-scan

(*CrysAlis PRO*; Agilent, 2013)

$T_{\min} = 0.899$, $T_{\max} = 1.000$

3225 measured reflections

1785 independent reflections

1374 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.026$

$\theta_{\max} = 27.5$ °, $\theta_{\min} = 3.6$ °

$h = -5 \rightarrow 4$

$k = -9 \rightarrow 14$

$l = -21 \rightarrow 23$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.044$

$wR(F^2) = 0.116$

$S = 1.06$

1785 reflections

111 parameters

0 restraints

Primary atom site location: structure-invariant direct methods

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + (0.048P)^2 + 0.110P]$

where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} < 0.001$

$$\Delta\rho_{\max} = 0.21 \text{ e } \text{\AA}^{-3}$$

$$\Delta\rho_{\min} = -0.24 \text{ e } \text{\AA}^{-3}$$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	1.1608 (4)	0.84580 (13)	0.32618 (8)	0.0170 (3)
C2	1.1137 (4)	0.96337 (13)	0.34798 (7)	0.0170 (3)
C3	0.9434 (4)	1.03957 (13)	0.29791 (8)	0.0178 (3)
C4	0.8238 (4)	1.00071 (13)	0.22693 (8)	0.0180 (3)
C5	0.8723 (4)	0.88435 (13)	0.20555 (8)	0.0166 (3)
C6	1.0425 (4)	0.80699 (13)	0.25538 (7)	0.0178 (3)
C7	1.3310 (4)	0.76401 (14)	0.37753 (8)	0.0193 (3)
C8	1.1889 (4)	1.11088 (14)	0.44323 (8)	0.0214 (4)
C9	0.5943 (4)	0.91476 (14)	0.08362 (8)	0.0227 (4)
H3	0.9077	1.1192	0.3120	0.021*
H4	0.7086	1.0541	0.1930	0.022*
H6	1.0782	0.7274	0.2410	0.021*
H8A	1.2963	1.1668	0.4099	0.032*
H8B	1.2895	1.1202	0.4945	0.032*
H8C	0.9421	1.1264	0.4425	0.032*
H9A	0.5317	0.8708	0.0375	0.034*
H9B	0.7498	0.9793	0.0723	0.034*
H9C	0.3872	0.9472	0.1041	0.034*
N1	1.4646 (4)	0.69750 (12)	0.41812 (7)	0.0264 (3)
O1	1.2456 (3)	0.99244 (9)	0.41788 (5)	0.0202 (3)
O2	0.7618 (3)	0.83694 (9)	0.13775 (5)	0.0219 (3)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0161 (7)	0.0162 (8)	0.0189 (7)	-0.0002 (6)	0.0038 (5)	0.0027 (6)
C2	0.0161 (8)	0.0185 (8)	0.0165 (7)	-0.0017 (6)	0.0025 (5)	0.0012 (6)
C3	0.0184 (8)	0.0147 (7)	0.0204 (7)	0.0000 (6)	0.0026 (6)	-0.0003 (6)
C4	0.0178 (8)	0.0170 (7)	0.0194 (7)	0.0008 (6)	0.0024 (6)	0.0039 (6)
C5	0.0157 (7)	0.0176 (8)	0.0165 (7)	-0.0028 (6)	0.0020 (5)	0.0008 (6)
C6	0.0169 (8)	0.0164 (7)	0.0206 (7)	-0.0007 (6)	0.0041 (6)	-0.0001 (6)
C7	0.0210 (8)	0.0174 (8)	0.0196 (7)	-0.0005 (7)	0.0030 (6)	-0.0018 (6)
C8	0.0260 (9)	0.0184 (8)	0.0196 (7)	0.0012 (7)	-0.0007 (6)	-0.0027 (6)
C9	0.0256 (9)	0.0229 (8)	0.0189 (7)	-0.0011 (7)	-0.0029 (6)	0.0031 (6)
N1	0.0334 (9)	0.0212 (7)	0.0243 (7)	0.0046 (6)	-0.0004 (6)	0.0003 (6)
O1	0.0258 (6)	0.0172 (6)	0.0171 (5)	0.0027 (5)	-0.0029 (4)	-0.0009 (4)
O2	0.0281 (6)	0.0196 (6)	0.0174 (5)	0.0000 (5)	-0.0033 (4)	0.0002 (4)

Geometric parameters (Å, °)

C2—C1	1.404 (2)	C8—H8C	0.9800
C2—C3	1.384 (2)	C8—H8B	0.9800
C3—C4	1.395 (2)	C8—H8A	0.9800
C3—H3	0.9500	C9—H9C	0.9800
C4—H4	0.9500	C9—H9B	0.9800
C5—C4	1.389 (2)	C9—H9A	0.9800
C5—C6	1.390 (2)	O1—C8	1.4381 (18)
C6—C1	1.391 (2)	O1—C2	1.3616 (17)
C6—H6	0.9500	O2—C9	1.4370 (18)
C7—N1	1.1492 (19)	O2—C5	1.3700 (17)
C7—C1	1.439 (2)		
C1—C6—H6	119.9	H8A—C8—H8B	109.5
C2—C1—C7	119.92 (13)	H8B—C8—H8C	109.5
C2—C3—C4	120.75 (14)	H9A—C9—H9C	109.5
C2—C3—H3	119.6	H9A—C9—H9B	109.5
C2—O1—C8	117.18 (11)	H9B—C9—H9C	109.5
C3—C4—H4	119.8	N1—C7—C1	179.12 (16)
C3—C2—C1	118.65 (13)	O1—C8—H8C	109.5
C4—C3—H3	119.6	O1—C8—H8B	109.5
C4—C5—C6	119.42 (13)	O1—C8—H8A	109.5
C5—C4—H4	119.8	O1—C2—C1	115.77 (13)
C5—C4—C3	120.35 (14)	O1—C2—C3	125.57 (14)
C5—C6—C1	120.16 (14)	O2—C9—H9C	109.5
C5—C6—H6	119.9	O2—C9—H9B	109.5
C5—O2—C9	117.39 (11)	O2—C9—H9A	109.5
C6—C1—C7	119.42 (13)	O2—C5—C4	125.03 (13)
C6—C1—C2	120.67 (14)	O2—C5—C6	115.55 (13)
H8A—C8—H8C	109.5		

Hydrogen-bond geometry (Å, °)

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
C8—H8B...O1 ⁱ	0.98	2.65	3.428 (2)	136
C8—H8B...N1 ⁱ	0.98	2.73	3.504 (2)	136
C9—H9B...N1 ⁱⁱ	0.98	2.71	3.640 (2)	158

Symmetry codes: (i) $-x+3, -y+2, -z+1$; (ii) $-x+5/2, y+1/2, -z+1/2$.