



RESEARCH ARTICLE

Open Access

Protective effect of magnesium on renal function in STZ-induced diabetic rats

Mohammad Reza Parvizi¹, Mohsen Parviz¹, Seyed Mohammad Tavangar², Nepton Soltani³, Mehri Kadkhodaei¹, Behjat Seifi¹, Yaser Azizi¹ and Mansoor Keshavarz^{1*}

Abstract

Background: Diabetic nephropathy is a serious complication of T1D (type one diabetes mellitus). Persistent hyperglycemia and subsequent hypomagnesemia is believed to develop kidney damage by activation of oxidative stress. We conducted this study to investigate the renoprotective effect of magnesium sulfate ($MgSO_4$) on renal histopathology and oxidative stress in diabetic rats.

Methods: The study included 70 male rats. The animals were divided into seven groups: control (CRL), control receiving $MgSO_4$ (CRL + Mg1 & CRL + Mg8), diabetic (DM1 & DM8) and diabetic receiving $MgSO_4$ (DM + Mg1 & DM + Mg8). Rats were given 20 mg/kg (i.p) Streptozocin (STZ) for 5 consecutive days in (MLD) multiple low doses to induce T1D. At day 10 treatment groups were received $MgSO_4$ (10 g/l) in drinking water, for 1 or 8 weeks. The blood glucose, BUN and creatinine levels were measured. Renal tissue levels of malondialdehyde (MDA) were measured by thiobarbituric acid (TBA) method to evaluate the oxidative stress. Renal histopathology was done using H & E staining method.

Results: Treatment with $MgSO_4$ significantly decreased the blood glucose in DM + Mg1 and DM + Mg8 groups as compared with DM1 and DM8. Magnesium treatment also decreased serum BUN and tissue level of MDA significantly in both short and long term treatment. The body weight loss and kidney weight to body weight ratio was improved by $MgSO_4$. Histological results showed there were no differences between DM and DM + Mg groups.

Conclusion: Our findings showed that diabetic nephropathy is associated with high blood glucose level and oxidative stress (significant increase in MDA level). The renal dysfunction and oxidative stress can be improved by magnesium sulfate administration. It is suggested that protection against development of diabetic nephropathy by $MgSO_4$ treatment involves changes in the blood glucose and oxidative stress.

Keywords: Diabetes mellitus type 1, Streptozocin, Diabetic nephropathy, Oxidative stress

Introduction

Diabetic nephropathy is a major long-term complication of Type 1 diabetes mellitus (T1D) [1-3]. It develops in more than of 40% of patients in spite of glucose control [4]. Oxidative stress has been considered to be a pathogenic factor for diabetic nephropathy [5]. Hyperglycemia is believed to activate oxidative stress resulting in proteinuria, basement membrane thickening, expansion of the mesangium, decline in filtration and nephromegaly followed by glomerular sclerosis [2,6]. It is suggested

that increased oxidative stress through reduction of plasma antioxidants and increased lipid peroxidation could intensify mesangial cells susceptibility to free radical injury [6,7]. Malondialdehyde (MDA) is a substance produced during polyunsaturated fatty acid peroxidation which has been detected in the serum of patients with T1D and correlate with the progression of disease [8-10]. MDA is known to have toxic influence on cell membrane structure. It can modulate signal transduction as well as modify proteins and DNA [11].

Magnesium (Mg) has received important attention for its potential in improving diabetes and its complications [12]. Mg deficiency has previously been proposed as a prominent factor leading to the pathogenesis of the diabetes

* Correspondence: mkesavarz@tums.ac.ir

¹Department of Physiology, School of Medicine, Tehran University of Medical Sciences, Tehran, Iran

Full list of author information is available at the end of the article

complications [13]. Hypomagnesemia has been implicated in the development of diabetic complications due to enhanced renal Mg excretion [6,13,14]. The study of Prabodh suggested that hypomagnesemia may be related to development of diabetic nephropathy [15]. Hypomagnesemia was found to be related to weak glycemic control and increased incidence of nephropathy and retinopathy [16]. The study of Srivastava indicated that $MgSO_4$ can improve oxidative stress by decreasing MDA generation in sodium-metavanadate induced lipid peroxidation [17]. However, the possibility that $MgSO_4$ could exert beneficial effects in improving diabetic renal damage has not been previously investigated. It seems that Mg as magnesium sulfate ($MgSO_4$) will be able to effectively limit oxidative stress. Thus this study was conducted to evaluate the renoprotective and antioxidant effects of oral magnesium administration on STZ- Induced diabetic rats in short and long term periods.

Materials and methods

Male Wistar rats, weighing 200 ± 20 g were used. Animals were kept at a constant temperature of $22 \pm 2^\circ C$ with fixed 12:12-h light-dark cycle. Animals were divided into seven groups ($n = 10$): Control (CRL), control treated by magnesium sulfate for one week (CRL + Mg1), control treated by magnesium sulfate for 8 weeks (CRL + Mg8), untreated diabetic for one week (DM1), untreated diabetic for 8 weeks (DM8), diabetic treated by magnesium sulfate for one week (DM + Mg1) and diabetic treated by magnesium sulfate for 8 weeks (DM + Mg8). All the experiments were approved by the Ethics committee of Tehran University of Medical Sciences (Tehran, Iran).

Diabetes induction

Diabetes were induced by intraperitoneal (i.p) injection of multiple low doses (20 mg/kg) of STZ (Sigma-Aldrich Inc., USA) for 5 consecutive days [18]. In 10th days, the blood glucose levels were determined using a glucometer (ACCU-CHEK Active, Germany) and animals with blood glucose levels above 250 mg/dl were considered to be diabetic [19]. Control rats were injected with the same volume of vehicle. Mg-treated rats were received 10 g/l of $MgSO_4$ added to the drinking water from 10th days [20]. Untreated groups of STZ- diabetic and control rats were given drinking water over the same period.

Blood sampling and biochemical assay

Blood samples were taken for glucose, creatinine, BUN, calcium and Mg levels measurements using a kit (Zistshimi, Tehran, Iran) and spectrophotometer (UV 3100, Shimadzu). Blood samples were centrifuged at 2000 g for 10 minutes at 4°C, and serums were used for biochemical assay. All analysis was performed in accordance with the

instructions provided by the manufacturers. Serum creatinine concentration was determined using Jaffe method [21] and BUN was determined using UV method by auto-analyzer (BT 3000).

Histological analysis

At the end of experiment, rats were euthanized with high-dose of ketamine. The right kidney of the animals in all groups were identified, resected, dried by tissue papers and weighed by digital scale on Sartorius balance to determine the change in the weight of organs with respect to their body weights. A change in kidney size was estimated by division the weight of the right kidney to total body weight [1]. The kidney was removed from each rat, put into 10% formalin and embedded in paraffin. Each sample was then cut into 5- μm sections with a microtome and deparaffined with xylene. Then sections are subjected to standard hematoxylin/eosin staining. The sections were observed under a light microscope at magnifications of 400x [22-24]. Determination of MDA in lipid peroxidation study was performed by thiobarbituric acid (TBA) method [25]. Briefly, the renal tissue was mixed with 2 volumes of 10% trichloroacetic acid (TCA) for protein precipitating. After centrifugation of the precipitate, supernatant is separated and reacted with TBA in boiling water for 10 min followed by cooling. The concentration of MDA was measured at 532 nm [26,27].

Statistical analysis

Results were expressed as mean \pm SEM. Differences among groups were evaluated by one-way analysis of variance (ANOVA) with Tukey post-hoc test. Statistical significance was achieved if the $p < 0.05$.


Results

Effects of magnesium sulfate administration on blood glucose

Before the experiment, there were no significant differences between blood glucose of groups (93.1 ± 5.1 vs 98.1 ± 3.1). At 10th days of study, blood glucose of diabetic animals significantly increased to opposite the non-diabetic ones (398.9 ± 15.3 vs 103.0 ± 6.2) [$F_{(3,36)} = 80.67$], ($p < 0.001$). Treatment with $MgSO_4$ significantly ($p < 0.01$) decreased the blood glucose in DM + Mg1 (230.4 ± 5.8 mg/dl) and DM + Mg8 (153.5 ± 12.6 mg/dl) [$F_{(3,36)} = 292.3$], ($p < 0.001$) in comparison with DM1 and DM8 respectively (Figure 1).

Change in body weight and kidney weight/body weight (KW/BW) ratio

There was significant reduction in body weight of DM1 compared to CRL + Mg1 group (209.4 ± 6.3 g vs. 228.5 ± 4.2 g) [$F_{(2,27)} = 1.696$] as well as DM8 compared with CRL + Mg8 group (194.7 ± 8.8 g vs. 230.8 ± 13.1 g) [$F_{(2,27)} =$



32.53] ($p < 0.0001$). Magnesium treatment prevented body weight loss in DM + Mg1 and DM + Mg8 compared with DM1 and DM8 significantly (219.1 ± 11.6 g vs. 194.7 ± 8.8 g) ($p < 0.01$) (Figure 2). In addition, diabetic rats had an increased kidney weight/body weight ratio, a marker for the body weight loss and renal size [28]. This ratio was significantly improved by treatment with $MgSO_4$ in DM + Mg8 groups (Figure 3).


Effects of magnesium sulfate administration on plasma level of Mg^{2+}

Plasma magnesium level in both DM1 and DM + Mg1 groups didn't show any changes in comparison with

control (2.0 ± 0.13 and 2.26 ± 0.12 mg/dl) [$F_{(3,36)} = 0.9309$]. 8 weeks after diabetes induction by STZ, magnesium level of plasma decreased in DM8 as compared to CRL + Mg8 groups. $MgSO_4$ treatment could increase the plasma level of magnesium significantly in DM + Mg8 as compared to DM8 groups (3.33 ± 0.3 vs 1.74 ± 0.4 mg/dl) [$F_{(3,36)} = 3.812$] $p < 0.05$ (Figure 4).

Effects of magnesium sulfate administration on plasma level of BUN and creatinine

Our results showed that STZ-induced diabetes caused a significant increase in serum BUN level in DM1 (41.37 ± 2.95 mg/dl) and DM8 (48.10 ± 3.79 mg/dl)






Figure 3 Effect of magnesium sulfate treatment on kidney weight/body weight ratio in control rats (CRL), control rats treated with magnesium sulfate (CRL + Mg), Untreated diabetic rats (DM) and diabetic rats treated with magnesium sulfate (DM + Mg) (10 g/l added in water) in 1 and 8 weeks ($n = 10$). Data are presented as mean \pm SE. * $p < 0.05$ vs. CRL, CRL + Mg8, # $p < 0.05$ vs DM8.

compared with CRL, CRL + Mg1 and CRL + Mg8 (24.77 ± 1.05 , 25.83 ± 1.5 and $22.3 \pm .66$ mg/dl) respectively [$F_{(3,36)} = 17.53$] ($p < 0.05$). Following treatment with magnesium sulfate serum BUN levels decreased in DM + Mg1 (26.3 ± 1.49 mg/dl) and DM + Mg8 (22.75 ± 1.68 mg/dl) groups, in comparison with DM1 (41.37 ± 2.95) and DM8 (48.10 ± 3.79) respectively [$F_{(3,36)} = 28.12$] ($p < 0.05$) (Figure 5).

After diabetes induction by STZ, a mild increase in creatinine level showed in 1 and 8 week groups (0.62 ± 0.018 and $0.64 \pm 0.13 \pm 0.03$ mg/dl) [$F_{(3,36)} = 1.971$] respectively. There were no significant differences in serum creatinine level between DM and DM + Mg in both weeks groups

(0.59 ± 0.02 and 0.58 ± 0.03 mg/dl) [$F_{(3,36)} = 2.368$] respectively (Figure 6).

Effects of magnesium sulfate treatment on MDA level in renal tissue

MDA was used as a marker of oxidative stress [29]. The renal content of MDA was higher levels significantly in DM1 and DM8 ($4.55 \pm .21$ and $4.72 \pm .31$ nmol/g protein) respectively in comparison with CRL ($2.17 \pm .05$ nmol/g protein) [$F_{(2,27)} = 34.94$]. Magnesium treatment attenuate renal content of MDA in DM + Mg1 and DM + Mg8 ($2.64 \pm .31$ and $3.12 \pm .51$ nmol/g protein) [$F_{(2,27)} = 12.13$] significantly compared to DM1 and DM8




Figure 4 Serum level of magnesium (Mg^{2+}) in CRL (control), CRL + Mg (control rats treated with $MgSO_4$), DM (diabetic) and DM + Mg (diabetic rats treated with $MgSO_4$) in 1 and 8 weeks after diabetes induction by STZ ($n = 10$). Data are presented as mean \pm SE. # $p < 0.05$ vs DM8.




Figure 5 Serum level of BUN in CRL (control), CRL + Mg (control rats treated with $MgSO_4$), DM (diabetic) and DM + Mg (diabetic rats treated with $MgSO_4$) in one and 8 weeks after diabetes induction by STZ ($n = 10$). Data are presented as mean \pm SE. *** $p < 0.001$ vs. CRL, CRL + Mg, ### $p < 0.001$ vs. DM.

respectively. However, there was no difference in renal content of MDA between DM + Mg1 and DM + Mg8 (Figure 7).

Histological study

Results of the renal tissue staining showed that the glomeruli of the control and diabetic groups were morphologically normal and that the treatment of the diabetic groups in 1 and 8 week did not have a clear effect on the structure of the glomeruli (Figure 8).

Discussion

Diabetes mellitus can cause serious health problems including macrovascular and microvascular complications

such as kidney failure, heart disease, stroke and etc., which affects the function of many organs [4]. One of them is injuries to the kidney tissue that results in renal dysfunction [30]. This study was designed to evaluate the renal complication of T1D in short and long term and probable protective effects of $MgSO_4$. We used the multiple low doses of STZ to induce T1D in rats according to previous reports [18,31]. Eight weeks after STZ-diabetes induction, some indexes of renal damage such as increasing BUN and elevation of renal MDA were appeared. Magnesium treatment could reverse renal function (decreasing BUN), decreasing MDA and improving hyperglycemia and kidney weight respect to body weight in diabetic rats.




Figure 6 Serum level of Creatinine in CRL (control), CRL + Mg (control rats treated with $MgSO_4$), DM (diabetic) and DM + Mg (diabetic rats treated with $MgSO_4$) in one and 8 weeks after diabetes induction by STZ ($n = 10$). Data are presented as mean \pm SE.




Figure 7 Kidney tissue level of MDA in CRL (control), and DM + Mg (diabetic rats treated with MgSO₄) in 1 and 8 weeks after diabetes induction by STZ (*n* = 10). Data are presented as mean ± SE. *** P < 0.001 vs CRL, ### P < 0.001 vs DM1, # P < 0.05 vs DM8.

Previous studies have been proved that treatment with MgSO₄ have beneficial effects on diabetes. It could improve some of diabetic complications such as hypertension [19,32], thermal hyperalgesia and structure of pancreatic islets [20,33]. Also MgSO₄ improves function of endothelium and restores hemodynamic and tubular function in postischemic rats [34]. With respect of Soltani study [20], we used magnesium supplementation (10 g/l) via drinking water to diabetic rats from 10th days after diabetes induction for short time (one week) or long time (eight weeks).

Our results showed that consumption of MgSO₄ in diabetic rats reversed the high blood glucose level. However, MgSO₄ couldn't return it to that control levels in

the treated diabetic rats. Magnesium plays an important role in the carbohydrate metabolism regulation [35]. We have previously showed that treatment by MgSO₄ could reduce the blood glucose in diabetic rats [20]. The effect of MgSO₄ on blood sugar levels may be related to the fact that it is a crucial cofactor for glucose transport and various enzymes involved in carbohydrate metabolism [36]. It has also been reported that oral MgSO₄ consumption in male obese rats improved glucose tolerance and delaying the progression of diabetes [37]. It is supposed that blood glucose levels can be decreased by Mg via increasing GLUT₄ mRNA expression in diabetic rats independent to insulin secretion [38]. Huang et al. showed




Figure 8 The images of kidney tissues (magnification x 400) in untreated diabetic rats (A & C) and diabetic treated by magnesium (B & D) in 1 and 8 weeks after diabetes induction by STZ. A: DM1, B: DM + Mg1, C: DM8, D: DM + Mg8.

that MgSO₄ increases the expression of GLUT₃ in the cortex and hippocampus of gerbil brains [39].

We showed that after one week MgSO₄ treatment (in drinking water), the plasma Mg levels in both DM1 and DM + Mg groups didn't show any significant changes in comparison with control group. In previous work, Soltani et al., showed that a significant reduction in Mg level in acute diabetic groups [32]. These discrepancy may be due to the dose of STZ and the model of diabetes induction. Because in this study we used MLD injection of STZ, but in the Soltani et al., study they used single high dose of STZ for diabetes induction [20]. In other hand, after few days following diabetes induction, rats showed polydipsia and polyuria along with diarrhea (in DM + Mg groups). Actually, in long term diabetes, ingestion of water and meal, reached near to control rats. It may be due to adaptation.

High Magnesium (as MgSO₄) ingestion has been reported to induce diarrhea. In short term diabetes, diarrhea and polyuria along with polydipsia can cause non-significant changes in magnesium level [39,40].

Our results showed that Mg administration to DM + Mg8, after eight weeks, could decrease the level of renal function markers such as BUN and creatinine in comparison to DM8. An elevation of BUN in DM8 groups may reflect decreasing in glomerular filtration rate (GFR) or it is probably due to increased degradation of proteins [41]. Avram et al., showed that there is a negative correlation between renal size and serum creatinine level [42]. So in our study, mildly elevation of creatinine may be associated with renomegaly as shown in this study.

Increasing BUN and plasma creatinine, has been reported as waste products of metabolism following kidney injury and they have known biomarkers of kidney degeneration [43]. In this study, the elevation of plasma BUN with hyperglycemia can be proposed as indicator for renal dysfunction [44]. Our data showed no significant changes in the creatinine level in diabetic animals versus CRL group. Increase in creatinine level occurs at the end stage of renal disease, and this is accompanied by histological alterations [45]. Kim et al., suggested that plasma BUN and creatinine is often measures together, but former is more sensitive marker for kidney injury [46].

Histological study also showed that there were no profound changes among the experimental groups. Eight weeks after induction of diabetes in DM8 groups' tissue didn't change histologically. It seems that it requires long time to cause an end stage renal disease with changes in creatinine level and histological changes. In previous study, it has been shown that it takes about 15 weeks to see profound histological changes [47]. So it may be due to insufficient time that we didn't see these changes.

Diabetes induction resulted in an increase in kidney MDA, which is an indicator of oxidative stress. Israa et al.,

revealed that enhancement of the oxidative stress as indicated by high MDA level is due to increase in blood glucose [48]. Metabolically, diabetes is characterized by diminished glucose utilization and increased lipid peroxidation, resulting in accumulation of MDA in renal tissue. The elevated level of MDA may be due to the poor antioxidant capacity of mesangial cells as a result of low glutathione (GSH) levels [49]. Evidences showed that in patients with T2D, the higher levels of lipid peroxidation and hypomagnesaemia are often reported [50]. Free radicals in DM cause peroxidative breakdown of phospholipids that lead to accumulation of MDA [51]. Agnieszka et al., have been shown that magnesium intake for 18 weeks could prevent lipid peroxidation stimulated by vanadium in hepatic tissue [52]. Thus, from our results it is seemed that magnesium may show an antioxidant activity through preventing lipid peroxidation in renal tissue that is indicated by a decrease in kidney MDA level. Ribeiro et al., demonstrated a negative correlation between magnesium and glucose levels as well as between magnesium and oxidative stress [3]. Magnesium deficiency enhances oxidative stress by increased production of free radicals and decreased of antioxidant defenses [53]. In this regard, histological alterations can be due to mildly proinflammatory invasion to renal tissues.

Hyperglycemia induces diabetic nephropathy via several biochemical pathways. Several mechanisms have revealed to depict the adverse effect of hyperglycemia, including protein kinase C (PKC) [48], mitogen-activated protein kinase (MAPK), polyol pathway, advanced glycation end products (AGE), aldose reductase activation [54] and oxidative stress [55]. In the other hand, it has been reported that hypomagnesemia is a novel predictor of end stage renal disease (ESRD) in patients with type 2 diabetic nephropathy [14]. So it is suggested that oxidative stress causing damage to mesangial cells and magnesium could reduce oxidative damages to mesangial cells as indicated by decreasing MDA production.

In addition, we assessed the effects of magnesium administration on the body weight and kidney weight to body weight ratio in diabetic rats. We found that induction of diabetes caused body weight loss. Decrease in body weight following STZ-induced diabetes notably in eight weeks groups, may be due to degradation of proteins as indicated by raising serum BUN and creatinine levels. Area under curve showed that in DM + Mg groups, Mg treatment caused body weight gain, especially in DM + Mg8 groups.

Also our results showed that KW/BW ratio was significantly elevated in untreated diabetic rats and decreased in DM + Mg groups. KW/BW ratio is a marker of renomegaly [56], its increment is an indicator of glomerular expansion due to diabetes. Several investigators have reported that KW/BW increases in DM animals [56]. Renet et al., showed that marked increase in the KW/BW, tubulo-interstitial fibrosis and fibronectin in STZ-induced

diabetic rats from 8 to 16 weeks [57]. Arozal et al., oxidative stress is associated with development of hypertrophy in diabetes [58]. Sharma et al., revealed that renal hypertrophy in T1D was related to overexpression of TGF- β 1 in the glomerular mesangial cells [59]. Our study further strengthens the notion that attenuation of kidney weight by magnesium, suggesting that Mg has the ability to protect the kidneys from oxidative injury. It may reverse kidney hypertrophy in STZ-diabetic rats.

Limitations of this study was to evaluate more precisely the oxidant and antioxidant markers for achieving better results, because of limitation in financial support. We haven't access to electron microscope for more precise evaluation of histological changes.

Conclusion

The current study determined that the increase in blood glucose level is related to the development of oxidative stress as indicated by high MDA level. Control of hyperglycemia by Mg supplementation can decreases oxidative damages as indicated by MDA and improves renal dysfunction via lowering of BUN and creatinine. However, further studies are required to clear the precise mechanism(s) involving for the protective effect of MgSO₄ against the diabetic nephropathy in experimental conditions.

Competing interest

The authors declare that they have no competing interests.

Authors' contribution

MRP performed experiments, data analysis and article writing. MP participated as advisor. MT participated in pathological data interpretation. NS participated as advisor. MK participated in pathological interpretation. BS participated data interpretation. YA participated in doing experiments and writing the article. MK participated as supervisor, statistical analysis and article writing. All authors read and approved the final manuscript.

Acknowledgment

This research was supported by Tehran University of Medical Sciences, Tehran, Iran.

Author details

¹Department of Physiology, School of Medicine, Tehran University of Medical Sciences, Tehran, Iran. ²Department of Pathology, School of Medicine, Tehran University of Medical Sciences, Tehran, Iran. ³Department of Physiology, School of Medicine, Hormozgan University of Medical Sciences, Bandar Abbas, Iran.

Received: 15 January 2014 Accepted: 23 July 2014

Published: 16 August 2014

References

- Lu HJ, Tzeng TF, Hsu JC, Kuo SH, Chang CH, Huang SY, Chang FY, Wu MC, Liu IM: An aqueous-ethanol extract of liriope spicata var. *prolifera* ameliorates diabetic nephropathy through suppression of renal inflammation. *Evid Based Complement Alternat Med* 2013, **2013**:201643. Epub 2013/09/13.
- Schena FP, Gesualdo L: Pathogenetic mechanisms of diabetic nephropathy. *J Am Soc Nephrol* 2005, **16**(Suppl 1):S30–S33. Epub 2005/06/07.
- Ribeiro MC, Avila DS, Barbosa NB, Meinerz DF, Waczuk EP, Hassan W, Rocha JB: Hydrochlorothiazide and high-fat diets reduce plasma magnesium levels and increase hepatic oxidative stress in rats. *Magnes Res* 2013, **26**(1):32–40. Epub 2013/05/10.
- Gross JL, de Azevedo MJ, Silveiro SP, Canani LH, Caramori ML, Zelmanovitz T: Diabetic nephropathy: diagnosis, prevention, and treatment. *Diabetes Care* 2005, **28**(1):164–176. Epub 2004/12/24.
- Kumawat M, Sharma TK, Singh I, Singh N, Ghalaut VS, Vardey SK, Shankar V: Antioxidant enzymes and lipid peroxidation in type 2 diabetes mellitus patients with and without nephropathy. *N Am J Med Sci* 2013, **5**(3):213–219. Epub 2013/04/30.
- Hans CP, Chaudhary DP, Bansal DD: Magnesium deficiency increases oxidative stress in rats. *Indian J Exp Biol* 2002, **40**(11):1275–1279. Epub 2003/09/19.
- Walti MK, Zimmermann MB, Spinias GA, Jacob S, Hurrell RF: Dietary magnesium intake in type 2 diabetes. *Eur J Clin Nutr* 2002, **56**(5):409–414. Epub 2002/05/10.
- Del Rio D, Stewart AJ, Pellegrini N: A review of recent studies on malondialdehyde as toxic molecule and biological marker of oxidative stress. *Nutr Metab Cardiovasc Dis* 2005, **15**(4):316–328. Epub 2005/08/02.
- Uchida K: Lipofuscin-like fluorophores originated from malondialdehyde. *Free Radic Res* 2006, **40**(12):1335–1338. Epub 2006/11/09.
- Kamper M, Tsipoukidi O, Chatzigeorgiou A, Lymeri M, Kamper EF: The antioxidant effect of angiotensin II receptor blocker, losartan, in streptozotocin-induced diabetic rats. *Transl Res* 2010, **156**(1):26–36. Epub 2010/07/14.
- Srivastava S, Chandrasekar B, Bhatnagar A, Prabhu SD: Lipid peroxidation-derived aldehydes and oxidative stress in the failing heart: role of aldose reductase. *Am J Physiol Heart Circ Physiol* 2002, **283**(6):H2612–H2619. Epub 2002/10/22.
- Li B, Tan Y, Sun W, Fu Y, Miao L, Cai L: The role of zinc in the prevention of diabetic cardiomyopathy and nephropathy. *Toxicol Mech Methods* 2013, **23**(1):27–33. Epub 2012/10/09.
- Kisters K, Gremmler B, Kozianka J, Hausberg M: Magnesium deficiency and diabetes mellitus. *Clin Nephrol* 2006, **65**(1):77–78. Epub 2006/01/25.
- Sakaguchi Y, Shoji T, Hayashi T, Suzuki A, Shimizu M, Mitsumoto K, Kawabata H, Niihata K, Okada N, Isaka Y, Rakugi H, Tsubakihara Y: Hypomagnesemia in type 2 diabetic nephropathy: a novel predictor of end-stage renal disease. *Diabetes Care* 2012, **35**(7):1591–1597. Epub 2012/04/14.
- Prabodh S, Prakash DS, Sudhakar G, Chowdary NV, Desai V, Shekhar R: Status of copper and magnesium levels in diabetic nephropathy cases: a case-control study from South India. *Biol Trace Elem Res* 2011, **142**(1):29–35. Epub 2010/06/17.
- Dasgupta A, Sarma D, Saikia UK: Hypomagnesemia in type 2 diabetes mellitus. *Indian J Endocrinol Metab* 2012, **16**(6):1000–1003. Epub 2012/12/12.
- Scibior A, Golebiowska D, Niedzwiecka I: Magnesium can protect against vanadium-induced lipid peroxidation in the hepatic tissue. *Oxid Med Cell Longev* 2013, **2013**:802734. Epub 2013/06/15.
- Maksimovic-Ivanic D, Trajkovic V, Miljkovic DJ, Mostarica Stojkovic M, Stosic-Grujicic S: Down-regulation of multiple low dose streptozotocin-induced diabetes by mycophenolate mofetil. *Clin Exp Immunol* 2002, **129**(2):214–223. Epub 2002/08/08.
- Soltani N, Keshavarz M, Sohanaki H, Zahedi Asl S, Dehpour AR: Relaxatory effect of magnesium on mesenteric vascular beds differs from normal and streptozotocin induced diabetic rats. *Eur J Pharmacol* 2005, **508**(1–3):177–181. Epub 2005/02/01.
- Soltani N, Keshavarz M, Minaii B, Mirershadi F, Zahedi Asl S, Dehpour AR: Effects of administration of oral magnesium on plasma glucose and pathological changes in the aorta and pancreas of diabetic rats. *Clin Exp Pharmacol Physiol* 2005, **32**(8):604–610. Epub 2005/08/27.
- Kojima N, Slaughter TN, Paige A, Kato S, Roman RJ, Williams JM: Comparison of the development diabetic induced renal disease in strains of Goto-Kakizaki rats. *J Diabetes Metab* 2013, Suppl 9(5). Epub 2013/12/10.
- Tesch GH, Allen TJ: Rodent models of streptozotocin-induced diabetic nephropathy. *Nephrology (Carlton)* 2007, **12**(3):261–266. Epub 2007/05/15.
- Cheng MF, Chen LJ, Cheng JT: Decrease of Klotho in the kidney of streptozotocin-induced diabetic rats. *J Biomed Biotechnol* 2010, **2010**:513853. Epub 2010/07/14.
- Baig MA, Gawali VB, Patil RR, Naik SR: Protective effect of herbomineral formulation (Dolabi) on early diabetic nephropathy in streptozotocin-induced diabetic rats. *J Nat Med* 2012, **66**(3):500–509. Epub 2011/11/26.
- Esterbauer H, Zollner H: Methods for determination of aldehydic lipid peroxidation products. *Free Radic Biol Med* 1989, **7**(2):197–203. Epub 1989/01/01.

26. Esterbauer H, Cheeseman KH: Determination of aldehydic lipid peroxidation products: malonaldehyde and 4-hydroxynonenal. *Methods Enzymol* 1990, **186**:407–421. Epub 1990/01/01.
27. Esterbauer H, Dieber-Rotheneder M, Waeg G, Puhl H, Tatzber F: Endogenous antioxidants and lipoprotein oxidation. *Biochem Soc Trans* 1990, **18**(6):1059–1061. Epub 1990/12/01.
28. Zafar M, Naeem-ul-Hassan Naqvi S: Effects of STZ-induced diabetes on the relative weights of kidney, liver and pancreas in albino rats: a comparative study. *Int J Morphol* 2010, **28**:135–142.
29. Pan HZ, Zhang L, Guo MY, Sui H, Li H, Wu WH, Qu NQ, Liang MH, Chang D: The oxidative stress status in diabetes mellitus and diabetic nephropathy. *Acta Diabetol* 2010, **47**(Suppl 1):71–76. Epub 2009/05/29.
30. Eid S, Maalouf R, Jaffa AA, Nassif J, Hamdy A, Rashid A, Ziyadeh FN, Eid AA: 20-HETE and EETs in diabetic nephropathy: a novel mechanistic pathway. *PLoS One* 2013, **8**(8):e70029. Epub 2013/08/13.
31. Cnop M, Welsh N, Jonas JC, Jorns A, Lenzen S, Eizirik DL: Mechanisms of pancreatic beta-cell death in type 1 and type 2 diabetes: many differences, few similarities. *Diabetes* 2005, **54**(Suppl 2):S97–S107. Epub 2005/11/25.
32. Soltani N, Keshavarz M, Sohanaki H, Dehpour AR, Zahedi Asl S: Oral magnesium administration prevents vascular complications in STZ-diabetic rats. *Life Sci* 2005, **76**(13):1455–1464. Epub 2005/02/01.
33. Kim DJ, Xun P, Liu K, Loria C, Yokota K, Jacobs DR Jr, He K: Magnesium intake in relation to systemic inflammation, insulin resistance, and the incidence of diabetes. *Diabetes Care* 2010, **33**(12):2604–2610. Epub 2010/09/03.
34. de Araujo M, Andrade L, Coimbra TM, Rodrigues AC Jr, Seguro AC: Magnesium supplementation combined with N-acetylcysteine protects against postischemic acute renal failure. *J Am Soc Nephrol* 2005, **16**(11):3339–3349. Epub 2005/09/24.
35. Anetor JL, Senjobi A, Ajose OA, Agbedana EO: Decreased serum magnesium and zinc levels: atherogenic implications in type-2 diabetes mellitus in Nigerians. *Nutr Health* 2002, **16**(4):291–300. Epub 2003/03/06.
36. Laires MJ, Monteiro CP, Bicho M: Role of cellular magnesium in health and human disease. *Front Biosci* 2004, **9**:262–276. Epub 2004/02/10.
37. Balon TW, Gu JL, Tokuyama Y, Jasman AP, Nadler JL: Magnesium supplementation reduces development of diabetes in a rat model of spontaneous NIDDM. *Am J Physiol* 1995, **269**(Pt 1):E745–E752. Epub 1995/10/01.
38. Solaimani H, Soltani N, Malekzadeh K, Sohrabipour S, Zhang N, Nasri S, Wang Q: Modulation of GLUT4 expression by oral administration of Mg (2+) to control sugar levels in STZ-induced diabetic rats. *Can J Physiol Pharmacol* 2014, **92**(6):438–444. Epub 2014/05/14.
39. Huang CY, Liou YF, Chung SY, Pai PY, Kan CB, Kuo CH, Tsai CH, Tsai FJ, Chen JL, Lin JY: Increased expression of glucose transporter 3 in gerbil brains following magnesium sulfate treatment and focal cerebral ischemic injury. *Cell Biochem Funct* 2010, **28**(4):313–320. Epub 2010/06/03.
40. Laurant P, Touyz RM, Schiffrin EL: Effect of magnesium on vascular tone and reactivity in pressurized mesenteric resistance arteries from spontaneously hypertensive rats. *Can J Physiol Pharmacol* 1997, **75**(4):293–300. Epub 1997/04/01.
41. Dabla PK: Renal function in diabetic nephropathy. *World J Diabetes* 2010, **1**(2):48–56. Epub 2011/05/04.
42. Avram MM, Hurtado H: Renal size and function in diabetic nephropathy. *Nephron* 1989, **52**(3):259–261. Epub 1989/01/01.
43. Kadkhodaee M, Mikaeili S, Zahmatkesh M, Golab F, Seifi B, Arab HA, Shams S, Mahdavi-Mazdeh M: Alteration of renal functional, oxidative stress and inflammatory indices following hepatic ischemia-reperfusion. *Gen Physiol Biophys* 2012, **31**(2):195–202. Epub 2012/07/12.
44. Trujillo J, Chirino YI, Molina-Jijon E, Anderica-Romero AC, Tapia E, Pedraza-Chaverri J: Renoprotective effect of the antioxidant curcumin: Recent findings. *Redox Biol* 2013, **1**(1):448–456. Epub 2013/11/06.
45. Sargin AK, Can B, Turan B: Comparative investigation of kidney mesangial cells from increased oxidative stress-induced diabetic rats by using different microscopy techniques. *Mol Cell Biochem* 2014, **390**(1-2):41–49. Epub 2013 Dec 29.
46. Kim MJ, Lim Y: Protective effect of short-term genistein supplementation on the early stage in diabetes-induced renal damage. *Mediators Inflamm* 2013, **2013**:510212. Epub 2013 Apr 29.
47. Yokozawa T, Yamabe N, Cho EJ, Nakagawa T, Oowada S: A study on the effects to diabetic nephropathy of Hachimi-ji-gan in rats. *Nephron Exp Nephrol* 2004, **97**(2):e38–e48. Epub 2004/06/26.
48. Israa FJA, Huda Arif J: Role of Antioxidant on Nephropathy in Alloxan Induced Diabetes in Rabbits. *Iraqi postgraduate Med J* 2009, **8**(4):398–402.
49. Giannoukakis N, Rudert WA, Trucco M, Robbins PD: Protection of human islets from the effects of interleukin-1beta by adenoviral gene transfer of an Ikappa B repressor. *J Biol Chem* 2000, **275**(47):36509–36513. Epub 2000/09/01.
50. Niranjan G, Mohanavalli V, Srinivasan AR, Ramesh R: Serum lipid peroxides and magnesium levels following three months of treatment with pioglitazone in patients with Type 2 Diabetes mellitus. *Diabetes Metab Syndr* 2013, **7**(1):35–37. Epub 2013/03/23.
51. Bhutia Y, Ghosh A, Sherpa ML, Pal R, Mohanta PK: Serum malondialdehyde level: Surrogate stress marker in the Sikkimese diabetics. *J Nat Sci Biol Med* 2011, **2**(1):107–112. Epub 2011/01/01.
52. Scibior A, Gołębierska D, Niedźwiecka I: Magnesium can protect against vanadium-induced lipid peroxidation in the hepatic tissue. *Oxid Med Cell Longev* 2013, **2013**:11.
53. Agarwal R, Iezhitsa IN, Agarwal P, Spasov AA: Mechanisms of cataractogenesis in the presence of magnesium deficiency. *Magnes Res* 2013, **26**(1):2–8. Epub 2013/05/28.
54. Dunlop M: Aldose reductase and the role of the polyol pathway in diabetic nephropathy. *Kidney Int Suppl* 2000, **77**:S3–S12. Epub 2000/09/21.
55. Suzuki D, Miyata T, Saotome N, Horie K, Inagi R, Yasuda Y, Uchida K, Izuhara Y, Yagame M, Sakai H, Kurokawa K: Immunohistochemical evidence for an increased oxidative stress and carbonyl modification of proteins in diabetic glomerular lesions. *J Am Soc Nephrol* 1999, **10**(4):822–832. Epub 1999/04/15.
56. Malatiali S, Francis I, Barac-Nieto M: Phlorizin prevents glomerular hyperfiltration but not hypertrophy in diabetic rats. *Exp Diabetes Res* 2008, **2008**:305403. Epub 2008/09/05.
57. Ren XJ, Guan GJ, Liu G, Zhang T, Liu GH: Effect of activin A on tubulointerstitial fibrosis in diabetic nephropathy. *Nephrology (Carlton)* 2009, **14**(3):311–320. Epub 2009/03/21.
58. Arozal W, Watanabe K, Veeraveedu PT, Ma M, Thandavarayan RA, Suzuki K, Tachikawa H, Kodama M, Aizawa Y: Effects of angiotensin receptor blocker on oxidative stress and cardio-renal function in streptozotocin-induced diabetic rats. *Biol Pharm Bull* 2009, **32**(8):1411–1416. Epub 2009/08/05.
59. Sharma K, Ziyadeh FN: Hyperglycemia and diabetic kidney disease. The case for transforming growth factor-beta as a key mediator. *Diabetes* 1995, **44**(10):1139–1146. Epub 1995/10/01.

doi:10.1186/s40200-014-0084-3

Cite this article as: Parvizi et al.: Protective effect of magnesium on renal function in STZ-induced diabetic rats. *Journal of Diabetes & Metabolic Disorders* 2014 **13**:84.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

