

Associations between the number of natural teeth and renal dysfunction

Hye Min Choi, MD, MS, PhD^a, Kyungdo Han, PhD^b, Yong Gyu Park, PhD^b, Jun-Beom Park, DDS, MSD, PhD^{c,*}

Abstract

The purpose of the present study was to investigate the association between the number of natural teeth and measures of kidney dysfunction, such as urinary albumin/creatinine ratio (ACR) and estimated glomerular filtration (eGFR) rate, using nationally representative data.

The data used were from the Korea National Health and Nutrition Examination Survey with cross-sectional design, which was conducted between 2011 and 2012; the sample analyzed in this study consisted of a total of 10,388 respondents, each of whom was 19 years or older and had no missing outcome variables. The association between the number of natural teeth and kidney function was assessed by multiple logistic regression and model was adjusted for age, sex, waist circumference, smoking, drinking, exercise, education, income, frequency of tooth brushing per day, diabetes, metabolic syndrome, urinary ACR, and eGFR.

The mean age, body mass index, and waist circumference were significantly higher among those with lower kidney function (urinary ACR ≥ 30 mg/g and eGFR < 60 mL/min/1.73m²). Urinary ACR and eGFR were associated with loss of natural teeth. As urinary ACR increased, the number of natural teeth decreased accordingly. Conversely, the number of natural teeth increased with an increase in eGFR.

This study showed that the number of natural teeth is inversely associated with the presence of kidney disease. Severity of tooth loss may be considered an independent risk indicator for kidney disease among Koreans. More epidemiological studies are warranted to investigate the role of tooth loss in kidney disease, to confirm this relationship and to test possible underlying mechanisms.

Abbreviations: ACR = albumin/creatinine ratio, eGFR = estimated glomerular filtration, KNHANES = Korean National Health and Nutrition Examination Survey.

Keywords: albuminuria, dentition, epidemiology, glomerula filtration rate, kidney diseases, oral health, tooth loss

1. Introduction

Observational studies have shown an association between poor oral health and systemic diseases.^[1,2] Periodontitis and pulp necrosis are important sources of systemic microinflammation in chronic kidney disease patients.^[3] It was suggested that periodontal disease is associated with increased risk of cardiovascular disease.^[4] In the previous study, severe periodon-

titis was associated with compromised glycemic control and periodontal treatment was associated with improvements in glycemic control in diabetic patients.^[5]

Chronic kidney diseases were shown to affect the oral health status of patients by inducing xerostomia, delayed eruption of teeth, calcification of root canals, and gingival hyperplasia.^[6] The incidence of periodontal disease (bleeding and calculus) was significantly higher among people with renal disease than among controls.^[7] Recently, it was reported that periodontitis was associated with increased prevalence of inflammatory markers in end-stage renal disease patients.^[8] Recent epidemiological studies have suggested that periodontitis is a major risk factor for renal failure.^[9,10] Within the authors' knowledge, there is no known report to evaluate the association between the number of natural teeth and eGFR with nationally representative data. Thus, this study was performed to investigate the association between the number of natural teeth and both urinary ACR and eGFR using nationally representative data.

2. Methods

2.1. The design of the study

The data from the Korea National Health and Nutrition Examination Survey (KNHANES), which was performed between 2011 and 2012, were used for the study. KNHANES is conducted to evaluate the health and nutritional status for the Korean people and KNHANES is an ongoing surveillance system in the Republic of Korea that assesses the health and nutritional status of Koreans, monitors trends in health risk factors and the

Editor: Pavlos Malindretos.

Funding: This research was supported by Basic Science Research Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Science, Information and Communication Technology & Future Planning (NRF-2014R1A1A1003106).

The authors report no conflicts of interest.

^a Department of Internal Medicine, Seonam University Myongji Hospital, Goyang, Gyeonggi-do, ^b Department of Biostatistics, ^c Department of Periodontics, College of Medicine, The Catholic University of Korea, Seoul, Republic of Korea.

* Correspondence: Jun-Beom Park, Department of Periodontics, Seoul St. Mary's Hospital, College of Medicine, The Catholic University of Korea, 222, Banpo-daero, Seocho-gu, Seoul 06591, Republic of Korea (e-mail: jbasoonis@yahoo.co.kr).

Copyright © 2016 the Author(s). Published by Wolters Kluwer Health, Inc. All rights reserved.

This is an open access article distributed under the Creative Commons Attribution License 4.0 (CCBY), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Medicine (2016) 95:34(e4681)

Received: 16 May 2016 / Received in final form: 26 July 2016 / Accepted: 30 July 2016

<http://dx.doi.org/10.1097/MD.0000000000004681>




Figure 1. Participant flow chart.

prevalence of major chronic diseases, and provides data for the development and evaluation of health policies and programs in Korea. The KNHANES uses a stratified and multistage probability-sampling design with a rolling survey-sampling model. The sample weights were used to calculate all the statistics in this survey. A total of 16,576 patients initially participated in the survey. The analysis in this study was confined to a total of 10,388 respondents who were 19 years or older and who had no missing values for the outcome variables (Fig. 1). This survey was reviewed and approved by the Institutional Review Board of the Korea Centers for Disease Control and Prevention.

2.2. Evaluation of kidney function

The level of kidney function is usually determined by urinary albumin/creatinine ratio (ACR) and by estimated glomerular filtration rate (eGFR).^[11,12] In this study, urinary ACR and eGFR were used for the categorization and evaluation of the presence of kidney diseases. When urinary ACR was 30 mg/g or greater, the patient was considered to have albuminuria.^[11]

The level of kidney function was also determined by eGFR using an abbreviated equation: $eGFR \text{ (mL/min/1.73m}^2\text{)} = 186.3 \times (\text{serum creatinine}^{-1.154}) \times (\text{age}^{-0.203})$; this result was then multiplied by the constant 0.742 if the patient was female.^[12] The equation was developed from the Modification of Diet in Renal Disease (MDRD) formula. Participants with $eGFR < 60 \text{ mL/min/1.73m}^2$ were considered to have kidney disease. For this study, chronic kidney disease was ascertained if a patient's eGFR was $< 60 \text{ mL/min/1.73m}^2$.^[13,14]

2.3. Anthropometric measurements

Trained staff members performed anthropometric measurements. Height and weight were measured to the nearest 0.1 cm and 0.1 kg, respectively. The participants wore light indoor clothing without shoes during the anthropometric measurements. Waist circumference was measured at the narrowest point between the iliac crest and the lower border of the rib cage.^[15,16] Body mass was divided by height squared to calculate body mass index.

2.4. Sociodemographic variables

Sociodemographic variables were obtained by the trained interviewers. The self-report questionnaire was used for the

smoking status: current smoker or not. Participants were also categorized into 2 groups by the number of drinking episodes (0 vs. 1 or more in the 1-month period before the interview).^[17]

Participants were regarded as regular physical exercisers if the participants performed vigorous exercise at least 3 times per week for at least 20 minutes per session or moderate exercise at least 5 times per week for at least 30 minutes per session.^[18] If the monthly house income was $< \$1092.40$, it was considered the lowest quartile. The participants were categorized by those with at least a high school degree and those with less education. Sleep duration and stress recognition were self-reported. Daily energy, fat, protein, and calcium intake were calculated based on the survey.

2.5. Biochemical measurements

Blood pressure measurements were performed using a standard mercury sphygmomanometer (Baumanometer; W.A. Baum Co, Inc, Copiague, NY). Systolic and diastolic blood pressures were measured 2 times at intervals of 5 minutes, and the analysis used the average values.

A blood sample was collected from each participant's antecubital vein after the individual had fasted for > 8 hours to measure concentrations of serum-fasting plasma glucose, glycated hemoglobin (HbA1c), total, low-density lipoprotein, high-density lipoprotein cholesterol, triglycerides; white blood cell count, and serum 25-hydroxyvitamin D. The blood samples were analyzed within 24 hours of transportation. Levels of serum-fasting plasma glucose, HbA1c, cholesterol, and triglycerides were measured with the analyzer (Hitachi Automatic Analyzer 7600, Hitachi, Tokyo, Japan) using enzymatic methods and commercially available kits (Daiichi, Tokyo, Japan).^[19] A gamma counter (1470 Wizard; PerkinElmer, Wallac, Turku, Finland) and a 25-hydroxyvitamin D ^{125}I radioimmunoassay kit (DiaSorin, Stillwater, MN, USA) were utilized to measure serum 25-hydroxyvitamin D levels.

2.6. Description of hypertension, diabetes and metabolic syndrome

If participant had a systolic blood pressure of $> 160 \text{ mmHg}$, a diastolic blood pressure of $> 90 \text{ mmHg}$, or currently used systemic antihypertensive drugs, the individuals were considered having hypertension.^[20] Diabetes was diagnosed when fasting blood sugar was $> 126 \text{ mg/dL}$ or when the individual was currently using antidiabetic medications.^[21] Metabolic syndrome was defined according to the criteria for Asians in the American Heart Association/National Heart, Lung, and Blood Institute Scientific Statement.^[22] Three or more of the following criteria must be met to be diagnosed with metabolic syndrome: waist circumference of at least 90 cm in men or 80 cm in women; fasting triglycerides of at least 150 mg/dL (or use of lipid-lowering medication); high-density lipoprotein cholesterol of $< 40 \text{ mg/dL}$ in men or 50 mg/dL in women (or use of medication); blood pressure of at least 130 mmHg systolic blood pressure and 85 mmHg diastolic pressure (or use of antihypertensive medication); and fasting blood glucose of at least 100 mg/dL (or use of antidiabetic medication).^[23]

2.7. Oral health behaviors

The time of day of patients' tooth brushing was noted as part of their oral health behavior and the frequency of daily tooth brushing was calculated from the oral health behavior.

Table 1**Baseline characteristics of study individuals according to kidney disease, as categorized by urinary ACR and eGFR.**

	Urinary ACR			eGFR		
	<30	≥30	<i>P</i> *	≥60	<60	<i>P</i> *
Unweighted, n	9528	860		10,053	335	
Age, y	45±0.3	58.1±0.8	<0.0001	45.4±0.3	69.4±0.8	<0.0001
Sex (male)	53.1 (0.5)	44.7 (2.0)	<0.0001	52.6 (0.5)	49.8 (3.3)	0.4055
Height, m	164.7±0.1	160.0±0.4	<0.0001	164.5±0.1	158.3±0.6	<0.0001
Weight, kg	64.6±0.2	64.4±0.6	0.7809	64.7±0.2	62.1±0.8	0.0011
Body mass index, kg/m ²	23.7±0.1	25±0.2	<0.0001	23.8±0.1	24.7±0.2	<0.0001
Waist circumference, cm	81.2±0.2	85.9±0.5	<0.0001	81.4±0.2	85.4±0.6	<0.0001
White blood cell count (×10 ³ cells/μL) [†]	5.86 (5.81–5.92)	6.17 (6.02–6.31)	<0.0001	5.87 (5.82–5.92)	6.38 (6.16–6.6)	<0.0001
Energy intake, kcal/day	2071.0±16.4	1826.3±39.2	<0.0001	2066±16.4	1547.0±43.2	<0.0001
Fat intake, g/day	18.9±0.2	14.7±0.4	<0.0001	18.7±0.2	12.2±0.4	<0.0001
Protein intake, g/day	14.8±0.1	13.8±0.2	<0.0001	14.8±0.1	12.6±0.2	<0.0001
Calcium intake, mg/day	517.3±5.5	460.4±15.9	0.0004	516.7±5.5	369.4±19.3	<0.0001
Systolic blood pressure	117.1±0.2	131±0.9	<0.0001	117.8±0.3	129.3±1.5	<0.0001
Diastolic blood pressure	76±0.2	79.3±0.6	<0.0001	76.3±0.2	73.9±1	0.015
Glucose	95.9±0.3	112.4±1.6	<0.0001	96.7±0.3	109.1±2.2	<0.0001
HbA1c	5.7±0.01	6.3±0.1	<0.0001	5.7±0.01	6.3±0.1	<0.0001
Serum 25-hydroxyvitamin D	17.1±0.1	17.5±0.3	0.1637	17.1±0.1	18.6±0.5	0.0008
Cholesterol	189.1±0.5	195±1.5	0.0002	189.6±0.5	184.4±2.9	0.0784
Low-density lipoprotein	111.1±0.4	112.5±1.4	0.3559	111.3±0.4	107.3±2.5	0.115
High-density lipoprotein	52.4±0.2	50.2±0.5	<0.0001	52.4±0.2	46.1±0.9	<0.0001
Triglyceride [‡]	109.3 (107.6–111.1)	139.1 (131.9v146.7)	<0.0001	110.7 (108.9–112.5)	131.1 (121.6–141.2)	<0.0001
Smoking (current)	24.9 (0.6)	19.6 (1.9)	0.0135	24.8 (0.6)	14.4 (2.8)	0.0032
Alcohol within 1 month	59.4 (0.7)	49.9 (2.4)	<0.0001	59.3 (0.7)	34.1 (3.4)	<0.0001
Exercise (yes)	18.9 (0.6)	18 (1.7)	0.6140	19 (0.5)	11.3 (2.2)	0.0044
Income (lowest quartile)	14 (0.6)	30.9 (2)	<0.0001	14.6 (0.6)	42 (3.3)	<0.0001
Education (high-school graduate or higher)	72.9 (0.7)	46.3 (2.4)	<0.0001	72 (0.8)	32.4 (3.4)	<0.0001
Hour of sleep	7.6±0.2	8.0±0.5	0.4526	7.7±0.2	7.6±0.7	0.9701
Stress	27.3 (0.6)	20.6 (1.7)	0.0005	27 (0.6)	17 (2.5)	0.0007
Hypertension	24.4 (0.6)	64.8 (2.4)	<0.0001	26 (0.7)	76.9 (2.7)	<0.0001
Diabetes	7.2 (0.3)	30.5 (2.1)	<0.0001	8.2 (0.4)	36.4 (3.7)	<0.0001
Metabolic syndrome	24.7 (0.6)	58.6 (2.3)	<0.0001	26.2 (0.6)	64.6 (3.3)	<0.0001
Frequency of toothbrushing per day			<0.0001			<0.0001
≤1	11.2 (0.4)	20.5 (2)		11.5 (0.4)	27.4 (3.2)	
2	43.8 (0.8)	42.7 (2.2)		43.8 (0.8)	41.5 (3.3)	
≥3	45 (0.8)	36.8 (2.2)		44.7 (0.8)	31.1 (3.1)	
Use of secondary oral products	57.9 (1.4)	42 (3)	<0.0001	57.3 (1.4)	36.2 (4.2)	<0.0001
Dental checkup within a year	25.8 (0.7)	19 (1.9)	0.0016	25.5 (0.6)	15 (2.5)	0.0004
Self-reported oral status (problematic)	46.4 (0.8)	50.6 (2.2)	0.0668	46.5 (0.8)	57.5 (3.5)	0.0018
Number of natural teeth	25±0.1	20.6±0.4	<0.0001	24.8±0.1	16.5±0.7	<0.0001

Data are presented as mean±standard error or percentage (standard error). ACR=albumin/creatinine ratio, eGFR=estimated glomerular filtration rate

* *P* values were obtained using an independent *t* test for continuous variables or a χ^2 test for categorical variables.[†] A log transformation was applied to the value, and the geometric mean (95% confidence interval) is shown.[‡] Geometric mean (95% confidence interval).

The survey also recorded patients' secondary oral product use, return dental checkup timing (e.g., within a year), and self-reported oral status.

In this study, the natural tooth was considered to be present if the permanent tooth was present or primary tooth was present. The total number of natural teeth was calculated from the examination. Training was provided to each examiner to minimize measurement errors during the examination as part of quality control.

2.8. Statistical analyses

The data were presented as mean±standard error or as a percent (standard error). To achieve normal distribution, a logarithmic transformation was performed when necessary. Either Student *t* test or a χ^2 test was used to investigate the differences as categorized by urinary ACR or eGFR. The association between

the number of natural teeth and kidney function was assessed by multiple logistic regression. Model 1 was adjusted for age, sex, and waist conference. Model 2 was adjusted for the same variables as Model 1 plus smoking, drinking, exercise, education, income, and frequency of tooth brushing per day. Model 3 was adjusted for the variables in Model 2 plus diabetes, metabolic syndrome, urinary ACR, and eGFR. The survey procedure of a statistical program (SAS version 9.2 for Windows, SAS Institute, Cary, NC) was used for statistical analysis to account for the complex sampling design. Two-sided *P* values of 0.05 were used for the statistical significance.

3. Results

Table 1 describes baseline characteristics of the study individuals according to kidney function, as categorized by urinary ACR and eGFR. Number of natural teeth was significantly lower among

Table 2**Subgroup analysis for each group in percentage (standard error).**

The number of natural teeth	Urinary ACR ≥ 30				eGFR < 60			
	≤ 19	20–27	28	P	≤ 19	20–27	28	P
Age, y								
19–64	12.8 (1.7)	6.0 (0.5)	3.1 (0.3)	<0.0001	2.5 (0.8)	0.8 (0.2)	0.4 (0.1)	<0.0001
65+	19.6 (1.3)	15.1 (1.5)	10.2 (2.7)	0.0063	11.8 (1.1)	9.4 (1.1)	2.7 (1.1)	0.002
Sex								
Male	15.9 (1.5)	6.3 (0.7)	2.8 (0.4)	<0.0001	8 (1.1)	2 (0.3)	0.6 (0.2)	<0.0001
Female	18 (1.6)	8.3 (0.7)	3.8 (0.4)	<0.0001	8.6 (1)	2.0 (0.3)	0.3 (0.1)	<0.0001
Body mass index								
<25	15.8 (1.2)	4.9 (0.5)	2.5 (0.3)	<0.0001	7.2 (0.8)	1.6 (0.2)	0.2 (0.1)	<0.0001
≥ 25	19.8 (2)	11.3 (1)	5.0 (0.7)	<0.0001	10.7 (1.7)	2.7 (0.4)	1.0 (0.3)	<0.0001
Waist circumference								
Male: <90 cm, Female: <80 cm	14.4 (1.4)	4.4 (0.4)	2.2 (0.3)	<0.0001	7.1 (0.9)	1.6 (0.2)	0.4 (0.1)	<0.0001
Male: ≥ 90 cm, Female: ≥ 80 cm	20.5 (1.7)	11.8 (1)	6.3 (0.9)	<0.0001	9.5 (1.2)	2.6 (0.4)	0.7 (0.2)	<0.0001
Hypertension								
No	8.0 (1.1)	3.6 (0.4)	2.2 (0.3)	<0.0001	5 (0.9)	0.5 (0.1)	0.1 (0)	<0.0001
Yes	23.9 (1.7)	14.9 (1.2)	9.4 (1.2)	<0.0001	10.4 (1.1)	4.8 (0.6)	2.6 (0.7)	<0.0001
Diabetes								
No	13.3 (1.3)	5.7 (0.5)	2.5 (0.3)	<0.0001	6.6 (0.8)	1.3 (0.2)	0.3 (0.1)	<0.0001
Yes	30.8 (2.9)	19.1 (2.2)	19 (2.9)	0.0011	13.4 (2.2)	7.3 (1.3)	2.6 (1.4)	0.0009
Metabolic syndrome								
No	12.7 (1.4)	4.1 (0.4)	2.0 (0.3)	<0.0001	5.7 (0.9)	1.0 (0.2)	0.2 (0.1)	<0.0001
Yes	22.2 (1.6)	13.7 (1.2)	9.0 (1.3)	<0.0001	10.5 (1.2)	4.1 (0.6)	1.5 (0.5)	<0.0001

ACR=albumin/creatinine ratio, eGFR=estimated glomerular filtration rate

those with lower kidney function (urinary ACR ≥ 30 mg/g and eGFR < 60 mL/min/1.73m²). The mean age, body mass index, and waist circumference were significantly higher among those with lower kidney function. Systolic and diastolic blood pressure, glucose, HbA1c, and triglycerides were also statistically higher among those with kidney diseases. High-density lipoprotein levels were significantly lower among those with kidney diseases. The percentage of individuals in the lowest-income quartile was higher among those with lower kidney function. The percentage of patients with at least a high school degree was lower among those with kidney disease. The prevalence of hypertension, diabetes, and metabolic syndrome was significantly higher among those with kidney disease. The subgroup analysis (categorized by number natural teeth for urinary ACR ≥ 30 mg/g and eGFR < 60 mL/min/1.73m²) is shown in Table 2. The number of natural teeth was lower among the participants with body mass index of ≥ 25 ($P < 0.01$). Similarly, the number of natural teeth was lower among the participants with hypertension, diabetes, or metabolic syndrome ($P < 0.01$).

Figure 2 shows the percentage and standard error for the participants, categorized by number of natural teeth and kidney status. As urinary ACR increased, the number of natural teeth decreased; conversely, the number of natural teeth increased as eGFR increased.

Table 3 shows the adjusted odds ratios and their 95% confidence intervals from the multiple logistic regression analyses for the individuals with kidney diseases (categorized by urinary ACR ≥ 30 mg/g or eGFR < 60 mL/min/1.73m²) according to the number of natural teeth. The odds ratios increased in the individuals with urinary ACR ≥ 30 mg/g. The adjusted odds ratios (with 95% confidence intervals) among participants with urinary ACR ≥ 30 mg/g were 1 [reference] for individuals with 28 natural teeth, 1.298 (0.989–1.703) for those with 20 to 27 teeth, and 1.758 (1.302–2.375) among those with ≤ 19 teeth. The adjusted odds ratios (with 95% confidence intervals) for participants with eGFR < 60 mL/min/1.73m² were 1 [reference] for individuals with 28 natural teeth, 1.329 (0.691–2.557) for those with 20 to 27 teeth, and 1.287 (0.659–2.513) for those with ≤ 19 teeth.




Figure 2. (A) Percentage and standard error for number of natural teeth, as categorized by kidney function using urinary ACR. (B) Percentage and standard error for number of natural teeth, as categorized by kidney function using eGFR.

Table 3

Adjusted odds ratios, 95% confidence intervals, and P values in multivariate logistic regression models based on number of natural teeth and kidney status in mean (95% confidence interval).

Number of natural teeth	Model 1	Model 2	Model 3*
		Urinary ACR ≥ 30	
28	1	1	1
20–27	1.329 (1.031–1.713)	1.337 (1.023–1.746)	1.298 (0.989–1.703)
≤ 19	1.825 (1.379–2.415)	1.727 (1.291–2.309)	1.758 (1.302–2.375)
P for trend	0.0001	0.0002	0.0003
Number of natural teeth	Model 1	Model 2	Model 3†
		eGFR < 60	
28	1	1	1
20–27	1.130 (0.643–1.989)	1.238 (0.675–2.271)	1.329 (0.691–2.557)
≤ 19	1.205 (0.681–2.132)	1.303 (0.686–2.476)	1.287 (0.659–2.513)
P for trend	0.5046	0.4406	0.5747

ACR = albumin/creatinine ratio, eGFR = estimated glomerular filtration rate. Model 1: adjusted for age, sex, and waist circumference. Model 2: Model 1 adjustments plus smoking, drinking, exercise, education, income, and frequency of tooth brushing per day.

* Model 3: Model 2 adjustments plus diabetes, metabolic syndrome, urinary eGFR.

† Model 3: Model 2 adjustments plus diabetes, metabolic syndrome, urinary ACR.

4. Discussion

This study clearly showed that the odd ratios for having fewer natural teeth tended to increase among the participants with lower kidney function (urinary ACR ≥ 30 mg/g and eGFR < 60 mL/min/1.73m²).

Kidney function may be determined using various methods.^[12,24–26] This study clearly showed that both urinary ACR and eGFR were associated with loss of natural teeth. The urine albumin test is used to evaluate kidney function and to screen people with chronic conditions, such as hypertension and diabetes, which put patients at an increased risk of kidney disease.^[27] Albumin is a protein that is present in high concentrations in the blood; it is not usually detected in the urine when the kidneys are functioning properly.^[28] Creatinine is a byproduct of muscle metabolism; its level in the urine is usually considered an indication of the urine concentration because it is normally released into the urine at a constant rate.^[29] Albuminuria has been shown to be related with periodontal disease.^[30] A previous report showed that decreased kidney function (characterized by low eGFR) may be associated with periodontitis.^[24] International recommendations suggest that the measurement of serum creatinine should be supplemented with eGFR using the MDRD study's equation.^[31,32] This previous report showed that the MDRD equation provided reasonably accurate GFR estimates in patients with chronic kidney disease.^[31] The lack of standardization of commercially available creatinine assays, which resulted in varying estimates of GFR, was considered to be problematic, and another study suggested the use of isotope dilution mass spectrometry without the requirement of standardization to the MDRD laboratory.^[32] Renal function may be estimated using the Chronic Kidney Disease Epidemiology Collaboration equation.^[25,26]

The mechanisms behind the association between kidney function and the number of natural teeth may be explained

partially by the following. Inflammation can be considered to be the intermediate factor between oral health behavior and systemic diseases.^[33] Poor oral health behavior may worsen inflammation; a previous report showed that poor oral hygiene was related to higher inflammatory markers.^[34,35] Periodontitis is a chronic inflammatory disease that results from a microbial infection within the subgingival dental plaque biofilm; the resulting inflammatory response may facilitate intravascular dissemination of microorganisms and their products throughout the body.^[36] This disease destroys tooth-supporting tissues and may lead to loss of teeth.^[37] Periodontitis leading to tooth loss has been labeled as an important potential risk factor for non-communicable diseases, including diabetes mellitus, cardiovascular diseases, pulmonary diseases, and osteoporosis.^[6] A higher percentage of periodontal disease among patients with renal disease was noted when these patients were compared with healthy individuals.^[7] The prevalence and severity of chronic periodontitis in the previous cohort of chronic kidney disease patients was markedly higher than it was in a geographically matched control population in Europe.^[38] In another report, poor oral health, which includes chronic periodontitis, was a common finding in patients undergoing hemodialysis, and chronic periodontitis was considered a continuous, reversible source of inflammation and this has a potential impact on mortality in patients undergoing hemodialysis.^[39] The authors suggested intervention trials to test the hypothesis that treatment of chronic periodontitis may improve survival in patients undergoing hemodialysis.^[39]

The most important strength of this study is that it is based on a nationally representative sample of Koreans. KNHANES is a nationwide survey of noninstitutionalized civilians; it uses sample participants and sample weights to represent the Korean population and considers survey nonresponse, complex survey design, and poststratification.^[40] However, some limitations should be considered regarding this study, as it used a cross-sectional design, which makes it difficult to determine whether there is a direct relationship between the exposure and the outcome.^[41] Another limitation of this study is that individuals' energy intake and oral health behavior were obtained self-reported and were under recall basis.^[42,43] Data regarding the inflammatory markers (such as C-reactive protein, tumor necrosis factor- α , and interleukin-6) were not available, so the study's explanation of inflammation and kidney disease is limited.^[44]

This study showed that the number of natural teeth is inversely associated with the presence of kidney disease. Among those with reduced kidney function (eGFR < 60 mL/min/1.73m²), the adjusted odds ratios for the group with 28 natural teeth were lower than the ratios for those with fewer natural teeth. Tooth loss may be considered an independent risk indicator for kidney disease in Korean people. More epidemiological studies are warranted to investigate the role of tooth loss in participants with kidney disease, to confirm this relationship, and to test possible underlying mechanisms.

Acknowledgments

The authors acknowledge the Korea Centers for Disease Control and Prevention for providing the data.

References

- Wilder RS, Bell KP, Phillips C, et al. Dentists' practice behaviors and perceived barriers regarding oral-systemic evidence: implications for education. *J Dent Educ* 2014;78:1252–62.

- [2] Paquette DW, Bell KP, Phillips C, et al. Dentists' knowledge and opinions of oral-systemic disease relationships: relevance to patient care and education. *J Dent Educ* 2015;79:626–35.
- [3] Niedzielska I, Chudek J, Kowol I, et al. The odontogenic-related microinflammation in patients with chronic kidney disease. *Ren Fail* 2014;36:883–8.
- [4] Madore F. Periodontal disease: a modifiable risk factor for cardiovascular disease in ESRD patients? *Kidney Int* 2009;75:672–4.
- [5] Casanova L, Hughes FJ, Preshaw PM. Diabetes and periodontal disease: a two-way relationship. *Br Dent J* 2014;217:433–7.
- [6] Wahid A, Chaudhry S, Ehsan A, et al. Bidirectional relationship between chronic kidney disease & periodontal disease. *Pak J Med Sci* 2013;29:211–5.
- [7] Tiwari V, Saxena V, Bhambhal A, et al. The oral health status of patients with renal disease in central India: a preliminary study. *J Ren Care* 2013;39:208–13.
- [8] Ariyamuthu VK, Nolph KD, Ringdahl BE. Periodontal disease in chronic kidney disease and end-stage renal disease patients: a review. *Cardiorenal Med* 2013;3:71–8.
- [9] Murakami M, Suzuki J, Yamazaki S, et al. High incidence of *Aggregatibacter actinomycetemcomitans* infection in patients with cerebral infarction and diabetic renal failure: a cross-sectional study. *BMC Infect Dis* 2013;13:557.
- [10] Fisher MA, Taylor GW, Shelton BJ, et al. Periodontal disease and other nontraditional risk factors for CKD. *Am J Kidney Dis* 2008;51:45–52.
- [11] Han SY, Hong JW, Noh JH, et al. Association of the estimated 24-h urinary sodium excretion with albuminuria in adult Koreans: the 2011 Korea National Health and Nutrition Examination Survey. *PLoS one* 2014;9:e109073.
- [12] Ahn JH, Yu JH, Ko SH, et al. Prevalence and determinants of diabetic nephropathy in Korea: Korea national health and nutrition examination survey. *Diabetes Metab J* 2014;38:109–19.
- [13] Park JI, Baek H, Jung HH. CKD and health-related quality of life: The Korea National Health and Nutrition Examination Survey. *Am J Kidney Dis* 2015.
- [14] So BH, Methven S, Hair MD, et al. Socio-economic status influences chronic kidney disease prevalence in primary care: a community-based cross-sectional analysis. *Nephrol Dial Transplant* 2015;30:1010–7.
- [15] Weisell RC. Body mass index as an indicator of obesity. *Asia Pac J Clin Nutr* 2002;11(Suppl 8):S681–4.
- [16] Oh SW, Shin SA, Yun YH, et al. Cut-off point of BMI and obesity-related comorbidities and mortality in middle-aged Koreans. *Obes Res* 2004;12:2031–40.
- [17] Agarwal DP. Cardioprotective effects of light-moderate consumption of alcohol: a review of putative mechanisms. *Alcohol Alcohol* 2002;37:409–15.
- [18] Oh JY, Yang YJ, Kim BS, et al. Validity and reliability of Korean version of International Physical Activity Questionnaire (IPAQ) short form. *J Korean Acad Fam Med* 2007;28:532–41.
- [19] Wallace TM, Levy JC, Matthews DR. Use and abuse of HOMA modeling. *Diabetes Care* 2004;27:1487–95.
- [20] Weber MA, Julius S, Kjeldsen SE, et al. Blood pressure dependent and independent effects of antihypertensive treatment on clinical events in the VALUE Trial. *Lancet* 2004;363:2049–51.
- [21] Jeon JY, Ko SH, Kwon HS, et al. Prevalence of diabetes and prediabetes according to fasting plasma glucose and HbA1c. *Diabetes Metab J* 2013;37:349–57.
- [22] Chun YH, Kim HR, Han K, et al. Total cholesterol and lipoprotein composition are associated with dry eye disease in Korean women. *Lipids Health Dis* 2013;12:84.
- [23] Chin SO, Rhee SY, Chon S, et al. Sarcopenia is independently associated with cardiovascular disease in older Korean adults: the Korea National Health and Nutrition Examination Survey (KNHANES) from 2009. *PLoS One* 2013;8:e60119.
- [24] Iwasaki M, Taylor GW, Sato M, et al. Cystatin C-based estimated glomerular filtration rate and periodontitis. *Gerodontology* 2016;33:328–34.
- [25] Choi HY, Joo DJ, Song MK, et al. The power of renal function estimation equations for predicting long-term kidney graft survival: a retrospective comparison of the chronic kidney disease epidemiology collaboration and the modification of diet in renal disease study equations. *Medicine (Baltimore)* 2016;95:e2682.
- [26] Terpos E, Christoulas D, Kastritis E, et al. The Chronic Kidney Disease Epidemiology Collaboration cystatin C (CKD-EPI-CysC) equation has an independent prognostic value for overall survival in newly diagnosed patients with symptomatic multiple myeloma; is it time to change from MDRD to CKD-EPI-CysC equations? *Eur J Haematol* 2013;91:347–55.
- [27] Martinez-Castelao A, Gorritz JL, Segura-de la Morena J, et al. Consensus document for the detection and management of chronic kidney disease. *Nefrologia* 2014;34:243–62.
- [28] Buser MC, Ingber SZ, Raines N, et al. Urinary and blood cadmium and lead and kidney function: NHANES 2007-2012. *Int J Hyg Environ Health* 2016;219:261–7.
- [29] Ix JH, de Boer IH, Wassel CL, et al. Urinary creatinine excretion rate and mortality in persons with coronary artery disease: the Heart and Soul Study. *Circulation* 2010;121:1295–303.
- [30] Salimi S, Ng N, Seliger SL, et al. Periodontal disease, renal dysfunction and heightened leukocytosis. *Nephron Clin Pract* 2014;128:107–14.
- [31] Levey AS, Coresh J, Greene T, et al. Using standardized serum creatinine values in the modification of diet in renal disease study equation for estimating glomerular filtration rate. *Ann Intern Med* 2006;145:247–54.
- [32] Vickery S, Stevens PE, Dalton RN, et al. Does the ID-MS traceable MDRD equation work and is it suitable for use with compensated Jaffe and enzymatic creatinine assays? *Nephrol Dial Transplant* 2006;21:2439–45.
- [33] de Oliveira C, Watt R, Hamer M. Toothbrushing, inflammation, and risk of cardiovascular disease: results from Scottish Health Survey. *BMJ (Clinical research ed)* 2010;340:c2451.
- [34] Akar H, Akar GC, Carrero JJ, et al. Systemic consequences of poor oral health in chronic kidney disease patients. *Clin J Am Soc Nephrol* 2011;6:218–26.
- [35] Li X, Kolltveit KM, Tronstad L, et al. Systemic diseases caused by oral infection. *Clin Microbiol Rev* 2000;13:547–58.
- [36] Mohangi GU, Singh-Rambirich S, Volchansky A. Periodontal disease: Mechanisms of infection and inflammation and possible impact on miscellaneous systemic diseases and conditions. *Sadj* 2013;68:464–7.
- [37] How KY, Song KP, Chan KG. *Porphyromonas gingivalis*: an overview of periodontopathic pathogen below the gum line. *Front Microbiol* 2016;7:53.
- [38] Sharma P, Dietrich T, Sidhu A, et al. The periodontal health component of the Renal Impairment In Secondary Care (RIISC) cohort study: a description of the rationale, methodology and initial baseline results. *J Clin Periodontol* 2014;41:653–61.
- [39] de Souza CM, Braosi AP, Luczyszyn SM, et al. Association among oral health parameters, periodontitis, and its treatment and mortality in patients undergoing hemodialysis. *J Periodontol* 2014;85:e169–78.
- [40] Han K, Ko Y, Park YG, et al. Associations between the number of natural teeth in postmenopausal women and duration of lactation: The 2010–2012 Korea National Health and Nutrition Examination Survey. *Maturitas* 2016;85:73–8.
- [41] Park JB, Han K, Park YG, et al. Association between alcohol consumption and periodontal disease: the 2008 to 2010 Korea National Health and Nutrition Examination Survey. *J Periodontol* 2014;85:1521–8.
- [42] Han K, Hwang E, Park JB. Association between Consumption of Coffee and the Prevalence of Periodontitis: The 2008–2010 Korea National Health and Nutrition Examination Survey. *PLoS One* 2016;11:e0158845.
- [43] Han K, Hwang E, Park JB. Excessive Consumption of Green Tea as a Risk Factor for Periodontal Disease among Korean Adults. *Nutrients* 2016;8.
- [44] Choi HM, Han K, Park YG, et al. Associations Among Oral Hygiene Behavior and Hypertension Prevalence and Control: The 2008 to 2010 Korea National Health and Nutrition Examination Survey. *J Periodontol* 2015;86:866–73.