

CORRECTION

Correction: Short-term anti-proteinuric effect of tacrolimus is not related to preservation of the glomerular filtration rate in IgA nephropathy: A 5-year follow-up study

Mi-yeon Yu, Yong-Chul Kim, Ho Suk Koo, Ho Jun Chin

The x-axis labels for [Fig 2D](#) are cut-off. The x-axis labels for [Fig 3B](#) are incorrect. Please see the correct [Fig 2](#) and [Fig 3](#) here.



OPEN ACCESS

Citation: Yu M-y, Kim Y-C, Koo HS, Chin HJ (2018) Correction: Short-term anti-proteinuric effect of tacrolimus is not related to preservation of the glomerular filtration rate in IgA nephropathy: A 5-year follow-up study. PLoS ONE 13(1): e0192266. <https://doi.org/10.1371/journal.pone.0192266>

Published: January 29, 2018

Copyright: © 2018 Yu et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.




Fig 2. The changes in UPCr (A, C) and eGFR (B, D) during the follow-up period after cessation of the clinical study. From left to right, each box-plot represents the follow-up periods of 0-weeks, 4-weeks, 8-weeks, 12-weeks and 16-weeks of the trial phase, and 12-weeks, 24-weeks, 52-weeks, and the final visit by October 2016 of the observational phase. A. The p-value for tests of between two groups was 0.130. B. The p-value for tests of between-subjects effects was 0.543. C. The p-value for tests of between-subjects effects was 0.488. Tac: Tacrolimus, RASi: renin-angiotensin-aldosterone system inhibitor, UPCr: urine protein to creatinine ratio, eGFR: estimated glomerular filtration rate by the equation of CKD-EPI. The p-value estimated by linear mixed effect model.

<https://doi.org/10.1371/journal.pone.0192266.g001>




Fig 3. The difference in outcome parameters between groups. Outcome parameters represent the followings; O1, decrease of time-averaged proteinuria from trial phase and to the observational phase; O2, remission of UPCr <0.2 g/g cr during observational phase; O3, rapid decline of eGFR ≥ 5 mL/min/1.73 m² during observational phase; O4, composite outcome of increase in serum cr level ($\geq 50\%$ from baseline) noted during observational phase or deterioration of renal function to end stage renal disease. Outcome parameters were compared by Chi-square test or Fisher's exact test according to the number of each cell in the tacrolimus group (A) and RASi group (B).

<https://doi.org/10.1371/journal.pone.0192266.g002>

Reference

1. Yu M-y, Kim Y-C, Koo HS, Chin HJ (2017) Short-term anti-proteinuric effect of tacrolimus is not related to preservation of the glomerular filtration rate in IgA nephropathy: A 5-year follow-up study. PLoS ONE 12(11): e0188375. <https://doi.org/10.1371/journal.pone.0188375> PMID: 29155873