



Published in final edited form as:

Nat Commun. ; 5: 5165. doi:10.1038/ncomms6165.

Suppression of the FOXM1 transcriptional program via novel small molecule inhibition

Michael V. Gormally^{1,2,3}, Thomas S. Dexheimer³, Giovanni Marsico², Deborah A. Sanders², Christopher Lowe¹, Dijana Matak-Vinkovi¹, Sam Michael³, Ajit Jadhav³, Ganesh Rai³, David J. Maloney³, Anton Simeonov³, and Shankar Balasubramanian^{1,2,*}

¹Department of Chemistry, University of Cambridge, Cambridge, CB2 1EW, UK

²Cancer Research UK, Cambridge Institute, Li Ka Shing Centre, Cambridge, CB2 0RE, UK

³National Center for Advancing Translational Sciences, NIH, Rockville, Maryland, 20850, USA

Abstract

The transcription factor FOXM1 binds to sequence-specific motifs on DNA (C/TAAACA) through its DNA binding domain (DBD), and activates proliferation- and differentiation-associated genes. Aberrant overexpression of FOXM1 is a key feature in oncogenesis and progression of many human cancers. Here — from a high-throughput screen applied to a library of 54,211 small molecules — we identify novel small molecule inhibitors of FOXM1 that block DNA binding. One of the identified compounds: FDI-6 (NCGC00099374) is characterized in depth and is shown to bind directly to FOXM1 protein, to displace FOXM1 from genomic targets in MCF-7 breast cancer cells, and induce concomitant transcriptional down-regulation. Global transcript profiling of MCF-7 cells by RNA-seq shows that FDI-6 specifically down regulates FOXM1-activated genes with FOXM1 occupancy confirmed by ChIP-seq. This small molecule mediated effect is selective for FOXM1-controlled genes with no effect on genes regulated by homologous forkhead family factors.

INTRODUCTION

Evidence is accumulating to implicate deregulation of transcription factor networks as a major pathogenic event in many human cancers¹⁻³. In healthy cells, transcription factors control the spatial and temporal conversion of DNA-encoded information into functional RNAs and proteins, thereby directing complex signaling pathways crucial to cellular function^{4,5}. In cancer, mutations in upstream regulators, aberrant gene amplification, and

Users may view, print, copy, and download text and data-mine the content in such documents, for the purposes of academic research, subject always to the full Conditions of use:http://www.nature.com/authors/editorial_policies/license.html#terms

*Correspondence: sb10031@cam.ac.uk.

Author Contributions

M.V.G., D.S., S.B., T.S.D., S.M., A.J. and A.S. designed research; M.V.G. and T.S.D. performed experiments; D. M.-V. performed mass spectrometry analysis; M.V.G., C.L., G.M., T.S.D., G.R., A.J., D.J.M., A.S., and S.B. analyzed data; M.V.G., S.B. wrote the paper, with edits from all authors.

Accession codes: RNAseq profiling GEO: GSE58626

Competing financial interests: The authors declare no competing financial interests.

further perturbations resulting from such changes may destabilize proper function of the transcription factor network and drive disease⁶⁻⁸. There is a need for tools to intervene directly with transcription factors to dissect how the complex feedback and regulatory mechanisms work in a healthy context, and to evaluate the potential for exploiting these targets therapeutically. Small molecule intervention is a particularly attractive avenue to address both of these points⁹⁻¹¹. First, ligands can be designed with favorable pharmacokinetic properties or modified to provide spatio-temporal delivery. Second, ligands tailored to specific domains or interfaces that minimize off-target effects may then be employed as tools to further interrogate the target's biological function. Unfortunately, transcription factors outside the nuclear receptor family lack substrate binding pockets, are often characterized by hydrophobic surfaces with few druggable regions and have historically proven difficult targets¹². Past efforts to perturb transcriptional pathways have therefore focused on known protein-protein interfaces or particular recognition elements on DNA. Notable examples include distamycin-inspired polyamides, which recognize the DNA minor groove in a sequence-specific manner and have been successfully applied to disrupt assembly of general transcription machinery at those loci¹³. Peptidomimetics, like the stapled alpha-helix BCL-2 domains (SAHBs) have generated protease resistant, cell-permeable tools to antagonize protein dimerization and induce selective activation of apoptotic pathways in cancer cells¹⁴. Such a strategy might be applied to disrupt transcription factor interactions with promoter DNA. Additionally, a recent publication reports a novel small molecule that directly binds to the oncogenic transcription factor ETV1, and inhibits its transcriptional activity¹⁵.

The transcription factor FOXM1 regulates a network of proliferation-associated genes critical to mitotic spindle assembly¹⁶, chromosome segregation¹⁷, and G2/M transition¹⁸, with depletion leading to cell cycle arrest. Importantly, aberrant up-regulation of FOXM1 has been shown to be a key driver of cancer progression and has been proposed as an initiating factor of oncogenesis^{3,19,20}. Furthermore, FOXM1 overexpression has been implicated in the development of chemotherapeutic resistance in human breast cancer^{21,22}, high protein levels correlate with poor clinical outcome^{23,24} and the endogenous level of FOXM1 has been suggested as a general diagnostic biomarker for cancer progression²⁵. Thus, inhibition of FOXM1 activity is an attractive goal for cancer therapy. FOXM1 functions as a gene-specific transcriptional activator by binding to DNA consensus sequences through a highly conserved and well-characterized DNA binding domain (DBD)²⁶. While such an interface has been recognized as an enticing target, the absence of small molecules known to bind FOXM1 makes structure-based rational design of drug candidates challenging²⁷.

Previously, others have shown that the natural product thiostrepton inhibits the transcriptional activity of FOXM1^{28,29}. We demonstrated that the ligand makes direct contact with FOXM1 and inhibits DNA binding in cells³⁰. However, thiostrepton is a promiscuous molecule with potent off-target effects most notably inhibition of the 20S proteasome³¹ and arrest of mitochondrial protein synthesis³². Thus thiostrepton cannot be used to elucidate the primary effects caused by the inhibition of FOXM1 binding to DNA³³. We designed and exploited a robust, biophysical high-throughput screening assay to identify inhibitors of FOXM1 that block DNA binding. We discovered an inhibitor that blocks

FOXM1 binding to DNA in human cancer cells and suppresses the transcription of genes under FOXM1 control.

RESULTS

FOXM1 DNA binding domain associates with consensus motif

To interrogate binding of FOXM1 to its consensus DNA duplex, we designed a fluorescence polarization (FP) assay (Fig. 1A). In this system association or dissociation of fluorophore-labeled DNA duplex to FOXM1 protein can be monitored as a function of linear and perpendicular polarization³⁴. We generated a 16 bp dsDNA probe containing repeats of the known FOXM1 recognition motif (T/CAAACA) and a single 6-carboxyfluorescein (6-FAM) label ligated to the 5' end of the forward strand. Recombinant FOXM1 DNA binding domain (DBD), a truncated fusion product containing only the winged helix domain (aa 222–360), was titrated with dsDNA probe and examined by FP, which demonstrated that FOXM1 DBD associates with the DNA probe in a dose dependent manner with $K_D = 224 \pm 30$ nM (Fig. 1B). This FOXM1 DBD construct has been previously identified as the minimal component of the canonical FOXM1-DNA binding mode²⁶. Notably, the dissociation constant for this truncated form of the protein for its 16 bp DNA consensus target is in good agreement with that of the full-length protein as determined by FP ($K_D = 300 \pm 64$ nM, Supplementary Fig. 1) or by electrophoretic mobility shift assay (EMSA)³⁰. Thus, we employed FOXM1 DBD to discover novel inhibitors of DNA binding.

Titration of a pre-bound complex of FOXM1 DBD and labeled dsDNA probe with increasing amounts of unlabeled dsDNA probe demonstrated the reversibility and specificity of the association (Fig. 1C). As expected, the IC_{50} of this displacement (57 ± 5 nM) was near the concentration of the labeled probe (50 nM), indicating that the 5'-carboxyfluorescein tag did not affect protein binding. Furthermore, thioestrepton, a previously identified inhibitor of FOXM1 DNA binding, disrupted the FOXM1 DBD-DNA interaction in gel shift assays with $IC_{50} = 45.0 \pm 16.4$ μ M (Supplementary Fig. 2), in good agreement with previously reported values using full length FOXM1, $IC_{50} = 42.6 \pm 1.4$ μ M³⁰.

Development of screening assay for inhibitors of FOXM1—DNA

To identify inhibitors of FOXM1 DNA binding, we developed a high throughput screen based on the perturbation of FOXM1 DBD—DNA complex measured by FP³⁵. Minor adjustments were made to the assay as presented above (Fig. 1C) to provide suitability for a high-throughput screening (HTS) protocol in 1536-well plate format. The assay buffer was supplemented with 0.01% Tween-20 and 0.1 mg mL⁻¹ bovine serum albumin to facilitate robotic dispensing, stabilize the FOXM1 DBD protein and minimize hydrophobic aggregation of material on plastic. An apparent K_D value of 158 nM was calculated for binding of FOXM1 DBD to the 16-bp DNA in 384-well plate format, showing good agreement with previous values from the initial FP runs (Supplementary Fig. 3A). The concentration of 400 nM FOXM1 DBD was chosen for subsequent experiments, which resulted in average window of 135 mP units between bound and unbound signals in 1536-well plate format (Supplementary Fig. 3B). Pipetting was simplified into two robotic dispense steps: 3 μ L 1.3X FOXM1 DBD + 1 μ L 4X DNA probe. In calibration runs with the

high throughput format, the signal-to-background ratio was approximately 3.2 and the Z' factor was 0.77. The Z' factor quantifies the statistical significance of the difference between positive and negative controls. A value above 0.5 indicates that more than 12 standard deviations separate the bound and unbound controls; therefore, 0.77 is well within the norm for a robust assay.

Identification of small molecule inhibitors of FOXM1—DNA

A collection of small molecules was screened at five concentrations ranging from 0.46 μM to a maximum of 115 μM . The screening library was comprised of a total of 54,211 compounds assembled at the National Center for Advancing Translational Sciences (NCATS). It consists of diverse drug-like molecules that provide a starting point for medicinal chemistry lead development. Compounds with annotated pharmacology including the NCGC Pharmaceutical Collection and the Tocris Biosciences diversity library were also included for potential repurposing³⁶. FOXM1 DBD was first dispensed into wells and compounds (23 nL at each concentration) were transferred with a Kalypsys 1536-pin tool. After 15 min pre-incubation, the DNA probe was dispensed and the mixture was incubated for a further 30 min before reading FP signal.

The assay reported consistently high Z' factors (average = 0.71) throughout all plates in the screen (Fig. 1D). Furthermore, a suitably low active (compounds with max inhibition >60%) hit rate of 0.32% suggested a useful screen and the existence of a druggable interaction. The normalized dose response data for all positive hits from the HTS are displayed as a waterfall plot in Fig. 1E with 170 actives shown in blue. All 170 actives from initial screening were checked for reproducibility and retested in the same assay, validating 86 of the original actives. Active molecules that exhibited poor reproducibility, bad dose response profile, non-specific DNA binding activity (hits in thiazole orange displacement counter screen), or were found to disrupt the FP assay (fluorescent artifacts or quenchers) were eliminated³⁷. Ultimately, 16 Forkhead Domain Inhibitory compounds (renamed FDI-(1-16), structures provided in Supplementary Fig. 4) were selected for further analysis.

Orthogonal EMSAs validate six active FDI compounds

EMSA was chosen as an orthogonal assay to further evaluate FDI-(1-16) in dose response. The binding of FOXM1 DBD is detected by EMSA ($K_D=301\pm 21$ nM, Fig. 2A) and is near the value reported by FP ($K_D=224\pm 30$ nM, Fig. 1B). To examine the effect of the inhibitors, preassembled complexes of FOXM1 DBD was incubated with increasing concentrations of ligand for 1.5 h at RT, target DNA was added, electrophoresis was performed, and DNA probe bands were visualized by fluorescence imaging. In this assay, 6 of the 16 FDI compounds exhibited dose-dependent inhibition of the FOXM1 DBD-DNA binding interaction (Fig. 2B, representative gels provided in Supplementary Fig. 6), confirming their activity in the primary FP screen. The remaining compounds were inactive below 200 μM ; these were presumed to be false positives in the FP assay and discarded. Values were also determined for 50% growth inhibition after 72 h ligand treatment (GI_{50}) in MCF-7 breast cancer cells, which highly express FOXM1 (Fig. 2B, inset and Supplementary Fig. 6). **FDI-6** was the most potent EMSA-validated compound, with a representative EMSA image shown in Fig. 2C. In addition, the concordance between GI_{50} and IC_{50} values for FDI-6 is

indicative of minimal off target activity. Furthermore, two compounds (**FDI-6** and **FDI-10**) exhibited potent IC_{50} values ($22.5 \pm 12.3 \mu\text{M}$ and $34.36 \pm 11.2 \mu\text{M}$, respectively) comparable to that of thiostrepton ($IC_{50} = 45.0 \pm 16.4 \mu\text{M}$, see Supplementary Fig. 2). It was noteworthy that the top three validated compounds by potency—**FDI-6**, **FDI-10**, and **FDI-11**—shared a common core structure (Fig. 2D) suggestive of a potential pharmacophore.

FDI-6 binds FOXM1 and displaces protein from target genes

Non-specific DNA recognition by the small molecule hits was already ruled out by the thiazole orange counter-screen (see earlier). It was then important to confirm that our shortlisted hits (**FDI-6**, **FDI-10**, and **FDI-11**) act via direct interaction with the protein, which we achieved by native mass spectrometry. FOXM1 protein is accurately detected by this method and exists as a dimer in solution, as is frequently observed for GST fusions³⁸. Hit compounds **FDI-6**, **FDI-10** and **FDI-11** were each found to bind directly to FOXM1 protein with a 1:1 stoichiometry (Figure 3A). We also tested FDI-9, a compound that was active in the primary FP assay, but failed to inhibit FOXM1-DNA by EMSA, and showed that, in contrast, it does not interact with the protein. This finding supports the use of EMSA as an orthogonal assay to flag artifacts and indirect inhibitors.

To determine whether an inhibitor of FOXM1-DNA in vitro could inhibit transcription factor binding in MCF-7 breast cancer cells, which highly express FOXM1³⁹, we selected the most potent ligand **FDI-6** for follow up experiments. Upon isolating the nuclear fractions of FDI-6 treated MCF-7 cells by centrifugation, western blots revealed that a 6 h dose of $20 \mu\text{M}$ **FDI-6**—approximately the biophysical IC_{50} of the ligand for disrupting the FOXM1-DNA interaction—decreased chromatin bound FOXM1 by nearly 50% while the overall amount of FOXM1 protein was unaffected (Fig. 3B&C). Importantly, and in contrast to reports for thiostrepton⁴⁰, **FDI-6** did not inhibit the 20S proteasome as determined by a standard proteasome assay (Supplementary Fig. 7).

The observation that **FDI-6** displaces FOXM1 from chromatin led us to initially investigate its effect on the gene targets *CCNB1* and *CDC25B*^{6,16}. Activation of these target genes has been attributed to the oncogenic potential of FOXM1, and depletion of FOXM1 reduced colony formation of acute myeloid leukemia cells due to inhibition of *CCNB1* and *CDC25B* expression⁴¹. Importantly, direct occupancy of FOXM1 at its consensus sequence within both *CCNB1* and *CDC25B* promoter regions activates transcription in reporter assays⁴². Thus, these were ideal targets to initially test the ability of **FDI-6** to disrupt FOXM1-DNA in the cell. To first examine FOXM1 occupancy in these regions, we performed chromatin immunoprecipitation (ChIP) with FOXM1 antibody followed by quantitative real time polymerase chain reaction (qRT-PCR) amplification of the bound DNA. This revealed a -0.68 and -0.64 log₂ fold decrease in FOXM1 occupancy at *CCNB1* and *CDC25B* gene promoters, respectively (Fig. 3D). The changes at specific genomic loci will not necessarily match the average global change quantitatively as there is a context-dependence for transcription factor binding^{43,44}. However, the magnitudes of changes observed here generally agree with those reported by the chromatin blotting experiments (Fig. 3B&C).

RNA-seq signature of FDI-6 associated with FOXM1 repression

As FOXM1 is a pure transcriptional activator^{45–47} we expect that its displacement from gene targets should lead to specific down-regulation of those gene transcripts. The ability of **FDI-6** to displace FOXM1 from two established genomic targets led us to investigate the global transcriptional effects of ligand treatment rigorously by genome-wide RNA sequencing (RNA-seq). We thus examined the effect of **FDI-6** treatment on RNA transcript levels in MCF-7 cells at 3, 6 and 9 h time points. The experiment was performed in triplicate at each of the three time-points and the untreated control. Fig. 4A shows the hierarchical clustering of gene expression profiles demonstrating that independent replicates at each time point group together and that their expression profiles show strong reproducibility. Furthermore, a substantial core set of 206 genes are found to be differentially expressed at each treatment time point, indicating agreement between samples (Fig. 4B, GEO: GSE58626 and differential expression is provided as Supplementary file). Fig. 4C shows a single replicate of differentially expressed genes after 3 h with up (red) and down (green) genes plotted by the logarithm of fold change (logFC, y-axis) and logarithm of counts per million reads (logCPM, x-axis).

We expected (i) specific down-regulation of FOXM1 gene targets, (ii) the activation of secondary and tertiary downstream responses to changes in the primary gene targets and (iii) non-FOXM1 specific, differentially expressed genes responding to the stress of ligand treatment. Table 1 presents the most significant ($p < 10^{-14}$) gene ontology terms for differentially expressed genes after ligand treatment. Importantly, cell cycle regulation and mitotic processes, of which FOXM1 is a known activator, are significantly down-regulated. After 6 h, *CCNB1* is decreased -0.86 (log₂ fold change, $P=1.9 \times 10^{-42}$) and *CDC25B* is decreased -0.36 ($P=4.3 \times 10^{-12}$). These values closely correlate with the fraction of FOXM1 that is displaced in ChIP-PCR (Fig. 3C) and also agree with down-regulation reported by separate qPCR validation runs after the same treatment (Supplementary Fig. 8). Many other established transcriptional targets of FOXM1²⁷ were also strongly down-regulated by **FDI-6** at each of the treatment time points. Measured at 6 h, these include *CCNA2* (-0.65 , $P=2.6 \times 10^{-27}$), *CDKN3* (-0.739 , $P=3.1 \times 10^{-13}$), *CENPA* (-0.61 , $P=8.3 \times 10^{-15}$), *KIF20A* (-1.22 , $P=1.1 \times 10^{-95}$), *NEK2* (-0.86 , $P=3.3 \times 10^{-16}$), *PLK1* (-0.92 , $P=3.2 \times 10^{-63}$), and *AURKA* (-0.83 , $P=1.4 \times 10^{-50}$) and compared in magnitude to changes seen following siRNA knockdown reported by Wonsey et al. (2005)¹⁶.

A FOXM1 inhibitor should exhibit this phenotype in any cell line expressing FOXM1. To rule out a cell-specific explanation for the response of ER-positive MCF-7 cells to **FDI-6**, we examined two additional patient-derived malignant adenocarcinoma cell lines commonly used in the literature: MDA-MB-231 ER-negative breast and PEO-1 ovarian cancer cells, which both highly express FOXM1^{39,48}. Both of these additional cell lines were sensitive to **FDI-6** in cell viability assays ($GI_{50} = 21.8 \mu\text{M}$ and $18.1 \mu\text{M}$, respectively) and exhibited comparable down-regulation of *CDC25B* after a 3 h treatment with the molecule (Fig. 5). This finding supports the proposed mechanism of action of **FDI-6** as an inhibitor of FOXM1 binding and suggests its utility across multiple cellular systems expressing the transcription factor.

FDI-6 effect is specific for genes with FOXM1 occupancy

A critical property of a *bona fide* FDI ligand, according to our model, would be the down-regulation of genes immediately downstream of FOXM1-occupied promoters. In contrast, genes that lack FOXM1 occupancy in the promoter should not be differentially expressed as a result of such a ligand, but may respond to downstream effects caused by changes in FOXM1-responsive genes. The identification of several established FOXM1 targets here supports **FDI-6** as a FOXM1 inhibitor. To further characterize the entire population of differentially expressed genes, we examined their correlation with FOXM1 promoter occupancy. Using the publicly available ENCODE FOXM1 ChIP-seq dataset in MCF-7 cells, we determined the abundance of verified FOXM1 protein binding sites within the up- and down-regulated sets of differentially expressed genes (Fig. 6A, top panel). This analysis revealed a highly significant ($P < 1 \times 10^{-16}$) 19-fold enrichment of FOXM1 binding sites exclusively in the down-regulated set. In contrast, the up-regulated differentially expressed genes were not significantly enriched ($P > 0.005$) for promoter FOXM1 binding sites over the background of genes not differentially expressed. Motif analysis of FOXM1 binding peaks reported the consensus sequence used to identify **FDI-6** originally (T/CAAACA, E-value = 2.5×10^{-62} , Supplementary Fig. 9), and ligand treatment specifically down-regulated these FOXM1 activated genes, providing strong evidence that the inhibitory effect of **FDI-6** on FOXM1 binding translates into cells.

We further applied this methodology to examine the specificity of **FDI-6** for FOXM1 over other Forkhead Box (FOX) proteins, which have highly homologous DBDs. We reanalyzed the RNAseq signature of **FDI-6** for enrichments in ChIP-seq binding peaks of all related forkhead factors for which available ENCODE datasets exist: FOXA1, FOXA2 and FOXP2 (Fig. 6A, middle three panels). Remarkably, despite the greater than 60% conservation among residues comprising the DBDs of FOX family transcription factors and the similarity of their consensus sequence motifs⁴⁹, genes differentially expressed after **FDI-6** treatment did not show enrichment over background for promoter occupancy of any other FOX transcription factors. This finding supports the specificity of the ligand for FOXM1 DBD over similar forkhead DBDs. Likewise, we show that there is no enrichment for sites of GATA1 occupancy (Fig. 6A, bottom panel). This latter comparison shows that the effect of the molecule is not associated with a general suppression of transcriptional activity, but is specific for the FOXM1 transcriptional program.

To further characterize the effects of **FDI-6** on the FOXM1 transcriptional program, we sorted sets of differentially expressed genes into groups based on their response profiles during the three treatment times. Our reasoning being that genes directly affected by FOXM1 occupancy would respond relatively quickly and consistently to displacement of FOXM1. This methodology, called time point cluster analysis, identified a set of 208 genes that are immediately down-regulated after **FDI-6** treatment (Cluster 5, Fig. 6B, cluster assignment of each gene is provided in Supplementary file). Inhibition of FOXM1-DNA should immediately down-regulate FOXM1-activated genes; thus, we expect that Cluster 5 should correlate best with promoter FOXM1 occupancy. Indeed, Cluster 5 is enriched 17-fold ($P = 5 \times 10^{-82}$) over the background for genes with promoter FOXM1 binding sites (Fig. 6C). Importantly, no other clusters exhibited significant enrichment ($P < 0.005$) of genes with

FOXM1 binding sites. Furthermore, each of the FOXM1 target genes discussed above is present within this cluster. This finding supports the notion that **FDI-6** induces a specific response of FOXM1 inhibition and that additional transcriptional responses represent downstream secondary effects of down-regulated primary FOXM1 targets or of non-specific ligand stress.

We remarked above that a number of the representative genes down-regulated by **FDI-6** were similarly decreased following siRNA knockdown of FOXM1¹⁶. We can explore this association across the entire transcriptome, as a highly specific compound should induce a similar differentially expressed gene signature to siRNA knockdown. Using an existing microarray profile of FOXM1 knockdown in BT-20 breast cancer cells (GEO accession: GSE2222) for comparison, we observe a significant ($P=6 \times 10^{-5}$) overlap with the **FDI-6** gene signature (Fig. 6D). This reveals a tendency for genes down-regulated by siRNA FOXM1 to be similarly down-regulated by the compound. In contrast, there is no significant ($P>0.05$) correlation between the genes up-regulated by the two treatments. Furthermore, the trend is even more pronounced within the subset of genes in Cluster 5, which are 3.7-fold ($P=2 \times 10^{-5}$) enriched for genes down-regulated by siRNA FOXM1. Lastly, it is a notable distinction that the small molecule acts directly on the DNA binding activity of FOXM1 and the phenotype arises after only 3 h treatment, in contrast to the 48 h incubation time needed to degrade FOXM1 protein by RNA interference. Taken together, these series of analyses show that **FDI-6** specifically inhibits FOXM1 association with consensus sequence DNA targets and prevents downstream transcriptional activation of these targets in cells, without inhibiting the related forkhead factors or perturbing FOXM1 protein level.

DISCUSSION

We have identified small molecule inhibitors that disrupt FOXM1-DNA interaction using an assay based on the fluorescence polarization response of a labeled consensus oligonucleotide bound to recombinant FOXM1 DBD protein. From a library of 54,211 structurally diverse ligands we identified six FDI compounds that were cross-validated in orthogonal binding assays. Notably, three of these compounds exhibit a common pharmacophore, suggesting a common structural mechanism. We further characterized the lead inhibitor NCGC00099374 (**FDI-6**) to show that it acts by targeting FOXM1 protein and broadly represses transcription of FOXM1-activated genes by displacing FOXM1 from chromatin. This activity is reflected throughout the transcriptome and is recapitulated by a gene ontology summary, which indicates that entire pathways known to be controlled by FOXM1 transactivation—mitosis, nuclear division, and cell cycle processes—are significantly inhibited by **FDI-6**. Taken together, our data support a direct mechanism of action whereby **FDI-6** blocks the binding of FOXM1 to target DNA, preventing the normal activation of FOXM1 target genes. We cannot absolutely rule out the possibility of a contribution from some unknown off-target effect(s) of FDI-6. However, it is a significant indication of the small molecule's specificity that despite the conserved DBD and a 71% overlap of ChIP-seq peaks for FOXM1 and FOXA1⁴⁴, genes repressed by **FDI-6** are exclusively targets of FOXM1 and not of FOXA1 or related forkhead factors.

An inhibitor with the ability to globally disrupt association of FOXM1 with DNA consensus sites in cells provides a valuable tool to probe FOXM1 occupancy. Previously, the ability to interrogate or intervene with the direct consequences of transcription factor binding has been limited. We previously reported that thiostrepton, a natural product antibiotic, binds directly to FOXM1 in cells and down-regulates target gene expression.³⁰ However, the use of thiostrepton is confounded by its inhibition of the 20S proteasome. Thiostrepton also induces significant toxicity ($GI_{50} = 2.7 \mu\text{M}$) well below the concentration needed to affect FOXM1 inhibition ($IC_{50} = 45.0 \pm 16.4 \mu\text{M}$). These peripheral activities of thiostrepton introduce some uncertainty about its dominant mechanism of action in cells and thus cannot be used to unambiguously probe FOXM1. In contrast, the novel inhibitor **FDI-6** is inactive against the proteasome and concordance between its biophysical IC_{50} and cellular GI_{50} values ($22.5 \pm 12.3 \mu\text{M}$ and $18.0 \pm 3.0 \mu\text{M}$, respectively) suggests it does not suffer from the off target effects of thiostrepton. The characteristics of smaller molecular size, greater aqueous solubility, and a scaffold amenable to future optimization are further advantages to the novel molecule.

We have also provided evidence for **FDI-6** mediated down-regulation of established FOXM1-controlled genes. By way of comparison, the same gene targets—*CDKN3* (−0.739, after 6 h **FDI-6**), *CENPA* (−0.61), *KIF20A* (−1.22), *NEK2* (−0.86), which all have strong FOXM1 binding sites in their promoters—were down-regulated by the same magnitude in a previous study employing qPCR after siRNA knockdown of FOXM1¹⁶. It is significant for its utility as a specific probe of FOXM1 occupancy that **FDI-6** stimulates a comparable transcriptional response after only 3 h and without ablating FOXM1 protein.

The genome-wide repression of the FOXM1 transcriptional program by **FDI-6** demonstrates a mechanistic proof-of-concept. At present, the application of FDI compounds to antagonize FOXM1-DNA provides a powerful means to probe the complex consequences of FOXM1 occupancy on transcription. Given that aberrant overexpression of FOXM1 protein is increasingly recognized as an initiating event in oncogenesis¹⁹, cancer progression⁵⁰ and therapeutic resistance^{21,22}, the capacity to inhibit FOXM1-DNA with small molecules has considerable potential which we will be exploring as part of future work.

METHODS

FOXM1 DBD protein preparation

FOXM1 DBD cDNA was cloned from the human full-length FOXM1c cDNA fragment (Origene Technologies). After PCR amplification with primers containing SgfI and MluI restriction sites, the products were gel purified and ligated in frame to the N terminus of GST into the linear pEX-N-GST vector (Origene Technologies) containing a T7 promoter, allowing for IPTG inducible expression of the fusion protein. Ligated plasmids were inserted into BL21/DE3 *E. Coli* cells by heat-induced transformation. Positive clones were selected by colony PCR screening and DNA sequencing confirmed successful insertion. Cells were grown at 37°C in LB media until reaching an optical density (OD) of 0.8 at which point, the temperature was decreased to 18°C and GST-FOXM1 DBD expression was induced by addition of 1 mM IPTG overnight. Culture conditions produced suitable amounts of protein (in excess of 2 mg L⁻¹ culture medium) that purified easily with no contaminant

or degradation product bands. Representative gel images of purified products are provided in Supplementary Fig. 10.

Fluorescence Polarization Assay

Setup runs of the fluorescence polarization (FP) assay were performed in 96-well plates and read on PHERAStar (BMG Labtech) with a 488/520 FP filter. The screening assay was scaled down successively to 384- and 1536-well plate formats and read on a ViewLux high-throughput charge-coupled device (CCD) imager (Perkin-Elmer, Wellesley, MA).

EMSA binding tests

Association of FOXM1 with 15 nM double stranded DNA consensus sequence oligo (forward strand: 5'-FAM- AAACAAACAAACAATC-3') was recorded by EMSA. Protein and dsDNA components were mixed in a 20 μ L reaction volume, incubated at RT for 30 min, and run directly on a 6% DNA retardation gel (Sigma) for 10 min at 40V and 20 min at 120V. The binding buffer consists of 20 mM Tris (pH 7.5), 100 mM KCl, 1 mM EDTA, 0.1 mM DTT, 10% vol/vol glycerol, 0.010 mg ml⁻¹ BSA. For displacement experiments, 2 μ L of 10X probe DNA was added to 18 μ L of FOXM1 DBD incubated with FDI compounds for 1.5 h ([FOXM1 DBD]_f = 750 nM, [DNA probe]_f = 15 nM). After 15 min samples were loaded directly and electrophoresis performed as above.

Cell Culture

MCF-7, MDA-MB-231, and PEO-1 cells were obtained from the European Collection of Animal Cell Cultures. MCF-7 and MDA-MB-231 cells were grown in Dulbecco's Modified Eagle Media (Sigma) supplemented with 10% fetal bovine serum (Sigma). PEO-1 cells were cultured in RPMI (Invitrogen) media also supplemented with 10% fetal bovine serum.

Antibodies

Rabbit polyclonal IgG FOXM1 antibody (C-20) was acquired from Santa Cruz Biotech (SC-502, lot# G2312). Actin was acquired from abcam (AC-15, lot GR91208-03, used 1:5000). Histone H3 antibody was acquired from Cell Signalling (4499S, lot 1, used 1:2000). LiCor Secondary antibodies were used for IR detection using goat-anti-mouse (680LT, lot C1109-02, used 1:20000) and goat-anti-rabbit (800CW, lot C30626-03, used 1:15000). Full gel images for each western blot are provided in Supplementary Fig. 11.

qRT-PCR primers

Quantitect primers (Quiagen) were used for qRT-PCR including ACTB (QT00095431), CCNB1 (QT00006615), CDC25B (QT00028350), and YWHAZ (QT00087962).

Native mass spectrometry

Mass spectra were recorded on a Synapt HDMS instrument (Waters UK Ltd., Manchester, UK). The concentration of FOXM1 was 4 μ mol dm⁻³ while the final concentration of compounds after adding to the protein was 50 μ mol dm⁻³. All measurements were carried out in a positive ion mode. Spraying conditions were: capillary voltage 1.7–1.8 kV, a cone voltage of 40–80 V, the source temperature 20°C, trap collision energy 12.0 V and transfer

collision energy 12.0 V with trap and transfer pressure 5.31×10^{-2} mbar, IMS pressure 5.02×10^{-1} mbar, TOF analyser pressure 1.2×10^{-6} mbar. External calibration of the spectra was achieved using caesium iodide at 100 mg ml^{-1} in water. Data acquisition and processing were performed using Micromass MassLynx 4.1.

20S Proteasome Inhibition

A 20S Proteasome Activity kit (Millipore) was used to determine compound effect on proteasome activity. The supplier provided standard protocol was followed. Briefly, an AMC peptide that is conditionally fluorescent upon proteasome cleavage was used to generate a standard curve to determine an appropriate concentration window for the assay. A 1:200 AMC concentration was determined to give suitable signal:noise response and used in the assay. Inhibitor (10 μM) or DMSO control, recombinant proteasome, and AMC substrate were mixed and incubated at 37°C for 90m. Fluorescence was then read on a Pherastar (BMG) using a 380/460 filter set. Inhibitory responses were determined normalized to DMSO control and compared to a known proteasome inhibitor MG-132.

Western blots

Western blot was carried out with total lysate and chromatin-isolated fractions from MCF-7 cells treated for 6 h with Compound 6 (40 μM) or DMSO control. For total lysate preparations, cells were grown and were harvested by scraping in PBS and pelleted by centrifugation. Cell pellets were lysed with 1 ml ice-cold $1 \times$ RIPA Buffer, with protease inhibitors (Roche). The lysate was agitated for 1 h at 4°C by end-to end rotation; the supernatant was collected following centrifugation and lysate concentration was measured using Pierce BCA Protein Assay kit following the manufacturer's protocol. Chromatin isolation protocol was adapted from Mendez et al.⁵¹ To isolate chromatin, cells were resuspended in Chromatin Isolation Buffer A (Table 6.1) and the cells were incubated for 5 min on ice. Nuclei were collected by low-speed centrifugation (4 m, 1,300 g at 4°C). Nuclei were washed once in Chromatin Isolation Buffer A, and then lysed in Chromatin Isolation Buffer B. Insoluble chromatin was collected by centrifugation (4 m, 1,700g at 4°C), washed once in Chromatin Isolation Buffer B, and centrifuged again under the same conditions. The final chromatin pellet (P3) was resuspended in Laemmli buffer and sonicated for 15 s in a Tekmar CV26 sonicator using a microtip at 25% amplitude. After harvesting, 20 μL of the recovered cell samples (total lysate or chromatin isolated fraction) were loaded onto 8–16% Tris-Glycine mini gels followed by electrophoresis at 150 V for 90 m. Rapid protein transfer was accomplished using the manufacture recommended 7 m protocol on the iBlot system (Invitrogen). The membrane was first washed in Odyssey block buffer (Li-Cor) three times (10 mL per wash). Primary actin antibody (total lysate control) or histone H3 antibody (chromatin isolation control) from Abcam were used at 1:5000 and 1:10000 respectively. The FOXM1 antibody used at 1:1000 dilution. Primary antibodies were incubated overnight at 4°C . Secondary antibodies were incubated at RT for 45 min. Goat anti-mouse IRDye-680 secondary antibody was used for actin and goat anti-rabbit IR Dye-800 for FOXM1. Blots were visualized on the Li-C or scanner and quantification of FOXM1 relative to β -actin (total lysate) or Histone H3 (chromatin isolation) was carried out for each lane.

Chromatin immunoprecipitation

MCF-7 cells were grown in 15 cm² dishes to 70% confluence prior to compound or DMSO control treatment for 6h. Cells were washed with PBS, cross-linked with 1% formaldehyde at RT for 10 min and quenched with 1/20 volume glycine (2.5 M). Cells were harvested by scraping into cold PBS containing protease inhibitors and cell pellets stored at -80 °C. Cell pellets were lysed sequentially with ChIP buffer I, buffer II and resuspended in buffer III (Table 5.2) containing protease inhibitors. Samples were sonicated on ice using a bioruptor UCD200 sonicator (Diagenode) for 15 min at high power with a pulse of 30 seconds on/30 seconds off. Samples then supplemented with Triton-X 100 to a final concentration of 1% and centrifuged at 20,000 × g for 10 min at 4°C. Protein-A dynabeads (Invitrogen) were washed with block solution (PBS plus 0.5% BSA) and incubated overnight with 3µg FOXM1 (Sc-502) antibody in block solution, followed by washing three times in block solution. Supernatants from cell lysates were incubated overnight with antibody capture beads, and then the beads were collected and washed 6 times with RIPA buffer II. Bound DNA was eluted by incubation at 65°C for 16 h in ChIP elution buffer and purified using a Qiaquick PCR clean-up kit. PCR was carried out using 2µl of purified DNA in a 20µl reaction with Power Sybr PCR mix (ABI), using the cycle conditions: 95 °C for 10 min followed by 40 cycles of 15 s at 95 °C, 30 s at 60 °C followed by generation of a dissociation curve. Primer sequences (5'-3') used are as follows: AAGAGCCCATCAGTTCCGCTTG (CDC25B promoter fwd), CCCATTTTACAGACCTGGACGC(CDC25B promoter rev), CGCGATCGCCCTGGAAACGCA (CCNB1 promoter fwd), CCCAGCAGAAACCAACAGCCGT (CCNB1 promoter rev), TGCCACACACCAGTGACTTT (CCND1 intergenic control fwd), ACAGCCAGAAGCTCCAAAAA (CCND1 intergenic control rev). Enrichment values were calculated in Excel by normalizing against Cyclin D1 intergenic control.

RNA Sequencing and analysis

Samples were processed following the standard TrueSeq (Illumina) protocol. Fastq files containing 100 single-end reads were processed with trim-galore (http://www.bioinformatics.babraham.ac.uk/projects/trim_galore/) to perform adapter trimming and low-quality reads filtering. The filtered reads were then aligned to the human genome (hg19) with tophat2⁵², and transcripts were assigned to gene and counted using htseq-count (<http://www.huber.embl.de/users/anders/HTSeq/doc/overview.html>). See Supplementary Table 1&2 for summary statistics.

Differential expression analysis

Differential expression analysis was performed using edgeR⁵³. The three replicates belonging to the different time points (3, 6, and 9 h) were tested for differential expression versus the control condition (0 h treatment), by using the exact test for differences between two groups of negative binomial counts (function exact Test in the R package 'edgeR'). This method computes the genewise difference of counts taking into account variability across replicates. By default, the method normalizes for the total library size by calculating counts per million of reads (CPM). Genes with a false discovery rate (FDR) below 0.01 were

considered significant for differential expression (Fig. 4C). For the overlap among differentially expressed genes shown in the Venn diagram (Fig. 4B), we considered only genes with $FDR < 0.01$ and absolute logarithm of fold change ($\log FC$) > 1 , i.e. genes up-regulated at least 2 folds or down-regulated at least 50%

Temporal clustering

The clustering of temporal profiles of gene expression was done using the R package ‘MFuzz’⁵⁴, which performs soft clustering of genes based on their expression values using the fuzzy c-means algorithm. We first selected only the 1552 genes having a substantial change in either one of the time points (i.e., $abs(\log FC) > 0.75$); we then run the fuzzy c-means algorithm with parameters 9 and 1.35, respectively the number of clusters and the fuzzification parameter. The temporal profiles used for clustering and shown in Fig. 6B are standardized such that the average expression value for each gene in the different time points is 0 and the standard deviation is 1. In the supporting .xls file “SI_diffexp_all” we report the assignment of each one of the 12873 genes to a temporal pattern, by calculating the maximal similarity (Pearson correlation coefficient) to the average profile of each of the 9 clusters previously identified.

Promoter binding analysis

We collected all peak files available at the website of the ENCODE project⁵⁵ (www.encodeproject.org) for all members of the FOX (Forkhead box) proteins plus GATA1 as an out-group. Peak files were calculated by the consortium using the peak caller MACS⁵⁶ on ChIPseq experiments done in duplicates, under standardized conditions. For the promoter binding analysis, 1/we took only peaks in common between the two duplicates by calculating the intersection of peaks with bedtools⁵⁷; 2/for each gene, we identified the promoter region as the genomic region 2000 bases upstream and 100 bases downstream of the transcription start site (tss); 3/we counted peaks inside the promoter regions; 4/we calculated the proportion of genes having at least one peak in the promoter region, dividing the genes in down-regulated (*down*), up-regulated (*up*) and not differentially expressed (*not de*); 5/we divided the three proportions so obtained by the respective value of the *not de* group for each data set separately; 6/the normalized proportion values of all data sets relative to the same transcription factor were averaged. For all the above steps, we considered as differentially expressed genes with $FDR < 0.01$ and $\log FC > 0.3$ (or < 0.3) at 3 h. P s determined using the non-parametric chi-squared test for proportions (function `prop.test`) implemented in the R software (<http://www.r-project.org>).

FOXM1 knockdown comparison

The categories “up_drug” and “down_drug” refer to genes up- ($\log FC > 0.5$) and down-regulated ($\log FC < -0.5$) after 3 hours treatment with FDI-6, as measured by RNAseq. The categories “up_kd” and “down_kd” refer to genes up- ($\log FC > 0.5$) and down-regulated ($\log FC < -0.5$) upon FOXM1 knockdown, as measured by Affymetrix expression array (<http://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE2222>). “cluster5_down” and “cluster5_up” refers to genes down-regulated ($\log FC < -0.5$) belonging to cluster 5 from the

temporal clustering analysis. *P*s determined using the non-parametric chi-squared test for proportions (function prop.test) implemented in the R software (<http://www.r-project.org>).

Motif analysis

Fasta sequences from the genomic intervals corresponding to peaks were analyzed by using the MEME-CHIP tool⁵⁸ from the MEME Suite for motif-based sequence analysis (<http://meme.nbcr.net/>), and enriched motifs were identified (Supplementary Fig. 9).

General chemistry experimental methods

Preparative purification was run on a Waters semi-preparative HPLC system using a Phenomenex Luna C18 (5 micron, 30 × 75 mm) at a flow rate of 45 mL min⁻¹. A gradient of 10% to 50% acetonitrile in water over 8 m (each containing 0.1% trifluoroacetic acid) was used as a mobile phase during the purification. Fraction collection was triggered by UV detection (220 nm). Analytical analysis was performed on an Agilent LC/MS (Agilent Technologies, Santa Clara, CA). Method 1: A 7 m gradient of 4% to 100% Acetonitrile (containing 0.025% trifluoroacetic acid) in water (containing 0.05% trifluoroacetic acid) was used with an 8 m run time at a flow rate of 1 mL min⁻¹. A Phenomenex Luna C18 column (3 micron, 3 × 75 mm) was used at a temperature of 50° C. Method 2: A 3 m gradient of 4% to 100% Acetonitrile (containing 0.025% trifluoroacetic acid) in water (containing 0.05% trifluoroacetic acid) was used with a 4.5 m run time at a flow rate of 1 mL min⁻¹. A Phenomenex Gemini Phenyl column (3 micron, 3 × 100 mm) was used at a temperature of 50° C. Purity was determined using an Agilent Diode Array Detector for both Method 1 and Method 2. Mass determination was performed using an Agilent 6130 mass spectrometer with electrospray ionization in the positive mode. ¹H NMR spectra were recorded on Varian 400 MHz spectrometer. Chemical shifts are reported in ppm with undeuterated DMSO-*d*₆ at 2.49 ppm as internal standard. High resolution mass spectrometry was recorded on Agilent 6210 Time-of-Flight LC/MS system. Confirmation of molecular formula was accomplished using electrospray ionization in the positive mode with the Agilent Masshunter software (version B.02).

Synthesis of 3-amino-*N*-(4-fluorophenyl)-6-(thiophen-2-yl)-4-(trifluoromethyl)thieno[2,3-*b*]pyridine-2-carboxamide TFA (FDI-6)

A mixture of 6-(thiophen-2-yl)-2-thioxo-4-(trifluoromethyl)-1,2-dihydropyridine-3-carbonitrile⁵⁹ (0.2 g, 0.699 mmol, 1 eq) and 2-chloro-*N*-(4-fluorophenyl)acetamide (0.138 g, 0.734 mmol, 1.05 eq) in ethanol (4 mL) was added potassium carbonate (0.241 g, 1.746 mmol). The reaction mixture was refluxed for 4 h then poured into ice water. The precipitate was collected by filtration and washed with water. The crude solid was taken up in 3 mL DMSO and purified on a preparative HPLC to obtain pure 3-amino-*N*-(4-fluorophenyl)-6-(thiophen-2-yl)-4-(trifluoromethyl)thieno[2,3-*b*]pyridine-2-carboxamide (**FDI-6**) as a TFA salt. LC-MS Retention Time: (method 1) = 7.081 min and (method 2) = 3.983 min; ¹H NMR (400 MHz, DMSO-*d*₆) δ 9.80 (s, 1H), 8.30 (s, 1H), 8.22 (dd, *J* = 3.8, 1.1 Hz, 1H), 7.85 (dd, *J* = 5.0, 1.1 Hz, 1H), 7.73–7.63 (m, 2H), 7.27 (dd, *J* = 5.0, 3.8 Hz, 1H), 7.24–7.14 (m, 2H), 6.77 (s, 2H); (101 MHz, DMSO-*d*₆) δ 163.5, 160.5, 159.8, 157.4, 152.3, 145.1, 142.0, 134.7,

134.7, 131.7, 131.2, 129.1, 129.1, 123.7, 123.6, 117.8, 115.2, 114.9, 112.6, 100.7; HRMS (ESI) m/z (M+H)⁺ calcd. for C₁₉H₁₂F₄N₃OS₂, 438.0352; found 438.367.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We thank Dr. Neil M. Bell for advice and assistance in protein purification. M.V.G. is supported by a Churchill Scholarship and an NIH OxCam Fellowship. G.M. is supported by a Wellcome Trust Investigator Award. S.B. is supported by a Wellcome Trust Senior Investigator Award. The authors gratefully acknowledge Cancer Research UK core funding. T.S.D., M.G., G.R., A.S., A.J. and D.J.M. were supported by the intramural research program of the National Center for Advancing Translational Sciences and the Molecular Libraries Initiative of the National Institutes of Health Common Fund (grant U54MH084681)

References

1. Darnell JE. Transcription factors as targets for cancer therapy. *Nat Rev Cancer*. 2002; 2:740–749. [PubMed: 12360277]
2. Libermann TA, Zerbini LF. Targeting transcription factors for cancer gene therapy. *Curr Gene Ther*. 2006; 6:17–33. [PubMed: 16475943]
3. Laoukili J, Stahl M, Medema RH. FoxM1: At the crossroads of ageing and cancer. *Biochimica et Biophysica Acta (BBA)- Reviews on Cancer*. 2007; 1775:92–102. [PubMed: 17014965]
4. Vaquerizas JM, Kummerfeld SK, Teichmann SA, Luscombe NM. A census of human transcription factors: function, expression and evolution. *Nature Reviews Genetics*. 2009; 10:252–263.
5. Spitz F, Furlong EEM. Transcription factors: from enhancer binding to developmental control. *Nature Reviews Genetics*. 2012; 13:613–626.
6. Bell D, et al. Integrated genomic analyses of ovarian carcinoma. *Nature*. 2011; 474:609–615. [PubMed: 21720365]
7. Szabova L, et al. Perturbation of Rb, p53, and Brca1 or Brca2 Cooperate in Inducing Metastatic Serous Epithelial Ovarian Cancer. *Cancer Research*. 2012; 72:4141–4153. [PubMed: 22617326]
8. Zhou J, et al. FOXM1 Modulates Cisplatin Sensitivity by Regulating EXO1 in Ovarian Cancer. *PLoS ONE*. 2014; 9:e96989. [PubMed: 24824601]
9. Koehler AN, Shamji AF, Schreiber SL. Discovery of an Inhibitor of a Transcription Factor Using Small Molecule Microarrays and Diversity-Oriented Synthesis. *J Am Chem Soc*. 2003; 125:8420–8421. [PubMed: 12848532]
10. Berg T. Inhibition of transcription factors with small organic molecules. *Current Opinion in Chemical Biology*. 2008; 12:464–471. [PubMed: 18706517]
11. Koehler AN. A complex task? Direct modulation of transcription factors with small molecules. *Current Opinion in Chemical Biology*. 2010; 14:331–340. [PubMed: 20395165]
12. Verdine GL, Walensky LD. The Challenge of Drugging Undruggable Targets in Cancer: Lessons Learned from Targeting BCL-2 Family Members. *Clinical Cancer Research*. 2007; 13:7264–7270. [PubMed: 18094406]
13. Dervan PB. Molecular recognition of DNA by small molecules. *Bioorg Med Chem*. 2001; 9:2215–2235. [PubMed: 11553460]
14. Walensky LD. Activation of Apoptosis in Vivo by a Hydrocarbon-Stapled BH3 Helix. *Science*. 2004; 305:1466–1470. [PubMed: 15353804]
15. Pop MS, et al. A Small Molecule That Binds and Inhibits the ETV1 Transcription Factor Oncoprotein. *Molecular Cancer Therapeutics*. 2014; 13:1492–1502. [PubMed: 24737027]
16. Wonsey DR, Follettie MT. Loss of the forkhead transcription factor FoxM1 causes centrosome amplification and mitotic catastrophe. *Cancer Research*. 2005; 65:5181–5189. [PubMed: 15958562]

17. Laoukili J, et al. FoxM1 is required for execution of the mitotic programme and chromosome stability. *Nat Cell Biol.* 2005; 7:126–136. [PubMed: 15654331]
18. McDonald D. FoxM1 dances with mitosis. *Nat Cell Biol.* 2005; 7:108–110. [PubMed: 15689977]
19. Teh, MT. *Stem Cells and Cancer Stem Cells: Therapeutic Applications in Disease and Injury.* Hayat, MA., editor. Vol. 3. Springer Verlag; 2011.
20. Koo CY, Muir KW, Lam EWF. FOXM1: From cancer initiation to progression and treatment. *BBA- Gene Regulatory Mechanisms.* 2012; 1819:28–37. [PubMed: 21978825]
21. Kwok JMM, et al. FOXM1 Confers Acquired Cisplatin Resistance in Breast Cancer Cells. *Molecular Cancer Research.* 2010; 8:24–34. [PubMed: 20068070]
22. Carr JR, Park HJ, Wang Z, Kiefer MM, Raychaudhuri P. FoxM1 Mediates Resistance to Herceptin and Paclitaxel. *Cancer Research.* 2010; 70:5054–5063. [PubMed: 20530690]
23. Carter SL, Eklund AC, Kohane IS, Harris LN, Szallasi Z. A signature of chromosomal instability inferred from gene expression profiles predicts clinical outcome in multiple human cancers. *Nat Genet.* 2006; 38:1043–1048. [PubMed: 16921376]
24. Priller M, et al. Expression of FoxM1 Is Required for the Proliferation of Medulloblastoma Cells and Indicates Worse Survival of Patients. *Clinical Cancer Research.* 2011; 17:6791–6801. [PubMed: 21918172]
25. Teh MT. Cells brainwashed by FOXM1: do they have potential as biomarkers of cancer? *Biomarkers Med.* 2012; 6:499–501.
26. Littler DR, et al. Structure of the FoxM1 DNA-recognition domain bound to a promoter sequence. *Nucleic Acids Research.* 2010; 38:4527–4538. [PubMed: 20360045]
27. Wang Z, et al. Forkhead box M1 transcription factor: A novel target for cancer therapy. *Cancer Treatment Reviews.* 2010; 36:151–156. [PubMed: 20022709]
28. Radhakrishnan SK, et al. Identification of a Chemical Inhibitor of the Oncogenic Transcription Factor Forkhead Box M1. *Cancer Research.* 2006; 66:9731–9735. [PubMed: 17018632]
29. Kwok JMM, et al. Thiostrepton selectively targets breast cancer cells through inhibition of forkhead box M1 expression. *Molecular Cancer Therapeutics.* 2008; 7:2022–2032. [PubMed: 18645012]
30. Hegde NS, Sanders DA, Rodriguez R, Balasubramanian S. The transcription factor FOXM1 is a cellular target of the natural product thiostrepton. *Nature Chemistry.* 2011; 3:725–731.
31. Halasi M, Gartel AL. A novel mode of FoxM1 regulation: positive auto-regulatory loop. *Cell Cycle.* 2009; 8:1966–1967. [PubMed: 19411834]
32. Zhang L, et al. Antibiotic susceptibility of mammalian mitochondrial translation. *FEBS Letters.* 2005; 579:6423–6427. [PubMed: 16271719]
33. Gartel AL. Thiostrepton, proteasome inhibitors and FOXM1. *Cell Cycle.* 2011; 10:4341–4342. [PubMed: 22134246]
34. Jameson DM, Ross JA. Fluorescence Polarization/Anisotropy in Diagnostics and Imaging. *Chem Rev.* 2010; 110:2685–2708. [PubMed: 20232898]
35. Simeonov A, et al. Fluorescence Spectroscopic Profiling of Compound Libraries. *J Med Chem.* 2008; 51:2363–2371. [PubMed: 18363325]
36. Huang R, et al. The NCGC Pharmaceutical Collection: A Comprehensive Resource of Clinically Approved Drugs Enabling Repurposing and Chemical Genomics. *Science Translational Medicine.* 2011; 3:80ps16–80ps16.
37. Jadhav A, et al. Quantitative Analyses of Aggregation, Autofluorescence, and Reactivity Artifacts in a Screen for Inhibitors of a Thiol Protease. *J Med Chem.* 2010; 53:37–51. [PubMed: 19908840]
38. Parker MW, Bello, Lo M, Federici G. Crystallization of glutathione S-transferase from human placenta. *J Mol Biol.* 1990; 213:221–222. [PubMed: 2342105]
39. Reinhold WC, et al. CellMiner: A Web-Based Suite of Genomic and Pharmacologic Tools to Explore Transcript and Drug Patterns in the NCI-60 Cell Line Set. *Cancer Research.* 2012; 72:3499–3511. [PubMed: 22802077]
40. Bhat UG, Halasi M, Gartel AL. FoxM1 Is a General Target for Proteasome Inhibitors. *PLoS ONE.* 2009; 4:e6593. [PubMed: 19672316]

41. Nakamura S, et al. The FOXM1 transcriptional factor promotes the proliferation of leukemia cells through modulation of cell cycle progression in acute myeloid leukemia. *Carcinogenesis*. 2010; 31:2012–2021. [PubMed: 20823107]
42. Leung TW, et al. Over-expression of FoxM1 stimulates cyclin B1 expression. *FEBS Letters*. 2001; 507:59–66. [PubMed: 11682060]
43. Chen X, et al. The Forkhead Transcription Factor FOXM1 Controls Cell Cycle-Dependent Gene Expression through an Atypical Chromatin Binding Mechanism. *Mol Cell Biol*. 2012; 33:227–236. [PubMed: 23109430]
44. Sanders DA, Ross-Innes CS, Beraldi D, Carroll JS, Balasubramanian S. Genome-wide mapping of FOXM1 binding reveals co binding with estrogen receptor alpha in breast cancer cells. *Genome Biol*. 2013; 14:R6. [PubMed: 23347430]
45. Wierstra I, Alves J. Despite its strong transactivation domain, transcription factor FOXM1c is kept almost inactive by two different inhibitory domains. *Biol Chem*. 2006; 387:963–976. [PubMed: 16913846]
46. Wierstra I, Alves J. FOXM1c transactivates the human c-myc promoter directly via the two TATA boxes P1 and P2. *FEBS Journal*. 2006; 273:4645–4667. [PubMed: 16965535]
47. Wierstra I, Alves J. FOXM1, a typical proliferation-associated transcription factor. *Biological Chemistry*. 2007; 388:1257–1274. [PubMed: 18020943]
48. Wiedemeyer WR. Reversing platinum resistance in high-grade serous ovarian carcinoma: targeting BRCA and the homologous recombination system. 2014:1–11.10.3389/fonc.2014.00034/abstract
49. Myatt SS, Lam EWF. The emerging roles of forkhead box (Fox) proteins in cancer. *Nat Rev Cancer*. 2007; 7:847–859. [PubMed: 17943136]
50. Balli D, et al. Foxm1 transcription factor is required for lung fibrosis and epithelial-to-mesenchymal transition. *The EMBO Journal*. 2013; 32:231–244. [PubMed: 23288041]
51. Méndez J, Stillman B. Chromatin association of human origin recognition complex, cdc6, and minichromosome maintenance proteins during the cell cycle: assembly of prereplication complexes in late mitosis. *Mol Cell Biol*. 2000; 20:8602–8612. [PubMed: 11046155]
52. Kim D, et al. TopHat2: accurate alignment of transcriptomes in the presence of insertions, deletions and gene fusions. *Genome Biol*. 2013; 14:R36. [PubMed: 23618408]
53. Robinson MD, McCarthy DJ, Smyth GK. edgeR: a Bioconductor package for differential expression analysis of digital gene expression data. *Bioinformatics*. 2009; 26:139–140. [PubMed: 19910308]
54. Kumar L, Futschik EM. Mfuzz: a software package for soft clustering of microarray data. *Bioinformatics*. 2007; 23:5–7. [PubMed: 18084642]
55. Consortium EP. The ENCODE (ENCyclopedia of DNA elements) project. *Science*. 2004; 306:636–640. [PubMed: 15499007]
56. Feng J, Liu T, Qin B, Zhang Y, Liu XS. Identifying ChIP-seq enrichment using MACS. *Nat Protoc*. 2012; 7:1728–1740. [PubMed: 22936215]
57. Quinlan AR, Hall IM. BEDTools: a flexible suite of utilities for comparing genomic features. *Bioinformatics*. 2010; 26:841–842. [PubMed: 20110278]
58. Machanick P, Bailey TL. MEME-ChIP: motif analysis of large DNA datasets. *Bioinformatics*. 2011; 27:1696–1697. [PubMed: 21486936]
59. Abdel-Monem MI, Mohamed OS, Bakhite EA. Fluorine-containing heterocycles: synthesis and some reactions of new 3-amino-2-functionalized-6-(2'-thienyl)-4-trifluoromethylthieno [2,3-b]pyridines. *Pharmazie*. 2001; 56:41–44. [PubMed: 11210666]




Figure 1. Design of FP assay to screen for inhibitors of FOXM1 DNA binding
(a) Schematic of FP assay principle. Polarized light excites the fluorophore labeled DNA consensus sequence probe. The proportion of polarized light recovered upon emission is related to the total mass of the probe, thus binding of probe and FOXM1 DBD can be monitored. **(b)** FP reports binding of FOXM1 DBD with tagged consensus DNA. **(c)** Unlabeled consensus DNA efficiently displaced tagged DNA in a dose dependent manner. A large signal window (>100 mP) between bound and unbound states indicated a robust assay. **(d)** Z' score for each plate in screen. Aggregate Z' score was 0.71, indicating a highly robust assay. **(e)** Waterfall plot showing dose-response curves for 170 active and 907 inconclusive hits in blue and black, respectively. Data are reported as average of replicates (n=3) and error bars indicate s.d. throughout the figure.




Figure 2. EMSA provides orthogonal biophysical validation of 16 lead compounds from FP screen

(a) Representative EMSA image shows the association of FOXM1 DBD with its fluorescently tagged DNA consensus. Binding curve for the FOXM1 DBD-DNA interaction was determined by quantification of EMSA replicates. (b) Quantification of EMSA bands confirmed six of the 20 lead compounds selected from the FP results inhibited FOXM1 DNA binding. In rank order of potencies: **FDI-6** > **FDI-10** > **FDI-11** > **FDI-4** > **FDI-2** > **FDI-7**. Representative gel images are provided in Supplementary concentrations in MCF-7 cells. (c) Representative Fig. 5. Each inset lists the IC_{50} as well as 72 h GI_{50} EMSA showing the inhibitory effect of **FDI-6** on the FOXM1 DBD-DNA complex. (d) Structures of **FDI-6**, **FDI-10**, and **FDI-11**, the most potent of the validated hit compounds, which all contain a common core structural element (highlighted in blue). Data are reported as average of replicates ($n=3$) and error bars indicate s.d. throughout figure.




Figure 3. Compound FDI-6 shows promising FOXM1-specific inhibitory effects in MCF-7 cells (a) FDI compounds were investigated for direct interaction with FOXM1 protein by nondenaturing nanoESI MS. EMSA validated hits **FDI-6** (yellow stars), **FDI-10** (green stars), and **FDI-11** (purple stars) associate with the protein in a 1:1 stoichiometric ratio. Negative control FDI-9 was identified as a false positive by EMSA and does not associate with the protein. Mass shifts indicating ligand association are indicated by red arrows. (b) Western blots on the whole cell and chromatin fractions revealed a specific displacement of DNA bound FOXM1 while total levels were unaffected. Total MCF-7 cell lysate and chromatin bound fractions were isolated after 6 h treatment with **FDI-6** or DMSO control. (c) Quantification of western blots revealed displacement of FOXM1 from DNA. Data were normalized to Actin (total lysate) and Histone H3 (chromatin fraction). (d) Targeted ChIP PCR revealed 6 h treatment with **FDI-6** decreased FOXM1 occupancy at known FOXM1 target genes. Refer to legend for c-d. Data are reported as average of replicates (n=3) and error bars indicate s.d. throughout the figure. In each case the statistical significance was determined by Student's t-test (*=P<0.05, **=P<0.01, ***=P<0.001).




Figure 4. Genome wide RNA-seq shows that FDI-6 treatment selectively down-regulates transcription of FOXM1 target genes

(a) Hierarchical clustering of gene expression profiles, showing the similarity of different treatment times. See Supplementary Table S2 for details of Euclidian distance between paired libraries. **(b)** Venn diagrams showing the overlap of differentially expressed genes at the different time points. **(c)** Global differential expression map of RNA-seq after 3 h treatment versus untreated. Y-axis: logarithm of fold change (logFC); X-axis: logarithm of counts per million of reads (logCPM). Red dots: up-regulated genes (n=1951); green dots: down-regulated genes (n=1552).




Figure 5. FDI-6 retains activity in MDA-MB-231 and PEO-1 adenocarcinoma cell lines
(a) 72 h treatment with FDI-6 shows dose dependent effect on cell growth in MDA-MB-231 and PEO-1 breast and ovarian cancer cell lines. **(b)** FDI-6 down-regulates CDC25B expression after 3 h treatment.




Figure 6. Differential expression signature of FDI-6 is associated with FOXM1 promoter occupancy and siRNA FOXM1
(a) Barplots showing the percentage of differentially expressed gene sets with a transcription factor occupancy in the promoter region (2 kb upstream of tss). Plots are shown for the closely related forkhead transcription factors FOXM1, FOXA1, FOXA2, FOXP2 as well as the transcriptional activator GATA1. For each plot, from left to right, bars represent down-regulated (down), up-regulated (up), and not differentially expressed (not de) gene sets. **(b)** Temporal cluster analysis grouping genes that show similar changes in expression after **FDI-6** treatment. **(c)** Barplot showing the percentage of genes in the different temporal clusters with a FOXM1 peak in the promoter region. **(d)** Barplot showing the fraction of genes that were up- and down-regulated by FDI-6 and were commonly differentially

expressed in an existing microarray dataset of FOXM1 siRNA knockdown. Error bars indicate s.e.m.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 1

Gene ontology signature of FDI-6

Significantly ($P < 10^{-14}$) over represented ontology terms for differentially expressed genes after 3 h FDI-6 treatment

DOWNREGULATED			UPREGULATED		
Gene Ontology Term	P-value	FDR	Gene Ontology Term	P-value	FDR
GO:0000278~mitotic cell cycle	1E-24	2E-21	GO:0031328~positive regulation of cellular biosynthetic process	2E-18	4E-15
GO:0000280~nuclear division	7E-23	1E-19	GO:0051173~positive regulation of nitrogen compound metabolic process	9E-18	2E-14
GO:0048285~organelle fission	2E-22	3E-19	GO:0010033~response to organic substance	2E-17	4E-14
GO:0022402~cell cycle process	1E-21	2E-18	GO:0010628~positive regulation of gene expression	2E-16	4E-13
GO:0007049~cell cycle	1E-19	2E-16	GO:0045935~positive regulation of nucleic acid metabolic process	1E-15	2E-12
GO:0015630~microtubule cytoskeleton	2E-17	2E-14	GO:0045941~positive regulation of transcription	1E-15	2E-12
GO:0005819~spindle	1E-16	1E-13	GO:0051254~positive regulation of RNA metabolic process	2E-14	4E-11
GO:0051301~cell division	4E-15	6E-12	GO:0042981~regulation of apoptosis	7E-14	1E-10