



Access this article online

Quick Response Code:



Website:

www.turkjemerged.com

DOI:

10.4103/tjem.tjem\_344\_21

# Comparison of pain and extent of anesthesia in digital blocks for isolated finger lacerations: A randomized controlled trial

Ali Jarragh\*, Ali Lari<sup>1</sup>, Waleed Burhamah<sup>2</sup>, Mohammed Alherz<sup>3</sup>, Abdullah Nouri<sup>1</sup>, Yahia Alshammari<sup>1</sup>, Ameer Al-Jasim<sup>4</sup>, Sulaiman AlRefai<sup>1</sup>, Naser Alnusif<sup>1</sup>

Department of Surgery, Kuwait University, <sup>1</sup>Department of Orthopedic Surgery, AlRazi Hospital, <sup>2</sup>Department of Plastic Surgery, AlBabtain Hospital, Al-Shuwaikh, Kuwait, <sup>3</sup>Department of Anatomy, Trinity College Dublin, Dublin, Ireland, <sup>4</sup>Department of Surgery, College of Medicine, University of Baghdad, Baghdad, Iraq  
\*Corresponding author

Submitted: 02-12-2021

Revised 1: 15-01-2022

Revised 2: 07-03-2022

Accepted: 08-03-2022

Published: 01-07-2022

ORCID:

AJ: 0000-0001-5392-2488

AL: 0000-0002-2070-6727

WB: 0000-0001-7707-2043

MA: 0000-0001-7641-9746

AN: 0000-0003-4251-5451

YA: 0000-0003-3507-1418

AA: 0000-0003-4125-9089

SA: 0000-0001-5348-9389

NA: 0000-0002-6801-9205

Address for  
correspondence:

Dr. Ali Jarragh,  
Department of Surgery,  
Kuwait University, Kuwait.  
E-mail: Ali.Jarragh@  
Ku.edu.kw

## Abstract:

**OBJECTIVES:** Digital injuries are among the most common presentations to the emergency department. In order to sufficiently examine and manage these injuries, adequate, prompt, and predictable anesthesia is essential. In this trial, we aim to primarily compare the degree of pain and anesthesia onset time between the two-injection dorsal block technique (TD) and the single-injection volar subcutaneous block (SV) technique. Further, we describe the temporal and anatomical effects of both techniques for an accurate delineation of the anesthetized regions.

**METHODS:** This is a single-center prospective randomized controlled trial involving patients presenting with isolated wounds to the fingers requiring primary repair under local anesthesia. Patients were randomized to either the SV or TD blocks. The primary outcome was procedure-related pain (Numerical Rating Scale). Further, we assessed the extent of anesthesia along with the anesthesia onset time.

**RESULTS:** A total of 100 patients were included in the final analysis, 50 on each arm of the study. The median pain score during injection was significantly higher in patients who received TD block than patients who received SV block (median [interquartile range] = 4 [2.25, 5.00] vs. 3.00 [2.00, 4.00], respectively,  $P=0.006$ ). However, anesthesia onset time was not statistically different among the groups ( $P=0.39$ ). The extent of anesthesia was more predictable in the dorsal block compared to the volar block.

**CONCLUSION:** The single-injection volar subcutaneous blocks are less painful with a similar anesthesia onset time. Injuries presenting in the proximal dorsal region may benefit from the two-injection dorsal blocks, given the anatomical differences and timely anesthesia of the region.

## Keywords:

Anesthetic techniques, digital nerve block, finger laceration, hand trauma, local anesthesia, nerve block

## Introduction

Digital injuries are among the most common presentations to the emergency department.<sup>[1]</sup> In order to sufficiently examine and manage these

injuries, adequate analgesia is commonly achieved either through local infiltration at the site of injury with a local anesthetic or by performing a digital nerve block. The latter represents the preferred technique, producing a less painful and a more predictable result.<sup>[2,3]</sup> Common among the various described digital nerve blocks,<sup>[4-7]</sup> are

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: WKHLRPMedknow\_reprints@wolterskluwer.com

**How to cite this article:** Jarragh A, Lari A, Burhamah W, Alherz M, Nouri A, Alshammari Y, et al. Comparison of pain and extent of anesthesia in digital blocks for isolated finger lacerations: A randomized controlled trial. Turk J Emerg Med 2022;22:125-30.

**Box-ED Section****What is already known on this study topic?**

- Finger lacerations are one of the most common presentations to the emergency department.
- Various digital block techniques are utilized in order to provide sufficient anesthesia for repair.

**How is this study structured?**

- This was a single-center, prospective randomized controlled trial that included data from 100 patients.
- The study compared two of the most common anesthetic techniques; single-injection volar subcutaneous block and the two-injection dorsal block.

**What does this study tell us?**

- The Single-injection volar subcutaneous blocks are less painful and provide a similar anesthesia onset time.
- Injuries to the dorsal region may benefit from the two-injection dorsal blocks, especially in the proximal dorsal region of the digit.

the traditional two-injection dorsal block technique (TD)<sup>[6]</sup> and the single-injection volar subcutaneous block (SV) technique.<sup>[7]</sup>

The single-injection volar subcutaneous block (SV) technique was first described by Harrison in 1991, requiring one subcutaneous injection at the volar side of the finger base.<sup>[7]</sup> Braun and Harris described the TD technique, whereby a local anesthetic is injected at each side of the finger base. For this technique, two separate dorsal injections are administered, one in each web space.<sup>[6]</sup>

We hypothesize that the SV technique is less painful than the TD technique. In this trial, we aim to primarily compare the degree of procedure-related pain and anesthesia onset time between the two block techniques. Further, we describe the temporal and anatomical effects of both techniques for an accurate delineation of the anesthetized regions.

## Methods

### Study design

This was a single-center, prospective randomized controlled trial. Informed consent was obtained from eligible patients prior to their inclusion in the study. The study was conducted in accordance with the 1964 Helsinki Declaration, approved by the Kuwaiti Ethical Committee board on the 6<sup>th</sup> of January, 2021 (UID 1742/2021), and registered on Research Registry (UIN: 6693). This study adheres to the appropriate CONSORT guidelines.

### Study population

One hundred ( $n = 100$ ) eligible participants over 18 years of age who presented with an isolated laceration to a single finger requiring primary repair under local anesthesia were included. The exclusion criteria included known allergies to local anesthesia, the presence of distracting concomitant injuries, multiple digital injuries, peripheral neuropathy, open fractures, finger infections, and tendon injuries. Injuries to the proximal dorsal aspect of the finger were excluded, as well as injuries to the thumb. The three volar anatomical regions of the finger were divided according to the proximal, middle, and distal joint creases, with the dorsal divisions corresponding to the volar. Further, patients with fractures, dislocations, and traumatic foreign bodies were also excluded. Finally, the included patients did not receive any form of intravenous or oral analgesia. The sample size calculation for this study was based on a two-tailed *t*-test for two independent groups (dorsal vs. volar blocks). A study with an effect size of 1 and a power of 90% required a total sample of 54 to test the association at a 5% alpha level. The power calculation was carried out using G\*Power 3.1.9.

### Randomization

Consecutive patients were selected in the emergency department of a tertiary orthopedic center if they met the inclusion criteria. Patients were block randomized to receive either the TD or SV techniques by computer-generated lists equally distributed to four trained providers. The four blocks corresponding to the number of the surgeons consisted of either 12 volar and 13 dorsal or 13 volar and 12 dorsal, making four blocks of 25 patients distributed to each surgeon. The order of the blocks and the patients within the block were randomized via a computer-generated sequence, which was concealed from the operator until a participant was enrolled.

### Techniques employed

The TD technique was performed by injecting 1.5 ml of 1% lidocaine without epinephrine at each lateral aspect of the base of the finger, just distal to the metacarpophalangeal joint, at the level of the web spaces. The SV technique was performed by injecting 3 ml of 1% lidocaine at the base of the finger at the proximal volar metacarpophalangeal crease. In all cases, a 27G needle was used.

### Outcome assessments

The primary outcome was pain measured using the Numerical Rating Scale (NRS), labeled 0 (painless) to 10 (worst pain). Measurements were taken across two stages, before the administration of local anesthesia (injury-related pain) and during administration (procedure-related pain). Patients were further evaluated for the anesthesia onset time and the extent of anesthesia according to each region of the digit. The anesthesia

onset time was assessed by using a needle prick in each of the six regions of the finger at 30-s intervals. The extent of anesthesia for each region of the finger was simultaneously recorded through the same technique, and the effect of anesthesia was obtained accordingly.

### Statistical analysis

Statistical analysis was performed using R 3.6.3. Counts and percentages were used to summarize categorical variables. The mean  $\pm$  standard deviation (standard deviation) or the median/interquartile range (IQR) was used to summarize the distribution of continuous normal and nonnormal variables, respectively. Unpaired *t*-test and Mann–Whitney test were used to assess the balance between groups at baseline for continuous normal and nonnormal variables, respectively. Chi-square test of independence was used to compare the distribution of categorical variables between patients who received TD and SV nerve blocks.

The primary analysis included comparing time to onset and final pain scores between patients who received TD and SV nerve blocks. The primary outcome analysis was performed using unpaired *t*-test and Mann–Whitney test for continuous normal and nonnormal variables, respectively. Hypothesis testing was performed at 5% levels of significance.

## Results

Between January 2021 and June 2021, a total of 100 patients were consecutively recruited and randomized to receive either TD (*n* = 50) or SV (*n* = 50) block [Figure 1]. Eight patients were excluded for not meeting inclusion criteria: two excluded on the basis of peripheral neuropathy, four excluded due to injury on the proximal dorsal aspect of the finger, and two excluded for the suspected presence of a foreign body.

The descriptive characteristics of our study population are displayed in Table 1. No statistical significance was observed between the distributions (*P* > 0.05) except the finger injured (*P* < 0.05). The majority of the injuries were distal (38%) and affected the volar aspect of the fingers (67%). Differences of age, baseline pain, sex, and injury type were not significant, thereby ensuring comparability across these groups.

The results of the primary outcomes are displayed in Tables 2 and 3 and Figure 2. The median procedure-related pain score was significantly higher in patients who received TD nerve block than patients who received SV nerve block (median [IQR] = 4 [2.25, 5.00] vs. 3.00 [2.00, 4.00], respectively). The difference in the procedure-related pain between the two groups was




Figure 1: CONSORT Flowchart displaying enrolment, allocation, follow up and analysis of participants

**Table 1: Distribution and between-group comparisons of participants according to patient and injury factors**

	All (n=100), n (%)	Dorsal (n=50), n (%)	Volar (n=50), n (%)	P
Surgeon				
A	25 (25.0)	12 (24.0)	13 (26.0)	0.984
B	25 (25.0)	13 (26.0)	12 (24.0)	
C	25 (25.0)	13 (26.0)	12 (24.0)	
D	25 (25.0)	12 (24.0)	13 (26.0)	
Sex				
Male	83 (83.0)	40 (80.0)	43 (86.0)	0.594
Female	17 (17.0)	10 (20.0)	7 (14.0)	
Age	34.5 (29.0-45.0)	37.5 (29.2-45.8)	33.0 (29.0-41.0)	0.265
Injury				
Dorsal	33 (33.0)	19 (38.0)	14 (28.0)	0.395
Volar	67 (67.0)	31 (62.0)	36 (72.0)	
Location				
Distal	45 (50.6)	23 (52.3)	22 (48.9)	0.466
Middle	23 (25.8)	9 (20.5)	14 (31.1)	
Multi-regional	21 (23.6)	12 (27.3)	9 (20.0)	
Injury-related pain	6.00 (4.00-7.00)	6.00 (5.00-7.00)	5.00 (4.00-6.75)	0.130
Side				
Left	47 (47.0)	21 (42.0)	26 (52.0)	0.423
Right	53 (53.0)	29 (58.0)	24 (48.0)	
Finger				
Index	38 (38.0)	16 (32.0)	22 (44.0)	0.002
Little	9 (9.00)	3 (6.00)	6 (12.0)	
Middle	38 (38.0)	17 (34.0)	21 (42.0)	
Ring	15 (15.0)	14 (28.0)	1 (2.00)	

**Table 2: The difference in the median procedure-related pain between groups**

	Dorsal (n=50)	Volar (n=50)	Δ 95% CI
Procedure-related pain	4.00 (2.25-5.00)	3.00 (2.00-4.00)	1 (0.25-1.75)

Continuous data were summarized as median (IQR). The difference and 95% CI for the difference between medians were computed using the approach suggested by Price and Bonnett.<sup>[8]</sup> IQR: Interquartile range, CI: Confidence interval, Δ: Delta

**Table 3: Anesthesia onset time between groups**

	Dorsal (n=50)	Volar (n=50)	P	Δ 95% CI
Anesthesia onset time (min)	2.00 (2.00-3.00)	2.00 (1.50-2.50)	0.385	0.13 (-0.17-0.43)

Continuous data were summarized as median (IQR). P values were obtained using Mann-Whitney test, Unpaired t-test was used to obtain the mean difference and 95% CI for the difference between groups. IQR: Interquartile range, CI: Confidence interval, Δ: Delta

1 (95% confidence interval [CI]: 0.25; 1.75). However, anesthesia onset time was not statistically different among the groups (P = 0.39).

Results showed that the difference in median NRS was statistically significant, as evidenced by the 95% CI that does not include zero.

The extent of anesthesia along with anesthesia onset time of each region is depicted in Figure 3. In the SV block, the anesthesia onset time is shortest in the volar region, followed by the middle and distal dorsal regions. The

proximal dorsal region took the longest time to become anesthetized. In the TD block, the proximal and dorsal regions were quickly anesthetized, followed by the middle and distal regions and finally the distal volar.

## Discussion

Sensory innervation of each digit is provided by four digital nerves entering the finger base bilaterally: two at the volar and two at the dorsal side. Targeting those nerves effectively while minimizing patient discomfort forms the basis of a desirable digital block. The technique of injecting the local anesthetic into the SC tissue above the flexor tendon sheath at the volar side of the hand was described by Low *et al.*<sup>[9]</sup> This technique is intended to reduce the risk of neurovascular injury that may be associated with the TD technique into the sides of the digit.

In this trial, we compared the two-injection dorsal block and single-injection volar subcutaneous block techniques by assessing procedure-related pain as well as the extent of anesthesia across the six regions of the finger. Our results showed that the mean pain score was significantly lower in patients who received a SV nerve block compared to patients who received TD nerve block. In similar trials, there is considerable heterogeneity related to the sample size, pain score used, severity of injury, type of injury, the comparison of blocks, and the local




Figure 2: Distribution of (A) procedure-related pain and (B) anesthesia onset time between groups. Dorsal; TD, Volar; SV

anesthesia used.<sup>[10-13]</sup> Yet despite this, there have been consistent results showing lower pain scores in the volar block,<sup>[12,14-17]</sup> despite statistical insignificance. Further, data suggest that patients prefer the SV injection.<sup>[14-16]</sup> Clement *et al.* and Cannon *et al.* both reported greater clinician satisfaction with the volar block, due to ease of administration and patient response.<sup>[12,15]</sup>

The time to onset was not statistically significant among our groups [Figure 3]. Our results showed a faster onset of anesthesia with the SV on the volar aspect. Okur *et al.* assessed time to complete anesthesia of the finger yielding similar results to our study.<sup>[18]</sup> While the TD more effectively anesthetized the proximal dorsal region, consistent with recent reports by Clement *et al.*<sup>[12]</sup> We found that the proximal dorsal region was eventually anesthetized by the volar block, albeit less timely, predictably, and effectively. Previous studies noted an increased risk of incomplete anesthesia over the proximal dorsal region, findings that were partially consistent with ours.<sup>[13,14]</sup> In addition, although we did not examine the duration of action for the TD technique, Afridi *et al.* reported a longer duration of action for the SV technique compared with the TD technique.<sup>[11]</sup>

As a limitation of our study, the various tangential techniques employed by some providers to reduce pain, which could have a significant impact and contribute to important variability, have not been examined. These include buffering the solution with sodium bicarbonate, smaller needle gauges, solutions at room temperature, pinching loose skin or creating “sensory noise” near the injection site, and injecting at 90°.<sup>[19-24]</sup> In addition, the study did not investigate patient and provider preference, a potential underlying element of which is the number of injections. However, confidence in our comparisons is achieved by the reasonably sized sample which removed sources of confounding by excluding complex presentations and including a single injury




Figure 3: Schematics depicting anaesthesia onset time measured at 30-second intervals in all participants. Values labelled according to their respective regions represent the median anaesthesia onset time in seconds. The grey colour gradient shows speed of anaesthesia from darkest (fastest) to lightest (slowest)

type. In addition, in an effort to accurately assess the extent of the temporal and regional effects of the blocks, we assessed variables at short 30-s intervals, compared with previous studies which employed 5- and 10-min intervals from the time of injection.<sup>[25,26]</sup>

## Conclusion

The single-injection volar subcutaneous blocks are less painful with a similar anesthesia onset time. The single-injection volar subcutaneous block is preferable in injuries to all regions of the digit except the proximal dorsal region. The two-injection dorsal blocks are ideal in the latter to ensure the complete and timely anesthesia of the region.

### Author contributions (CRediT)

Ali Jarragh: Conceptualization, writing original draft, and methodology.

Ali Lari: Conceptualization, writing original draft, presentation, and methodology.

Waleed Burhamah: Data curation and writing original draft.

Mohammed Alherz: Writing review and editing and formal analysis.

Abdullah Nouri: Data curation and editing.

Yahia Alshammari: Data collection and editing.

Ameer Al-Jasim: Formal analysis and resources.

Sulaiman Alrefai: Data collection and editing.

Naser Alnusif: Review, conceptualization, and supervision.

### Conflicts of interest

None declared.

### Consent to participate

Written consent was obtained from all included participants.

### Ethical approval

Ethical approval was obtained from the Ethical Committee Board at the Ministry of Health - 06/01/2021 (UID 1742/2021).

### Funding

None.

## References

- Ootes D, Lambers KT, Ring DC. The epidemiology of upper extremity injuries presenting to the emergency department in the United States. *Hand (N Y)* 2012;7:18-22.
- Robson AK, Bloom PA. Suturing of digital lacerations: Digital block or local infiltration? *Ann R Coll Surg Engl* 1990;72:360-1.
- Chale S, Singer AJ, Marchini S, McBride MJ, Kennedy D. Digital versus local anesthesia for finger lacerations: A randomized controlled trial. *Acad Emerg Med* 2006;13:1046-50.
- Myers S. Intermetacarpal Blocks; 2008. Available from: <http://www.myhand.com.au/health-professionals/injection-techniques/intermetacarpal-blocks>. [Last accessed on 2015 Dec 06].
- Hart RG, Fernandes FA, Kutz JE. Transthecal digital block: An underutilized technique in the ED. *Am J Emerg Med* 2005;23:340-2.
- Braun H, Harris ML, editors. *Operations on the extremities. In: Local Anesthesia: Its Scientific Basis and Practical Use.* 2<sup>nd</sup> ed. New York: Lea & Febiger; 1924. p. 366-7.
- Harbison S. "Transthecal digital block: Flexor tendon sheath used for anaesthetic infusion". *J Hand Surg Am* 1991;16:957.
- Bonett DG, Price RM. Statistical inference for a linear function of medians: Confidence intervals, hypothesis testing, and sample size requirements. *Psychol Methods* 2002;7:370-83.
- Low CK, Vartany A, Engstrom JW, Poncelet A, Diao E. Comparison of transthecal and subcutaneous single-injection digital block techniques. *J Hand Surg Am* 1997;22:901-5.
- Vinycomb TI, Sahhar LJ. Comparison of local anesthetics for digital nerve blocks: A systematic review. *J Hand Surg Am* 2014;39:744-51.e5.
- Afridi RA, Masood T, Ahmed E, Obaidullah AM, Alvi HF. Comparison of the efficacy of single volar subcutaneous digital block and the dorsal two injections block. *J Ayub Med Coll Abbottabad* 2014;26:88-91.
- Clement P, Doomen L, van Hooft M, Hessels R. Regional anaesthesia on the finger: Traditional dorsal digital nerve block versus subcutaneous volar nerve block, a randomized controlled trial. *Injury* 2021;52:883-8.
- Yin ZG, Zhang JB, Kan SL, Wang P. A comparison of traditional digital blocks and single subcutaneous palmar injection blocks at the base of the finger and a meta-analysis of the digital block trials. *J Hand Surg Br* 2006;31:547-55.
- Williams JG, Lalonde DH. Randomized comparison of the single-injection volar subcutaneous block and the two-injection dorsal block for digital anesthesia. *Plast Reconstr Surg* 2006;118:1195-200.
- Cannon B, Chan L, Rowlinson JS, Baker M, Clancy M. Digital anaesthesia: One injection or two? *Emerg Med J* 2010;27:533-6.
- Hung VS, Bodavula VK, Dubin NH. Digital anaesthesia: Comparison of the efficacy and pain associated with three digital nerve block techniques. *J Hand Surg Br* 2005;30:581-4.
- Martires KJ, Malbasa CL, Bordeaux JS. A randomized controlled crossover trial: Lidocaine injected at a 90-degree angle causes less pain than lidocaine injected at a 45-degree angle. *J Am Acad Dermatol* 2011;65:1231-3.
- Okur OM, Şener A, Kavaklı HŞ, Çelik GK, Doğan NÖ, İçme F, et al. Two injection digital block versus single subcutaneous palmar injection block for finger lacerations. *Eur J Trauma Emerg Surg* 2017;43:863-8.
- Joukhadar N, Lalonde D. How to minimize the pain of local anesthetic injection for wide awake surgery. *Plast Reconstr Surg Glob Open* 2021;9:e3730.
- Strazar AR, Leynes PG, Lalonde DH. Minimizing the pain of local anesthesia injection. *Plast Reconstr Surg* 2013;132:675-84.
- Cepeda MS, Tzortzopoulou A, Thackrey M, Hudcova J, Arora Gandhi P, Schumann R. Adjusting the pH of lidocaine for reducing pain on injection. *Cochrane Database Syst Rev* 2010;12:CD006581.
- McKay W, Morris R, Mushlin P. Sodium bicarbonate attenuates pain on skin infiltration with lidocaine, with or without epinephrine. *Anesth Analg* 1987;66:572-4.
- Arendt-Nielsen L, Egekvist H, Bjerring P. Pain following controlled cutaneous insertion of needles with different diameters. *Somatosens Mot Res* 2006;23:37-43.
- Hogan ME, vanderVaart S, Perampaladas K, Machado M, Einarson TR, Taddio A. Systematic review and meta-analysis of the effect of warming local anesthetics on injection pain. *Ann Emerg Med* 2011;58:86-98.e1.
- Cousins MJ, Bridenbaugh PO. *Neural Blockade in Clinical Anesthesia and Pain Medicine.* Philadelphia: Lippincott Williams & Wilkins; 2009. p. 94-100.
- Roberts JR, Hedges JR. *Clinical Procedures in Emergency Medicine. Section V: Anesthetic and Analgesic Techniques.* Philadelphia: Elsevier Health Sciences; 2014. p. 530.