













Impaired myocardial reserve underlies reduced exercise capacity and heart rate recovery in preterm-born young adults

Odaro J. Huckstep ^{1,2†}, Holger Burchert ^{1†}, Wilby Williamson ¹, Fernando Telles ¹, Cheryl M.J. Tan ¹, Mariane Bertagnolli ^{1,3}, Linda Arnold ¹, Afifah Mohamed ^{1,4}, Kenny McCormick ⁵, Henner Hanssen ⁶, Paul Leeson ¹, and Adam J. Lewandowski ^{1*}

¹Oxford Cardiovascular Clinical Research Facility, Division of Cardiovascular Medicine, Radcliffe Department of Medicine, University of Oxford, Headley Way, John Radcliffe Hospital, Oxford OX39DU, UK; ²Department of Biology, United States Air Force Academy, 2355 Faculty Drive, Suite 2P389, Colorado 80840, USA; ³Hospital Sacré-Coeur Research Center, CIUSSS du Nord-de-l'Île-de-Montréal, 5400 Boulevard Gouin Ouest, Montreal, Quebec H4J1C5, Canada; ⁴Department of Diagnostic Imaging and Radiotherapy, Faculty of Health Sciences, Universiti Kebangsaan Malaysia, Jalan Raja Muda Abdul Aziz, Kuala Lumpur 50300, Malaysia; ⁵Department of Paediatrics, Headley Way, John Radcliffe Hospital, Oxford OX39DU, UK; and; ⁶Department of Sport, Exercise and Health, University of Basel, Birsstrasse 320B, Basel 4052, Switzerland

Received 17 December 2019; editorial decision 11 March 2020; accepted 12 March 2020; online publish-ahead-of-print 17 April 2020

Aims

We tested the hypothesis that the known reduction in myocardial functional reserve in preterm-born young adults is an independent predictor of exercise capacity (peak VO_2) and heart rate recovery (HRR).

Methods and results

We recruited 101 normotensive young adults ($n = 47$ born preterm; 32.8 ± 3.2 weeks' gestation and $n = 54$ term-born controls). Peak VO_2 was determined by cardiopulmonary exercise testing (CPET), and lung function assessed using spirometry. Percentage predicted values were then calculated. HRR was defined as the decrease from peak HR to 1 min (HRR₁) and 2 min of recovery (HRR₂). Four-chamber echocardiography views were acquired at rest and exercise at 40% and 60% of CPET peak power. Change in left ventricular ejection fraction from rest to each work intensity was calculated (EF Δ 40% and EF Δ 60%) to estimate myocardial functional reserve. Peak VO_2 and per cent of predicted peak VO_2 were lower in preterm-born young adults compared with controls (33.6 ± 8.6 vs. 40.1 ± 9.0 mL/kg/min, $P = 0.003$ and $94\% \pm 20\%$ vs. $108\% \pm 25\%$, $P = 0.001$). HRR₁ was similar between groups. HRR₂ decreased less in preterm-born young adults compared with controls (-36 ± 13 vs. -43 ± 11 b.p.m., $P = 0.039$). In young adults born preterm, but not in controls, EF Δ 40% and EF Δ 60% correlated with per cent of predicted peak VO_2 ($r^2 = 0.430$, $P = 0.015$ and $r^2 = 0.345$, $P = 0.021$). Similarly, EF Δ 60% correlated with HRR₁ and HRR₂ only in those born preterm ($r^2 = 0.611$, $P = 0.002$ and $r^2 = 0.663$, $P = 0.001$).

Conclusions

Impaired myocardial functional reserve underlies reductions in peak VO_2 and HRR in young adults born moderate-preterm. Peak VO_2 and HRR may aid risk stratification and treatment monitoring in this population.

Keywords

preterm birth • prematurity • exercise capacity • heart rate recovery • echocardiography • cardiac reserve

Introduction

Prematurity is the leading cause of under-five child mortality,¹ though more than 90% of those born preterm now survive into adulthood due to modern advances in pre- and post-natal care. Nevertheless,

those born preterm suffer an increased prevalence of respiratory, cardiovascular, and neurological diseases in later life.¹ Cardiovascular sequelae related to preterm birth include hypertension, early heart failure, and ischaemic heart disease, as well as increased mortality from cardiovascular disorders.^{2–4}

*Corresponding author. Tel: +44(0)1865221392; Fax: +44(0)1865572840. E-mail: adam.lewandowski@cardiov.ox.ac.uk

[†]These authors contributed equally to this work

© The Author(s) 2020. Published by Oxford University Press on behalf of the European Society of Cardiology.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

The rate of maximum oxygen uptake during exercise (peak VO_2) and heart rate recovery (HRR) are recognized as independent predictors of morbidity and all-cause mortality.^{5,6} Individuals born preterm have been reported to have lower peak VO_2 ⁷ and have recently been shown to have impaired HRR,^{8–10} though the underlying mechanisms are unknown. Besides metabolic and cardiac alterations, interrupted pulmonary development, which normally continues late into the third trimester, is generally considered to contribute to potential exercise deficits.⁷ However, 80–85% of preterm births fall within the range of moderate to late prematurity (32–36 weeks gestational age) where acute neonatal respiratory distress syndrome is rarely seen.¹¹ This variety of alterations, their association to gestational age, and the fact that impaired peak VO_2 and HRR are seen in a range of conditions¹² limits risk stratification and compromises the interpretation of impaired peak VO_2 and HRR in individuals with a history of preterm birth.

Our previous investigations identified an altered cardiac phenotype in preterm-born young adults, even in those born moderate to late preterm.^{13,14} These cardiac alterations include an impaired increase in left ventricular (LV) ejection fraction (EF) between rest and light to moderate exercise intensity, suggestive of a reduced myocardial functional reserve.¹⁵ We therefore tested the hypothesis that this reduction in myocardial functional reserve in preterm-born young adults is an independent predictor of peak VO_2 and HRR.

Methods

Study design

We completed the Young Adult Cardiovascular Health sTudy (YACHT); an observational, case-control study with a primary outcome measure of cardiopulmonary exercise capacity to investigate cardiovascular structure and function, along with cardiopulmonary capacity and exercise stress response in preterm- and term-born young adults. Participants completed a detailed multimodal set of study measures including 24-h ambulatory blood pressure monitoring, cardiopulmonary exercise testing (CPET), and cardiac imaging. Ethical approval for YACHT was granted by the South Central Berkshire Research Ethics Committee (14/SC/0275). Study registration was completed via www.clinicaltrials.gov (NCT02103231).

Study population

A total of 149 participants aged 18–40 years were recruited through purposive active and passive recruitment as previously described.¹⁵ In total, 101 non-hypertensive young adult participants (47 preterm-born, 54 term-born) meeting the inclusion and exclusion criteria were included in this analysis.¹⁵ Cohort characteristics are provided in *Table 1*.

Study visit

Participants were instructed to fast overnight for 12 h but to drink plenty of water prior to attending a study visit at the University of Oxford Centre for Clinical Magnetic Resonance (OCMR) and Oxford Cardiovascular Clinical Research Facility (CCRF) in the John Radcliffe Hospital in Oxford, UK. All measurements were completed by trained study investigators and clinical staff, including height, weight, venepuncture, and blood pressure as previously described (see [Supplementary material online](#)).¹⁵

Spirometry

Spirometry was performed according to standard guidelines by using the Cortex Metalyzer 3B (Metalyzer 3B, Cortex Biophysik GmbH, Leipzig, Germany).¹⁶ The spirometer underwent calibration checks of the flow

and volume before each test using a Futuremed's 3.0-L calibration syringe (Futuremed, CA, USA). The participants were seated and guided through the test wearing the nose-clips with the lips tightly sealed around the mouthpiece. A minimum of three acceptable forced expiratory manoeuvres were made for every participant. To ensure consistency and high quality of the spirometry data, visual checks of the waveforms were done before the highest values from the data were selected and recorded on a clinical research form. The parameters recorded included forced expiratory volume in 1 s (FEV_1) and forced vital capacity (FVC), as well as per cent of predicted values according to Quanjer *et al.*¹⁷

CPET and HRR

Following spirometry, participants completed a peak CPET on a seated stationary cycle ergometer (Ergoline GmbH, Bitz, Germany) using a validated incremental protocol with respiratory gases collected and measured using the same Metalyzer as used during spirometry. Heart rate was recorded using continuous electrocardiogram (ECG) monitoring and blood pressure was recorded every 4 min using a manual mercury sphygmomanometer (ACCOSON Freestyle, Essex, UK). Participants maintained a rate of 60 revolutions per minute during the test, which began with one quiescent minute of resting measurements followed by a 2-min warm-up with a 20 watt workload. After the warm-up period, workload increased to 35 watts. To normalize test duration to ~8–12 min, participants who reported higher activity or fitness levels had their workload increased to 75 watts after the warm-up period. Workload then incremented by 15 watts each minute and participants cycled continuously until exhaustion prevented them from maintaining at least 50 revolutions per minute or safety termination criteria were met.¹⁸ Participants then completed a 2-min cool-down period at 50 watts and the revolutions per minute of their preference. Per cent of predicted peak VO_2 was calculated based on the Wassermann weight algorithm.¹⁹

Stress echocardiography

After completing CPET, participants had a 30-min recovery period during which they were offered water and a cereal bar. Sub-maximal exercise stress echocardiography was then performed on study subjects for 3 min at 40% and 60% of their peak exercise wattage achieved during CPET testing as previously described.¹⁵ In brief, after 2 min at each prescribed workload and with steady-state heart rate verified, apical four-chamber views optimized for the LV and pulsed wave Doppler imaging of mitral inflow were collected at each stage using either a Philips iE33 or Philips EPIQ 7C (Philips Healthcare, Surrey, UK) cardiology ultrasound machine with an X5-1 xMATRIX array transducer to compare with resting echocardiography measures as previously described.¹⁵ Participants were given a 3-min rest period between each stage. Offline image analysis was only performed on images where heart rate was within 5% of maximum achieved steady-state heart rate for a given stage.

Data processing

Echocardiography analysis

Echocardiography analysis was completed offline in accordance with standard guidelines as previously described (see [Supplementary material online](#), Methods).¹⁵ Resting LVEF was calculated via the bi-plane Modified Simpson's method using apical four-chamber and apical two-chamber views while sub-maximal exercise LVEF at 40% and 60% were calculated by the single-plane Modified Simpson's method using apical four-chamber views. EF change was calculated from rest to 40% and 60% exercise intensity ($\text{EF}\Delta 40\%$ and $\text{EF}\Delta 60\%$). Intraclass correlation coefficients for LV ejection fraction at rest and under physiological stress at 40% and 60% for intra/inter-observer variability were 0.89/0.85, 0.90/0.89, and 0.89/0.90, respectively.

Table 1 Cohort clinical characteristics

Group characteristics	Preterm-born young adults (n = 47)	Term-born young adults (n = 54)	P-value
Demographics and anthropometrics			
Age (years)	22.7 ± 3.04	23.6 ± 3.8	0.239
Gestational age (weeks)	32.8 ± 3.23	39.5 ± 1.37	<0.001
32–36 weeks, n (%)	38 (80.9)	0 (0.0)	
28–31 weeks, n (%)	5 (10.6)	0 (0.0)	
<28 weeks, n (%)	4 (8.5)	0 (0.0)	
Male, n (%)	14 (30)	26 (48)	0.061
Birth weight (grams)	1916 ± 806	3390 ± 424	<0.001
Gestational hypertension, n (%)	8 (17.0)	0 (0.0)	0.002
Small for gestational age, n (%)	2 (4.3)	0 (0.0)	0.214
Height (cm)	167 ± 9	175 ± 10	<0.001
BMI (kg/m ²)	23.3 ± 4.5	22.7 ± 2.7	0.401
Biochemistry			
Total cholesterol (mmol/L)	4.72 ± 0.65	4.18 ± 0.77	0.001
HDL (mmol/L)	1.49 ± 0.31	1.47 ± 0.26	0.882
LDL (mmol/L)	2.80 ± 0.71	2.32 ± 0.60	0.001
Triglycerides (mmol/L)	1.12 ± 0.66	0.87 ± 0.36	0.031
High sensitivity CRP (mg/L)	1.57 ± 2.42	1.14 ± 1.96	0.412
Glucose (mmol/L)	5.02 ± 0.41	4.82 ± 0.51	0.030
Insulin (pmol/L)	51.1 ± 29.0	35.8 ± 29.4	0.012
Insulin resistance	0.96 ± 0.54	0.68 ± 0.59	0.020
Brachial blood pressure (mmHg)			
Resting systolic	119 ± 9	115 ± 8	0.014
Resting diastolic	70 ± 8	66 ± 5	0.014
Average 24h systolic	119 ± 6	119 ± 8	0.385
Average 24h diastolic	71 ± 5	69 ± 5	0.065
Cardiac function			
Heart rate (b.p.m.)	75 ± 13	61 ± 12	<0.001
Ejection fraction (%)	63.1 ± 5.4	63.4 ± 4.7	0.624
Stroke index (mL/m ²)	29.6 ± 6.4	32.2 ± 7.8	0.173
Pulmonary function			
FVC (% of predicted)	109 ± 15.2	109 ± 12.0	0.917
FEV ₁ /FVC (%)	81.4 ± 6.4	81.6 ± 8.1	0.795

Group characteristics presented as mean ± SD. Biochemistry, blood pressure, cardiac function, and pulmonary function *P*-values adjusted for sex. Insulin resistance was calculated using the Homeostatic Model Assessment Index (HOMA) calculator (www.dtu.ox.ac.uk/homacalculator/). Bold *P* values are statistically significant (*P* < 0.05). BMI, body mass index; FEV₁, forced expiratory volume in 1 s; FVC, forced vital capacity; HDL, high density lipoprotein; LDL, low-density lipoprotein.

Heart rate recovery

The ECG of each participant after the cessation of exercise was inspected by two researchers for valid readings at 1 and 2 min after exercise. Heart rate was smoothed within the Metalyzer Software (MetaSoft Studio, Cortex Biophysik GmbH, Leipzig, Germany) using the moving average (time interval) set to 15 s so that an average from the data points in this specified time interval (around the supporting point) was formed. The heart rates at 1 and 2 min after the cessation of exercise were then identified and extracted for each participant. HRR was defined as the absolute drop in heart rate from peak heart rate to 1 min (HRR₁) and 2 min after exercise (HRR₂).

Statistical analysis


Statistical analysis was performed using SPSS Version 25. Shapiro–Wilk testing and visual inspection were used to assess normality of variable distribution. Direct, between-group comparisons were performed using

independent-samples Student's *t*-tests for normally distributed data and Mann–Whitney and Kruskal–Wallis tests for non-normally distributed data. Between-group comparisons were adjusted for sex. Linear regression modelling was completed using forced entry with missing data excluded pairwise. Our final study numbers *n* = 47 vs. *n* = 54 allowed us to detect a 0.77-standard deviation difference between groups for study measures, powered at 80% with *P* = 0.05. *P*-values < 0.05 were considered statistically significant.

Results

Clinical characteristics

There were no differences in body mass index or age, but between-group sex distribution approached significance with a 30% male pre-term group vs. a 48% male term-born group, *P* = 0.061. Between-



group comparisons were, therefore, adjusted for sex throughout. Resting LVEF and stroke index measured by echocardiography were similar between groups. Baseline pulmonary function measured by spirometry was also similar between groups with no significant differences in per cent of predicted FVC or FEV_1/FVC . Baseline group characteristics are provided in *Table 1*.

Exercise characteristics

Reduced aerobic exercise capacity and HRR in preterm-born young adults

Peak cardiopulmonary exercise capacity was lower in preterm-born young adults, who on average, achieved $94\% \pm 20\%$ of their predicted peak VO_2 vs. $108\% \pm 25\%$ in term-born controls, $P=0.001$ (*Figure 1*). Peak VO_2 measured in mL/kg/min was also lower in those born preterm (33.6 ± 8.6 vs. 40.1 ± 9.0 mL/kg/min, $P=0.003$). Heart rate at the ventilatory anaerobic threshold (VAT) was higher in the preterm group (132 ± 18 vs. 124 ± 17 beats/min, $P=0.038$) but peak achieved heart rate was similar between groups. Ventilatory reserve at peak ventilation rate was similar between preterm-born and term-born individuals ($23.3\% \pm 17.8\%$ vs. $21.2\% \pm 18.3\%$ of maximum minute ventilatory volume, respectively, $P=0.513$) (*Table 2*). As shown in *Figure 2*, HRR_1 was similar between groups (-21 ± 9 vs. -24 ± 9 b.p.m., $P=0.457$) and remained similar after additional adjustment for per cent of predicted peak VO_2 ($P=0.492$). HRR_2 was slower in

Table 2 Exercise characteristics

	Preterm-born young adults ($n=47$)	Term-born young adults ($n=54$)	P-value
Exercise performance			
Per cent of predicted peak VO_2	94 ± 20	108 ± 25	0.001
Peak VO_2 (mL/kg/min)	33.6 ± 8.6	40.1 ± 9.0	0.003
OUE slope	2.39 ± 0.73	3.05 ± 0.89	0.001
VE/ VCO_2 slope	28.1 ± 3.3	25.9 ± 3.8	0.011
Peak RER	1.19 ± 0.06	1.19 ± 0.06	0.738
Cardiovascular function			
Heart rate at VAT (b.p.m.)	132 ± 18	124 ± 17	0.038
Heart rate at 60% intensity (b.p.m.)	159 ± 11	151 ± 14	0.055
Peak HR achieved (b.p.m.)	189 ± 9	186 ± 11	0.218
HRR_1 (b.p.m.)	-21 ± 9	-24 ± 9	0.457
HRR_2 (b.p.m.)	-36 ± 13	-43 ± 11	0.039
Systolic blood pressure at 60% intensity (mmHg)	146 ± 16	149 ± 14	0.567
Diastolic blood pressure at 60% intensity (mmHg)	81 ± 10	77 ± 7	0.139
Peak systolic blood pressure (mmHg)	165 ± 22	167 ± 21	0.885
Peak diastolic blood pressure (mmHg)	89 ± 11	82 ± 10	0.002
Peak pulse pressure (mmHg)	76 ± 23	86 ± 23	0.168
Pulmonary function			
Breaths per minute at peak VO_2	44.3 ± 6.7	47.4 ± 11.8	0.188
Ventilatory reserve at peak (%)	23.3 ± 17.8	21.2 ± 18.3	0.513
Objective measure of physical activity			
MVPA (h/week)	14.2 ± 5.8	14.9 ± 6.1	0.633
VPA (h/week)	0.9 ± 0.9	1.2 ± 1.2	0.146

Exercise characteristics presented as mean \pm SD. P-values adjusted for sex. Bold P values are statistically significant ($P<0.05$).

HRR, heart rate recovery; MVPA, moderate to vigorous physical activity; OUE Slope, oxygen uptake efficiency slope; RER, respiratory exchange ratio; VAT, ventilatory anaerobic threshold; VE/ VCO_2 , minute ventilation/carbon dioxide production; VO_2 , oxygen uptake; VPA, vigorous physical activity.




Figure 2 Heart rate recovery at one and two minutes post exercise. There was no significant difference in heart rate recovery after one minute (HRR1) between preterm-born young adults in blue ($n = 40$) and term-born controls in green ($n = 46$). Heart rate recovery after two minutes (HRR2) was significantly reduced in preterm-born young adults ($n = 39$) compared to term-born controls ($n = 46$). Boxplots represent min to max.

preterm-born young adults (-36 ± 13 vs. -43 ± 11 b.p.m., $P = 0.039$), but no significant difference was found after additional adjustment for per cent of predicted peak VO_2 ($P = 0.813$). Exercise characteristics are presented in [Table 2](#).

Change in EF during exercise predicts exercise capacity and HRR in preterm-born young adults

Bivariate correlations were carried out to investigate whether EFA, pulmonary, and physical activity measures were associated with per cent of predicted peak VO_2 and HRR. As shown in [Supplementary data](#) online, [Table S1](#), EFA $\Delta 40\%$ as well as EFA $\Delta 60\%$ were correlated with per cent of predicted peak VO_2 only in preterm-born young adults ($r^2 = 0.430$, $P = 0.015$ and $r^2 = 0.345$, $P = 0.021$, respectively) ([Figure 3A](#) and [C](#)). EFA $\Delta 60\%$ was correlated with HRR $_1$ and HRR $_2$ only in preterm-born young adults ($r^2 = 0.611$, $P = 0.002$ and $r^2 = 0.663$, $P = 0.001$, respectively) ([Figure 4A](#) and [C](#)).

Multivariable analysis for confounding effects of pulmonary capacity and dynamic function in preterm-born subjects

To determine if the relationships between EFA and per cent of predicted peak VO_2 as well as HRR in preterm-born subjects were




Figure 3 Increase in LVEF from baseline to 40% and 60% as a predictor of peak VO_2 . Change in LVEF from rest to 40% exercise intensity (EFA $\Delta 40\%$) was predictive of percent of predicted peak VO_2 in the preterm-born group (Panel A: $r^2 = 0.430$, $P = 0.015$, $n = 13$), but not in full-term born controls (Panel C: $r^2 < 0.001$, $P = 0.953$, $n = 28$). Change in LVEF from rest to 60% exercise intensity (EFA $\Delta 60\%$) was similarly predictive of percent of predicted peak VO_2 in the preterm-born group (Panel B: $r^2 = 0.345$, $P = 0.021$, $n = 15$), but not in full-term controls (Panel D: $r^2 = 0.003$, $P = 0.766$, $n = 30$).




Figure 4 Change in LVEF from rest to 60% exercise intensity as a predictor of heart rate recovery. Change in LVEF from rest to 60% exercise intensity (EFΔ60%) was predictive of heart rate recovery at 1 and 2 min (HRR₁ and HRR₂, respectively) post-exercise in preterm-born young adults (Panel A: $r^2 = 0.611$, $P = 0.002$, $n = 13$ and Panel B: $r^2 = 0.663$, $P = 0.001$, $n = 13$, respectively) but not in term-born young adults (Panel C: $r^2 = 0.068$, $P = 0.230$, $n = 23$ and Panel D: $r^2 = 0.003$, $P = 0.816$, $n = 23$, respectively).

significantly influenced by pulmonary volume or dynamic pulmonary function, we completed multivariable linear regression analyses. Based on bivariable regression analyses, significant variables were carried forward to a multivariable model. To predict per cent of predicted peak VO_2 in the preterm group, we included per cent predicted FVC, ventilatory reserve at peak VO_2 , and EFΔ40% and then EFΔ60% in two separate models. EFΔ40% and EFΔ60% remained the strongest predictors of per cent of predicted peak VO_2 (beta = 0.538, $P = 0.084$ and beta = 0.506, $P = 0.152$) (Table 3). For the prediction of HRR₁ and HRR₂ in the preterm group, we included per cent of predicted FVC and EFΔ60% in the model. EFΔ60% remained the strongest predictor of HRR₁ (beta = -1.030, $P = 0.002$) and HRR₂ (beta = -0.948, $P = 0.003$).

Discussion

This study reports for the first time that normotensive, moderately preterm-born young adults exhibited reduced aerobic exercise capacity and slower HRR. These reductions were strongly related to their level of impairment in LV systolic function when going from rest to mild and moderate levels

of exercise intensity. In addition, mechanical pulmonary function in this moderately preterm group was similar to controls, and exercise LV systolic response remained the strongest predictor of exercise capacity and HRR in preterm-born young adults after adjusting for pulmonary volume and dynamic function.

Impaired preterm myocardial functional reserve underlies reduced exercise capacity

Several diseases including cardiac amyloidosis,²⁰ diabetes,²¹ metabolic syndrome,²² and heart failure of various aetiologies^{23,24} are associated with impaired cardiac contractility during exercise with overall exercise capacity subsequently reduced. However, the mechanisms responsible for diminished increases in exercise EF are understood to vary in each instance, including poor coronary blood flow resulting in challenged sub-endocardial perfusion²⁰ and dysfunctional myocardial energetics with reduced adenosine triphosphate production.^{21,22} Due to the structural remodelling with thicker myocardial walls in the LV in those born preterm,¹³ they may have greater susceptibility to subtle, subclinical deficits in coronary blood flow, though this

Table 3 Multiple linear regression models of per cent of predicted peak VO₂, HRR₁, and HRR₂ in preterm-born young adults

Preterm-born young adults (n = 47)				
	B	95%CI	Beta	P-value
Per cent of predicted peak VO ₂				
Per cent of predicted FVC	0.176	-0.634 to 0.983	0.132	0.634
Vent. reserve (%)	-0.234	-0.883 to 0.415	-0.205	0.435
EFA40% (%)	1.269	-0.208 to 2.746	0.538	0.084
Per cent of predicted peak VO ₂				
Per cent of predicted FVC	0.031	-0.905 to 0.967	0.023	0.943
Vent. reserve (%)	-0.270	-0.886 to 0.345	-0.236	0.354
EFA60% (%)	1.162	-0.498 to 2.823	0.506	0.152
HRR ₁ (b.p.m.)				
Per cent of predicted FVC	0.227	-0.106 to 0.561	0.367	0.159
EFA60% (%)	-1.099	-1.673 to -0.525	-1.030	0.002
HRR ₂ (b.p.m.)				
Per cent of predicted FVC	0.170	-0.293 to 0.633	0.198	0.432
EFA60% (%)	-1.405	-2.203 to -0.608	-0.948	0.003

Bold P values are statistically significant ($P < 0.05$). FVC represents forced vital capacity; EFA40%, ejection fraction change from rest to 40% exercise intensity; EFA60%, ejection fraction change from baseline to 60% exercise intensity; B, unstandardized regression coefficient.

would require further study to assess. In this study, however, we did observe elevated low-density lipoprotein cholesterol, triglycerides, plasma glucose, plasma insulin, and insulin resistance in the preterm group compared with controls.¹⁵ Other previous studies of preterm-born individuals have reported similar findings of impaired glucose and lipid metabolism in varied preterm-born study groups.^{13,25} Accordingly, while myocardial energetics were not assessed in this work, the presence of subclinical alterations in glucose and lipid metabolism suggests that research into the role of myocardial energetics as a contributing factor to the impaired systolic response to exercise in preterm-born subjects is warranted.

Pulmonary physiology does not explain reduced preterm exercise capacity

We observed that impaired LV systolic response to exercise stress was associated with diminished exercise capacity in normotensive preterm-born young adults. However, because lung development continues into the third trimester, pulmonary function is often considered a risk area for deficits and vulnerability in preterm-born infants. In particular, the saccular and alveolar phases of development may be interrupted by premature delivery. In spite of this, spirometric values in our preterm-born group were similar to controls. Interestingly, this comports with a growing body of evidence that moderate to late preterm birth may be less impacting of pulmonary capacity later in life than once thought.²⁶

In this study, the similar pulmonary volume and dynamic function between study groups and results from the multivariable analysis indicate that change in EF from rest to exercise, rather than mechanical ventilation, predicted exercise capacity. The elevated VE/VCO₂ slope in the preterm group indicates that exercise was limited by cardiac function. Such elevations most commonly occur secondary to deficits in cardiac performance, which drive a compensatory response of increased ventilation.²⁷ This ventilation perfusion mismatch is

common in heart failure, where exercise VE/VCO₂ slopes often exceed a value of 30. In the otherwise healthy, preterm-born young adults in this study, the presence of this compensatory increased ventilatory response relative to CO₂ production and O₂ uptake, combined with typical ventilatory reserve margin (23.3%) are consistent with cardiac limited exercise performance.²⁷

Systolic myocardial impairment and delayed heart rate recovery

In this article, we have shown that HRR is reduced at 2 min after exercise in normotensive, moderately preterm-born young adults and that this associated with impaired systolic function during exercise rather than pulmonary function. The metaboreflex is well known for its effect on blood pressure but the effect on post-exercise heart rate remains controversial.²⁸ However, it seems plausible that systolic myocardial impairment in the preterm-born young adults results in hypoperfusion of metabolite eliminating organs, and thus the clearance and deactivation of the metaboreflex are delayed. Furthermore, delayed HRR is commonly seen in heart failure populations⁶ and metaboreflex modulation plays a key role in the 'muscle hypothesis'.²⁹ As Crisafulli³⁰ explains, the normal haemodynamic response to metaboreflex activation is, among other factors, an increase in arterial blood pressure achieved by increased cardiac contractility boosting stroke volume to maintain adequate muscle perfusion. In absence of normal contractile reserves, exaggerated peripheral vasoconstriction is required to sustain pressure, which increases systemic vascular resistance.³⁰ The increased afterload further impairs stroke volume which exacerbates muscle perfusion deficits and ultimately produces early fatigue. Whether such a mechanism exists in conjunction with preterm birth would require detailed future research. However, Crump *et al.*³ recently reported an association of preterm birth with ischaemic heart disease even in those born moderately preterm, which further suggests this mechanism might exist.

To our knowledge, apart from one study comparing HRR after sub-maximal exercise,¹⁰ only two other studies investigated HRR after peak exercise testing. The first was conducted on healthy very preterm-born adolescents⁹ and the second on healthy very preterm-born young adults.⁸ In contrast to our study, they found reductions in HRR after the first minute. Furthermore, HRR remained lower in the very preterm young adults of the second study after adjusting for maximal oxygen consumption to control for fitness level, which was not the case for the moderately preterm-born individuals in our study. Whether this discrepancy indicates more severe or additional impairments in more premature populations remains to be determined as methodological differences (active vs. passive recovery) hinder direct comparisons.³¹ Interestingly, the mentioned second study also found that hypoxia did not further slow-down HRR in very preterm young adult individuals compared with controls. This result may support the muscle hypothesis, as compensatory increases in stroke volume or vasoconstriction, to counter the oxygen drop, may already be working at their physiological limits to compensate for the reduced contractile reserve. Although HRR is widely used as a prognostic marker,⁶ the underlying autonomic mechanisms, especially in preterm populations, are incompletely understood, and thus further research is needed to elucidate autonomic function contributions to differences in HRR.

Clinical implications

Overall, the moderately premature gestational age of the preterm-born group makes our results more broadly relevant, given that this reflects the demographic of the majority of preterm births.³² The reduced association between physical activity and exercise capacity in the preterm group suggests that prematurity may alter cardiovascular adaptation to aerobic training and require tailored approaches to lifestyle intervention, though randomized control trials will be needed to assess this. The finding that a lower myocardial functional reserve seen in those young adults born preterm explains a significant proportion of the lower peak VO_2 and slower HRR adds further clinical relevance to the altered preterm cardiac phenotype. In addition, peak VO_2 and HRR in this population may be important and beneficial for future risk stratification and treatment monitoring, acting as surrogate measures of reduced myocardial functional reserve. Continued investigation and longitudinal follow-up will be necessary to more fully understand the clinical implications of these findings.

Study limitations

We designed YACHT as an observational study around a single, full-day study visit. As the exercise stress testing came towards the end of the visit in order not to confound other measures, not all individuals were willing to continue with the sub-maximal exercise testing and echocardiography stress testing component. Also, due to challenges with imaging quality, not all echocardiography scans were of sufficient quality for analysis. Finally, the overall sample size was modest with a lower percentage of males in the preterm cohort, and therefore, analyses were adjusted for sex where appropriate. Nevertheless, the dataset was sufficiently powered to make between-group comparisons for the YACHT primary outcome of cardiopulmonary aerobic fitness as well as LVEF at rest, 40%, and 60% of maximal exercise capacity using echocardiography.

There was a wide gestational age range in our participants with the majority of our preterm-born individuals born moderate to late preterm, which reflects the demographic of the general preterm population.³² Although this makes our findings more relevant to a larger proportion of the population, larger studies will be needed to fully explore to what extent the severity of the LV systolic response to exercise is altered specifically in very and extreme preterm-born adults, who are at greater risk of perinatal complications. Finally, while pulmonary function and objective measurement of physical activity (accelerometry) were included in this analysis, direct assessment of pulmonary gas diffusion and broader ranging lifestyle characteristics, including historical physical activity profile, were not assessed.

Conclusions

Preterm-born young adults have reduced exercise capacity and HRR, which relate to the preterm functional phenotype of reduced LV systolic response to physical exercise. Although below average, exercise capacity was within the normal range and did not reflect acute clinical deficits. Further research is needed to develop a more complete understanding of cardiopulmonary exercise capacity and function in the preterm-born population, as well as which mechanisms underlie reduced peak oxygen consumption and HRR in preterm individuals of varying degrees of prematurity.

Supplementary data

Supplementary data are available at *European Heart Journal - Cardiovascular Imaging* online.

Acknowledgements

The authors acknowledge support from the Oxford BHF Centre of Research Excellence and National Institute for Health Research (NIHR) Oxford Biomedical Research Centre. The views expressed in this article are those of the author and do not reflect the official policy or position of the United States Air Force, Department of Defense, or the U.S. Government.

Funding

Study funding was provided by a British Heart Foundation (BHF) project grant (PG/13/58/30397). A.J.L. is funded by a BHF Research Fellowship (FS/18/3/33292) and O.J.H. received funding from the U.S. Air Force Institute of Technology.

Conflict of interest: none declared.

References

1. Liu L, Oza S, Hogan D, Perin J, Rudan I, Lawn JE et al. Global, regional, and national causes of child mortality in 2000-13, with projections to inform post-2015 priorities: an updated systematic analysis. *Lancet* 2015;**385**:430-40.
2. Carr H, Cnattingius S, Granath F, Ludvigsson JF, Edstedt Bonamy A-K. Preterm birth and risk of heart failure up to early adulthood. *J Am Coll Cardiol* 2017;**69**: 2634-42.
3. Crump C, Howell EA, Stroustrup A, McLaughlin MA, Sundquist J, Sundquist K. Association of preterm birth with risk of ischemic heart disease in adulthood. *JAMA Pediatr* 2019;**173**:736-43.
4. Crump C, Sundquist K, Sundquist J, Winkleby MA. Gestational age at birth and mortality in young adulthood. *JAMA* 2011;**306**:1233-40.

5. Myers J, Prakash M, Froelicher V, Do D, Partington S, Atwood JE. Exercise capacity and mortality among men referred for exercise testing. *N Engl J Med* 2002;**346**:793–801.
6. Cole CR, Blackstone EH, Pashkow FJ, Snader CE, Lauer MS. Heart-rate recovery immediately after exercise as a predictor of mortality. *N Engl J Med* 1999;**341**:1351–7.
7. Edwards MO, Kotecha SJ, Lowe J, Watkins WJ, Henderson AJ, Kotecha S. Effect of preterm birth on exercise capacity: a systematic review and meta-analysis. *Pediatr Pulmonol* 2015;**50**:293–301.
8. Haraldsdottir K, Watson AM, Beshish AG, Pegelow DF, Palta M, Tetri LH et al. Heart rate recovery after maximal exercise is impaired in healthy young adults born preterm. *Eur J Appl Physiol* 2019;**119**:857–66.
9. Haraldsdottir K, Watson AM, Goss KN, Beshish AG, Pegelow DF, Palta M et al. Impaired autonomic function in adolescents born preterm. *Physiol Rep* 2018;**6**:e13620.
10. Karvonen R, Sipola M, Kiviniemi AM, Tikanmaki M, Jarvelin MR, Eriksson JG et al. Postexercise heart rate recovery in adults born preterm. *J Pediatr* 2019;**214**:89–95.
11. Lovering AT, Elliott JE, Laurie SS, Beasley KM, Gust CE, Mangum TS et al. Ventilatory and sensory responses in adult survivors of preterm birth and bronchopulmonary dysplasia with reduced exercise capacity. *Annals ATS* 2014;**11**:1528–37.
12. Pierpont GL, Adabag S, Yannopoulos D. Pathophysiology of exercise heart rate recovery: a comprehensive analysis. *Ann Noninvasive Electrocardiol* 2013;**18**:107–17.
13. Lewandowski AJ, Augustine D, Lamata P, Davis EF, Lazdam M, Francis J et al. Preterm heart in adult life: cardiovascular magnetic resonance reveals distinct differences in left ventricular mass, geometry, and function. *Circulation* 2013;**127**:197–206.
14. Lewandowski AJ, Bradlow WM, Augustine D, Davis EF, Francis J, Singhal A et al. Right ventricular systolic dysfunction in young adults born preterm. *Circulation* 2013;**128**:713–20.
15. Huckstep OJ, Williamson W, Telles F, Burchert H, Bertagnolli M, Herdman C et al. Physiological stress elicits impaired left ventricular function in preterm-born adults. *J Am Coll Cardiol* 2018;**71**:1347–56.
16. Bush A, Cramer D. Guidelines for the measurement of respiratory function. *Respir Med* 1994;**88**:798.
17. Quanjer PH, Tammeling GJ, Cotes JE, Pedersen OF, Peslin R, Yernault JC. Lung volumes and forced ventilatory flows. Report working party standardization of lung function tests, European Community for steel and coal. Official statement of the European Respiratory Society. *Eur Respir J Suppl* 1993;**16**:5–40.
18. Fletcher GF, Ades PA, Kligfield P, Arena R, Balady GJ, Bittner VA et al. Exercise standards for testing and training: a scientific statement from the American Heart Association. *Circulation* 2013;**128**:873–934.
19. Wasserman K. *Principles of Exercise Testing and Interpretation: Including Pathophysiology and Clinical Applications*. 4th ed. Philadelphia, PA and London: Lippincott Williams & Wilkins; 2005. xvi, p585.
20. Clemmensen TS, Eiskjær H, Mølgaard H, Larsen AH, Soerensen J, Andersen NF et al. Abnormal coronary flow velocity reserve and decreased myocardial contractile reserve are main factors in relation to physical exercise capacity in cardiac amyloidosis. *J Am Soc Echocardiogr* 2018;**31**:71–8.
21. Wahl MP, Scalzo RL, Regensteiner JG, Reusch J. Mechanisms of aerobic exercise impairment in diabetes: a narrative review. *Front Endocrinol (Lausanne)* 2018;**9**:181.
22. Fournier SB, Reger BL, Donley DA, Bonner DE, Warden BE, Gharib W et al. Exercise reveals impairments in left ventricular systolic function in patients with metabolic syndrome. *Exp Physiol* 2014;**99**:149–63.
23. Martens P, Verbrugge FH, Nijst P, Dupont M, Mullens W. Limited contractile reserve contributes to poor peak exercise capacity in iron-deficient heart failure. *Eur J Heart Fail* 2018;**20**:806–8.
24. Abudiyab MM, Redfield MM, Melenovsky V, Olson TP, Kass DA, Johnson BD et al. Cardiac output response to exercise in relation to metabolic demand in heart failure with preserved ejection fraction. *Eur J Heart Fail* 2013;**15**:776–85.
25. Parkinson JR, Hyde MJ, Gale C, Santhakumaran S, Modi N. Preterm birth and the metabolic syndrome in adult life: a systematic review and meta-analysis. *Pediatrics* 2013;**131**:e1240–63.
26. Kotecha SJ, Watkins WJ, Paranjothy S, Dunstan FD, Henderson AJ, Kotecha S. Effect of late preterm birth on longitudinal lung spirometry in school age children and adolescents. *Thorax* 2012;**67**:54–61.
27. Balady GJ, Arena R, Sietsema K, Myers J, Coke L, Fletcher GF et al.; Council on Epidemiology and Prevention. Clinician's Guide to cardiopulmonary exercise testing in adults: a scientific statement from the American Heart Association. *Circulation* 2010;**122**:191–225.
28. Michael S, Graham KS, Davis G. Cardiac autonomic responses during exercise and post-exercise recovery using heart rate variability and systolic time intervals—a review. *Front Physiol* 2017;**8**:301.
29. Coats AJ, Clark AL, Piepoli M, Volterrani M, Poole-Wilson PA. Symptoms and quality of life in heart failure: the muscle hypothesis. *Br Heart J* 1994;**72**(2 Suppl):S36–9.
30. Crisafulli A. The impact of cardiovascular diseases on cardiovascular regulation during exercise in humans: studies on metaboreflex activation elicited by the post-exercise muscle ischemia method. *Curr Cardiol Rev* 2017;**13**:293–300.
31. Carter R, Watenpugh DE, Wasmund WL, Wasmund SL, Smith ML. Muscle pump and central command during recovery from exercise in humans. *J Appl Physiol* 1999;**87**:1463–9.
32. Kaczmarczyk K, Wiszomirska I, Sturzmowicz M, Magiera A, Błażkiewicz M. Are preterm-born survivors at risk of long-term respiratory disease? *Theor Adv Respir Dis* 2017;**11**:277–87.