

Case report

Open Access

Extraabdominal fibromatosis in retroperitoneal space

Akira Kikkawa¹, Akira Kido^{*2,3}, Tsukasa Kumai³ and Toru Hoshida²

Address: ¹Department of Urology, Hannna Central Hospital, Nara, Japan, ²Divison of Orthopedics and Traumatology, Medical Center for Emergency and Critical Care, Nara Prefectural Nara Hospital, Nara, Japan and ³Department of Orthopedic Surgery, Nara Medical University, Nara, Japan

Email: Akira Kikkawa - hanna-ch@m3.kcn.ne.jp; Akira Kido* - akirakid@narmed-u.ac.jp; Tsukasa Kumai - kumakumat@aol.com; Toru Hoshida - thoshida@narmed-u.ac.jp

* Corresponding author

Published: 03 October 2004

Received: 25 May 2004

World Journal of Surgical Oncology 2004, 2:33 doi:10.1186/1477-7819-2-33

Accepted: 03 October 2004

This article is available from: <http://www.wjso.com/content/2/1/33>

© 2004 Kikkawa et al; licensee BioMed Central Ltd.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Background: Fibromatosis or desmoid tumor covers a broad spectrum of benign fibrous tissue proliferations. It is characterized by infiltrative growth and a tendency towards recurrence; however, unlike sarcoma, it never metastasizes.

Case presentation: We report on a case of extraabdominal fibromatosis originating from the retroperitoneal space in a 43-year-old woman. Seven years earlier she had undergone ureterolysis and ureteroureterostomy for ureteral obstruction. Computed tomography revealed a tumor between the iliocostalis and the psoas muscle. Histopathological evaluation revealed uniform proliferation of spindle cells, with a moderate amount of collagen fibers, suggesting extraabdominal fibromatosis (desmoid tumor). The tumor was surgically resected, and since then, the patient has remained asymptomatic without any restrictions of daily living activities and without any signs of tumor recurrence during the two-year follow-up.

Conclusions: Complete resection is the treatment of choice. Adjuvant therapy using non steroidal anti-inflammatory agents, tamoxifen, interferon, anti-neoplastic agents, and radiotherapy, either alone or in combination finds application for unresectable or recurrent cases.

Background

The term "fibromatosis" covers a broad spectrum of benign fibrous tissue proliferations, the biological behavior of which is similar to both benign fibrous lesions and fibrosarcoma. Like fibrosarcoma, fibromatosis is characterized by infiltrative growth and a tendency towards recurrence; however, unlike sarcoma, it never develops metastasis [1]. Therefore, the most important strategy is to prevent direct invasion into adjacent tissues. Extraabdominal fibromatosis principally originates from the connective tissue of muscles and the overlying fascia or aponeurosis. It may occur in a variety of anatomical locations, including the muscles of the shoulder, the chest

wall and back, thigh, and head and neck. However, solitary occurrence is rare in retroperitoneal space [1,2]. Here, we report on a case of extraabdominal fibromatosis in the retroperitoneum. Resection was successfully performed, and the patient has been tumor-free for two years after surgery.

Case presentation

A 43-year-old woman with a history of schizophrenia since 1982, and a history of hospitalization to help the patient to acquire social communication abilities, at the age of 23 presented with slight pain on her left flank and back. In 1995, she was treated with ureterolysis and



Figure 1

Contrast enhanced computed tomography showing the tumor location between iliocostalis and psoas muscles in retroperitoneal space.

ureteroureterostomy because of left-sided ureteral obstruction. Histological evaluation of the biopsy revealed benign fibrous tissue proliferations; however, no further evaluation and surgical excision was planned as her mental state was deteriorating. She was put on regular follow-up with computed tomography (CT) scans. In May 2002, she was referred from the psychiatric hospital to our Department of Urology, as the tumor tended to grow. CT scan with contrast enhancement revealed a tumor located between iliocostalis and psoas muscles in retroperitoneal space. The peripheral part of the tumor was enhanced, while the central part did not. The left paravertebral muscles around the tumor showed atrophy. The medial margin of the tumor was deformed by a left transverse process of the second lumbar spine, suggesting invasive behavior (Figure 1). Coronal magnetic resonance imaging (MRI) demonstrated the tumor to be located beneath the left kidney. The central part of the tumor was found to be of iso/low intensity in the T1-weighted phase and of hetero-

geneously high intensity in the T2-weighted phase. The marginal part showed very low intensity in both phases (Figure 2). In May 2002, needle biopsy was performed and revealed that the tumor consisted of well-proliferated spindle cells rich in collagen fibers, an observation that was inconsistent with the histological pictures made in 1995. In June 2002, resection was performed using a paraspinal approach. Although the tumor strongly adhered to adjacent tissues, including the urinary tract and peritoneum, it was marginally resected, including paravertebral muscles and part of the spine. As the kidney was less affected by the tumor, ablation posed no problem. On gross examination, the cut surface appeared homogeneously gray and glossy (Figure 3). Histologically, a uniform proliferation of spindle cells with a moderate amount of collagen fibers led to a diagnosis of extraabdominal fibromatosis in the retroperitoneal space (Figure 4). No adjuvant treatment was given and during the two years of




Figure 2
Coronal magnetic resonance imaging demonstrating the tumor beneath the left kidney.




Figure 3
Gross appearance of the resected tumor. The cut surface homogeneously appears gray and glossy.

follow-up, the patient has remained asymptomatic, with no restrictions of daily living. There were no clues as to recurrence of the tumor in computed tomography.

Discussion

Extraabdominal fibromatosis may occur in a variety of anatomic locations; the principle sites of the involvement are the shoulder, chest wall and back, thigh and head and neck. Origin of extraabdominal fibromatosis from any mesenchymal tissue is now well recognized [1,2]. Several authors have reported retroperitoneal fibromatosis in patients with familial adenomatous polyposis (Gardner

syndrome) [2,4-6], however, solitary occurrence of fibromatosis is very rarely reported [1,2,7-9].

Our patient did not have a family history and upper gastrointestinal endoscopy, colonoscopy, or ophthalmoscopy were normal suggesting that our patient may be negative for the syndrome. The exact histological origin of the tumor remains to be verified. The findings of CT suggested an origin from paravertebral muscles. Interestingly, this assumption was corroborated by a computed tomography performed in April 2000, which revealed that the previous tumor was located intramuscularly




Figure 4

Photomicrograph showing the tumor composed of uniform spindled proliferation with moderate account of collagen fibers (hematoxylin and eosin $\times 400$).

Principally, complete resection is the therapy of choice for this type of tumors [10]. Adjuvant therapy using non steroidal anti-inflammatory drugs (NSAIDs), tamoxifen, interferon, anti-neoplastic agents, radiation, and a combinations of these, have been reported for cases that are difficult to resect [1], the exact benefit offered by them is not known due to thin literature. Radiation therapy is accepted as an effective treatment after incomplete resection [11,12]. Recently, preoperative radiotherapy was reported to be useful for the local control [13].

In our case the tumor detection was delayed because the psychiatric status of our patient which has been unstable for several years. As wide resection of the tumor reduces

the risk of recurrence, an early diagnosis is required for this type of tumor, which is difficult as most of these patients are asymptomatic. While the silent area contains several vital organs, extraabdominal fibromatosis should be considered for the differential diagnosis for such a lesion.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

AKik and AKid performed the operation, are responsible for the clinical work and helped with the preparation of the manuscript. TK is the orthopedic consultant and

helped with the preparation and editing of the manuscript. TH coordinated and drafted the manuscript. All authors read and approved the final manuscript.

Acknowledgements

Written consent was obtained from the patient for publication. We thank Mr. Bernd Wuesthoff for editing the manuscript.

References

1. Weiss SW, Goldblum JR: **Fibromatosis**. *Soft tissue tumors* Edited by: Weiss SW, Goldblum JR. St. Louis, Mosby; 2001:309-346.
2. Campanacci M, Enneking WF: **Aggressive fibromatosis**. *Bone and soft tissue tumors* Edited by: Campanacci M, Enneking WF. Padova (Italy): Springer Verlag; 1999:925-936.
3. Cleeve LK: **Retroperitoneal Fibrosis, Tumors, and Cysts**. *Urologic Surgery* Edited by: Glenn JF. Philadelphia, Lippincott; 1991:404-413.
4. Gurbuz AK, Giardiello FM, Petersen GM, Krush AJ, Offerhaus GJ, Booker SV, Kerr MC, Hamilton SR: **Desmoid tumors in familial adenomatous polyposis**. *Gut* 1994, **35**:377-381.
5. Heiskanen I, Jarvinen HJ: **Occurrence of desmoid tumors in familial adenomatous polyposis and results of treatment**. *Int J Colorectal Dis* 1996, **11**:157-162.
6. Rodriguez-Bigas MA, Mahoney MC, Karakousis CP, Petrelli NJ: **Desmoid tumors in patients with familial adenomatous polyposis**. *Cancer* 1994, **74**:1270-1274.
7. Goldblum J, Fletcher JA: **Desmoid-type fibromatosis**. *World Health Organization classification, pathology, and genetics of tumors of soft tissue and bone* Edited by: Fletcher CDM, Unni KK, Mertens F. Lyon, IARC press; 2002:83-84.
8. Pignatti G, Barbanti-Brodano G, Ferrari D, Gherlinzoni F, Bertoni F, Bacchini P, Barbieri E, Giunti A, Campanacci M: **Extraabdominal desmoid tumor. A study of 83 cases**. *Clin Orthop* 2000, **375**:207-213.
9. Douglas PJ, Antonio NG, Ivy PA: **Local control of extra-abdominal desmoid tumors**. *J Bone Joint Surg Am* 1996, **78**:848-854.
10. Overhaus M, Decker P, Fischer HP, Textor HJ, Hirner A: **Desmoid tumors of the abdominal wall: A case report**. *World J Surg Oncol* 2003, **1**:11 [<http://www.wjso.com/content/1/1/11>].
11. Karakousis CP, Mayordomo J, Zografos GC, Driscoll DL: **Desmoid tumors of the trunk and extremity**. *Cancer* 1993, **72**:1637-1641.
12. Schulz-Ertner D, Zierhut D, Mende U, Harms W, Branitzki P, Wannenmacher M: **The role of radiation therapy in the management of desmoid tumors**. *Strahlenther Onkol* 2002, **178**:78-83.
13. O'Dea FJ, Wunder j, Bell RS, Griffin AM, Catton C, O'Sullivan B: **Pre-operative radiation therapy is effective in the treatment of fibromatosis**. *Clin Orthop* 2003, **415**:19-24.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

