

MEETING ABSTRACT

Open Access

Cutaneous vascular & sudomotor responses to heat-stress in smokers & non-smokers

Nicole E Moyen, Hannah A Anderson, Jenna M Burchfield, Matthew A Tucker, Melina A Gonzalez, Forrest B Robinson, Matthew S Ganio*

From 15th International Conference on Environmental Ergonomics (ICEE XV) Portsmouth, UK. 28 June - 3 July 2015

Introduction

As approximately one billion people worldwide are chronic smokers [1] it is important to determine smokers' thermoregulatory responses to heat-stress. Although local maximal vasodilation may be attenuated in smokers [2], skin blood flow responses during whole-body heat stress are unknown. Moreover, it is unknown if sweat rate is altered in smokers; theoretically the binding of nicotine to nicotinic acetylcholine receptors [2] may initiate an earlier onset of sweating during whole-body heat stress compared to non-smokers [3]. The purpose of this study was to compare cutaneous vascular and sudomotor responses to whole-body passive heat-stress between smokers and non-smokers.

Methods

Nine male chronic smokers [SMK; 10 (6) cigarettes/day for 11.8 (9.5) y; 26 (8) y; 177.7 (6.6) cm; 80.6 ± 21.1 kg] and 13 male non-smokers [N-SMK; 28 (9) y; 177.6 (6.8) cm; 77.2 (8.2) kg] were matched for age, height, body mass, and exercise habits (all $p > 0.05$). Subjects were passively heated via water-perfused suits until gastrointestinal temperature (T_{gi}) increased 1.5 °C. Local sweat rate (LSR) via ventilated capsule and cutaneous vasomotor activity (CVC) via Laser Doppler on the forearm were continuously recorded; blood pressure, heart rate, sweat gland activation (SGA), sweat gland output (SGO), T_{gi} , and mean-weighted skin temperature (T_{sk}) were taken at baseline and each 0.5 °C T_{gi} increase. LSR and CVC onsets and sensitivities were calculated with mean body temperature (T_b) = 0.9* T_{gi} + 0.1* T_{sk} [4].

Table 1 Mean (SD) CVC and LSR parameters on the forearm for SMK and N-SMK during passive heat stress

	Measurement	Smokers	Non-smokers
CVC	CVC onset (ΔT_b from baseline, °C)	0.31 (0.12)	0.61 (0.21)*
	CVC plateau (% of max)	68.4 (27.4)	68.4 (21.6)
	CVC sensitivity ($\Delta\%$ max per °C ΔT_b)	82.5 (46.2)	58.9 (23.3)
LSR	LSR onset (ΔT_b from baseline, °C)	0.35 (0.14)	0.52 (0.19)*
	LSR plateau ($\text{mg}\cdot\text{cm}^{-2}\cdot\text{min}^{-1}$)	0.79 (0.26)	1.00 (0.13)*
	LSR sensitivity ($\Delta\text{mg}\cdot\text{cm}^{-2}\cdot\text{min}^{-1}$ per °C ΔT_b)	0.60 (0.40)	0.63 (0.21)

*Significant difference between groups ($p < 0.05$).

Results

No differences existed between SMK and N-SMK for T_{gi} , T_{sk} , T_b , heart rate, mean arterial pressure, LSR, CVC, and SGA with each 0.5 °C T_{gi} increase (all $p > 0.05$). Overall, SGO tended to be lower in SMK than N-SMK [SMK = 5.94 (3.49) vs. N-SMK = 8.94 (3.99) $\mu\text{g}\cdot\text{gland}^{-1}\cdot\text{min}^{-1}$; $p = 0.08$].

Discussion

Smokers' CVC and LSR onsets occurred at an earlier T_b than non-smokers, possibly because heat stress enhances nicotine kinetics (*i.e.* binding of nicotine to nicotinic acetylcholine receptors; [2,3]). The lower LSR at plateau during whole-body heating might indicate a thermoregulatory impairment in young smokers, and is likely a result of decreased sweat gland output and not activation.

* Correspondence: msganio@uark.edu

Human Performance Laboratory, Department of Health, Human Performance, and Recreation, University of Arkansas, Fayetteville, AR, USA

Conclusion

Compared to non-smokers, smokers had an earlier onset but similar sensitivity (i.e. increase in response per increase in T_b) for sweating/cutaneous vasodilation. These data suggest that overall, most young chronic smokers' thermo-regulatory responses to whole-body passive heat stress are not impaired.

Published: 14 September 2015

References

1. Alwan A: **Global status report on noncommunicable diseases 2010.** World Health Organization; 2011.
2. Kilaru S, Frangos SG, Chen AH, Gortler D, Dhadwal AK, Araim O, Sumpio BE: **Nicotine: a review of its role in atherosclerosis.** *J Am Coll Surg* 2001, **193**(5):538-546.
3. Ogawa T: **Local effect of skin temperature on threshold concentration of sudorific agents.** *J Appl Physiol* 1970, **28**:18-22.
4. Stolwijk JA: **A mathematical model of physiological temperature regulation in man.** *National Aeronautics and Space Administration* 1971.

doi:10.1186/2046-7648-4-S1-A98

Cite this article as: Moyen *et al.*: Cutaneous vascular & sudomotor responses to heat-stress in smokers & non-smokers. *Extreme Physiology & Medicine* 2015 **4**(Suppl 1):A98.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

