

Research article

Open Access

Immunity related genes in dipterans share common enrichment of AT-rich motifs in their 5' regulatory regions that are potentially involved in nucleosome formation

Jesus Hernandez-Romano^{1,5}, Francisco J Carlos-Rivera², Heladia Salgado³, Hector Lamadrid-Figueroa⁴, Veronica Valverde-Garduño¹, Mario H Rodriguez¹ and Jesus Martinez-Barnetche*¹

Address: ¹Centro de Investigación sobre Enfermedades Infecciosas, Instituto Nacional de Salud Pública, Av. Universidad 655, Col Sta Maria Ahuacatitlan, CP 62508, Cuernavaca, Morelos, México, ²Independent Consultant on Computational Sciences, Cuernavaca, Morelos, México, ³Centro de Ciencias Genómicas, Universidad Nacional Autónoma de México, Av Universidad s/n Col Chamilpa 62210, Cuernavaca, Morelos, México, ⁴Dirección de Estadística, Centro de Investigación en Evaluación y Encuestas (CIEE), Instituto Nacional de Salud Pública, Av Universidad 655, Col Sta Maria Ahuacatitlan, CP 62508, Cuernavaca, Morelos, México and ⁵Ingeniería en Biotecnología, Universidad Politécnica del Estado de Morelos, Boulevard Cuauhnahuac 566, Col Lomas del Texcal, Jiutepec, Morelos, México

Email: Jesus Hernandez-Romano - jesushromano@hotmail.com; Francisco J Carlos-Rivera - javier@nediam.com.mx; Heladia Salgado - heladia@ccg.unam.mx; Hector Lamadrid-Figueroa - hlamadrid@insp.mx; Veronica Valverde-Garduño - vvalverde@insp.mx; Mario H Rodriguez - mhenry@insp.mx; Jesus Martinez-Barnetche* - jmbarnet@insp.mx

* Corresponding author

Published: 9 July 2008

Received: 11 March 2008

BMC Genomics 2008, 9:326 doi:10.1186/1471-2164-9-326

Accepted: 9 July 2008

This article is available from: <http://www.biomedcentral.com/1471-2164/9/326>

© 2008 Hernandez-Romano et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Background: Understanding the transcriptional regulation mechanisms in response to environmental challenges is of fundamental importance in biology. Transcription factors associated to response elements and the chromatin structure had proven to play important roles in gene expression regulation. We have analyzed promoter regions of dipteran genes induced in response to immune challenge, in search for particular sequence patterns involved in their transcriptional regulation.

Results: 5' upstream regions of *D. melanogaster* and *A. gambiae* immunity-induced genes and their corresponding orthologous genes in 11 non-melanogaster drosophilid species and *Ae. aegypti* share enrichment in AT-rich short motifs. AT-rich motifs are associated with nucleosome formation as predicted by two different algorithms. In *A. gambiae* and *D. melanogaster*, many immunity genes 5' upstream sequences also showed NFκB response elements, located within 500 bp from the transcription start site. In *A. gambiae*, the frequency of ATAA motif near the NFκB response elements was increased, suggesting a functional link between nucleosome formation/remodelling and NFκB regulation of transcription.

Conclusion: AT-rich motif enrichment in 5' upstream sequences in *A. gambiae*, *Ae. aegypti* and the *Drosophila* genus immunity genes suggests a particular pattern of nucleosome formation/chromatin organization. The co-occurrence of such motifs with the NFκB response elements suggests that these sequence signatures may be functionally involved in transcriptional activation during dipteran immune response. AT-rich motif enrichment in regulatory regions in this group of co-regulated genes could represent an evolutionary constrained signature in dipterans and perhaps other distantly species.

Background

Organismal complexity is dependent on the network that regulates gene expression, rather than the number of genes in its genome [1-3]. Thus, one of the biggest challenges in postgenomic research is understand the regulatory mechanisms controlling location, timing and intensity of gene expression.

Organisms are permanently sensing changes in their environment. Environmental agents activate cellular signaling pathways that lead to a rapid expression of specific genes to respond to changes. These pathways transmit their signal to specific transcription factors (TFs) which gain access to response elements (REs) located in promoter and enhancer regions of the corresponding gene [2] resulting in transcriptional activation. In eukaryotes these protein-DNA interactions occur in the context of a chromatin template within the cell nucleus. The fundamental unit of chromatin is the nucleosome, composed by a segment of 146 base pairs of double stranded DNA wrapped around a core of histone proteins [4]. Initially, nucleosomes were regarded as structures required for the packing of long DNA molecules into the cellular nucleus [5], but it is now clear that chromatin structure plays a central role in the regulation of gene expression [6-9]. At least three mechanisms have been proposed for the active role of chromatin in transcriptional regulation. First, by preventing TF binding to its cognate RE as revealed by the pioneering studies in the expression of PHO5 gene in response to phosphate starvation [10]. Secondly, wrapping DNA in nucleosomes may promote transcription by allowing closely adjacent RE access to their cognate TF [11,12]. Third, nucleosomes may approximate distant regulatory elements, as it occurs in the alcohol-dehydrogenase (Adh) promoter region of *Drosophila* [13].

Nucleosomes are located in preferred positions with respect to DNA sequence [14-21]. It has been shown that on a statistical level, groups of experimentally obtained nucleosomal sequences display periodicity in the occurrence of dinucleotides such as GG, TA, TG, and TT [14,15,20] or trinucleotides such as VWG ([G/C/A] [A/T]G) [19]. This periodicity tends to occur approximately every 10 bp, coinciding with one turn of the DNA chain and confers better bending properties required for wrapping DNA around the histone core. However, this periodicity is difficult to identify on individual nucleosomal sequences due to a low signal/noise ratio. The non-random distribution of nucleosomes suggests that some DNA sequences are more likely to form stable nucleosomes, and therefore nucleosome forming sequences could be predicted using computational methods based on the sequence features identified so far [20,22].

Immune responses are inducible phenomena resulting from a close relationship between the environment, pathogen signal detection systems and the gene expression machinery [23]. Upon pathogen recognition, several transduction pathways are activated leading to the activation of TFs that induce gene expression [24]. In *Drosophila melanogaster*, the Toll and Imd pathways converge in the activation of the NFκB/Rel-related TFs, Dif and Relish, respectively, which bind to NFκB REs located in the 5' upstream regions of antimicrobial peptide genes, thus promoting their transcription [25].

Understanding the transcriptional regulation mechanisms during insect immune response is of fundamental interest in biology, but also could provide the rational basis for developing strategies to control vector borne diseases. In this work, we describe that immunity genes induced upon immune challenge in *D. melanogaster* and *Anopheles gambiae*, the main African malaria vector, share an enrichment of AT-rich motifs in their 5' regulatory regions. Enrichment of AT-rich motifs was also observed in 10 additional non-melanogaster *Drosophila* species and *Aedes aegypti* immunity orthologs. These motifs are different to REs in terms of statistical frequency and length. Their occurrence correlates with predicted nucleosomal positions [18,20,22], suggesting that AT-rich motifs may be involved in chromatinization and transcriptional regulation of immunity related transcriptional gene modules in these insects.

Results

Regulatory regions of immunity-related co-expressed genes of *Anopheles gambiae* and *Drosophila melanogaster* induced upon immune challenge are enriched in AT-rich specific DNA motifs

We used public available and author provided microarray databases [26,27], coupled to bioinformatics analysis tools for regulatory sequences, to identify sequence patterns potentially involved in transcriptional regulation operating during immune response in *D. melanogaster* and *A. gambiae*.

Microarray data describing the temporal transcriptional profile for 13,196 *D. melanogaster* genes [27], and 2,883 *A. gambiae* genes [26] in response to various immune challenges were used to select genes with the following expression profiles: immunity induced, repressed and non-modified (Figure 1). The *Drosophila* microarray data were obtained using Oregon^R adult males challenged with *Escherichia coli* and *Micrococcus luteus* [27]. The *Anopheles* microarray data were obtained using the *A. gambiae* cell line 4a-3B challenged with several bacteria species or microbial products [26]. Gene groups were selected according to 1) gene ontology and Interpro assignments as well as induction after an immune challenge, 2) genes




Figure 1
Expression profiles of immunity, non-modified and repressed genes of *A. gambiae* (A) and *D. melanogaster* (B). In red color are shown the immunity (Im) induced genes, in black the non-modified (NM) genes and green the down-regulated (DR) genes. The expression values are $\log_2(f_2/f_1)$, as reported by Dimopoulos [26] and De Gregorio [27]. C) and D) are examples of co-expressed genes of *A. gambiae* in response to *P. aeruginosa* and *E. coli*, respectively. 4a-3B cells of *A. gambiae* were challenged with nine microbial stimuli and analyzed at 1, 4, 8, 12, 18 and 24 hours post-challenge (except for *P. aeruginosa*, for which the 8 h sample was omitted). For *D. melanogaster*, adult males were challenged by pricking the thorax with a needle dipped into a bacterial culture of *Escherichia coli* and *Micrococcus luteus* and analyzed at 1.5, 3, 6, 12 and 24 hours after challenge (HAC).

that were not modified upon immune challenges, and 3) down-regulated genes upon immune challenges. Two additional groups were added as controls: 4) randomly selected genes and 5) computer-randomly generated

sequences (artificial). Table 1 presents the number of analyzed genes in each group and Tables 2, 3, 4, 5, 6, 7 list the Ensembl or AnEST ID, gene description, and chromosomal location for the genes used in the analysis. The

Table 1: Number of 5' upstream sequences analyzed according expression pattern and functional class.

Organism	Immunity up-regulated	Down-regulated	Non modified	Random genes	Artificial
<i>Anopheles gambiae</i>	18	16	19	20	20
<i>Drosophila melanogaster</i>	36	29	32	30	30

Groups of *A. gambiae* and *D. melanogaster* genes selected for *in silico* analysis for searching potential DNA regulatory motifs. 2500 nucleotides were analyzed.

expression profiles of the three gene groups (immunity induced, repressed and non-modified) of both species are shown in Figure 1.

To investigate whether the 5' regulatory regions of immunity-related genes shared common DNA motifs, 2500 bp 5' upstream sequences (5'-US) were recovered using Biomart, of Ensembl [28] and analyzed for statistically overrepresented motifs of 2 to 8 nucleotides in length, using Oligo-Analysis, which is based on binomial distribution [29]. The background oligonucleotide frequencies were estimated calculating the relative frequencies of all possible oligonucleotides (ranging from 2 to 8 bp) within the 5'-US of 2500 bp of length of 13,166 *A. gambiae* or 13,172 *D. melanogaster* genes. Oligonucleotide occurrences were counted for each group of 5'-US and their statistical significance was estimated on the basis of the background frequencies. The significance index (sigocc) reflects the degree of overrepresentation of each motif on a logarithmic scale [30].

Analysis of 5'-US of *A. gambiae* immunity genes showed that the main motifs statistically overrepresented were those of 2, 3 and 4 nucleotides in length (Table 8). Similar to *A. gambiae*, *D. melanogaster* 5'-US of immunity genes showed an enrichment of 2, 3 and 4 letter motifs (Table 8 and Tables 9, 10, 11, 12, 13, 14). Motifs of these length in 5'-US of immunity genes showed higher sig_{occ} than those obtained in 5'-US of down-regulated and non-modified genes in both insects. For *A. gambiae* immunity genes, the highest score for 4 pb motifs was of 11.26 (ATAA), versus 2.59 (AAAA) and 2.46 (CGAC) in down-regulated and non-modified genes, respectively (Table 9). Motifs of 3 bp and 2 bp in length also present the highest sig_{occ} in immunity induced *A. gambiae* genes: 6.11 for the AAA motif in immunity genes, versus 0.77 for the same motif in down-regulated genes, without 3 bp motif in non-modified genes (Table 11); 20.99 for AA motif in immunity genes, versus 0.72 and 1.73 for CG in down-regulated and non-modified genes, respectively (Table 13). Similar results for 2, 3 and 4 bp motifs were obtained in *D. melanogaster* 5'-US (Tables 10, 12 and 14). These observations underline the high overrepresentation of certain motifs in 5'-US of

Table 2: Immunity-related genes from *A. gambiae*.

Gene ID*	Description	Chromosome Name	Band
AGAP000182	Serine protease domain (IPR001314)	X	4A
AGAP000694	Antimicrobial peptide Cecropin (IPR000875)	X	1C
AGAP001212	Peptidoglycan recognition protein long class (PGRP-LB)	2R	7A
AGAP002457	Chitin binding domain, Glycoside hydrolase (IPR002557, IPR001579)	2R	12B
AGAP003960	Serine protease domain, Gastrulation defective precursor (IPR001314, IPR001254)	2R	17A
AGAP004455	Gram negative binding protein subgroup B (GNBPB1)	2R	19A
AGAP005246	Serpin	2L	21E
AGAP005848	Fibrinogen domain (IPR002181)	2L	23A
AGAP006348	LRR Toll	2L	24B
AGAP007036	Leucine-rich repeat (IPR001611)	2L	26C
TCLAG170406	Cactus	3R	29C
TCLAG170981	Phosphotyrosine interaction region (IPR006020)	3R	30E
AGAP008645	Putative infection responsive (Gambicin)	3R	31A
TCLAG168975	Dopa Decarboxylase isoform I	3R	33B
AGAP010133	Scavenger receptor class B Croquemort type	3R	37A
AGAP010816	Thioester-Containing Protein (TEP3)	3L	39C
AGAP010833	CLIP-domain serine protease subfamily B (CLIPB14) (IPR001254, IPR006604)	3L	39C
AGAP011197	Fibrinogen	3L	41D

*According to Ensembl [28] and AnoEST [67].

The Ensemble Gene ID or AnoEST cluster ID, description, chromosome and band location are showed. For genes without description, InterPro annotation is showed.

Table 3: Immunity-related genes from *D. melanogaster*.

Ensembl Gene ID	Description	Chromosome Name	Band
CG10146	Attacin-A precursor	2R	51C
CG10794	Diptericin B	2R	55F
CG10810	Drosomycin precursor	3L	63D
CG10812	drosomycin-5	3L	63D
CG10816	Drosocin precursor	2R	51C
CG11331	Serpin-27A CG11331-PA	2L	26F
CG11709	Peptidoglycan-recognition protein-SA precursor (Protein semmelweis)	X	10C
CG11992	Nuclear factor NF-kappa-B p110 subunit (Relish protein) (Rel-p110)	3R	85C
CG12763	Diptericin precursor	2R	55F
CG1365	Cecropin-A1/A2 precursor	3R	99E
CG1367	Cecropin-A1/A2 precursor	3R	99E
CG1373	Cecropin-C precursor	3R	99E
CG1385	Defensin precursor	2R	46D
CG14704	Peptidoglycan-recognition protein-LB precursor	3R	86E
CG14745	Peptidoglycan-recognition protein-SC2 precursor	2R	44E
CG15066	Immune-induced peptide 23 precursor (DIM-23)	2R	55C
CG18106	Immune-induced peptide 2 precursor (DIM-2)	2R	55C
CG18108	Immune-induced peptide 1 precursor (DIM-1)	2R	55C
CG18279	Immune-induced peptides precursor (DIM-10; DIM-12; DIM-13; DIM-24)	2R	50A
CG18372	Attacin-B precursor	2R	51C
CG1857	Necrotic	2R	42F
CG1878	Cecropin-B precursor	3R	99E
CG4432	Peptidoglycan-recognition protein-LC	3L	67B
CG4437	Peptidoglycan-recognition protein-LF (PGRP-like protein)	3L	67B
CG4740	Attacin-C precursor	2R	50A
CG5490	Protein toll precursor	3R	97D
CG5848	NF-kappa-B inhibitor cactus.	2L	35F
CG5974	Probable serine/threonine-protein kinase pelle	3R	97E
CG6134	Protein spaetzle precursor	3R	97E
CG6667	Embryonic polarity protein dorsal	2L	36C
CG6794	Dorsal-related immunity factor Dif	2L	36C
CG7496	Peptidoglycan-recognition protein-SD precursor.	3L	66A
CG7629	Attacin-D	3R	90B
CG8175	Metchnikowin precursor	2R	52A
CG8846	Thor CG8846-PA	2L	23F
CG9681	Peptidoglycan-recognition protein-SBI precursor	3L	73C

these insect immunity genes. Interestingly, the motifs sequences with the highest scores in 5'-US of immunity genes were the same in both organisms: TA, AA, AT, AAA and ATAA (Table 8).

The sigocc for the ATAA motif in 2500 pb 5'-US of *A. gambiae* immunity genes was 11.26, indicating that this four base-pairs motif is expected to occur in one of 1011.26 groups with similar numbers of sequences of the same length of random sequences. In comparison, the ATAA motif was not present in non-modified and down-regulated genes of *A. gambiae* (Table 15 and Table 9). The ATAA motif in 5'-US *D. melanogaster* immunity genes had a sigocc of 14.21, versus 2.2 in down-regulated genes, and was absent in non-modified genes (Table 15 and Table 10). Similar results were obtained for the conserved motifs AAA, AA, AT and TA. These results indicate that sta-

tistical overrepresentation of these motifs is specific of immunity genes 5'-US (Table 15).

In agreement to the Oligo-Analysis, bootstrapping analysis using POBO [31] confirmed that the average occurrence of the conserved motifs TA, AA, AT, AAA and ATAA, was significantly higher ($p < 0.0001$) in immunity promoters compared to the whole genome, the non-modified genes, down-regulated genes and the random sequence sets for both insects (Figure 2).

As a consequence of AT-rich motifs over-representation, a slight increase in AT% content was observed in 5'-US immunity genes, the average AT% for *D. melanogaster* immunity genes was of 60.4%, versus 57.6% and 57.2% for non-modified and down-regulated genes, respectively, these differences were not significant ($p > 0.001$), indicat-

Table 4: Non-modified genes from *A. gambiae*.

Gene ID*	Description	Chromosome Name	Band
AGAP000120	Cytoskeleton-associated proteins (CAP-Gly)	X	4B
AGAP000528	HMG-I and HMG-Y, DNA-binding Basic-leucine zipper (bZIP) transcription factor	X	2B
AGAP012290	Transporter	X	2B
AGAP000551	Dehydrogenase, E1 component, Transketolase, central region	X	2B
AGAP000855	No description	X	5A
AGAP002378	Fumarate lyase, Delta crystallin	2R	12B
AGAP003119	Translation initiation factor 4C (1A)	2R	14B
AGAP004295	DTW domain	2R	18D
AGAP005032	Calreticulin/calnexin	2L	21A
AGAP005429	WD-40 repeat	2L	22B
AGAP005558	Peptidase M16, C-terminal	2L	22C
AGAP006084	Antifreeze protein, type I, Ubiquitin system component Cue	2L	23C
AGAP006729	Domain of unknown function DUF1907	2L	25C
AGAP009156	BRICHOS	3R	33C
AGAP009547	Endoplasmic reticulum targeting sequence, Torsin	3R	34C
TCLAG188271	Neutral zinc metalloproteases, zinc-binding region signature	X	1D
TCLAG158277	Leucine aminopeptidase-related (PTHR11963)	2R	15D
TCLAG153172	RNA 3'-terminal phosphate cyclase, insert region	2L	20C
TCLAG187751	Malic oxidoreductase	X	4A

*According to Ensembl [28] and AnEST [67].

ing that AT-rich motif over-representation was not due to a significant increase in AT % that could lead to a random AT-rich motifs enrichment. Similar results were obtained for *A. gambiae* (Figure 3).

5' upstream regions of immunity-related orthologous genes in the genus *Drosophila* and in *Aedes aegypti* are also enriched with AT-rich motifs

In order to determine if other related dipteran species share the same motifs identified in *A. gambiae* and *D. melanogaster*, 5'-US of orthologous immunity genes of other *Drosophila* species and another *Culicidae* family member, *Aedes aegypti* were analyzed, regardless their transcriptional profile. Orthologous genes from those initially selected from *D. melanogaster* expression profiles were selected from the recently sequenced eleven *Drosophila* species [32,33]. Tables 16 and 17 show the list of orthologous genes present in each *Drosophila* species and *Ae. aegypti*. 5'-US from these genes were screened with Oligo-Analysis. Figure 4 shows the results obtained for the 4bp motifs TATA, AAAA, ATAA, AAAT and TTAA, which were statistically over-represented in the majority of 5'-US of immunity orthologous genes of 12 *Drosophila* species, *A. gambiae* and *Ae. aegypti*. We observed some phylogenetic correlations for some motifs. The most prevalent motif and with the highest sig_{occ} scores was the TATA motif, which was within the best ranked for most *Drosophila* species (9/12), but absent in *Anopheles* and *Aedes*. The AAAA motif was also highly ranked among some *Drosophila* species but not in mosquitoes. The ATAA motif was highly ranked in *D. melanogaster*, *D. simulans* and *A. gambiae*, but absent in other *Drosophilas* and *Ae. aegypti*. Finally, motifs

such as TTAA and AAAT were highly ranked in mosquitoes only. Intriguingly, *D. persimilis* did not show any enrichment of AT-rich motifs at all, regardless its close genetic distance to species displaying clear AT-rich motif enrichment. It should be noted that Sig_{occ} values for the non-melanogaster sequences and *Ae. aegypti* were considerably lower than those observed in *A. gambiae* and *D. melanogaster* (Table 18. See additional file 1).

AT-rich tetrads are associated with high nucleosomal potential

Once observed that AT-rich motif enrichment was a general feature of 5'-US of immunity genes in several dipteran species, we evaluated the association of some of these motifs with predicted nucleosomal sites. Experimentally stable nucleosomes in mouse have AT-rich motifs, including the AA [20], TA [14,20], TATA and ATAA motifs [18]. The ATAA motif include the three 2 bp motifs statistically over-represented in immunity genes (AA, TA and AT) of *A. gambiae* and *D. melanogaster*. Taking into account the highly conserved nucleosomal structure and given that the ATAA motif was enriched in both *D. melanogaster* and *A. gambiae*, as well as in other *Drosophila* species, we hypothesized that the ATAA motif could also participate in nucleosome formation in dipteran immune response genes. Some algorithms have been developed to predict the chromatin structure from sequence [20,22,34]. The RECON algorithm uses experimentally determined nucleosomal sequences coupled to Monte Carlo methods and discriminant analysis of dinucleotide frequencies [22]. It searches for a partition of non-overlapping regions in the nucleosomal sequences that provides the maximal value

Table 5: Non-modified genes from *D. melanogaster*

Ensembl Gene ID	Description	Chromosome Name	Band
CG10062	Peptidase M28	2R	56D
CG10203	Eggshell protein, Zinc finger, CCHC-type, RNA-binding region RNP-1 (RNA recognition motif)	2L	27C
CG11489	Serine/threonine protein kinase, active site	3L	79D
CG12929	No description	2R	45F
CG13046	No description	3L	72D
CG13230	No description	2R	47D
CG13329	Histone H3, Dopamine D4 receptor	2R	50A
CG14212	HAD-superfamily subfamily IB hydrolase, hypothetical 1	X	18D
CG14218	Vinculin/alpha-catenin	X	18D
CG14332	No description	3R	90A
CG14962	Zinc finger, C2H2-type	3L	63B
CG15016	Probable mitochondrial 28S ribosomal protein S6	3L	64B
CG15458	Putative 60S ribosomal protein L33	X	19D
CG15526	No description	3R	99D
CG15824	No description	2L	21E
CG17105	Eggshell protein	2L	32A
CG2076	BAX inhibitor related (PTHR23291)	X	10A
CG3723	Peptidase, cysteine peptidase active site, Dynein heavy chain, N-terminal, ATPase associated with various cellular activities	3R	93E
CG4148	Zinc finger, C2H2-type, HMG-I and HMG-Y, DNA-binding	2L	35D
CG4550	Opsin, KiSS-1 peptide receptor	3R	92B
CG4733	Calcium-binding EF-hand, Antifreeze protein, type I	3R	92B
CG5179	Cyclin-dependent kinase 9	2R	58F
CG5242	mitochondrial ribosomal protein L40	3R	86E
CG5675	Aminoacyl-tRNA synthetase, class I, Phosphotyrosine interaction region	X	16B
CG6253	60S ribosomal protein L14	3L	66D
CG6745	tRNA pseudouridine synthase D, TruD	3L	66D
CG6982	Claudin tight junction protein, Voltage-dependent calcium channel gamma	3R	94C
CG7173	Protease inhibitor, Kazal-type	3L	78D
CG7283	60S ribosomal protein L10a-2	3L	68E
CG9029	Orphan nuclear receptor, HMR type,	2L	26A
CG9091	Probable 60S ribosomal protein L37-A	X	13B
CG9961	Phosphoglycerate kinase	2L	23A

Table 6: Down-regulated genes from *A. gambiae*.

Gene ID*	Description	Chromosome Name	Band
AGAP000385	no description	X	2C
AGAP000654	Ribosomal protein S30, ubiquitin	X	1C
AGAP000655	Ribosomal protein S11	X	1C
AGAP000719	Adenosylhomocysteinase	X	1A
AGAP000862	Glycoside hydrolase	X	5A
AGAP001604	Putative Tyr/Ser/Thr phosphatase	2R	8D
AGAP002499	Probable methylmalonate-semialdehyde dehydrogenase, Mitochondrial precursor	2R	12C
AGAP003521	Alpha-2-macroglobulin RAP, C-terminal	2R	15D
AGAP004993	EGF-like, laminin	2L	20D
AGAP006782	ADP, ATP carrier protein 1 (ADP/ATP translocase 1)	2L	25D
AGAP007406	Elongation factor 1 alpha	2L	27D
AGAP007524	ST7	2L	28A
AGAP008914	no description	3R	32C
TCLAG186387	no description	X	3D
AGAP000720	no description	X	1A
TCLAG168991	no description	3R	33B

*According to Ensembl [28] and AnoEST [67].

Table 7: Down-regulated genes from *D. melanogaster*.

Ensembl Gene ID	Description	Chromosome Name	Band
CG10467	Aldose 1-epimerase	3L	65A
CG10475	Peptidase S1 and S6, chymotrypsin	3L	65A
CG10621	Homocysteine S-methyltransferase	2L	37B
CG11236	D-amino acid oxidase	2L	27B
CG11854	Hormone binding, cysteine peptidase active site	3R	96C
CG12351	Trypsin delta/gamma precursor	2R	47F
CG12385	Trypsin theta precursor	2R	47F
CG12387	Trypsin zeta precursor	2R	47F
CG15096	Sugar transporter superfamily	2R	55F
CG15231	Immune-induced peptide 4 precursor (DIM-4)	2R	57B
CG18030	Peptidase S1 and S6, chymotrypsin/Hap	3R	99F
CG18179	Peptidase S1 and S6, chymotrypsin/Hap	3L	67C
CG18180	Peptidase S1 and S6, chymotrypsin/Hap	3L	67C
CG18255	Stretchin-Mlck, isoform E	2R	52D
CG4178	Larval serum protein I beta chain precursor (Hexamerin I beta)	2L	21E
CG4812	Peptidase S1 and S6, chymotrypsin/Hap	2R	50A
CG4950	Leucine-rich repeat	3L	72D
CG5932	Esterase/lipase/thioesterase	3L	77C
CG6580	Peptidase S1 and S6, chymotrypsin/Hap	3L	65A
CG7214	No description	2L	28C
CG7216	Adult cuticle protein I precursor (dACP-I)	2L	28C
CG7459	Copper transporter 1B	3R	84F
CG8562	Peptidase M14, carboxypeptidase A	3L	65F
CG8579	Peptidase S1 and S6, chymotrypsin/Hap	2R	44E
CG8867	Peptidase S1 and S6, chymotrypsin/Hap	2L	25B
CG8869	Peptidase S1 and S6, chymotrypsin/Hap	2L	25B
CG9259	No description	2L	39A
CG9466	Glycoside hydrolase, family 38	2L	29F
CG9468	Glycoside hydrolase, family 38	2L	29F

of the Mahalanobis distance that discriminates between nucleosomal and non-nucleosomal sequences. In this way, RECON determines the probability that a sequence forms nucleosomes and assign a nucleosomal potential value to each nucleotide according to the context of the sequence in which the nucleotide is immersed. Positive values of nucleosomal potential correspond to reliable predictions of nucleosome formation sites with a confidence level of $p < 0.05$ ($\alpha = 0.05$), nucleosomal potential of +1 corresponds to the best predictions.

Using RECON, ATAA motifs were preferentially associated to positive values of nucleosomal potential in all the biological groups, both in *A. gambiae* and *D. melanogaster* (Figure 5), supporting a possible role for this motif in nucleosome formation. From Figure 5 is evident that ATAA is associated with positive nucleosomal potential values independently of the group of biological 5'-US analyzed. As expected, immunity, down-regulated, non-modified and random selected 5'-US of *A. gambiae* and *D. melanogaster*, all have ATAA motifs, however, 5'-US of immunity genes have a significant increased number of ATAA motifs. The frequency of ATAA associated with positive nucleosomal potential in immunity 5'-US is higher

than non-modified ($p = 0.001$ both in *A. gambiae* and *D. melanogaster*), down-regulated ($p = 0.006$ in *A. gambiae* and $p = 0.012$ in *D. melanogaster*), random ($p = 0.005$ in *A. gambiae* and $p < 0.001$ in *D. melanogaster*) and non-biological (artificial) sequences ($p < 0.001$ in *A. gambiae* and *D. melanogaster*). Analyzing ATAA distribution per group of genes, more than 70% and 80% of all ATAA motifs in *A. gambiae* and *D. melanogaster*, respectively, were located within regions of positive nucleosomal potential values ($p < 0.001$), indicating a possible role of this motif in nucleosome formation.

Additional information derived from Figure 5 is that the combination of RECON with Oligo-Analysis results allows detection of a property inherent to biological sequences. The ATAA distribution with respect to nucleosomal potential values was utterly different between biological and non-biological (artificial) sequences for both insects. On the one hand, biological sequences had more ATAA than non-biological sequences (1065 ATAA motifs in 30 random biological sequences versus 613 in 30 random non-biological sequences). The majority of the biological ATAA motifs were associated to positive nucleosomal potential (872/1065 or 81.9% motifs in 30

Table 8: Motif overrepresentation in 5' upstream sequences (2,500 bp) of immunity-induced genes.

Organism	Motif length (bp)	Motif	sig _{occ}
Anopheles gambiae	2	AA	20.99
		TA	14.39
		AT	13.25
	3	AAA	6.11
		TCA	2.84
		AAC	2.17
	4	ATAA	11.26
		AATA	8.82
		TTAA	4.62
	5	AATAA	9.37
		ATAAA	5.1
		AATAC	3.7
	6	CAATAC	2.53
		AATAAT	2.19
		ATAATG	1.5
		AAAATAA	1.3
	7	ATTATTA	0.41
AAATAAA		0.33	
Drosophila melanogaster	2	AA	37.43
		TA	15.64
		AT	14.8
	3	AAA	22.3
		GAA	1.45
		AAG	0.95
	4	AAAA	16.61
		ATAA	14.21
		TAAA	9.59
	5	TAAAA	10.75
		AAAAT	8.29
		ATAAA	7.49
	6	TGATAA	5.97
		TAAAAA	5.64
		ATAAAA	5.36
	7	GAAAAAC	2.73
		TTAAAAA	2.39
CTTATCA		2.03	

First three motifs statistically overrepresented found with Oligo-Analysis for each motif length, in *A. gambiae* and *D. melanogaster* upregulated immunity genes. In bold are indicated motifs with high sig_{occ} conserved in both insects.

random biological sequences versus 208/613 or 33.9% in 30 random non-biological sequences). Additionally, the non-biological sequences presented an inverse distribution of ATAA motifs, with more ATAA motifs associated to negative values of nucleosomal potential (289/613 or 47.2% ATAA negatives versus 208/613 or 33.9% ATAA positives in 30 random non-biological sequences).

We also evaluated the association of ATAA and TATA motifs in *D. ananassae*, representing the non-melanogaster subgroup; *D. pseudoobscura* of the *obscura* group and *D. grimshawi*, the most phylogenetically distant species belonging to the Hawaiian *Drosophila*, as well as the

TTAA and AAAT motif in *Ae. aegypti* (non-drosophilid dipteran), with nucleosomal potential calculated by the RECON algorithm. As shown in Figure 6, both motifs in the drosophilid species analyzed are clearly associated to positive nucleosomal potential values ($p < 0.001$; Figure 6A–B). Similarly, both motifs analyzed in *Ae. aegypti* were associated to high nucleosomal potential values and the difference between AT-rich motifs with positive nucleosomal potential and AT-rich motifs with other values was statistically significant ($p < 0.001$) (Figure 6C).

ATAA motifs correlate with high Nucleosomal Occupancy p values (pNO)

Segal and col. [20] recently reported an algorithm to predict nucleosome positions that takes into account sequence composition and thermodynamic properties. Using a collection of nucleosome bound DNA sequences from yeast, chicken or human, they constructed probabilistic models that represent the DNA sequence preferences for nucleosome formation and assign a p value to each nucleotide of the analyzed sequence; this value indicates the probability that the position is occupied by a nucleosome (p of Nucleosomal Occupancy, pNO).

Applying the three models to all the groups of sequences analyzed from *A. gambiae* and *D. melanogaster*, we found that the ATAA motifs were associated with high pNO values when the sequences were analyzed using the yeast model (Figure 7). Distribution of ATAA was very similar to that obtained using RECON, showing a coincidence between two independent methods to predict the association of ATAA with nucleosomal positions. All the biological groups of analyzed sequences presented ATAA motifs associated largely with pNO > 0.8, however, immunity genes had a significant increased number of ATAA associated with pNO > 0.8 values in relation to other biological sequences ($p < 0.05$, except between immunity and down-regulated genes of *D. melanogaster*, where $p = 0.155$).

More than 85% of all ATAA motifs found in 5'-US of *A. gambiae* and *D. melanogaster* were associated with pNO > 0.8 ($p < 0.001$). For *A. gambiae* immunity genes, 88% (476/541) of ATAA motifs were associated with pNO > 0.8, a similar distribution was obtained for the other *A. gambiae* gene groups (Figure 7A). The difference between ATAA associated with pNO > 0.8 and ATAA associated with pNO < 0.5 or undefined values was statistically significant ($p < 0.001$), showing a clear correlation between ATAA and high values of pNO. In a similar way, 85.9% (1340/1560) of ATAA motifs in 5'-US *D. melanogaster* immunity genes had pNO > 0.8, with a significant difference with regard to ATAA with pNO < 0.5 or undefined values ($p < 0.001$), the distribution of ATAA in the other *D. melanogaster* groups of genes also was statistically significant ($p < 0.001$) (Figure 7B).

Table 9: Motifs of 4 bp reported by Oligo-Analysis in 5'-US (2500 pb) of *A. gambiae* genes.

Over-expressed immunity genes		Down-regulated genes		Non-modified genes	
Motif	sig _{occ}	Motif	sig _{occ}	Motif	sig _{occ}
ataa	11.26	aaaa	2.59	cgac	2.46
aata	8.82	aaat	1.38	ccgc	0.93
ttaa	4.62	caga	0.88	cggc	0.93
tcaa	4.02	cgcc	0.23	acgg	0.33
taaa	3.78			ccga	0.18
aatg	3.41				
atta	2.92				
aact	2.47				
atca	2.44				
atac	2.09				
aaat	1.77				
gtta	1.14				
tata	1.01				
attc	0.91				
acat	0.82				
aaaa	0.57				
aaag	0.37				
cata	0.36				
aaga	0.32				
gata	0.31				
atga	0.23				
ctta	0.18				
aaca	0.15				
attg	0.11				
atag	0.10				
taca	0.10				

The combination of Oligo-Analysis and pNO results also revealed a difference between biological and non-biological sequences. Thirty randomly selected biological sequences of *D. melanogaster* had 90% (959/1065) of ATAA motifs associated with pNO > 0.8 versus 55.1% (338/613) of ATAA motifs in 30 non-biological sequences, showing again a non-random distribution of biological ATAA motif and tagging it as part of a potential nucleosomal code (Figure 7B).

Surprisingly no ATAA motifs were found with pNO values between 0.8 and 0.5 (values that define the "medium p value" range, see methods) in any of the other gene groups analyzed in *A. gambiae* and *D. melanogaster* (Figure 7).

We evaluated the association to probability of nucleosomal occupancy (pNO) [20] of ATAA and TATA motifs in *D. ananassae*, *D. pseudoobscura* and *D. grimshawi*; and the TTAA and AAAT motif in *Ae. aegypti*. As shown in Figure 8, both motifs in the drosophilid species analyzed are clearly associated to pNO > 0.8 values ($p < 0.001$, Figures 8A–B). Similarly, both motifs analyzed in *Ae. aegypti* were associated to pNO > 0.8 values ($p < 0.001$, Figure 8C).

Taken together, we found a consistent tendency, demonstrated by two independent methods, showing that the AT-rich motif enrichment within a specific sequence context might favour nucleosome formation in immune genes of a wide variety of dipteran species.

A. gambiae and D. melanogaster 5'-US of immunity genes have NF κ B response elements located in the first 500 pb of their 5'-US

NF κ B transcription factors in both insects and vertebrates are involved in immune gene expression regulation [23,25]. Using MEME [35], NF κ B REs were identified in the 5'-US of immunity genes, but not in 5'-US of down-regulated, non-modified and random genes. Moreover, the NF κ B REs were enriched within the first 500 bp of the 5'-US of immunity genes, both in *A. gambiae* and *D. melanogaster* (Figure 9). In order to investigate if there is a functional and physical association between ATAA motifs and NF κ B motifs, the distribution of ATAA motifs was analysed with respect to the transcription initiation site (TIS). Although ATAA motifs were distributed along the whole sequence in all gene groups, the highest frequencies were found to be located within the -251 to -500 interval in immunity genes in both insects (Figure 10). We further

Table 10: Motifs of 4 bp reported by Oligo-Analysis in 5'-US (2500 pb) of *D. melanogaster* genes.

Immunity-induced genes		Down-regulated genes		Non-modified genes	
Motif	sig _{occ}	Motif	sig _{occ}	Motif	sig _{occ}
aaaa	16.61	ataa	2.20	caaa	0.65
ataa	14.21	ctaa	2.16	aaaa	0.31
taaa	9.59	ccag	1.03	aaac	0.18
aaat	9.15	ccca	1.00		
aata	7.66	atca	0.89		
tata	5.80	atgg	0.58		
gaaa	5.62	gata	0.48		
aatt	5.44	cgga	0.46		
atat	4.16	agat	0.38		
ttaa	3.28	taga	0.34		
atta	2.47	aatc	0.31		
aaag	2.34	gacc	0.18		
aatc	2.04	gaca	0.01		
aaac	1.70				
aaga	1.64				
tcaa	1.08				
agaa	0.98				
atca	0.66				
gtaa	0.63				
attc	0.52				
atga	0.26				
acaa	0.12				
caaa	0.02				

quantified ATAA frequency within ± 200 bp from the NF κ B site of immunity, NM and DR genes in both insects. In the case of *A. gambiae*, the frequency of ATAA motifs around NF κ B REs is significantly higher in immunity genes compared to both non-modified ($p < 0.01$) and down-regulated genes ($p = 0.048$). However, in the case of *D. melanogaster* we did not find significant differences in the ATAA frequency in relation to NF κ B RE among the gene groups ($p > 0.1$), although the tendency was equal to that of *A. gambiae* (Figure 11). The association between NF κ B REs to AT-rich motifs and possibly nucleosomes of

immunity genes of *A. gambiae* and *D. melanogaster* may function as a specific link between the chromatin structure and the remodelling machinery needed for the expression of immune response genes.

Discussion

In this work, we have documented that AT-rich motifs are over-represented in 5' upstream regions of immunity genes of mosquitoes and drosophilids. We documented also that the position of the AT-rich motifs is associated to nucleosomes as predicted by two different algorithms for

Table 11: Motifs of 3 bp reported by Oligo-Analysis in 5'-UR (2500 pb) of *A. gambiae* genes.

Over-expressed immunity genes		Down-regulated genes		Non-modified genes	
Motif	sig _{occ}	Motif	sig _{occ}	Motif	sig _{occ}
aaa	6.11	aaa	0.77	none	
tca	2.84	gac	0.05		
aac	2.17				
caa	1.59				
taa	1.43				
ata	0.72				
act	0.58				
aat	0.44				
aca	0.36				

Table 12: Motifs of 3 bp reported by Oligo-Analysis in 5'-US (2500 pb) of *D. melanogaster* genes.

Immunity-induced genes		Down-regulated genes		Non-modified genes	
Motif	sig _{occ}	Motif	sig _{occ}	Motif	sig _{occ}
aaa	22.3	cca	2.39	caa	1.7
gaa	1.45	acc	0.71	aaa	1.42
aag	0.95				
tca	0.89				

nucleosome positioning, pointing out to a possible role of this motif in the transcriptional regulation of these functionally related genes through modification of chromatin structure involving nucleosome positioning.

Previous reports have found that sequences that form extremely stable nucleosomes are enriched with AT motifs referred as TATA boxes, which in many cases included the ATAA motif [18]. When we correlated the positions of this motif with the output of two different algorithms that predict nucleosome positions [20,22], we found that this motif correlates almost exclusively with positions with a high probability to form nucleosomes, suggesting that the ATAA motif enrichment is a DNA sequence pattern associated to nucleosome formation in these functionally related immunity genes. The conservation of enrichment of AT-rich motifs in 5'-US of immunity-related genes of *Drosophilidae* and *Culicidae*, which diverged 250 million years ago [36], suggest that this common feature may be the result of evolutionary constrained epigenetic mechanism of transcriptional regulation in immune-responsive genes in dipterans. More studies are required to define if this could be part of a more general mechanism of regulation in metazoans. The case of *D. persimilis* represents a caveat for our attempt generalize the implications of our findings, however, we cannot exclude that the current status of the annotation of such genomes may affect the results.

There are two conflicting views about nucleosome formation: one establishes that nucleosomes can potentially be

formed anywhere in the genome regardless the sequence and therefore, it is not possible to predict sites for nucleosome formation [4]. The other proposes that nucleosomes are associated to certain DNA sequences or sequence patterns that have an effect on the bending properties of DNA during nucleosome formation [14,18]. This point of view has been gaining support in recent years due to the documentation of a great variability in the bending potential of DNA sequences [14,15,37] and therefore their capacity to form nucleosomes [18,38,39].

Two of the three two-letter motifs statistically over-represented in immunity promoters of *A. gambiae* and *D. melanogaster*, TA and AA, have been previously associated to nucleosome formation in human, yeast, chicken and mouse [14,15,20,21]. Additionally, the ATAA motif, which is associated to nucleosome positions, and includes the three motifs containing two letters with the highest scores in both organisms (AA, TA and AT), have been found in sequences that form stable nucleosomes [18]. Thus, AT-rich motifs in 5'-US regions of mosquitoes and drosophilid immunity genes could participate in the transcriptional regulation of genes induced by immune challenges in a different way to the typical response elements. In contrast to response elements, which can be functional single or in pairs in a promoter region, the AT-rich motifs are statistically enriched, with several copies distributed in a diffuse pattern through the promoter regions, suggesting its involvement in nucleosome formation. This diffuse sequence pattern of AT-rich motifs, different to discreet patterns displayed by response elements, represents a new

Table 13: Motifs of 2 bp reported by Oligo-Analysis in 5'-US (2500 pb) of *A. gambiae* genes.

Over-expressed immunity genes		Down-regulated genes		Non-modified genes	
Motif	sig _{occ}	Motif	sig _{occ}	Motif	sig _{occ}
aa	20.99	cg	0.72	cg	1.73
ta	14.39			cc	1.46
at	13.25				

Table 14: Motifs of 2 bp reported by Oligo-Analysis in 5'-US (2500 pb) of *D. melanogaster* genes.

Immunity-induced genes		Down-regulated genes		Non-modified genes	
Motif	sig _{occ}	Motif	sig _{occ}	Motif	sig _{occ}
aa	37.43	at	3.27	aa	2.97
ta	15.64	ta	2.25	at	0.11
at	14.8	cc	0.15		

insight on the role of DNA sequence context in transcriptional regulation.

Several reports have documented that genes with similar functions share similar nucleosomal occupancy patterns. Levitsky and col. [34], using the RECON algorithm to analyze distinct functional types of human promoters, found that tissue-specific gene promoters present higher nucleosomal potential than genes commonly expressed in many tissues (housekeeping genes). Segal and col. [20], using the Nucleosome Position Prediction algorithm to analyze different kinds of genomic sequences and gene sets biologically related, found that nucleosome occupancy varies depending of the analyzed genomic location type, and that groups of genes functionally related can be classified on the basis of their profiles of nucleosome occupancy in the open reading frames and intergenic regions. Recently, Lee and col. [40], using Hidden Markov Models to analyze experimentally obtained nucleosomes, also found a correlation between function and nucleosome occupancy. Each of these reports used a different method to analyze data sequences, and all found that nucleosomal sequences follow a distinctive pattern associated to the functionality of the genes.

In relation to RECON [22] and Nucleosome Positioning Prediction [20], it is important to note that none of these programs search for *a priori* defined motifs, the input for

both programs are biological nucleosomal sequences from which information is extracted.

It has been shown that gene expression co-regulation is highly conserved in eukaryotes, for example, *Saccharomyces cerevisiae* and *Caenorhabditis elegans*, which diverged 1500 million years ago, still share a group of co-regulated genes [41], so it is plausible that *Drosophilidae* and *Culicidae* which diverged only 250 million years ago also share groups of functionally related co-regulated genes. The enrichment of AT-rich motifs in groups of co-regulated genes involved in immune response could provide the basis for developing new tools for the identification of different functional gene modules based on the compositional context of non-coding regulatory DNA. However, the high sig_{occ} observed in manually curated datasets compared to the low sig_{occ} observed in automatically annotated datasets highlights the importance of accurate TIS for regulatory region analysis.

Insect immunity relies on innate defense mechanisms to combat pathogens. In *D. melanogaster*, the Imd and Toll pathways lead to the activation of Rel/NFκB transcription factors that control a substantial proportion of the transcriptionally modified genes in response to pathogen infection [42]. Many components of these pathways are conserved in *A. gambiae* and *Ae. aegypti* [43] and are also remarkably conserved in innate immunity signaling pathways in mammals (TLR and TNF-R signaling pathways,

Table 15: Overrepresentation of AT rich motifs is specific of immunity-induced genes

Organism	Group of genes	AA score	TA score	AT score	AAA score	ATAA score
<i>Anopheles gambiae</i>	Immunity induced	20.99	14.39	13.25	6.11	11.26
	Down-regulated	---	---	---	0.77	---
	Non-modified	---	---	---	---	---
	Random genes	0.42	---	---	2.51	---
	Artificial	---	12.11	---	---	---
<i>Drosophila melanogaster</i>	Immunity induced	37.43	15.64	14.80	22.30	14.21
	Down-regulated	---	2.25	3.27	---	2.2
	Non-modified	2.97	---	0.11	1.42	---
	Random genes	1.9	---	---	0.7	---
	Artificial	---	---	---	---	---

The overrepresented motifs are specific of immunity-related gene promoters of *Anopheles gambiae* and *Drosophila melanogaster*.




Figure 2

Bootstrap to verify the ATAA enrichment in 5'-US regions of immunity-induced genes of *A. gambiae* and *D. melanogaster*. In the vertical axis are the number of pseudoclusters having a given occurrence of the motif, in the horizontal axis are indicated the number of occurrences of the motif in pseudoclusters. The red curve represents ATAA enrichment in pseudoclusters generated from the complete genome of *A. gambiae* (**A**) or *D. melanogaster* (**B**), the blue curve represents ATAA enrichment in immunity-induced genes, the yellow and green curve represents ATAA enrichment in the non-modified genes group. Similar results were obtained for the TA, AA, AT and AAA motifs in both insects and versus the down-regulated genes.




Figure 3
Percentage of AT between the different groups of sequences of *A. gambiae* (A) and *D. melanogaster* (B).
 The difference of AT% between groups is not statistically significant ($p > 0.001$), therefore, the enrichment of ATAA motif is not due to a bias in sequence composition. NM: non-modified genes, DR: down-regulated genes. Standard deviation is showed.

respectively) [33,44-47]. The set of induced genes in both insects described here belong to the same functional group and many of them have NF κ B response elements within 500 pb upstream from the predicted transcription start site, the same location where functionally important NF κ B REs have been found in these and other insects [48-52]. Our findings indicate that besides being regulated by

NF κ B, the enrichment with the ATAA motif constitutes a particular pattern of chromatin structure involved in transcriptional regulation of these genes.

Interestingly, NF κ B transcription factors bind to their response elements even if they are packaged in a nucleosome [53]. Once bound to their response elements, NF κ B transcription factors can recruit chromatin remodeling complexes to expose other response elements and allow the formation of the initiation complex [54]. In the vertebrate immune system, chromatin structure is critical to establish Th1-Th2 differentiation through the action of specific transcription factor as GATA-3 and T-bet [55], and several cytokines possess nucleosomes located in their promoters which need to be removed to allow gene expression [56-59]. Thus, epigenetic phenomena such as histone modification (altered nucleosome conformation) [60] or remodeling of chromatin (change of nucleosome position) [61] are commonly a required step to achieve gene expression in response to external stimuli.

Based on the obtained results and previously reported information, we propose a model in which a subgroup of insect immunity genes remains silent in absence of an immune challenge due to nucleosome formation in their 5'-US regions. The presence of these nucleosomes occludes the access of transcription factors to REs involved in gene expression. After an immune challenge, the Toll and/or Imd pathways are activated which in turn lead to activation of Rel/NF κ B transcription factors, which are translocated to the nucleus and bind to their NF κ B REs and recruit chromatin modifying/remodeling factors that release DNA from nucleosomes allowing its interaction with the transcriptional machinery.

Functionally related genes could harbor in their regulatory region a regulatory code represented by the combination of REs plus, in some cases, particular short motifs associated to chromatin structure. This regulatory code functions like a lock, genes that need to be co-expressed will share the same lock, represented by REs organized in a similar way, or by specific REs associated to motifs that confer a distinctive chromatin structure. Cells are continuously sensing its environment and responding to adapt. The regulatory state of the cell, defined by the presence and state of activity of transcription factors [3], also changes continuously; this regulatory state represents the "key" needed to open the proposed lock. The active transcription factors present in a given time in the cell, determines the form of the "key" for the lock, and therefore, the class of promoters that will be open or closed. In the case of the immune genes studied here, we have identified evidence that is compatible with a potential regulatory unit involving chromatin structure (associated with ATAA), Rel/NF κ B transcription factors and NF κ B response ele-

Table 16: List of Drosophilid orthologous immunity genes used for over-representation analysis of AT-rich tetrads in 5' upstream regions (2000 bp)

Subgenus <i>Sophophora</i>						Subgenus <i>Drosophila</i>					
<i>melanogaster group</i>						<i>obscura group</i>	<i>willistoni group</i>	<i>virilis group</i>	<i>repleta group</i>	<i>Hawaiiian</i>	
<i>melanogaster</i>	<i>simulans</i>	<i>sechellia</i>	<i>yakuba</i>	<i>erecta</i>	<i>ananassae</i>	<i>pseudobscura</i>	<i>persimilis</i>	<i>willistoni</i>	<i>virilis</i>	<i>mojavensis</i>	<i>grimshawi</i>
FBgn0012042	<i>Dsim</i> \AttA	<i>Dsec</i> \AttA	<i>Dyak</i> \AttA			<i>Dpse</i> \GAI0109					
FBgn0030310	<i>Dsim</i> \PGR P-SA	<i>Dsec</i> \GM13088	<i>Dyak</i> \PGR P-SA			<i>Dpse</i> \GAI1152	<i>Dper</i> \GL26887	<i>Dwil</i> \GK25449	<i>Dvir</i> \GJ15950	<i>Dmojl</i> \GII6473	
FBgn0000276	<i>Dsim</i> \CecA1	<i>Dsec</i> \CecA1				<i>Dpse</i> \Cec1					
FBgn0000279	<i>Dsim</i> \CecC	<i>Dsec</i> \CecC	<i>Dyak</i> \CecC	<i>Dere</i> \CecC		<i>Dpse</i> \CecV					
FBgn0041581	<i>Dsim</i> \AttB					<i>Dpse</i> \GAI4910					
FBgn0000277			<i>Dyak</i> \CecA2			<i>Dpse</i> \Cec11			<i>Dvir</i> \Cec2A		
FBgn0025583						<i>Dpse</i> \GAI4796					
FBgn0011274	<i>Dsim</i> \Dif	<i>Dsec</i> \GM17131	<i>Dyak</i> \Dif	<i>Dere</i> \GG21748	<i>Dana</i> \GF15105	<i>Dpse</i> \GAI9867	<i>Dper</i> \GL18755	<i>Dwil</i> \GK24830	<i>Dvir</i> \GJ16233	<i>Dmojl</i> \GII6552	<i>Dgri</i> \GHI0078
FBgn0004240	<i>Dsim</i> \GD11417	<i>Dsec</i> \GM21923	<i>Dyak</i> \Dpt	<i>Dere</i> \GG21934	<i>Dana</i> \GF11124	<i>Dpse</i> \GAI1797	<i>Dper</i> \GL11494	<i>Dwil</i> \GK20931	<i>Dvir</i> \GJ19915	<i>Dmojl</i> \GII0150	<i>Dgri</i> \GH22131
FBgn0000462	<i>Dsim</i> \dl	<i>Dsec</i> \GM17128	<i>Dyak</i> \dl	<i>Dere</i> \GG21746	<i>Dana</i> \GF15103	<i>Dpse</i> \GAI9765	<i>Dper</i> \GL18753	<i>Dwil</i> \GK10525	<i>Dvir</i> \GJ16232	<i>Dmojl</i> \GII6541	<i>Dgri</i> \GHI0077
FBgn0035434	<i>Dsim</i> \dro5	<i>Dsec</i> \GM14562	<i>Dyak</i> \dro5	<i>Dere</i> \dro5							
FBgn0035976	<i>Dsim</i> \GD14166	<i>Dsec</i> \GM25129	<i>Dyak</i> \GE20818	<i>Dere</i> \GG15356	<i>Dana</i> \GF10680	<i>Dpse</i> \GA29295	<i>Dper</i> \GL18427	<i>Dwil</i> \GK23800	<i>Dvir</i> \GJ13383	<i>Dmojl</i> \GII2108	<i>Dgri</i> \GHI5170
FBgn0035977	<i>Dsim</i> \GD14168	<i>Dsec</i> \GM25131	<i>Dyak</i> \GE20821	<i>Dere</i> \GG15359	<i>Dana</i> \GF10681	<i>Dpse</i> \GAI8183	<i>Dper</i> \GL18430	<i>Dwil</i> \GK23822	<i>Dvir</i> \GJ13386	<i>Dmojl</i> \GII2111	<i>Dgri</i> \GHI5173
FBgn0034328		<i>Dsec</i> \GM19911	<i>Dyak</i> \GE13917	<i>Dere</i> \GG20976	<i>Dana</i> \GF12769	<i>Dpse</i> \GA24273	<i>Dper</i> \GL16707	<i>Dwil</i> \GK23237	<i>Dvir</i> \GJ22454	<i>Dmojl</i> \GII9391	<i>Dgri</i> \GH20222
FBgn0003717	<i>Dsim</i> \TI	<i>Dsec</i> \GM10345	<i>Dyak</i> \TI	<i>Dere</i> \GG11504	<i>Dana</i> \GF16456	<i>Dpse</i> \TI		<i>Dwil</i> \GK13544	<i>Dvir</i> \TI	<i>Dmojl</i> \GII2147	<i>Dgri</i> \GHI4238
FBgn0014018	<i>Dsim</i> \Rel	<i>Dsec</i> \Rel	<i>Dyak</i> \Rel	<i>Dere</i> \GG17396	<i>Dana</i> \GF18430	<i>Dpse</i> \GAI1317	<i>Dper</i> \GL12490	<i>Dwil</i> \GK14061	<i>Dvir</i> \GJ23481	<i>Dmojl</i> \GII0193	<i>Dgri</i> \GHI8237
FBgn0010381	<i>Dsim</i> \Drs	<i>Dsec</i> \GM14569	<i>Dyak</i> \GE21361	<i>Dere</i> \GG15135							
FBgn0043575	<i>Dsim</i> \PGR P-SC2	<i>Dsec</i> \GM21061	<i>Dyak</i> \GE19223	<i>Dere</i> \GG23381		<i>Dpse</i> \GAI3217		<i>Dwil</i> \GK21737	<i>Dvir</i> \GJ21836	<i>Dmojl</i> \GII8809	
FBgn0034329						<i>Dpse</i> \GAI4798					
FBgn0034407	<i>Dsim</i> \GD11419	<i>Dsec</i> \GM21924	<i>Dyak</i> \DptB	<i>Dere</i> \GG21935	<i>Dana</i> \GF11125	<i>Dpse</i> \GAI0563	<i>Dper</i> \GL11495	<i>Dwil</i> \GK20932	<i>Dvir</i> \GJ19917	<i>Dmojl</i> \GII9362	<i>Dgri</i> \GH22132
FBgn0010388	<i>Dsim</i> \Dro	<i>Dsec</i> \GM21566	<i>Dyak</i> \GE13605	<i>Dere</i> \GG20474	<i>Dana</i> \GF11338	<i>Dpse</i> \GAI0577	<i>Dper</i> \GL11457	<i>Dwil</i> \GK19342			
FBgn0028990	<i>Dsim</i> \GD22555	<i>Dsec</i> \GM14001	<i>Dyak</i> \GE18418	<i>Dere</i> \GG23599	<i>Dana</i> \GF14458	<i>Dpse</i> \GAI0926	<i>Dper</i> \GL25590	<i>Dwil</i> \GK18897	<i>Dvir</i> \GJ21412	<i>Dmojl</i> \GII4139	<i>Dgri</i> \GHI3574
FBgn0010385	<i>Dsim</i> \Def	<i>Dsec</i> \GM20557	<i>Dyak</i> \GE21830	<i>Dere</i> \GG25236	<i>Dana</i> \GF12914	<i>Dpse</i> \GAI2570	<i>Dper</i> \GL10559	<i>Dwil</i> \GK21361	<i>Dvir</i> \GJ22479	<i>Dmojl</i> \GII9416	<i>Dgri</i> \GH21581
FBgn0037906	<i>Dsim</i> \GD15111	<i>Dsec</i> \GM26096	<i>Dyak</i> \GE24618	<i>Dere</i> \GG17219	<i>Dana</i> \GF16643	<i>Dpse</i> \GAI3189	<i>Dper</i> \GL12268	<i>Dwil</i> \GK11791	<i>Dvir</i> \GJ10054	<i>Dmojl</i> \GII2456	<i>Dgri</i> \GHI5357
FBgn0033835						<i>Dpse</i> \GAI4871					
FBgn0002930	<i>Dsim</i> \nec	<i>Dsec</i> \GM20812	<i>Dyak</i> \nec	<i>Dere</i> \GG10764	<i>Dana</i> \GF12243	<i>Dpse</i> \GAI4995	<i>Dper</i> \GL10947	<i>Dwil</i> \GK21526	<i>Dvir</i> \GJ14938	<i>Dmojl</i> \GII0049	<i>Dgri</i> \GH20376
FBgn0000278	<i>Dsim</i> \CecB	<i>Dsec</i> \CecB	<i>Dyak</i> \GE10866	<i>Dere</i> \GG11950	<i>Dana</i> \GF16190						

Table 16: List of Drosophilid orthologous immunity genes used for over-representation analysis of AT-rich tetrads in 5' upstream regions (2000 bp) (Continued)

FBgn0041 579	<i>Dsim\AttC</i>	<u>Dsec\GM</u> <u>21465</u>	<u>Dyak\GE</u> <u>12538</u>	<u>Dere\GG</u> <u>20378</u>	<u>Dana\GF</u> <u>13689</u>			<u>Dwil\GK</u> <u>17822</u>	<u>Dvir\GJ21</u> <u>173</u>	<u>Dgri\GH2</u> <u>1628</u>	
FBgn0000 250	<i>Dsim\cact</i>	<u>Dsec\GM</u> <u>17910</u>	<u>Dyak\cact</u>	<u>Dere\GG</u> <u>24172</u>	<u>Dana\GF</u> <u>15324</u>	<i>Dpse\GA1</i> 9176	<i>Dper\GL</i> 14084	<u>Dwil\GK</u> <u>23715</u>	<u>Dvir\GJ17</u> <u>162</u>	<u>Dmojl\GI1</u> <u>4683</u>	<u>Dgri\GH1</u> <u>1429</u>
FBgn0010 441	<i>Dsim\pll</i>	<u>Dsec\GM</u> <u>10117</u>	<u>Dyak\pll</u>	<u>Dere\GG</u> <u>12125</u>	<u>Dana\GF</u> <u>18719</u>	<i>Dpse\GA1</i> 9272	<i>Dper\GL</i> 23050	<u>Dwil\GK</u> <u>22593</u>	<u>Dvir\GJ10</u> <u>484</u>	<u>Dmojl\GI2</u> <u>2298</u>	<u>Dgri\GH1</u> <u>6651</u>
FBgn0035 806	<i>Dsim\PGR</i> P-SD		<u>Dyak\PGR</u> P-SD	<u>Dere\GG</u> <u>14444</u>	<u>Dana\GF</u> <u>10655</u>	<i>Dpse\GA2</i> 0392	<i>Dper\GL</i> <u>26612</u>	<u>Dwil\GK</u> <u>23884</u>	<u>Dvir\GJ11</u> <u>481</u>	<u>Dmojl\GI1</u> <u>2249</u>	<u>Dgri\GH1</u> <u>4703</u>
FBgn0038 530	<u>Dsim\GD</u>	<u>Dsec\GM</u> <u>15251</u>	<u>Dyak\AttD</u>	<u>Dere\GG</u> <u>22395</u>	<u>Dana\GF</u> <u>17819</u>	<i>Dpse\GA2</i> 0491	<i>Dper\GL</i> <u>21518</u>	<u>Dwil\GK</u> <u>12085</u>	<u>Dvir\GJ22</u> <u>662</u>	<u>Dmojl\GI2</u> <u>4598</u>	<u>Dgri\GH1</u> <u>8099</u>
FBgn0014 865	<i>Dsim\Mtk</i>	<u>Dsec\GM</u> <u>21609</u>	<u>Dyak\GE</u> <u>11702</u>	<u>Dere\GG</u> <u>20517</u>		<i>Dpse\GA2</i> 0868		<u>Dwil\GK</u> <u>19217</u>			
FBgn0022 073	<i>Dsim\Thor</i>	<u>Dsec\GM</u> <u>11143</u>	<u>Dyak\Thor</u>	<u>Dere\GG</u> <u>24948</u>	<u>Dana\GF</u> <u>15239</u>	<i>Dpse\GA2</i> 1364	<i>Dper\GL</i> <u>18656</u>	<u>Dwil\GK</u> <u>18436</u>	<u>Dvir\GJ17</u> <u>238</u>	<u>Dmojl\GI1</u> <u>5007</u>	<u>Dgri\GH1</u> <u>0214</u>
FBgn0043 578	<i>Dsim\PGR</i> P-SB1	<u>Dsec\GM</u> <u>24370</u>	<u>Dyak\GE</u> <u>22193</u>	<u>Dere\GG</u> <u>15854</u>	<u>Dana\GF</u> <u>10506</u>	<i>Dpse\GA2</i> 1961	<i>Dper\GL</i> <u>13224</u>	<u>Dwil\GK</u> <u>17343</u>	<u>Dvir\GJ12</u> <u>160</u>	<u>Dmojl\GI1</u> <u>1935</u>	<u>Dgri\GH1</u> <u>5225</u>

Underlined: Genome-wide drosophilid orthologs.

Italic: Curated drosophilid ortholog

ments. Other regulatory codes could exist involving any one of these components, in addition to others.

The role of chromatin structure in gene expression regulation during immune response of insects remains poorly explored. This work provides a first insight into this complex regulatory mechanism potentially shared by immune genes of *drosophilidae* and *culicidae*.

Conclusion

Immunity genes of *A. gambiae*, *Ae. aegypti*, *D. melanogaster* and many other Drosophilid species share a common enrichment of AT-rich motifs in their 5'-US regions. AT-rich motifs are frequently associated to bioinformatic nucleosome positioning predictions, suggesting their participation in a particular nucleosome organization involved in transcriptional regulation of an immunity co-regulated module. Many of these regulatory regions also have NFκB response elements within the first 500 bp 5'

from the transcription start site. These two features suggest that the mechanism of transcriptional regulation of immune response genes in dipterans are conserved and might occur through modifications in chromatin structure of promoter regions mediated by NFκB-dependent recruitment of remodeling factors. Our findings suggest that AT-rich motif enrichment in regulatory regions in this group of co-regulated genes could represent an evolutionary constrained signature in dipterans and perhaps other species, despite their evolutionary distance.

Methods

Gene selection criteria

Microarray data for *A. gambiae* immune response was kindly provided by the author [26]. Microarray data from *D. melanogaster* immune response [27] was downloaded from [62]. Analysis of expression profiles was conducted using the TMEV version 3.1 module of TM4 microarray software suite [63].

Table 17: Aedes aegypti orthologous immunity genes used for over-representation analysis of AT-rich motifs in 5'-US (2500 bp)

Ensembl Gene ID	Description
AAEL000621	antibacterial peptide, putative
AAEL000709	developmental protein cactus
AAEL001794	macroglobulin/complement
AAEL001802	macroglobulin/complement
AAEL002972	brain chitinase and chia
AAEL003889	gram-negative bacteria binding protein
AAEL004522	Orthologous of putative infection responsive short peptide
AAEL005787	serine protease, putative
AAEL007765	serine protease inhibitor 4, serpin-4
AAEL008646	fibrinogen and fibronectin
AAEL009423	cd36 antigen
AAEL009520	Leucine rich domain
AAEL010171	peptidoglycan recognition protein sb2
AAEL010737	aromatic amino acid decarboxylase
AAEL014238	aromatic amino acid decarboxylase




Figure 4
Four bp AT-rich motif over-representation analysis in 5' US of dipteran immunity related induced genes and their corresponding phylogenetic relationships based on [32]. Each AT-rich motif was plotted based on the rank obtained according to the corresponding Sig_{Occ} for each genome.

Three gene clusters were selected for each species by hierarchical clustering [64] from the microarray databases. For *A. gambiae*, gene selection criteria were as follows: **Induced immunity-related genes:** genes associated with immunity based on the protein structural features, and with expression values in $\log_2(f_2/f_1) > 0$ in at least 7/10 immunological challenges and 3/6 points for each challenge. **Non-modified genes:** genes with an expression mean of $\log_2(f_2/f_1) \pm 0.10$ and a standard deviation of ± 0.15 in 9/10 immunological challenges. **Down-regulated**

genes: genes with an expression level of $\log_2(f_2/f_1) \leq -0.5855$ (a repression level of at least 1.5 times with respect to control cells) in at least 4/10 immune challenges and in 4/6 times for each challenge, and with a maximal expression level of $\log_2(f_2/f_1) < 0.6785$ in only one point per challenge (a maximal expression level less than 1.6 times with respect to control cells in only one point). For *D. melanogaster*, gene definitions were as follows: **Immunity related genes:** genes associated with immunity based on Gene Ontology classification (GO:0006952, defense




Figure 5
ATAA motifs are associated preferentially with regions of positive nucleosomal potential in *A. gambiae* (A) and *D. melanogaster* (B). ATAA motifs are present in all the groups of sequences analyzed, however 5'-US of immunity genes have more ATAA per sequence. ATAA distribution differ between biological and non-biological (artificial) sequences, which have less ATAA motifs and are associated primarily with negative values of nucleosomal potential. Standard deviation is showed. (NM:non-modified, DR:down-regulated, Random: random selected genes, Artificial: computer-derived non-biological sequences).




Figure 6
AT-rich motifs in the family *Drosophilidae* and *Culicidae* are associated preferentially with regions of positive nucleosomal potential calculated by the RECON algorithm[22]. ATAA (A) and TATA (B) motifs are associated with positive nucleosomal potential in *D. ananassae*, *D. pseudoobscura* and *D. grimshawi* ($p < 0.001$), which belong to different groups. Over-represented TTAA and AAAT motifs in *Ae. aegypti* are also associated preferentially with regions of positive nucleosomal potential (C, $p < 0.001$). Standard deviation is showed.

response, biological process), and with expression values in $\log_2(f_2/f_1) > 0$ in 6/6 time points of bacterial challenge. **Non-modified genes:** genes with an expression mean of $\log_2(f_2/f_1) \pm 0.1$ and a standard deviation of ± 0.07 in 6/6 time points of bacterial challenge. **Down-regulated genes:** genes with an expression level of $\log_2(f_2/f_1) \leq -0.5855$ (a repression level of at least 1.5 times with respect to control) in 6/6 times of the bacterial challenge. Two additional groups were included in the analysis: **Random genes:** two groups of 20 and 30 genes were randomly

selected from the *A. gambiae* and *D. melanogaster* genomes, respectively, using the "Random Gene Selection" tool of RSA-Tools [65]. **Random sequences (artificial):** two groups of 20 and 30 random non-biological sequences were generated using the "Random DNA sequence" tool of Sequence Manipulation Suite, version 2 [66], using this tool we generated random sequences with equal proportions of each nucleotide (~ 0.25).




Figure 7
ATAA motifs are associated with high p values of nucleosomal occupancy (pNO), in *A. gambiae* (A) and *D. melanogaster* (B), evaluated with the software "Nucleosomes Positioning", reported by Segal and col.[20], when the yeast model is applied. None ATAA motif was associated with pNO values between 0.8 and 0.5. Non-biological sequences (artificial) show an ATAA distribution different to biological sequences. Note the similarity between this figure and figure 5. Standard deviation is showed.





Figure 8
AT-rich motifs in *Drosophilidae* and *Culicidae* families are associated with high p values of nucleosomal occupancy (pNO), calculated according to Segal, et al [20] using the yeast model for *D. ananassae*, *D. pseudoobscura* and *D. grimshawi* (A and B) and *Ae. aegypti* (C). pNO is plotted in relation to its association with ATAA (A) and TTA (B) or TTA and AAAT (C) motif frequency. In every species, the AT-rich motif analyzed was associated to high pNO. Standard deviation is showed.

Using these definitions, for *A. gambiae*, we first selected the expression profiles and the associated gene was then identified. To verify the annotated transcription initiation site, each gene prediction was manually curated by two approaches: The first was by aligning corresponding EST clusters obtained from AnoEST [67] and UNIGENE [68] to the *A. gambiae* genome (AgamP3, Ensembl release 45, Jun 2007) using BLAST in the ENSEMBL genome browser [28]. The second was based on manual verification of the presence of either TATA-box, Initiator sequence (Inr) or downstream promoter element (DPE) [69]. For *Drosophila*, the gene ID was included in the microarray database.

Once the gene associated with each profile was identified, the 5' regulatory regions were recovered for *D. melanogaster* (BDGP4.3) and *A. gambiae* (AgamP3, Ensembl release 45, Jun 2007) genes using Ensembl's data mining tool Biomart [70].

An additional set of 5' upstream 2000 bp sequences from the 12 *Drosophila* species derived from the 12 *Drosophila* genome project [32] retrieved from [71], release R1.1 for *D. virilis* (23 sequences), R1.2 for *D. ananassae* (23 sequences), *D. erecta* (27 sequences), *D. grimshawi* (21 sequences), *D. mojavensis* (21 sequences), *D. persimilis* (20 sequences), *D. sechellia* (28 sequences), *D. simulans* (29






Figure 10
ATAA motifs are located predominantly in the -251 to -500 region of immunity genes in *A. gambiae* and *D. melanogaster*, showing a coincidence with regions where NF κ B RE are more abundant.

sequences), *D. yakuba* (27 sequences) and *D. willistoni* (24 sequences), R2.2 for *D. pseudoobscura* (27 sequences) and R5.7 for *D. melanogaster* (35 sequences) and 15 *Aedes aegypti* [72] 2500 bp sequences retrieved from Biomart [70] (AAEGL1) were also included for motif over-representation analysis. Selection of the *Drosophila* sequences was done based on orthology to *D. melanogaster* immunity gene data set described in Table 16, according to Fly-Base annotations, but not expression data. For *Ae. aegypti*, genes were also selected based on one to one orthology to immunity genes in *A. gambiae* according to Table 17.

Statistically overrepresented DNA motifs

To identify statistically overrepresented DNA motifs in 5' DNA regulatory regions of selected genes, we used the Oligo-Analysis program [29] searching for DNA motifs of 2 to 8 nucleotides of length in 5' upstream regions of 2500 or 2000 nucleotides. For the analysis, we created our own expected frequency tables for each motif length, using 5' upstream regions of 2500 nucleotides length corresponding to 13,172 genes of *D. melanogaster*; 13,166 genes of *A. gambiae*, and 16,691 in *Ae. aegypti*. A similar approach was used for the 11 additional *Drosophila* species using pre-computed 2000 bp upstream 5' sequences. The obtained expected frequency tables were used to estimate the expected number of occurrences for each oligonucleotide in induced, down-regulated, non-modified, random biological and random no-biological sets of sequences. The analyzed sequences were aligned to detect and avoid duplication between sequences, and duplicated regions larger than 40 nucleotides inside a sequence were removed. Also, to prevent a bias due to self-overlapping, a non-overlapping mode was adopted. The detection of overrepresented oligonucleotides was based on an estimation of the significance of the observed occurrences (O_{occ}). For each oligonucleotide, the p value (P_{occ}) was calculated on the basis of the binomial distribution. Because the analysis comprise multiple tests (256 in the case of tetranucleotides), the possibility exists that even low p values appeared by chance. To correct for such a multitesting effect, the p values were multiplied by the number of oligonucleotides. This correction results in an expected value (E_{occ}). The significance index [$sig_{occ} = -\log(E_{occ})$] reflects the degree of overrepresentation for each oligonucleotide in a logarithmic scale [30].

The motifs overrepresentation identified with Oligo-Analysis, was verified using POBO [31], which uses bootstrap to verify the statistical overrepresentation of a given motif.

Nucleosome positioning prediction

To predict regions of nucleosomal occupancy in the sequences of the distinct groups of *A. gambiae* and *D. melanogaster* 5'-US, we used two programs: RECON [22], which assigns a nucleosomal potential value at each posi-

tion in a sequence using sliding windows of 160 pb based on statistical distribution of dinucleotide frequencies, and Nucleosome Position Prediction [20], which predicts nucleosomal positioning using probabilistic and thermodynamic models, assigning a p value of nucleosomal occupancy (pNO) to each position of a sequence. For this last program, we analyzed 5'-US using yeast, chicken and human models, and both published and working versions of the program. The length of the 5'-US analyzed was of 2500 bp for *A. gambiae*, *D. melanogaster* and *Ae. aegypti* and 2000 bp for non-melanogaster *Drosophila* species. For RECON, we used 2660 bp that comprised the 2500 bp promoter, flanked by 80 bp, in order to recover nucleosomal potential values for all promoter positions.

Analysis of motifs position regarding predicted nucleosomal regions

To associate the results obtained with the program Oligo-Analysis and those obtained with RECON and Nucleosome Position Prediction, perl scripts were written to automatically associate motifs coordinates (obtained with DNA-Pattern, [65] with tables containing "Nucleosomal potential" values (obtained with RECON) or "p of nucleosomal occupancy (pNO)" values (obtained with Nucleosome Position Prediction).

To determine if there was a correlation between the findings of Oligo-analysis and RECON, using perl scripts, the corresponding value of nucleosomal potential obtained with RECON was assigned to each position of the ATAA motif, having four values for each motif, on basis to these four values each ATAA motif was classified as: 1) positive, if the four positions of ATAA were positive; 2) negative, if the four positions were negative; 3) mixed, if at least one position was of opposite sign to the others, and 4) undefined, if at least one position was an N. Once classified, the distribution of ATAA motifs between these four categories was statistically evaluated.

Similarly, to determine if there was a correlation between the findings of Oligo-analysis and Nucleosome Position Prediction, using perl scripts, the corresponding pNO value obtained with the algorithm "Nucleosome Position Prediction" was assigned to each position of the ATAA motif, having again four values for each motif, one per each position. Based on these four values, each ATAA motif was classified as: 1) high p value motif if the four positions had occupancy p values higher than 0.8, 2) medium p value motif if the four positions had occupancy p values between 0.8 and 0.5, 3) low p value motif if the four positions had occupancy p values below of 0.5, and 4) undefined if at least one value were not belonging to the same range of values. Additionally, given that this software uses yeast, chicken and human models, and have a working and a published version, data generated with

each model and each version were analyzed. This program does not accept sequences with Ns, therefore, in some cases the number of sequences analyzed by group of 5' upstream sequences was slightly smaller. For *A. gambiae*: 16 immunity, 12 down-regulated, 17 random genes and 17 non-modified. For *D. melanogaster*, only the group of non-modified genes was modified, from 32 to 31 sequences. For the other *Drosophila* species and *Ae. aegypti* the number of analysed sequences was the same for both programs.

Analysis of 5'-US with alignment matrices

Using alignment matrices constructed on NFκB REs identified in immunity 5'-US of *A. gambiae* and *D. melanogaster* by MEME [35] (Figure 12), a matrix-based search was carried out in the different groups of genes using the PATSER algorithm [73], searching for NFκB REs. A lower threshold estimation of 5.0 was assigned.

Statistical analysis

In order to compare the distribution of AT-rich motifs associated with positive nucleosomal potential values among groups of genes, we fitted ordinary least squares regression models with robust standard errors, with the number of positive AT-rich motifs as the dependent variable and dummy variables of the corresponding group of genes as predictors for all dipteran species. We fitted similar models to compare among groups of genes the distribution of AT-rich motifs associated with pNO > 0.8, and the AT% difference between groups of genes. Additionally, we compared the number of AT-rich motifs within all types of nucleotide sequences, either associated with pos-

itive vs. negative values of nucleosomal potential or pNO > 0.8 vs. pNO < 0.5, by use of paired Student's T tests.

To evaluate the distribution of NFκB REs throughout the promoter regions, comparisons were done using Poisson regression with the count of NFκB RE as response variable and the group of genes as independent variable. The models support the χ^2 goodness-of-fit test, when the model was not supported, a Kruskal-Wallis non-parametric test was done. Ordinary least squares regression analysis was performed to compare the counts of ATAA in the vicinity (± 200 bp) of NFκB motifs by type of gene group (immunity, non-modified and down-regulated), in both *A. gambiae* and *D. melanogaster*.

Authors' contributions

JH-R participated in the design, data acquisition, data analysis and interpretation and wrote the manuscript. HS, FJC-R wrote perl scripts for data acquisition. HL-F performed statistical analysis. VV-G participated in data interpretation as well as revising the manuscript. MHR participated in conception, data analysis and interpretation as well as revising the manuscript. JM-B participated in conception and design, data acquisition and analysis, interpretation and drafted the manuscript.

Additional material

Additional file 1

Contains Table 18. 4 bp motif enrichment in 5'-US of Immunity genes of the *Drosophila* genus and *Aedes aegypti*.

Click here for file

[<http://www.biomedcentral.com/content/supplementary/1471-2164-9-326-S1.pdf>]

A)

A	0	0	1	12	4	0	1	4
C	2	0	0	2	1	0	12	11
G	14	5	13	0	0	0	2	1
T	0	11	2	2	11	16	1	0

B)

A	8	0	0	0	18	3	0	0	0	0	5	16
C	0	0	0	4	0	3	0	14	24	23	17	3
G	11	26	18	21	0	0	4	1	0	0	0	0
T	7	0	8	1	8	20	22	11	2	3	4	7

Figure 12
Alignment matrices for NFκB motifs identified by MEME[34] in immunity genes of *A. gambiae* (A) and *D. melanogaster* (B).

Acknowledgements

Conacyt PhD scholarship I59376 to JHR. SEP-Conacyt P44797 Young Researcher award to JMB. Maritza Solano for technical assistance; Mara Tellez Rojo, Julio Collado Vides, Mario Zurita and Jaques van Helden and the two reviewers for helpful discussions and suggestions.

References

- Carroll SB: **Genetics and the making of *Homo sapiens***. *Nature* 2003, **422**:849-857.
- Levine M, Tjian R: **Transcription regulation and animal diversity**. *Nature* 2003, **424**:147-151.
- Davidson EH: *The regulatory genome* Canada: Academic Press; 2006.
- Khorasanizadeh S: **The nucleosome: from genomic organization to genomic regulation**. *Cell* 2004, **116**:259-272.
- Kornberg RD: **Chromatin structure: a repeating unit of histones and DNA**. *Science* 1974, **184**:868-871.
- Wu C: **Chromatin remodelling and the control of gene expression**. *J Biol Chem* 1997, **272**:28171-28174.
- Narlikar GJ, Fan HY, Kingston RE: **Cooperation between complexes that regulate chromatin structure and transcription**. *Cell* 2002, **108**:475-487.
- Smale ST, Fisher AG: **Chromatin structure and gene regulation in the immune system**. *Ann Rev Immunol* 2002, **20**:427-462.

9. Schulze SR, Wallrath LL: **Gene regulation by chromatin structure: paradigms established in *Drosophila melanogaster*.** *Annu Rev Entomol* 2007, **52**:171-192.
10. Schmid A, Fascher KD, Hörz W: **Nucleosome disruption at the yeast PHO5 promoter upon PHO5 induction occurs in the absence of DNA replication.** *Cell* 1992, **71**:853-864.
11. Lewin B: **Chromatin and gene expression: constant questions, but changing answers.** *Cell* 1994, **79**:397-406.
12. Beato M, Eisfeld K: **Transcription factor access to chromatin.** *Nucleic Acids Res* 1997, **25**:3559-3563.
13. Jackson JR, Benyajati C: **DNA-histone interactions are sufficient to position a single nucleosome juxtaposing *Drosophila* Adh adult enhancer and distal promoter.** *Nucleic Acids Res* 1993, **21**:957-967.
14. Trifonov EN, Sussman JL: **The pitch of chromatin DNA is reflected in its nucleotide sequence.** *Proc Natl Acad Sci* 1980, **77**:3816-3820.
15. Satchwell SC, Drew HR, Travers AA: **Sequence periodicities in chicken nucleosome core DNA.** *J Mol Biol* 1986, **191**:659-675.
16. Shrader TE, Crothers DM: **Artificial nucleosome positioning sequences.** *Proc Natl Acad Sci* 1989, **86**:7418-7422.
17. Ioshikhes I, Bolshoy A, Derenshteyn K, Borodovsky M, Trifonov EN: **Nucleosome DNA sequence pattern revealed by multiple alignment of experimentally mapped sequences.** *J Mol Biol* 1996, **262**:129-139.
18. Widlund HR, Cao H, Simonsson S, Magnusson E, Simonsson T, Nielsen PE, Kahn JD, Crothers DM, Kubista DM: **Identification and characterization of genomic nucleosome-positioning sequences.** *J Mol Biol* 1997, **267**:807-817.
19. Stein A, Bina M: **A signal encoded in vertebrate DNA that influences nucleosome positioning and alignment.** *Nucleic Acids Res* 1999, **27**:848-853.
20. Segal E, Fondufe-Mittendorf Y, Chen L, Thastrom A, Field Y, Moore IK, Wang JZ, Widom J: **A genomic code for nucleosome positioning.** *Nature* 2006, **442**:772-778.
21. Johnson SM, Tan FJ, McCullough HL, Riordan DP, Fire AZ: **Flexibility and constraint in the nucleosome core landscape of *Caenorhabditis elegans* chromatin.** *Genome Res* 2006, **16**:1505-1516.
22. Levitsky VG: **RECON: a program for prediction of nucleosome formation potential.** *Nuc Acids Res* 2004, **32**:W346-W349.
23. Kimbrell DA, Beutler B: **The evolution and genetics of innate immunity.** *Nat Rev Genet* 2001, **2**:256-267.
24. Hernández-Romano J, Martínez-Barnetteche J, Rodríguez-López MH: **Transcriptional regulation of immune related genes in insects: Insights into the genomics of *Anopheles* immune response.** In *Genes, Genomes and Genomics Volume 1*. Edited by: Thangadurai D, Tang W, Pullaiah T. New Delhi: Regency Publications; 2006:98-130.
25. Ferrandon D, Imler JL, Hetru C, Hoffmann JA: **The *Drosophila* systemic immune response: sensing and signalling during bacterial and fungal infections.** *Nat Rev Immunol* 2007, **7**:862-874.
26. Dimopoulos G, Christophides GK, Meister S, Schultz J, White KP, Barillas-Mury C, Kafatos FC: **Genome expression analysis of *Anopheles gambiae*: responses to injury, bacterial challenge, and malaria infection.** *Proc Natl Acad Sci* 2002, **99**:8814-8819.
27. De Gregorio E, Spellman PT, Rubin GM, Lemaitre B: **Genome-wide analysis of the *Drosophila* immune response by using oligonucleotide microarrays.** *Proc Natl Acad Sci* 2001, **98**:12590-12595.
28. Birney E, Andrews D, Caccamo M, Chen Y, Clarke L, Coates G, Cox T, Cunningham F, Curwen V, Cutts T, Down T, Durbin R, Fernandez-Suarez XM, Flicek P, Graf S, Hammond M, Herrero J, Howe K, Iyer V, Jekosch K, Kahari A, Kasprzyk A, Keefe D, Kokocinski F, Kulesha E, London D, Longden I, Melsopp C, Meidl P, Overduin B, Parker A, Proctor G, Prlic A, Rae M, Rios D, Redmond S, Schuster M, Sealy I, Searle S, Severin J, Slater G, Smedley D, Smith J, Stabenau A, Stalker J, Trevanion S, Ureta-Vidal A, Vogel J, White S, Woodwork C, Hubbard TJ: **Ensembl 2006.** *Nucleic Acids Res* 2006, **34**:D556-561.
29. van Helden, André B, Collado-Vides J: **Extracting regulatory sites from the upstream region of yeast genes by computational analysis of oligonucleotide frequencies.** *J Mol Biol* 1998, **281**:827-842.
30. Hulzink RJ, Weerdesteyn H, Croes AF, Gerats T, van Herpen MM, van Helden J: **In silico identification of putative regulatory sequence elements in the 5'-untranslated region of genes that are expressed during male gametogenesis.** *Plant Physiol* 2003, **132**:75-83.
31. Kankainen M, Holm L: **POBO, transcription binding site verification with bootstrapping.** *Nucleic Acid Res* 2004, **32**:W222-W229.
32. Drosophila 12 Genomes Consortium, Clark AG, Eisen MB, Smith DR, Bergman CM, Oliver B, Markow TA, Kaufman TC, Kellis M, Gelbart W, Iyer VN, Pollard DA, Sackton TB, Larracuent AM, Singh ND, Abad JP, Abt DN, Adryan B, Aguade M, Akashi H, Anderson WW, Aquadro CF, Ardell DH, Arguello R, Artieri CG, Barbash DA, Barker D, Barsanti P, Batterham P, Batzoglou S, Begun D, Bhutkar A, Blanco E, Bosak SA, Bradley RK, Brand AD, Brent MR, Brooks AN, Brown RH, Butlin RK, Caggese C, Calvi BR, Bernardo de Carvalho A, Caspi A, Castrezana S, Celniker SE, Chang JL, Chapple C, Chatterji S, Chinwalla A, Civetta A, Clifton SW, Comeron JM, Costello JC, Coyne JA, Daub J, David RG, Delcher AL, Delehaunty K, Do CB, Ebling H, Edwards K, Eickbush T, Evans JD, Filipitski A, Findeiss S, Freyhult E, Fulton L, Fulton R, Garcia AC, Gardiner A, Garfield DA, Garvin BE, Gibson G, Gilbert D, Gnerre S, Godfrey J, Good R, Gotea V, Gravely B, Greenberg AJ, Griffiths-Jones S, Gross S, Guigo R, Gustafson EA, Haerty W, Hahn MW, Halligan DL, Halpern AL, Halter GM, Han MV, Heger A, Hillier L, Hinrichs AS, Holmes I, Hoskins RA, Hubisz MJ, Hultmark D, Huntley MA, Jaffe DB, Jagadeeshan S, Jeck WR, Johnson J, Jones CD, Jordan WC, Karpen GH, Kataoka E, Keightley PD, Kheradpour P, Kirkness EF, Koerich LB, Kristiansen K, Kudrna D, Kulathinal RJ, Kumar S, Kwok R, Lander E, Langley CH, Lapoint R, Lazzaro BP, Lee SJ, Levesque L, Li R, Lin CF, Lin MF, Lindblad-Toh K, Llopart A, Long M, Low L, Lozovsky E, Lu J, Luo M, Machado CA, Makalowski W, Marzo M, Matsuda M, Matzkin L, McAllister B, McBride CS, McKernan B, McKernan K, Mendez-Lago M, Minx P, Mollenhauer MU, Montooth K, Mount SM, Mu X, Myers E, Negre B, Newfield S, Nielsen R, Noor MA, O'Grady P, Pachter L, Papacait M, Parisi MJ, Parisi M, Parts L, Pedersen JS, Pesole G, Phillipy AM, Ponting CP, Pop M, Porcelli D, Powell JR, Prohaska S, Pruitt K, Puig M, Quesneville H, Ram KR, Rand D, Rasmussen MD, Reed LK, Reenan R, Reily A, Remington KA, Rieger TT, Ritchie MG, Robin C, Rogers YH, Rohde C, Rozas J, Rubenfield MJ, Ruiz A, Russo S, Salzberg SL, Sanchez-Gracia A, Saranga DJ, Sato H, Schaeffer SW, Schatz MC, Schlenke T, Schwartz R, Segarra C, Singh RS, Sirot L, Sirot M, Sisneros NB, Smith CD, Smith TF, Spieth J, Stage DE, Stark A, Stephan W, Strausberg RL, Strempel S, Sturgill D, Sutton G, Sutton GG, Tao W, Teichmann S, Tobarri YN, Tomimura Y, Tsolas JM, Valente VL, Venter E, Venter JC, Vicario S, Vieira FG, Vilella AJ, Villasante A, Walenz B, Wang J, Wasserman M, Watts T, Wilson D, Wilson RK, Wing RA, Wolfner MF, Wong A, Wong GK, Wu CI, Wu G, Yamamoto D, Yang HP, Yang SP, Yorke JA, Yoshida K, Zdobnov E, Zhang P, Zhang Y, Zimin AV, Baldwin J, Abdouelleil A, Abdulkadir J, Abebe A, Abera B, Abreu J, Acer SC, Aftuck L, Alexander A, An P, Anderson E, Anderson S, Arachi H, Azer M, Bachantsang P, Barry A, Bayul T, Berlin A, Bessette D, Bloom T, Blye J, Boguslavskiy L, Bonnet C, Boukhgalter B, Bourzgui I, Brown A, Cahill P, Channer S, Cheshtsang Y, Chuda L, Citroen M, Collymore A, Cooke P, Costello M, D'Acio K, Daza R, De Haan G, DeGray S, DeMaso C, Dhargay N, Dooley K, Dooley E, Doricent M, Dorje P, Dorjee K, Dupes A, Elong R, Falk J, Farina A, Faro S, Ferguson D, Fisher S, Foley CD, Franke A, Friedrich D, Gadbois L, Gearin G, Gearin CR, Giannoukos G, Goode T, Graham J, Grandbois E, Grewal S, Gyaltsen K, Hafez N, Hagos B, Hall J, Henson C, Hollinger A, Honan T, Huard MD, Hughes L, Hurhula B, Husby ME, Kamat A, Kanga B, Kashin S, Khazanovich D, Kisner P, Lance K, Lara M, Lee W, Lennon N, Letendre F, LeVine R, Lipovsky A, Liu X, Liu J, Liu S, Lokysang T, Lokysang Y, Lubonja R, Lui A, MacDonald P, Magnalis V, Maru K, Matthews C, McCusker W, McDonough S, Mehta T, Meldrim J, Meneus L, Mihai O, Mihalev A, Mihova T, Mittelman R, Mlenga V, Montmayeur A, Mulrain L, Navidi A, Naylor J, Negash T, Nguyen T, Nguyen N, Nicol R, Norbu C, Norbu N, Novod N, O'Neill B, Osman S, Markiewicz E, Oyono OL, Patti C, Phunkhang P, Pierre F, Priest M, Raghuraman S, Rege F, Reyes R, Rise C, Rogov P, Ross K, Ryan E, Settipalli S, Shea T, Sherpa N, Shi L, Shih D, Sparrow T, Spaulding J, Stalker J, Stange-Thomann N, Stavropoulos S, Stone C, Strader C, Tesfaye S, Thomson T, Thoulutsang Y, Thoulutsang D, Topham K, Topping I, Tsamla T, Vassiliev H, Vo A, Wangchuk T, Wangdi T, Weiland M, Wilkinson J, Wilson A, Yadav S, Young G, Yu Q, Zembek L, Zhong D, Zimmer A, Zwirko Z, Jaffe DB, Alvarez P, Brockman WW, Butler J, Chin C, Gnerre S, Grabherr M, Kleber M, Mauceli E, MacCallum I: **Evolution of genes and genomes on the *Drosophila* phylogeny.** *Nature* 2007, **8**:203-218.

33. Richards S, Liu Y, Bettencourt BR, Hradecky P, Letovsky S, Nielsen R, Thornton K, Hubisz MJ, Chen R, Meisel RP, Couronne O, Hua S, Smith MA, Zhang P, Liu J, Bussemaker HJ, van Batenburg MF, Howells SL, Scherer SE, Sodergren E, Matthews BB, Crosby MA, Schroeder AJ, Ortiz-Barrientos D, Rives CM, Metzker ML, Muzny DM, Scott G, Steffen D, Wheeler DA, Worley KC, Havlak P, Durbin KJ, Egan A, Gill R, Hume J, Morgan MB, Miner G, Hamilton C, Huang Y, Waldron L, Verduzco D, Clerc-Blankenburg KP, Dubchak I, Noor MA, Anderson W, White KP, Clark AG, Schaeffer SW, Gelbart W, Weinstock GM, Gibbs RA: **Comparative genome sequencing of *Drosophila pseudoobscura*: chromosomal, gene, and cis-element evolution.** *Genome Res* 2005, **15**:1-18.
34. Levitsky VG, Podkolodnaya OA, Kolchanov NA, Podkolodny NL: **Nucleosome formation potential of eukaryotic DNA: calculation and promoters analysis.** *Bioinformatics* 2001, **17**:998-1010.
35. Bailey TL, Elkan C: **The value of prior knowledge in discovering motifs with MEME.** In *Proceedings of the third international conference on intelligent systems for molecular biology* AAAI Press, Menlo Park, CA; 1995:21-29.
36. Zdobnov EM, von Mering C, Letunic I, Torrents D, Suyama M, Copley RR, Christophides GK, Thomasova D, Holt RA, Subramanian GM, Mueller HM, Dimopoulos G, Law JH, Wells MA, Birney E, Charlab R, Halpern AL, Kokoza E, Kraft CL, Lai Z, Lewis S, Louis C, Barillas-Mury C, Nusskern D, Rubin GM, Salzberg SL, Sutton GG, Topalis P, Wides R, Wincker P, Yandell M, Collins FH, Ribeiro J, Gelbart WM, Kafatos FC, Bork P: **Comparative genome and proteome analysis of *Anopheles gambiae* and *Drosophila melanogaster*.** *Science* 2002, **298**:149-159.
37. Widom J: **Role of DNA sequence in nucleosome stability and dynamics.** *Q Rev Biophys* 2001, **34**:269-324.
38. Anderson JD, Widom J: **Poly(dA-dT) promoter elements increase the equilibrium accessibility of nucleosomal DNA target sites.** *Mol Cell Biol* 2001, **21**:3830-3839.
39. Sekinger EA, Moqtaderi Z, Struhl K: **Intrinsic histone-DNA interactions and low nucleosome density are important for preferential accessibility of promoter regions in yeast.** *Mol Cell* 2005, **18**:735-748.
40. Lee W, Tillo D, Bray N, Morse RH, Davis RW, Hughes TR, Nislow C: **A high-resolution atlas of nucleosome occupancy in yeast.** *Nat Genet* 2007, **39**:1235-1244.
41. Snel B, van Noort V, Huynen MA: **Gene co-regulation is highly conserved in the evolution of eukaryotes and prokaryotes.** *Nucleic Acids Res* 2004, **32**:4725-4731.
42. De Gregorio E, Spellman PT, Tzou P, Rubin GM, Lemaitre B: **The Toll and Imd pathways are the major regulators of the immune system response in *Drosophila*.** *EMBO J* 2002, **21**:2568-2579.
43. Waterhouse RM, Kriventseva EV, Meister S, Xi Z, Alvarez KS, Bartholomay LC, Barillas-Mury C, Bian G, Blandin S, Christensen BM, Dong Y, Jiang H, Kanost MR, Koutsos AC, Levashina EA, Li J, Ligoxygakis P, Maccallum RM, Mayhew GF, Mendes A, Michel K, Osta MA, Paskewitz S, Shin SW, Vlachou D, Wang L, Wei W, Zheng L, Zou Z, Severson DW, Raikhel AS, Kafatos FC, Dimopoulos G, Zdobnov EM, Christophides GK: **Evolutionary dynamics of immune-related genes and pathways in disease-vector mosquitoes.** *Science* 2007, **316**:1738-1743.
44. Hoffmann JA, Kafatos FC, Janeway CA, Ezekowitz RAB: **Phylogenetic perspectives in innate immunity.** *Science* 1999, **284**:1313-1318.
45. Hoffmann JA, Reichhart JM: ***Drosophila* innate immunity: an evolutionary perspective.** *Nat Immunol* 2002, **3**:121-126.
46. Zheng L, Zhang L, Lin H, McIntosh MT, Malacrida AR: **Toll-like receptors in invertebrate innate immunity.** *ISJ* 2005, **2**:105-113.
47. Dziarski R, Gupta D: **Mammalian PGRPs: novel antibacterial proteins.** *Cell Microbiol* 2006, **8**:1059-1069.
48. Sun SC, Lindstrom I, Lee JY, Faye I: **Structure and expression of the attacin genes in *Hyalophora cecropia*.** *Eur J Biochem* 1991, **196**:247-254.
49. Kadalayil L, Petersen U, Engström Y: **Adjacent GATA and κB-like motifs regulate the expression of a *Drosophila* immune gene.** *Nuc Acids Res* 1997, **25**:1233-1239.
50. Eggleston P, Lu W, Zhao Y: **Genomic organization and immune regulation of the defensin gene from the mosquito, *Anopheles gambiae*.** *Insect Mol Biol* 2000, **9**:481-490.
51. Zheng XL, Zheng AL: **Genomic organization and regulation of three cecropin genes in *Anopheles gambiae*.** *Insect Mol Biol* 2002, **11**:517-525.
52. Senger K, Armstrong GW, Rowell WJ, Kwan JM, Markstein M, Levine M: **Immunity regulatory DNAs share common organizational features in *Drosophila*.** *Mol Cell* 2004, **13**:19-32.
53. Angelov D, Lenouvel F, Hans F, Müller CW, Bouvet P, Bednar J, Moudrianakis EN, Cadet J, Dimitrov S: **The histone octamer is invisible when NF-κappaB binds to the nucleosome.** *J Biol Chem* 2004, **279**:42374-42382.
54. Hoberg JE, Yeung F, Mayo MW: **SMRT derepression by the IκappaB kinase alpha: a prerequisite to NF-κappaB transcription and survival.** *Mol Cell* 2004, **16**:245-255.
55. Murphy KM, Reiner SL: **The lineage decisions of helper T cells.** *Nat Rev Immunol* 2002, **2**:933-944.
56. Weinmann AS, Plevy SE, Smale ST: **Rapid and selective remodeling of a positioned nucleosome during the induction of IL-12 p40 transcription.** *Immunity* 1999, **11**:665-675.
57. Agalioti T, Lomvardas S, Parekh B, Yie J, Maniatis T, Thanos D: **Ordered recruitment of chromatin modifying and general transcription factors to the IFN-β promoter.** *Cell* 2000, **103**:667-678.
58. Attema JL, Reeves R, Murray V, Levichkin I, Temple MD, Tremethick DJ, Shannon MF: **The human IL-2 promoter can assemble a positioned nucleosome that becomes remodeled upon T cell activation.** *J Immunol* 2002, **169**:2466-2476.
59. Holloway AF, Rao S, Chen X, Shannon MF: **Changes in chromatin accessibility across the GM-CSF promoter upon T cell activation are dependent on nuclear factor kappaB proteins.** *J Exp Med* 2003, **197**:413-23.
60. Strahl BD, Allis CD: **The language of covalent histone modifications.** *Nature* 2000, **403**:41-45.
61. Saha A, Wittmeyer J, Cairns BR: **Chromatin remodelling: the industrial revolution of DNA around histones.** *Nature Rev Mol Cell Biol* 2006, **7**:437-447.
62. **Genome-wide Gene Expression Patterns of *Drosophila* in Response to Immune Challenge** [<http://www.fruitfly.org/expression/immunity/data.shtml>]
63. Saeed AI, Sharov V, White J, Li J, Liang W, Bhagabati N, Braisted J, Klapa M, Currier T, Thiagarajan M, Sturn A, Snuffin M, Rezantsev A, Popov D, Ryltsov A, Kostukovich E, Borisovsky I, Liu Z, Vinsavich A, Trush V, Quackenbush J: **TM4: a free, open-source system for microarray data management and analysis.** *Biotechniques* 2003, **34**:374-378.
64. Eisen MB, Spellman PT, Brown PO, Botstein D: **Cluster analysis and display of genome-wide expression patterns.** *Proc Natl Acad Sci* 1998, **95**:14863-14868.
65. **Regulatory Sequence Analysis Tools** [<http://rsat.scmbb.ulb.ac.be/rsat/>]
66. **Sequence Manipulation Suite** [http://bioinformatics.org/sms2/random_dna.html]
67. Kriventseva EV, Koutsos AC, Blass C, Kafatos FC, Christophides GK, Zdobnov EM: **AnoEST: toward a *gambiae* functional genomics.** *Genome Res* 2005, **15**:893-899.
68. Wheeler DL, Church DM, Federhen S, Lash AE, Madden TL, Pontius JU, Schuler GD, Schriml LM, Sequeira E, Tatusova TA, Wagner L: **Database resources of the National Center for Biotechnology.** *Nucleic Acids Res* 2003, **31**:28-33.
69. Burke TW, Kadonaga JT: **The downstream core promoter element, DPE, is conserved from *Drosophila* to humans and is recognized by TAFII60 of *Drosophila*.** *Genes Dev* 1997, **11**:3020-3031.
70. Durinck S, Moreau Y, Kasprzyk A, Davis S, De Moor B, Brazma A, Huber W: **BioMart and Bioconductor: a powerful link between biological databases and microarray data analysis.** *Bioinformatics* 2005, **21**:3439-3440.
71. **Flybase: a database of *Drosophila* genes and genomes** [<http://www.flybase.org>]
72. Nene V, Wortman JR, Lawson D, Haas B, Kodira C, Tu ZJ, Loftus B, Xi Z, Megy K, Grabherr M, Ren Q, Zdobnov EM, Lobo NF, Campbell KS, Brown SE, Bonaldo MF, Zhu J, Sinkins SP, Hogenkamp DG, Amedeo P, Arensburg P, Atkinson PV, Bidwell S, Biedler J, Birney E, Bruggner RV, Costas J, Coy MR, Crabtree J, Crawford M, Debruyne B, Decaprio D, Eglmeier K, Eisenstadt E, El-Dorry H, Gelbart WM, Gomes SL, Hammond M, Hannick LI, Hogan JR, Holmes MH, Jaffe D, Johnston JS, Kennedy RC, Koo H, Kravitz S, Kriventseva EV, Kulp D,

Labutti K, Lee E, Li S, Lovin DD, Mao C, Mauceli E, Menck CF, Miller JR, Montgomery P, Mori A, Nascimento AL, Naveira HF, Nusbaum C, O'leary S, Orvis J, Pertea M, Quesneville H, Reidenbach KR, Rogers YH, Roth CW, Schneider JR, Schatz M, Shumway M, Stanke M, Stinson EO, Tubio JM, Vanzee JP, Verjovski-Almeida S, Werner D, White O, Wyder S, Zeng Q, Zhao Q, Zhao Y, Hill CA, Raikhel AS, Soares MB, Knudson DL, Lee NH, Galagan J, Salzberg SL, Paulsen IT, Dimopoulos G, Collins FH, Birren B, Fraser-Liggett CM, Severson DW: **Genome sequence of *Aedes aegypti*, a major arbovirus vector.** *Science* 2007, **316**:1718-1723.

73. Hertz GZ, Stormo GD: **Identifying DNA and protein patterns with statistically significant alignments of multiple sequences.** *Bioinformatics* 1999, **15**:563-577.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

