




Complete Genome Sequence of Enterotoxigenic *Escherichia coli* Podophage LL11

Matthew Theodore,^a Lauren Lessor,^a Chandler O'Leary,^a  Rohit Kongari,^a  Jason Gill,^a Mei Liu^a

^aCenter for Phage Technology, Texas A&M University, College Station, Texas, USA

ABSTRACT Enterotoxigenic *Escherichia coli* (EPEC) is an opportunistic pathogen that commonly causes foodborne illness. Study of bacteriophages against this pathogen could be useful to develop alternative treatment approaches. Here, we present the complete genome sequence of LL11, a T7-like podophage that infects EPEC.

Enterotoxigenic *Escherichia coli* (EPEC) is a global health concern due to its ability to cause traveler's diarrhea, as well as the increase in the emergence of antibiotic-resistant strains (1–3). The study of bacteriophages that infect EPEC provides insight into alternative therapeutic and preventative approaches (4, 5). Here, we present the complete genome sequence of podophage LL11, which infects EPEC.

Podophage LL11 was isolated using a clinical EPEC isolate from a municipal wastewater treatment plant in College Station, TX, in 2011. Host bacteria were cultured on LB broth or agar (Difco) at 37°C with aeration. The phage was cultured and propagated by the soft-agar overlay method (6). It was identified as a podophage using negative-stain transmission electron microscopy performed at the Texas A&M University Microscopy and Imaging Center as described previously (7). Phage genomic DNA was prepared using a modified Promega Wizard DNA cleanup kit protocol as described previously (7). Phage LL11 DNA was prepared using the GS FLX Titanium general DNA library preparation kit and sequenced by FLX Titanium 454 pyrosequencing at the Emory GRA Genome Center (Emory University, GA); trimmed FLX Titanium sequence reads were assembled into a single contig at 96.3-fold coverage using Newbler 2.5.3 (454 Life Sciences) with default settings. Contig completion was confirmed by PCR using primers (5'-AGCAATGCCTTGCCTAAG-3', 5'-AGTCGTATTCGTCTGGTTAAAG-3') facing away from the center of the assembled contig and by Sanger sequencing of the resulting product, with the contig sequence manually corrected to match the resulting Sanger sequencing read. GLIMMER 3.0 (8) and MetaGeneAnnotator 1.0 (9) were used to predict protein coding genes, with manual correction for appropriate gene starts, and tRNA genes were predicted with ARAGORN 2.36 (10). Rho-independent termination sites were identified via TransTermHP (<http://transterm.cbcb.umd.edu/>). Sequence similarity searches were conducted by BLASTp 2.2.28 (11) against the NCBI nonredundant (nr), UniProt Swiss-Prot (12), and TrEMBL databases. InterProScan 5.15-54.0 (13), LipOP (14), and TMHMM 2.0 (15) were used to predict protein function. All analyses were conducted at default settings via the CPT Galaxy (16) and WebApollo (17) interfaces (<https://cpt.tamu.edu>).

The complete genome of LL11 is 44,185 bp and is most similar to those of the phages K1-5 (GenBank accession no. [NC_008152](https://ncbi.nlm.nih.gov/nuccore/NC_008152)) and K1E ([AM084415](https://ncbi.nlm.nih.gov/nuccore/AM084415)) (18) with 74% and 57% overall similarity at the nucleotide level, respectively, as determined by BLASTn. LL11 has a GC content of 45.1% and contains 55 protein coding genes, most of which had homologues in phage T7 ([NC_001604](https://ncbi.nlm.nih.gov/nuccore/NC_001604)). The annotated genes include those responsible for morphogenesis, such as the tail spike, tail protein, major capsid protein, portal protein, and scaffolding protein, genes involved in DNA replication, such as DNA polymerase, RNA polymerase, and DNA primase, lysis genes, such as a class-II

Citation Theodore M, Lessor L, O'Leary C, Kongari R, Gill J, Liu M. 2019. Complete genome sequence of enterotoxigenic *Escherichia coli* podophage LL11. *Microbiol Resour Announc* 8:e00693-19. <https://doi.org/10.1128/MRA.00693-19>.

Editor Catherine Putonti, Loyola University Chicago

Copyright © 2019 Theodore et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Mei Liu, meiliu@tamu.edu.

Received 10 June 2019

Accepted 14 July 2019

Published 8 August 2019

holin, endolysin with a conserved lysozyme domain, and a separated spanin pair. Similarly to phage K1-5 (AY370674), LL11 also has two predicted tail fiber proteins related to tail fiber proteins of K1-5 (NC_008152), with one identical to K5 lyase (YP_654147) and the other distantly related to K1 endosialidase (YP_654148). These tail fiber proteins could allow LL11 to infect multiple host strains using different host receptor proteins in a manner similar to that of phage K1-5 (19).

Data availability. The genome sequence of phage LL11 was submitted to GenBank under accession no. MH729818. The associated BioProject, SRA, and BioSample accession numbers are PRJNA222858, SRR9214597, and SAMN11975578, respectively.

ACKNOWLEDGMENTS

We thank John Deaton of Deerland Enzymes for provision of pathogenic *E. coli* strains. This work was supported by Deerland Enzymes, Inc., and by funding from the National Science Foundation (awards EF-0949351 and DBI-1565146). Additional support came from the Center for Phage Technology (CPT), an Initial University Multidisciplinary Research Initiative supported by Texas A&M University and Texas AgriLife, and from the Department of Biochemistry and Biophysics at Texas A&M University.

We are grateful for the advice and support of the CPT staff and the Texas A&M University Microscopy and Imaging Center.

This announcement was prepared in partial fulfillment of the requirements for BICH464 Phage Genomics, an undergraduate course at Texas A&M University.

REFERENCES

- Croxen MA, Law RJ, Scholz R, Keeney KM, Wlodarska M, Finlay BB. 2013. Recent advances in understanding enteric pathogenic *Escherichia coli*. Clin Microbiol Rev 26:822–880. <https://doi.org/10.1128/CMR.00022-13>.
- Lim JY, Yoon J, Hovde CJ. 2010. A brief overview of *Escherichia coli* O157:H7 and its plasmid O157. J Microbiol Biotechnol 20:5–14.
- Sadeghabadi AF, Ajami A, Fadaei R, Zandieh M, Heidari E, Sadeghi M, Ataei B, Hoseini SG. 2014. Widespread antibiotic resistance of diarrheagenic *Escherichia coli* and *Shigella* species. J Res Med Sci 19:551–555.
- Brüssow H. 2005. Phage therapy: the *Escherichia coli* experience. Microbiology 151:2133–2140. <https://doi.org/10.1099/mic.0.27849-0>.
- Pouillot F, Chomton M, Blois H, Courroux C, Noelig J, Bidet P, Bingen E, Bonacorsi S. 2012. Efficacy of bacteriophage therapy in experimental sepsis and meningitis caused by a clone O25b:H4–ST131 *Escherichia coli* strain producing CTX-M-15. Antimicrob Agents Chemother 56:3568. <https://doi.org/10.1128/AAC.06330-11>.
- Adams MK. 1959. Bacteriophages. Interscience Publishers, Inc., New York, NY.
- Gill JJ, Berry JD, Russell WK, Lessor L, Escobar-Garcia DA, Hernandez D, Kane A, Keene J, Maddox M, Martin R, Mohan S, Thorn AM, Russell DH, Young R. 2012. The *Caulobacter crescentus* phage phiCbK: genomics of a canonical phage. BMC Genomics 13:542. <https://doi.org/10.1186/1471-2164-13-542>.
- Delcher AL, Harmon D, Kasif S, White O, Salzberg SL. 1999. Improved microbial gene identification with GLIMMER. Nucleic Acids Res 27:4636–4641. <https://doi.org/10.1093/nar/27.23.4636>.
- Noguchi H, Taniguchi T, Itoh T. 2008. MetaGeneAnnotator: detecting species-specific patterns of ribosomal binding site for precise gene prediction in anonymous prokaryotic and phage genomes. DNA Res 15:387–396. <https://doi.org/10.1093/dnares/dsn027>.
- Laslett D, Canback B. 2004. ARAGORN, a program to detect tRNA genes and tmRNA genes in nucleotide sequences. Nucleic Acids Res 32:11–16. <https://doi.org/10.1093/nar/gkh152>.
- Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, Madden TL. 2009. BLAST+: architecture and applications. BMC Bioinformatics 10:421. <https://doi.org/10.1186/1471-2105-10-421>.
- UniProt Consortium T. 2018. UniProt: the universal protein knowledgebase. Nucleic Acids Res 46:2699. <https://doi.org/10.1093/nar/gky092>.
- Jones P, Binns D, Chang HY, Fraser M, Li W, McAnulla C, McWilliam H, Maslen J, Mitchell A, Nuka G, Pesseat S, Quinn AF, Sangrador-Vegas A, Scheremetjew M, Yong SY, Lopez R, Hunter S. 2014. InterProScan 5: genome-scale protein function classification. Bioinformatics 30:1236–1240. <https://doi.org/10.1093/bioinformatics/btu031>.
- Juncker AS, Willenbrock H, Von Heijne G, Brunak S, Nielsen H, Krogh A. 2003. Prediction of lipoprotein signal peptides in Gram-negative bacteria. Protein Sci 12:1652–1662. <https://doi.org/10.1110/ps.0303703>.
- Krogh A, Larsson B, von Heijne G, Sonnhammer EL. 2001. Predicting transmembrane protein topology with a hidden Markov model: application to complete genomes. J Mol Biol 305:567–580. <https://doi.org/10.1006/jmbi.2000.4315>.
- Cock PJ, Gruning BA, Paszkiewicz K, Pritchard L. 2013. Galaxy tools and workflows for sequence analysis with applications in molecular plant pathology. PeerJ 1:e167. <https://doi.org/10.7717/peerj.167>.
- Lee E, Helt GA, Reese JT, Munoz-Torres MC, Childers CP, Buels RM, Stein L, Holmes IH, Elsik CG, Lewis SE. 2013. Web Apollo: a web-based genomic annotation editing platform. Genome Biol 14:R93. <https://doi.org/10.1186/gb-2013-14-8-r93>.
- Stummeyer K, Schwarzer D, Claus H, Vogel U, Gerardy-Schahn R, Mühlenhoff M. 2006. Evolution of bacteriophages infecting encapsulated bacteria: lessons from *Escherichia coli* K1-specific phages. Mol Microbiol 60:1123–1135. <https://doi.org/10.1111/j.1365-2958.2006.05173.x>.
- Scholl D, Rogers S, Adhya S, Merrill CR. 2001. Bacteriophage K1-5 encodes two different tail fiber proteins, allowing it to infect and replicate on both K1 and K5 strains of *Escherichia coli*. J Virol 75:2509–2515. <https://doi.org/10.1128/JVI.75.6.2509-2515.2001>.