

CORRECTION


Correction: Early onset hyperuricemia is a prognostic marker for kidney graft failure: Propensity score matching analysis in a Korean multicenter cohort

Miyeun Han, Jung Pyo Lee, Seokwoo Park, Yunmi Kim, Yong Chul Kim, Curie Ahn, Duck Jong Han, Jongwon Ha, In Mok Jung, Chun Soo Lim, Yon Su Kim, Young Hoon Kim, Yun Kyu Oh

The affiliation for the last author is incorrect. Yun Kyu Oh is not affiliated with #5 but with #2 Department of Surgery, Seoul National University Boramae Medical Center, Seoul, Korea.

Reference

1. Han M, Lee JP, Park S, Kim Y, Kim YC, Ahn C, et al. (2017) Early onset hyperuricemia is a prognostic marker for kidney graft failure: Propensity score matching analysis in a Korean multicenter cohort. PLoS ONE 12(5): e0176786. doi:[10.1371/journal.pone.0176786](https://doi.org/10.1371/journal.pone.0176786) PMID: [28467476](https://pubmed.ncbi.nlm.nih.gov/28467476/)



OPEN ACCESS

Citation: Han M, Lee JP, Park S, Kim Y, Kim YC, Ahn C, et al. (2017) Correction: Early onset hyperuricemia is a prognostic marker for kidney graft failure: Propensity score matching analysis in a Korean multicenter cohort. PLoS ONE 12(6): e0179779. <https://doi.org/10.1371/journal.pone.0179779>

Published: June 12, 2017

Copyright: © 2017 Han et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.