

Meeting abstract

Open Access

T Cell transformation by herpesvirus saimiri requires STAT5 pathways

D Lengfelder*, B Vogel, ED Mazumder, B Scholz, E Heck, M Schmidt, I Müller-Fleckenstein, B Fleckenstein, B Biesinger and A Ensser

Address: Friedrich-Alexander Universität Erlangen-Nürnberg, Institut für Klinische und Molekulare Virologie, Erlangen, Germany

* Corresponding author

from 12th Joint Meeting of the Signal Transduction Society (STS). Signal Transduction: Receptors, Mediators and Genes Weimar, Germany. 29–31 October 2008

Published: 26 February 2009

Cell Communication and Signaling 2009, **7**(Suppl 1):A80 doi:10.1186/1478-811X-7-S1-A80

This abstract is available from: <http://www.biosignaling.com/content/7/S1/A80>

© 2009 Lengfelder et al; licensee BioMed Central Ltd.

Aberrant growth stimulation by Signal Transducers and Activators of Transcription (STAT) is implicated in human carcinogenesis. The viral effector oncoprotein Tip, which binds to and is phosphorylated by the T-cell lymphocyte kinase Lck, is crucial in the activation of the cellular STAT5 signalling pathway in Herpesvirus-transformed T-cell lines, a model for malignant T-cell proliferation. While Tip-Lck interaction is required for transformation by HVS, mutation of Tip tyrosine residues had distinct effects. Two defined mutations of Tip can direct the phosphorylation and activation of either STAT3 or STAT5 by Lck. Tip Y114 mutation to phenylalanine (TipY114F) abolished the constitutive STAT3 activation observed in HVS-wildtype transformed T cells. Conversely, TipY114F enhanced the efficiency of human T cell transformation in absence of exogenous interleukin-2 (IL-2). In contrast, mutation of the major phosphorylation site, Y127, in Tip is compatible with viral transformation only when IL-2 is supplemented. This growth factor requirement correlated with STAT5 activation by Lck and Tip. Our current work focuses on the synergy among Tip, Lck, STAT5 and JAK family kinases. Interaction analysis will address whether Tip or the Y127F mutant mediate interactions between STAT5 and Lck or other Src family members. The role of the JAK-STAT pathway was elucidated using specific JAK inhibitors. We assume that, for viral transformation, Tip preferentially targets and reprograms the STAT5 pathway, which is central for T-cell growth and homeostasis. Comparable modulation of the STAT5 signalling may be involved in other forms of malignancy and cell differentiation; their

analysis may therefore be a useful model for the development of new therapeutics.