

ORAL PRESENTATION

Open Access

Aortic valve replacement in octogenarians: performance analysis of EuroSCORE II

D Hernandez-Vaquero*, R Diaz, JC Llosa

From 23rd World Congress of the World Society of Cardio-Thoracic Surgeons
Split, Croatia. 12-15 September 2013

Background

EuroSCORE has been proposed to identify patients at high risk for surgical aortic valve replacement and estimate for them the risk-benefit of percutaneous implantation. However, it has been clearly demonstrated that this system overestimates mortality, particularly in the elderly population undergoing surgery for aortic valve replacement. An actualized model called EuroSCORE II has been recently proposed to predict mortality after cardiac surgery but it has not been validated yet in the elderly population undergoing aortic valve replacement. We aimed to validate the new risk model in an octogenarian population who underwent surgery for aortic valve replacement in our center.

Methods

From March 2005 to March 2013, 378 consecutive octogenarian patients underwent aortic valve replacement in our center. Calibration (Hosmer-Lemeshow test and risk-adjusted mortality ratio) and discrimination (area under the receiver-operating characteristic curve) were calculated to assess the capacity of EuroSCORE II to predict 30-day mortality. Moreover, the same analysis was made in a subgroup of patients who underwent isolated aortic valve replacement.

Results

Observed mortality was 8,7% in the overall population and the predicted mortality by EuroSCORE II was 6,31% so that it underestimated the real mortality rate. However, the general calibration assessed by Hosmer-Lemeshow test was not bad ($p=0,15$) and the discriminatory power was very good (C -Statistic= 0,90). Assessing calibration by quartiles of risk, an adequate predictive power was

observed in the first three quartiles but not in the fourth (Hosmer-Lemeshow, $p=0,032$; risk adjusted mortality ratio=1,93). Similar results were obtained in the subgroup of patients who underwent isolated replacement.

Conclusions

EuroSCORE II is a good model for predicting mortality after surgery for aortic valve replacement in octogenarians. However, this system underestimates the mortality rate in the highest quartile of risk.

Published: 11 September 2013

doi:10.1186/1749-8090-8-S1-O102

Cite this article as: Hernandez-Vaquero et al.: Aortic valve replacement in octogenarians: performance analysis of EuroSCORE II. *Journal of Cardiothoracic Surgery* 2013 **8**(Suppl 1):O102.

Submit your next manuscript to BioMed Central
and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit



* Correspondence: dhvaquero@gmail.com

Cardiac Surgery Department, Central University Hospital of Asturias, Oviedo, Spain