

POSTER PRESENTATION

Open Access

Cardiovascular magnetic resonance myocardial feature tracking predicts severity of wall motion abnormalities following acute coronary syndromes

Kalpa De Silva^{1*}, Asela Bandara¹, Andreas Schuster², Pablo Lamata³, Roy Jogiya², Shazia T Hussain², Kaleab N Asrress¹, Nic Smith², Michael Marber¹, Eike Nagel², Simon Redwood¹, Divaka Perera¹, Sven Plein^{4,2}

From 16th Annual SCMR Scientific Sessions
San Francisco, CA, USA. 31 January - 3 February 2013

Background

Left ventricular wall motion assessment following acute myocardial infarction (AMI) allows prediction of functional recovery. Current methods are based upon visual assessment, with inherent operator variability. CMR myocardial feature tracking (CMR-FT) is a recently introduced technique for tissue voxel motion tracking on standard steady-state free precession (SSFP) images to derive circumferential and radial myocardial mechanics. We sought to determine whether CMR-FT could be used as a quantitative measure of wall motion assessment and predict recovery in function following AMI.

Methods

Patients presenting with non-ST elevation myocardial infarction (NSTEMI) were studied using a 3 Tesla Phillips Achieva system with Multi-transmit[®] technology and a 32-channel receiver coil, immediately prior to, and 3-months following, percutaneous coronary revascularization (PCI). Cine images were acquired using a SSFP cine technique, with analysis performed on the left-ventricular (LV) short-axis stack, covering the LV from apex to base. The acquisition pulse sequence provided a typical spatial resolution of 1.8 x 1.8 x 8 mm with a 2mm inter-slice gap and a temporal resolution of 50 frames per second. A 5-point wall motion scoring system was used to determine regional wall motion abnormalities (RWMA), with derivation of an indexed wall motion score (WMSI) specific to

the region of infarction, according to the AHA 16-segment model. Myocardial strain parameters were derived following automated endo- and epi-cardial wall motion tracking of the SSFP cine short axis stack using dedicated software (Tomtec[™], Germany). Pre-PCI LV short axis circumferential (Ecc) and radial (Err) peak strains were related to WMSI using a Pearson correlation analysis. Visual and quantitative analyses were performed by two blinded reviewers.

Results

18 patients (Mean age 57±9 years, 78% male) with a mean Troponin T and LV ejection fraction (LVEF) of 1.6 ±1.1µg/L and 56±8% respectively completed the scanning protocol. Pre-PCI CMR-FT strain parameters were derived in all cases. Mean index WMSI was 2.57±0.03 reducing to 1.98±0.03 at follow-up (p=0.004). Mean index Ecc and Err were -23.1±15.5 and 21.0±15.5 respectively. Neither Ecc nor Err correlated with global LVEF (R=0.10, p=0.69 and R=-0.03, p=0.89, respectively). However, Ecc (R=0.51, p=0.04) and Err (R=-0.50, p=0.04) showed a linear and inverse linear correlation, respectively, with the regional WMSI. In addition, index Ecc correlated with the 3-month WMSI (R=0.55, p=0.03).

Conclusions

CMR-FT may be used to quantify severity of wall motion following an ACS. Circumferential strain parameters are more accurate in predicting infarct related wall motion abnormalities and recovery in function.

¹Cardiovascular Division, St. Thomas' Hospital, King's College London, London, UK

Full list of author information is available at the end of the article

Funding

Heart Research UK.

Author details

¹Cardiovascular Division, St. Thomas' Hospital, King's College London, London, UK. ²Imaging Sciences, St. Thomas' Hospital, King's College London, London, UK. ³Department of Computer Science, University of Oxford, Oxford, UK. ⁴Multidisciplinary Cardiovascular Research Centre, Leeds Institute of Genetics, Health and Therapeutics, Leeds, UK.

Published: 30 January 2013

doi:10.1186/1532-429X-15-S1-P200

Cite this article as: De Silva et al.: Cardiovascular magnetic resonance myocardial feature tracking predicts severity of wall motion abnormalities following acute coronary syndromes. *Journal of Cardiovascular Magnetic Resonance* 2013 **15**(Suppl 1):P200.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

