

Original Article



Population attributable fraction of indicators for musculoskeletal diseases: a cross-sectional study of fishers in Korea

Jaehoo Lee ^{1,2}, Bohyun Sim ^{1,2}, Bonggyun Ju ^{1,2}, Chul Gab Lee ¹,
Ki-Soo Park ^{3,4}, Mi-Ji Kim ^{3,4}, Jeong Ho Kim ^{5,6}, Kunhyung Kim ^{5,6}, and
Hansoo Song ^{1,2*}

OPEN ACCESS

Received: Apr 22, 2022
1st Revised: Jul 26, 2022
2nd Revised: Aug 10, 2022
Accepted: Aug 12, 2022
Published online: Sep 7, 2022

*Correspondence:

Hansoo Song

Department of Occupational and
Environmental Medicine, Chosun University
Hospital, 365 Pilmun-daero, Dong-gu,
Gwangju 61453, Korea.
Email: oemsong@chosun.ac.kr

Copyright © 2022 Korean Society of
Occupational & Environmental Medicine
This is an Open Access article distributed
under the terms of the Creative Commons
Attribution Non-Commercial License (<https://creativecommons.org/licenses/by-nc/4.0/>)
which permits unrestricted non-commercial
use, distribution, and reproduction in any
medium, provided the original work is properly
cited.

ORCID iDs

Jaehoo Lee
<https://orcid.org/0000-0001-7365-8952>
Bohyun Sim
<https://orcid.org/0000-0001-6996-018X>
Bonggyun Ju
<https://orcid.org/0000-0003-0433-8554>
Chul Gab Lee
<https://orcid.org/0000-0001-7017-2623>
Ki-Soo Park
<https://orcid.org/0000-0001-5571-3639>
Mi-Ji Kim
<https://orcid.org/0000-0002-8646-832X>
Jeong Ho Kim
<https://orcid.org/0000-0003-3876-2724>

¹Department of Occupational and Environmental Medicine, Chosun University Hospital, Gwangju, Korea

²Center for Fishermen's Safety and Health, Chosun University Hospital, Gwangju, Korea

³Department of Preventive Medicine, College of Medicine and Institute of Health Science, Gyeongsang National University, Jinju, Korea

⁴Center for Fishermen's Safety and Health, Gyeongsang National University Hospital, Jinju, Korea

⁵Department of Occupational and Environmental Medicine & Institute of Environmental and Occupational Medicine, Inje University Busan Paik Hospital, Busan, Korea

⁶Center for Fishermen's Safety and Health, Inje University Busan Paik Hospital, Busan, Korea

ABSTRACT



Background: The musculoskeletal disease (MSD) burden is an important health problem among Korean fishers. We aimed to investigate the indicators of the prevalence of MSD and contributions of significant indicators to MSD in Korean fishers.

Methods: This cross-section study included 927 fishers (male, 371; female, 556) aged 40 to 79 years who were enrolled from 3 fishery safety and health centers. The outcome variable was one-year prevalence of MSD in 5 body parts (the neck, shoulder, hand, back, and knee). Independent variables were sex, age, educational attainment, household income, job classification, employment xlink:type, hazardous working environment (cold, heat, and noise), ergonomic risk by the 5 body parts, anxiety disorder, depression, hypertension, diabetes, and hyperlipidemia. The adjusted odds ratio of MSDs by the 5 body parts were calculated using multiple logistic regression analysis. We computed the population attributable fraction (PAF) for each indicators of MSDs using binary regression models.

Results: The one-year prevalence of MSD in the neck, shoulder, hand, back, and knee was 7.8%, 17.8%, 7.8%, 27.2%, and 16.2% in males vs. 16.4%, 28.1%, 23.0%, 38.7%, and 30.0% in females, respectively. The ergonomic risk PAF according to the body parts ranged from 22.8%–59.6% in males and 22.8%–50.3% in female. Mental diseases showed a significant PAF for all body parts only among female (PAF 9.1%–21.4%). Cold exposure showed a significant PAF for the neck, shoulder, and hand MSD only among female (25.6%–26.8%). Age was not a significant indicator except for the knee MSD among female.

Conclusions: Ergonomic risk contributed majorly as indicators of MSDs in both sexes of fishers. Mental disease and cold exposure were indicators of MSDs only among female fishers. This information may be important for determining priority risk groups for the prevention of work-related MSD among Korean fishers.

Keywords: Ergonomic risk; Fisher; Musculoskeletal disease; Population attributable fraction

Kunhyung Kim <https://orcid.org/0000-0001-9060-7233>Hansoo Song <https://orcid.org/0000-0001-6000-1572>

Abbreviations

CI: confidence interval; KOSHA: Korea Occupational Safety and Health Agency; MSD: musculoskeletal disease; PAF: population attributable fraction; PAR: population attributable risk; OR: odds ratio.

Funding

This study was supported by research funds from the Ministry of Oceans and Fisheries Affairs of the Republic of Korea.

Competing interests

The authors declare that they have no competing interests.

Author contributions

Conceptualization: Song H. Data curation: Song H, Lee J. Formal analysis: Song H. Funding acquisition: Song H. Investigation: Song H, Lee CG, Lee J, Sim B, Joo B, Park KS, Kim MJ, Kim JH, Kim K. Methodology: Song H, Lee CG. Software: Lee J. Validation: Song H, Lee J. Visualization: Song H. Writing - original draft: Lee J. Writing - review & editing: Song H, Park KS, Kim J, Lee CG.

BACKGROUND

Although musculoskeletal disease (MSD) is non-fatal, it is an important public health problem because it accounts for a high proportion of the years lived with disability in South Korea.¹ The direct and indirect costs of MSD was 9.0% of the Korean gross domestic product, and highest among the 20–64 years old working group.² Therefore occupation-based approach is important for preventing MSD. Manufacturing, construction, hotel and restaurants, education, wholesale and retail, public administration and defense,³ and agricultural/fishery workers⁴ are high-risk work sector groups for MSD. However, the MSD among small-scale self-employed agricultural and fishery workers has received less attention compared to that among employed workers. In particular, in the fishery industry, MSD is relatively unknown, and the related research is few. However, some previous studies have shown that fisher's MSD are very serious. In Sweden, the United States, United Kingdom, and Sri Lanka, the prevalence of low back pain of 50%–80% has been reported.⁵ A cohort study of MSD in Danish fishermen, using hospitalization data, reported a high incidence of knee arthrosis, thoracolumbar disc disease, carpal tunnel syndrome.⁶ In our previous study, the prevalence of collapsed lumbar disc in fishers aged 40–69 years was 23.7%.⁷ Fishers include those working in fishing vessel fishery, aquaculture, diving, and seafood processing. The fishers' exposures vary in ergonomic stress, including net or line pulling, heavy manual lifting, and repetitive hand exertion. In addition, fishery is known as a very high-risk occupation with fatal and non-fatal injuries. Most of the previous studies on MSDs have focused only on relative risk, and few studies have revealed how much disease can be prevented when risk factors are removed.^{8,9} Therefore, this study aimed to determine the indicators of MSD among fishers, an occupation at high-risk of MSD, and identify the association between major indicators and MSD. Furthermore, we calculated the extent to which significant indicators contribute to MSD, to determine the priority of prevention.

METHODS

Subjects

In this cross-sectional study, 957 fishers from 3 fishery safety and health centers funded by Ministry of Oceans and Fisheries voluntarily participated in a fisher's health survey, from June 2018 to August 2020. The inclusion criteria were subjects aged 40–79 years whose occupation could be identified as fishery. Accordingly, 22, 3, 3, and 2 subjects aged < 40 years, > 80 years, with unclear occupations, and insufficient questionnaire details, respectively, were excluded. Finally, a total of 927 subjects were included in the study.

Variables

Data were collected through a questionnaire and medical examination. The dependent variable, MSD in 5 body parts (the neck, shoulders, hands, back, and knees), was defined as an experience of pain of > 1 week or > 1 per month, moderate pain intensity or higher, and an experience of medical care during the past 1 year using the musculoskeletal symptom questionnaire developed by the Korea Occupational Safety and Health Agency (KOSHA).¹⁰ Independent variables included sex, age, household income, employment type, job classification, cold exposure,¹¹ noise exposure,¹¹ heat exposure,¹¹ ergonomic risk in the 5 body parts, mental disease (depression, anxiety) obtained by questionnaire (**Supplementary Table 1**), and chronic diseases (hypertension, diabetes mellitus, hyperlipidemia) obtained by medical examination and medication history. Household income was divided into quartiles.

Employment was divided into owner (self-employed), owner's household members, and employees. Cold, heat, and noise exposure were divided into either no exposure or 3 additional categories by counting the number of days of > 2 hours daily exposure, over the past year (**Supplementary Data 1**). Job classification were as follows: aquaculture (shellfish, fish, and seaweeds); and fishing vessel fishery (with trap gill net, long line, Haenyeo [Korean female sea divers], and oyster shucker). For comparison between job classifications, aquaculture (shellfish), with a higher income level and fishery mechanization rate, was used as the reference. Depression was defined using Patient Health Questionnaire-9 (cut-off point 10).¹² Anxiety disorder was defined using the Generalized Anxiety Disorder 7-item (cut-off point 6).¹³ Ergonomic risk was investigated using a structured face-to-face questionnaire. Ergonomic risk factors composed of the intensity of the task, average daily working hours, and number of working days during the recent year. We identified ergonomic risk for each of 5 body parts (the neck, shoulders, hands, back, knees). A total ergonomic risk score was calculated by the following formula: A Score (point) = The Intensity of the Task (mild = 0.5, moderate = 1.0, severe = 1.5) × Average Daily Working Hours (< 2 hours = 1, 2–3 hours = 2, 4–5 hours = 3, 6–7 hours = 4, ≥ 8 hours = 5) × number of working days during the recent year. A score of ≥ 250 was defined as a significant musculoskeletal burden task. A 250-point score implies that the fishers had continuously performed an average of 2 hours of work per day for the last year. Hypertension was defined based on a systolic blood pressure ≥ 140 mmHg or diastolic blood pressure ≥ 90 mmHg. Diabetes mellitus was defined based on hemoglobin A1c ≥ 6.5% or fasting blood glucose ≥ 126 mg/dL. Hyperlipidemia was defined based on a low-density lipoprotein cholesterol level ≥ 150 mg/dL or triglyceride ≥ 500 mg/dL or total cholesterol minus high-density lipoprotein ≥ 190 mg/dL. Patients with hypertension, diabetes mellitus, and hyperlipidemia included those who had been on medications.

Statistical analysis

A χ^2 test was performed to evaluate the univariate association between ergonomic risk and MSDs. To find valuable and clear indicators for comparative analysis by MSDs xlink:type and sex, odds ratios (ORs) and 95% confidence intervals (CIs) were calculated for model 1 (adjusted for sex and age) and model 2 (adjusted for major variables) using binary multiple logistic regression analysis. To assess the population attributable risk (PAR) and 95% CI of significant indicators, we used STATA module as described by Newson to compute PARs from binary regression models.¹⁴ PAR was calculated as the difference in prevalence between scenario-0 (when exposed to all harmful factors) and scenario-1 (when each of the harmful factors was excluded). Furthermore, the prevalence was calculated for each condition, and PAR was obtained from the difference. The population attributable fraction (PAF) of each indicators of MSDs was calculated as follows.

$$PAF = \frac{PAR \text{ Exposed}}{\text{The Prevalence in Total Subjects}} \times 100$$

For statistical analysis, STATA 16 version (StataCorp LLC., College Station, TX, USA) was used.

Ethics statement

The protocol of this study was reviewed and approved by the Institutional Review Board of Chosun University Hospital (approval No. CHOSUN 2018-5-014). Informed consent was obtained in writing from the study subjects.

RESULTS

Table 1 shows the characteristics of the study subjects. A total of 927 subjects (mean age: 61.1, standard deviation: 8.0 years, with no significant difference by sex), 371 (40%)

Table 1. Characteristic of subjects

Variables		Male (n = 371, 40.0)	Female (n = 556, 60.0)	Total	p-value
Age (yr)	40–49	50 (13.5)	55 (9.9)	105 (11.3)	0.052
	50–59	112 (30.2)	212 (38.1)	324 (35.0)	
	60–69	160 (43.1)	228 (41.0)	388 (41.9)	
	70–79	49 (13.2)	61 (11.0)	110 (11.9)	
	Mean ± SD	60.2 ± 8.4	60.0 ± 7.8	61.1 ± 8.0	
Educational attainment	High	176 (47.4)	158 (28.4)	334 (36.0)	< 0.001
	Middle	119 (32.1)	161 (29.0)	280 (30.2)	
	Elementary	76 (20.5)	237 (42.6)	313 (33.8)	
Household income (10,000 won)	< 1,500	63 (17.0)	147 (26.4)	210 (22.7)	< 0.001
	1,500–2,399	66 (17.8)	131 (23.6)	197 (21.3)	
	2,400–4,000	94 (25.3)	139 (25.0)	233 (25.1)	
	> 4,000	128 (34.5)	93 (16.7)	221 (23.8)	
	No answer	20 (5.4)	46 (8.3)	66 (7.1)	
Employment xlink:type	Owner	287 (77.4)	145 (26.1)	432 (46.6)	< 0.001
	Household member	28 (7.5)	215 (38.7)	243 (26.2)	
	Employee	56 (15.1)	196 (35.3)	252 (27.2)	
Job classification ^a	A	51 (13.7)	51 (9.2)	102 (11.0)	< 0.001
	B	14 (3.8)	17 (3.1)	31 (3.3)	
	C	82 (22.1)	62 (11.2)	144 (15.5)	
	D	137 (36.9)	114 (20.5)	251 (27.1)	
	E	62 (16.7)	42 (7.6)	104 (11.2)	
	F	0 (0.0)	98 (17.6)	98 (10.6)	
	G	25 (6.7)	172 (30.9)	197 (21.3)	
Cold exposure ^b (days/1 year)	No	115 (31.0)	212 (38.1)	327 (35.3)	0.003
	1–60	151 (40.7)	167 (30.0)	318 (34.3)	
	61–119	66 (17.8)	95 (17.1)	161 (17.4)	
	≥ 120	39 (10.5)	82 (14.7)	121 (13.1)	
Noise exposure ^c (days/1 year)	No	238 (64.2)	371 (66.7)	609 (65.7)	0.012
	1–60	66 (17.8)	60 (10.8)	126 (13.6)	
	61–199	46 (12.4)	93 (16.7)	139 (15.0)	
	≥ 200	21 (5.7)	32 (5.8)	53 (5.7)	
Heat exposure ^d (days/1 year)	No	157 (42.5)	276 (63.7)	433 (46.9)	0.126
	1–29	92 (24.9)	112 (54.9)	204 (22.1)	
	30–89	68 (18.4)	87 (56.1)	155 (16.8)	
	≥ 90	52 (14.1)	80 (60.6)	132 (14.3)	
Chronic disease ^e	No	138 (39.3)	218 (40.1)	356 (39.8)	0.821
	Yes	213 (60.7)	326 (59.9)	539 (60.2)	
Hypertension	No	217 (61.8)	335 (61.6)	552 (61.7)	0.942
	Yes	134 (38.2)	209 (38.4)	343 (38.3)	
Diabetes mellitus	No	272 (77.5)	473 (86.9)	745 (83.2)	< 0.001
	Yes	79 (22.5)	71 (13.1)	150 (16.8)	
Hyperlipidemia	No	281 (80.1)	418 (76.8)	699 (78.1)	0.256
	Yes	70 (19.9)	126 (23.2)	196 (21.9)	
Mental disease ^f	No	338 (91.1)	463 (83.3)	801 (86.4)	0.001
	Yes	33 (8.9)	93 (16.7)	126 (13.6)	
Depression	No	354 (95.4)	524 (94.2)	878 (94.7)	0.434
	Yes	17 (4.6)	32 (5.8)	49 (5.3)	
Anxiety disorder	No	341 (91.9)	469 (84.4)	810 (87.4)	0.001
	Yes	30 (8.1)	87 (15.6)	117 (12.6)	

Values are presented as number (%). The p-value by χ^2 test.

^aA: Aquaculture (shellfish), B: Aquaculture (fish), C: Aquaculture (seaweeds), D: Fishing vessel fishery with trap gill net, E: Fishing vessel fishery with long line, F: Haenyeo (Korean female sea diver), G: Oyster shucker. ^bLow temperatures, whether indoors or outdoors. ^cNoise that is loud enough to raise your voice when speaking to others. ^dTemperatures high enough to make you sweat even when you are not working. ^eIf patient has hypertension and/or diabetes mellitus. ^fIf the patient has depression and/or anxiety disorder.

male, and 556 (60%) female were included in this analysis. By sex, there was a statistically significant difference in educational attainment, highest among high school male (47.4%) and elementary school female (42.6%); household income, with the highest of 40 million won among male (34.5%) than female (16.7%); employment xlink:type, highest among male owners (77.4%) and female household members (38.7%); and by job classification, Haenyeo were all female while the female proportion among oyster shucker was 87.3%. In the hazardous working environment, males are more exposed to cold (69.0%), heat (57.5%), and noise (35.8%) than their female counterparts. By sex, male and female differed significantly in diabetes mellitus (22.5% vs. 13.1%) and depression prevalence (8.9% vs. 16.7%), respectively.

Table 2 shows the prevalence of MSDs by the 5 body parts according to each variable. Female showed a higher prevalence of MSDs in all body parts than male ($p < 0.05$ by χ^2 test). By age, significant difference was observed in the knees MSD prevalence only ($p < 0.05$ by χ^2 test), which was lowest in those aged 40–49 years (17.1%) and highest in those aged 60–69 years (27.3%). The prevalence of neck, back, and knees MSD decreased at higher educational attainment. As household income increased, the prevalence of MSDs decreased in all body parts except the neck. By employment xlink:type, the prevalence was lowest among owners in all body parts. As the ergonomic risk increased, the prevalence of MSD in all body parts increased significantly. As the duration of exposure to cold increased, the prevalence of MSD in the hands, back, and knees increased significantly. The effects of cold exposure may have been confounded by outdoor work.

Therefore, we tried to confirm whether cold exposure was related to musculoskeletal disorders through sensitivity analysis including heat exposure. However, there was no dose-response association of noise and heat exposure with MSD. Chronic diseases including hypertension, diabetes mellitus, and hyperlipidemia were not associated with MSDs. On the contrary, the prevalence of the back and knees MSD was lower in patients with diabetes mellitus. Mental diseases including depression and anxiety disorder significantly increased the prevalence of MSDs in all body parts. We presented the results by sex in **Supplementary Tables 2 and 3**, respectively.

Table 3 shows the age and sex adjusted ORs and 95% CIs for MSDs of the 5 body parts using binary logistic regression (model 1). Educational attainment showed a significant relationship with back MSD. Fishers with lower educational attainment had higher OR of MSDs in the back. According to the job classification, neck MSD had significantly higher ORs in aquaculture (fish), fishing vessel fishery with long line, and oyster shucker, in hand MSD. Knees MSD had significantly higher OR in aquaculture (seaweeds) and fishing vessel fishery with long line. In particular, the fishing vessel fishery with trap gill net and Haenyeo showed a low OR overall. Ergonomic risk point grade showed a dose-response relationship with MSDs in all body parts. Mental diseases showed a significant association with MSDs in all body parts. Cold exposure showed an association with MSDs in neck and hands MSD. We presented the results by sex in **Supplementary Tables 4 and 5**, respectively.

Table 4 shows the adjusted OR by ergonomic risk, age, educational attainment, mental diseases, and cold exposure for MSDs in the relevant body parts by sex, respectively (model 2). In both sexes, significant OR occurred in all body parts with ergonomic risk. The adjusted OR of the highest ergonomic risk point grade (≥ 750 points) in male occurred with the neck 6.28 (95% CI: 1.92–20.51); shoulders 6.18 (95% CI: 2.83–13.46); hands 4.05 (95% CI: 1.29–12.71); back 5.86 (95% CI: 2.83–12.10); and knees 3.07 (95% CI: 1.35–6.97). The OR of

Indicators for musculoskeletal diseases among fishers in Korea

Table 2. The association between variables and musculoskeletal disease of 5 body parts

Variables		Neck		Shoulder		Hand		Back		Knee	
		%	p-value	%	p-value	%	p-value	%	p-value	%	p-value
Sex	Male	7.8	< 0.001	17.8	< 0.001	7.8	< 0.001	27.2	< 0.001	16.2	< 0.001
	Female	16.4		28.1		23.0		38.7		30.0	
Age (years)	40–49	13.3	0.197	24.8	0.753	21.9	0.141	23.8	0.077	17.1	0.039
	50–59	13.9		23.8		17.3		32.7		22.8	
	60–69	13.9		24.5		16.0		39.2		27.3	
	70–79	6.4		21.8		14.5		30.0		26.4	
Education	High	12.9	0.255	23.7	0.309	17.1	0.709	25.4	< 0.001	18.6	< 0.001
	Middle	9.6		20.7		15.4		35.4		23.9	
	Elementary	16.0		27.2		18.2		42.2		31.3	
Household income (10,000 won)	< 1,500	16.7	0.011	26.7	< 0.001	17.6	0.004	43.3	< 0.001	32.9	< 0.001
	1,500–2,399	14.2		27.4		21.8		38.6		26.4	
	2,400–4,000	11.2		25.8		19.3		32.6		22.7	
	> 4,000	13.1		21.7		13.1		31.7		23.1	
	No answer	3.0		6.1		4.5		4.5		3.0	
Employment	Owner	10.2	0.024	20.4	0.049	11.8	< 0.001	30.1	0.015	21.1	0.069
	Member	13.2		25.9		20.6		41.2		28.4	
	Employee	17.5		28.2		22.2		34.1		26.6	
Job classification ^a	A	11.8	0.001	21.6	0.005	9.8	< 0.001	34.3	0.001	20.6	< 0.001
	B	29.0		29.0		38.7		45.2		19.4	
	C	17.4		27.8		16.0		40.3		32.6	
	D	6.4		14.7		11.2		23.9		15.1	
	E	14.4		29.8		18.3		46.2		32.7	
	F	11.2		26.5		15.3		35.7		31.6	
	G	16.2		28.9		25.4		33.5		25.4	
Ergonomic risk by each body parts	< 250	7.0	< 0.001	16.3	< 0.001	8.8	< 0.001	19.8	< 0.001	19.7	< 0.001
	250–499	14.4		28.7		19.8		35.4		25.0	
	500–749	20.0		23.4		21.6		47.3		29.1	
	≥ 750	27.5		44.3		30.9		53.8		40.7	
Cold exposure (days/1 year)	No	11.6	0.053	22.3	0.030	15.3	0.004	36.4	0.246	25.7	0.127
	1–60	11.3		20.8		14.2		26.4		18.9	
	61–119	14.9		28.6		17.4		39.8		26.7	
	≥ 120	18.2		30.6		28.1		40.5		33.1	
Noise exposure (days/1 year)	No	12.6	0.686	23.6	0.686	17.1	0.618	35.5	0.091	25.6	0.471
	1–60	16.7		28.6		17.5		37.3		22.2	
	61–199	12.9		23.0		18.0		26.6		20.1	
	≥ 200	7.5		18.9		11.3		30.2		28.3	
Heat exposure (days/1 year)	No	12.5	0.432	24.2	0.760	17.1	0.324	26.3	0.212	26.3	0.468
	1–29	11.3		21.6		13.7		25.0		17.6	
	30–89	16.1		25.8		16.8		28.4		21.3	
	≥ 90	13.6		25.0		22.0		48.5		33.3	
Chronic disease ^b	No	12.1	0.427	24.4	0.913	16.6	0.791	34.6	0.721	21.9	0.148
	Yes	13.9		24.1		17.3		33.4		26.2	
Hypertension	No	13.4	0.804	25.5	0.250	17.6	0.551	33.9	0.986	23.7	0.515
	Yes	12.8		22.2		16.0		33.8		25.7	
Diabetes mellitus	No	14.0	0.127	25.2	0.124	17.6	0.286	35.3	0.041	25.8	0.043
	Yes	9.3		19.3		14.0		26.7		18.0	
Hyperlipidemia	No	12.3	0.141	24.0	0.780	16.7	0.712	34.0	0.817	23.3	0.131
	Yes	16.3		25.0		17.9		33.2		28.6	
Mental disease ^c	No	10.4	< 0.001	20.6	< 0.001	14.5	< 0.001	31.2	< 0.001	22.1	< 0.001
	Yes	29.4		45.2		32.5		52.4		39.7	
Depression	No	12.1	0.001	22.8	< 0.001	16.2	0.009	33.3	0.024	23.9	0.088
	Yes	28.6		44.9		30.6		49.0		34.7	
Anxiety disorder	No	10.4	< 0.001	20.6	< 0.001	14.6	< 0.001	31.1	< 0.001	22.0	< 0.001
	Yes	30.8		47.0		33.3		54.7		41.9	
Total		12.9		23.9		16.9		34.1		24.5	

The *p*-value by χ^2 test in employment xlink:type, job classification, chronic disease, hypertension, diabetes mellitus, hyperlipidemia, mental disease, depression, anxiety disorder. The *p*-value by Cochran-Armitage trend test in age, educational attainment, household income, cold exposure, noise exposure, heat exposure. ^aA: Aquaculture (shellfish), B: Aquaculture (fish), C: Aquaculture (seaweeds), D: Fishing vessel fishery with trap gill net, E: Fishing vessel fishery with long line, F: Haenyeo (Korean female sea diver), G: Oyster shucker. ^bIf patient has hypertension and/or diabetes mellitus. ^cIf patient has depression and/or anxiety disorder.

Indicators for musculoskeletal diseases among fishers in Korea

Table 3. Age, sex adjusted multiple logistic regression analysis (model 1)

Variable	Neck		Shoulder		Hand		Back		Knee	
	OR	95% CI	OR	95% CI	OR	95% CI	OR	95% CI	OR	95% CI
Educational attainment										
High	1.00		1.00		1.00		1.00		1.00	
Middle	0.22	0.40–1.24	0.80	0.52–1.23	0.88	0.53–1.45	1.46	0.99–2.16	1.17	0.75–1.81
Elementary	0.40	0.72–2.27	1.06	0.67–1.67	0.90	0.53–1.56	1.93	1.26–2.94	1.46	0.92–2.34
Household income										
< 1,500	1.00		1.00		1.00		1.00		1.00	
1,500–2,399	0.80	0.46–1.39	1.09	0.69–1.70	1.36	0.82–2.26	0.89	0.59–1.33	0.83	0.53–1.29
2,400–4,000	0.62	0.35–1.09	1.04	0.67–1.63	1.21	0.72–2.02	0.74	0.50–1.12	0.76	0.49–1.19
> 4,000	0.85	0.46–1.54	0.94	0.57–1.54	0.89	0.49–1.62	0.86	0.56–1.35	1.01	0.62–1.64
No answer	0.17	0.04–0.74	0.17	0.06–0.50	0.21	0.06–0.71	0.06	0.02–0.21	0.06	0.01–0.25
Employment xlink:type										
Owner	1.00		1.00		1.00		1.00		1.00	
Member	0.76	0.43–1.33	1.02	0.66–1.58	1.07	0.65–1.76	1.32	0.89–1.95	1.07	0.70–1.65
Employee	1.14	0.68–1.90	1.20	0.80–1.81	1.31	0.82–2.12	0.98	0.67–1.42	1.01	0.67–1.54
Job classification^a										
A	1.00		1.00		1.00		1.00		1.00	
B	2.90	1.06–7.97	1.41	0.56–3.55	6.68	2.41–18.5	1.24	0.53–2.88	0.73	0.26–2.07
C	1.70	0.80–3.63	1.42	0.77–2.61	1.91	0.85–4.32	1.25	0.73–2.16	1.88	1.02–3.48
D	0.54	0.24–1.20	0.61	0.34–1.23	1.22	0.56–2.70	0.51	0.30–0.86	0.58	0.31–1.08
E	1.36	0.59–3.11	1.63	0.86–3.10	2.59	1.11–6.04	1.60	0.90–2.86	1.95	1.02–3.75
F	0.77	0.29–2.11	0.89	0.41–1.91	0.87	0.32–2.35	0.55	0.28–1.09	0.76	0.36–1.62
G	1.05	0.50–2.20	1.17	0.65–2.10	2.23	1.05–4.74	0.64	0.38–1.10	0.84	0.46–1.54
Ergonomic risk										
< 250	1.00		1.00		1.00		1.00		1.00	
250–499	2.00	1.18–3.40	2.04	1.38–3.01	2.34	1.42–3.85	2.17	1.50–3.15	1.32	0.89–1.96
500–749	2.94	1.63–5.30	1.46	0.86–2.47	2.75	1.54–4.91	3.76	2.43–5.83	1.89	1.11–3.20
≥ 750	4.44	2.57–7.66	4.09	2.64–6.32	4.09	2.53–6.61	4.77	3.16–7.18	3.06	1.98–4.74
Noise exposure (days/1 year)										
No	1.00		1.00		1.00		1.00		1.00	
1–60	1.44	0.84–2.47	1.40	0.90–2.19	1.15	0.68–1.96	1.22	0.81–1.84	0.99	0.61–1.59
61–199	0.92	0.53–1.61	0.93	0.60–1.45	1.00	0.61–1.64	0.62	0.41–0.95	0.71	0.45–1.13
≥ 200	0.55	0.19–1.58	0.76	0.37–1.57	0.62	0.25–1.51	0.81	0.44–1.51	1.22	0.64–2.32
Cold exposure (days/1 year)										
No	1.00		1.00		1.00		1.00		1.00	
1–60	1.14	0.70–1.88	0.99	0.68–1.45	1.06	0.68–1.66	0.67	0.47–0.95	0.72	0.49–1.06
61–199	1.54	0.88–2.70	1.48	0.95–2.30	1.32	0.78–2.24	1.17	0.78–1.74	1.04	0.67–1.62
≥ 120	1.80	1.01–3.22	1.54	0.96–2.47	2.26	1.35–3.78	1.17	0.76–1.81	1.36	0.86–2.17
Heat exposure (days/1 year)										
No	1.00		1.00		1.00		1.00		1.00	
1–29	1.06	0.62–1.80	0.91	0.61–1.37	0.85	0.52–1.39	0.63	0.43–0.93	0.62	0.41–0.96
30–89	1.52	0.90–2.56	1.15	0.75–1.76	1.08	0.65–1.78	0.74	0.49–1.10	0.79	0.51–1.24
≥ 90	1.28	0.71–2.30	1.08	0.68–1.72	1.54	0.93–2.55	1.68	1.12–2.53	1.36	0.88–2.10
Chronic disease^b										
No	1.00		1.00		1.00		1.00		1.00	
Yes	1.22	0.81–1.83	1.00	0.73–1.37	1.10	0.76–1.60	0.94	0.70–1.25	1.25	0.91–1.73
Mental disease^c										
No	1.00		1.00		1.00		1.00		1.00	
Yes	3.25	2.06–5.12	3.00	2.02–4.46	2.60	1.68–4.02	2.18	1.48–3.21	2.06	1.38–3.08

OR: odds ratio; CI: confidence interval.

^aA: Aquaculture (shellfish), B: Aquaculture (fish), C: Aquaculture (seaweeds), D: Fishing vessel fishery with trap gill net, E: Fishing vessel fishery with long line, F: Haenyeo (Korean female sea diver), G: Oyster shucker. ^bIf patient has hypertension and/or diabetes mellitus. ^cIf patient has depression and/or anxiety disorder.

the highest ergonomic risk point grade (≥ 750 points) in female occurred with the neck 3.41 (95% CI: 1.77–6.57); shoulder 3.28 (95% CI: 1.87–5.77); hands 4.69 (95% CI: 2.64–8.34); back 4.06 (95% CI: 2.40–6.88); and knees 2.87 (95% CI: 1.66–4.96). The OR of mental diseases in female occurred with the neck 3.56 (95% CI: 2.07–6.12); shoulder 3.18 (95% CI: 1.96–5.15); hands 3.05 (95% CI: 1.83–5.10); back 2.58 (95% CI: 1.58–4.23); and knees 2.10

Indicators for musculoskeletal diseases among fishers in Korea

Table 4. Adjusted OR of major 5 potential indicators by multiple logistic regression by sex (model 2)

Variables	Neck		Shoulder		Hand		Back		Knee	
	OR	95% CI	OR	95% CI	OR	95% CI	OR	95% CI	OR	95% CI
Male										
Age (years)										
40–49	1.00		1.00		1.00		1.00		1.00	
50–59	0.73	0.19–2.83	0.83	0.31–2.21	0.57	0.17–1.89	0.69	0.30–1.62	0.42	0.15–1.17
60–69	1.00	0.22–4.52	1.04	0.36–2.98	0.31	0.07–1.30	0.72	0.29–1.77	0.41	0.14–1.21
≥ 70	0.52	0.06–4.23	1.46	0.39–5.44	1.65	0.34–7.95	0.98	0.33–2.97	0.64	0.18–2.27
Educational attainment										
High	1.00		1.00		1.00		1.00		1.00	
Middle	1.26	0.38–4.18	1.10	0.51–2.37	0.68	0.56–5.04	0.67	0.85–3.29	2.35	1.03–5.37
Elementary	2.72	0.80–9.30	1.03	0.43–2.45	0.90	0.23–3.57	1.86	0.87–3.95	3.13	1.25–7.84
Ergonomic risk										
< 250	1.00		1.00		1.00		1.00		1.00	
250–499	4.35	1.45–13.06	2.88	1.44–5.79	4.62	1.70–12.57	3.07	1.63–5.77	1.19	0.53–2.66
500–749	7.02	2.17–22.77	1.97	0.65–5.94	2.70	0.77–9.55	6.21	3.01–12.81	2.29	0.96–5.49
≥ 750	6.28	1.92–20.51	6.18	2.83–13.46	4.05	1.29–12.71	5.86	2.83–12.10	3.07	1.35–6.97
Cold exposure (days/1 year)										
No	1.00		1.00		1.00		1.00		1.00	
1–60	1.22	0.48–3.07	0.65	0.32–1.28	1.02	0.39–2.68	0.62	0.34–1.13	0.50	0.24–1.04
61–119	0.31	0.06–1.55	0.91	0.41–2.05	0.65	0.18–2.38	0.87	0.42–1.77	0.73	0.31–1.69
≥ 120	1.17	0.32–4.29	0.71	0.27–1.92	1.30	0.39–4.39	0.90	0.39–2.07	1.37	0.57–3.31
Mental disease ^a										
No	1.00		1.00		1.00		1.00		1.00	
Yes	1.38	0.40–4.73	2.01	0.85–4.77	1.51	0.40–5.75	1.08	0.45–2.55	1.14	0.42–3.12
Female										
Age (years)										
40–49	1.00		1.00		1.00		1.00		1.00	
50–59	0.96	0.40–2.28	0.64	0.31–1.33	0.60	0.28–1.28	1.49	0.68–3.25	2.09	0.90–4.84
60–69	0.83	0.31–2.24	0.59	0.26–1.35	0.61	0.26–1.44	1.85	0.79–4.33	3.17	1.27–7.91
≥ 70	0.45	0.11–1.76	0.53	0.19–1.49	0.49	0.16–1.53	1.21	0.43–3.42	2.77	0.94–8.22
Educational attainment										
High	1.00		1.00		1.00		1.00		1.00	
Middle	0.59	0.29–1.23	0.85	0.47–1.55	0.79	0.41–1.49	1.45	0.83–2.54	0.76	0.43–1.35
Elementary	1.01	0.49–2.08	1.12	0.60–2.10	0.94	0.48–1.83	1.68	0.94–3.01	0.87	0.48–1.56
Ergonomic risk										
< 250	1.00		1.00		1.00		1.00		1.00	
250–499	1.92	1.01–3.68	1.78	1.09–2.93	1.87	1.03–3.38	1.80	1.11–2.92	1.28	0.80–2.07
500–749	2.46	1.17–5.16	2.12	0.64–2.26	2.60	1.30–5.21	2.78	1.57–4.93	1.41	0.71–2.82
≥ 750	3.41	1.77–6.57	3.28	1.87–5.77	4.69	2.64–8.34	4.06	2.40–6.88	2.87	1.66–4.96
Cold exposure (days/1 year)										
No	1.00		1.00		1.00		1.00		1.00	
1–60	1.53	0.80–2.90	1.67	1.01–2.77	1.60	0.92–2.78	1.02	0.64–1.62	1.04	0.64–1.69
61–119	2.27	1.15–4.49	2.04	1.15–3.62	1.75	0.94–3.28	1.64	0.95–2.81	1.23	0.71–2.12
≥ 120	2.11	1.04–4.29	2.49	1.38–4.51	3.31	1.77–6.20	1.61	0.92–2.81	1.45	0.82–2.57
Mental disease ^a										
No	1.00		1.00		1.00		1.00		1.00	
Yes	3.56	2.07–6.12	3.18	1.96–5.15	3.05	1.83–5.10	2.58	1.58–4.23	2.10	1.31–3.39

OR: odds ratio; CI: confidence interval.

^aIf the patient has depression and/or anxiety disorder.

(95% CI: 1.31–3.39). The OR of cold exposure in female occurred with the neck 2.11 (95% CI: 1.04–4.29); shoulder 2.49 (95% CI: 1.38–4.51); and hands 3.31 (95% CI: 1.77–6.20). Knee MSD showed a significant association with educational attainment in male and age in female. MSDs showed no significant association with age, mental health, and cold exposure in male; and educational attainment in female.

Indicators for musculoskeletal diseases among fishers in Korea

Table 5. PAR and PAF of indicators for 5 body parts musculoskeletal diseases by sex

Variables	Neck		Shoulder		Hand		Back		Knee	
	% or %p	95% CI	% or %p	95% CI	% or %p	95% CI	% or %p	95% CI	% or %p	95% CI
Male										
All	7.8	5.6–10.8	17.8	14.4–21.8	7.8	5.6–10.9	27.2	23.2–31.6	16.2	12.9–20.1
PAR										
Age	-1.1	-9.0–6.8	0.1	-10.5–10.6	-4.2	-13.7–5.3	-4.6	-17.4–8.4	-10.5	-24.6–4.0
Educational attainment	1.9	-0.8–4.7	0.5	-4.6–5.5	0.9	-2.3–4.2	5.1	0.0–10.2	5.9	1.7–10.1
Ergonomic risk	4.7	2.2–7.1	7.1	3.9–10.4	3.8	1.4–6.2	13.7	9.3–18.0	3.7	0.2–7.1
Cold exposure	-0.4	-4.1–3.4	-3.0	-8.7–2.8	-0.3	-3.7–3.2	-3.8	-10.2–2.6	-3.3	-8.9–2.2
Mental disease	0.2	-0.8–1.3	1.0	-0.3–2.4	-0.2	-4.5–4.1	0.1	-1.2–1.5	0.2	-1.1–1.4
PAF										
Educational attainment							18.8	0.0–32.3	36.4	13.2–50.2
Ergonomic risk	59.6	39.5–65.6	40.2	27.2–47.5	48.6	26.0–56.5	50.3	40.1–57.1	22.8	1.9–35.4
Female										
All	16.4	13.7–19.5	28.1	24.7–31.7	23.0	19.9–26.5	38.7	35.0–42.5	30.0	26.5–33.8
PAR										
Age	-2.1	-12.9–8.7	-8.9	-22.3–4.9	-8.1	-21.4–5.5	8.8	-4.9–22.0	14.4	4.5–23.9
Educational attainment	-1.7	-7.5–4.2	0.1	-6.9–7.1	-1.5	-8.6–5.6	6.9	-0.7–14.4	-2.9	-10.9–5.2
Ergonomic risk	5.8	2.6–9.1	6.8	2.7–10.8	9.7	5.7–13.6	13.0	7.8–18.2	4.8	1.4–8.2
Cold exposure	4.4	0.9–7.9	7.2	2.8–11.6	6.1	2.2–10.0	3.4	-1.5–8.2	2.0	-2.7–6.8
Mental disease	3.5	1.8–5.2	4.2	2.4–6.0	3.4	1.7–5.2	3.5	1.7–5.4	2.8	0.9–4.6
PAF										
Age									48.0	17.0–70.7
Ergonomic risk	35.6	18.8–46.6	24.1	11.1–34.0	42.2	28.9–51.5	33.7	22.2–42.8	15.9	5.2–24.1
Cold exposure	26.8	6.6–40.5	25.6	13.3–36.6	26.1	11.1–37.7				
Mental disease	21.4	13.3–26.5	14.9	9.6–18.8	15.0	8.6–19.6	9.1	4.8–12.6	9.2	3.5–13.6

All: prevalence (%) of each musculoskeletal disease and 95% confidence interval. PAR: %p and 95% confidence interval. PAF: proportion (%) and 95% confidence interval.

PAF: population attributable fraction; PAR: population attributable risk.

Table 5 shows the PAR and PAF and their 95% CIs for MSDs in the relevant body parts, including the indicators according to sex, respectively. PAF of age in female was 48.0% (95% CI: 17.0–70.7) in the knees. PAF of educational attainment in male was 36.4% (95% CI: 13.2–50.2) in knee. PAF of ergonomic risk in male was 59.6% (95% CI: 2.2–7.1) in the neck, 40.2% (95% CI: 3.9–10.4) in shoulders, 48.6% (95% CI: 26.0–56.5) in hands, 50.3% (95% CI: 40.1–57.1) in back, and 22.8% (95% CI: 1.9–35.4) in knees. PAF of ergonomic risk in female was 35.6% (95% CI: 18.8–46.6) in neck, 24.1% (95% CI: 11.1–34.0) in shoulders, 42.2 (95% CI: 28.9–51.5) in hands, 33.7% (95% CI: 22.2–42.8) in back, and 15.9% (95% CI: 5.2–24.1) in knees. PAF of cold exposure in female was 26.8% (95% CI: 6.6–40.5) in neck, 25.6% (95% CI: 13.3–36.6) in shoulders, and 26.1 (95% CI: 11.1–37.7) in hand. PAF of mental diseases in female was 21.4% (95% CI: 13.3–26.5) in neck, 14.9% (95% CI: 9.6–18.8) in shoulders, 15.0 (95% CI: 8.6–19.6) in hands, 9.1% (95% CI: 4.8–12.6) in back, and 9.2% (95% CI: 3.5–13.6) in knees.

Fig. 1 shows the PAF of MSDs among fishers by sex. The colored bars refer to modifiable factors. In male with MSDs, a contribution of 40.2%–69.1% was shown according to body parts. In Female with MSDs, a contribution of 42.8%–83.8% was shown.

DISCUSSION

In sex and age-adjusted multiple logistic regression model, significant indicators associated with MSDs were educational attainment, job classification, ergonomic risk, cold exposure, heat exposure, and mental diseases (**Table 4**). In the final binary logistic regression and PAF estimation by sex, heat exposure had some significant results but was excluded because the




Fig. 1. Population attributable fraction of indicators for MSD among fishers by sex. Subject number of MSD by sex (male, female): Neck (29, 91) Shoulder (66, 156) Hands (29, 128) Back (101, 215) Knees (60, 167). Colored bar: modifiable indicators. Grey bar: non-modifiable indicators. MSD: musculoskeletal disease.

OR of MSDs increased in the non-exposed group. We included ergonomic risk instead of job classification because 2 variables overlapped as ergonomic hazards and even Haenyeo included only female.

The major contributing factor of MSD in fishers was ergonomic risk. In our study, PAF according to body parts ranged from 22.8%–59.6% and 22.8%–50.3% in male and female, respectively. These results are similar to those of previous studies. Punnett et al.⁹ reported the attributable proportions of low back pain in ergonomic stress between 21%–41% depending on the region, and were higher in the less developed regions. This regional differences is explained by the labor force participation rate and population distribution of occupations, especially the proportion of farmers.¹⁵ Nambiema et al.⁸ showed an incident upper extremity musculoskeletal disorders associated with high physical exertion (PAF 30%) and working with arms above the shoulder level (PAF 7%), in a prospective study using the French Cosali cohort.

Mental diseases showed a significant PAF for MSDs in the neck, shoulder, hand, back, and knee regions in female only (PAF 9.1%–21.4% by body parts). Chronic pain is an important factor in the development of depression, and the coexistence of chronic pain and depression is commonly observed.^{9,16,17} Recent studies have shown that chronic pain and depression share similar neurological changes. For example, a study reported that the damage to dopamine activity in the limbic areas reduces the utilization of monoamine neurotransmitters.¹⁸ Another study showed that depression in the early stage of pain increases the possibility of prolonged pain.¹⁹ we recognized the reciprocal relationship between MSDs and mental diseases. Interestingly, we found sex difference in this relationship. In a study of sex differences in upper extremity MSD, female responded more strongly to psychological factors.²⁰

Interestingly, we found gender differences in this relationship. A study by Treaster et al.²⁰ on gender differences in upper extremity MSD showed that women responded more strongly to psychological factors. Higher pain sensitivity in female has been a consistent result of many studies.²¹ Zhang et al.²² analyzed gender differences in pain sensitivity in a functional MRI

study of 450 healthy subjects. In this study, female had a lower pain threshold and a greater fear response to pain than male. The study presented sex differences in the volumes of amygdala subnuclei, which regulate responses to anxiety and fear.²²

In this study, cold exposure showed a significant PAF for upper extremity MSD including the neck, shoulder, and hand only among female (25.6%–26.8%). Fishing has been reported as the most hazardous occupation, due to exposure to cold stress.²³ Pienimäki²⁴ explains that the occurrence of musculoskeletal symptoms increases as the duration of cold exposure increases. Working in a cold environment resulted in an elevated risk of OR: 2.2 (95% CI: 1.30–3.72) among male and OR: 1.6 (95% CI: 0.78–3.31) among female. Exposure to cold environment combined with repetitive work may have negative effect on muscle function, and fatigue may lead to strain injuries.^{25,26} The cause of the sex difference in the impact of cold exposure is unclear, and related studies are insufficient. However, Kaikaew et al.²⁷ showed that female perceive the experience of cold and onset of shivering faster than male in an experimental study including voluntary subjects.

Interestingly, Age was not a significant variable of MSDs in fishers except with the knee MSD among female. In this study, MSD was defined based on health care visits and subjective symptoms, not based on objective findings such as radiological degenerative signs or disability. Fishers with severe degenerative MSD could adapt by avoiding physical work that induces pain. In heavy physical occupations such as fishing, having MSD with disability are the main reasons for early retirement. Paradoxically, health care use may be higher among younger fishers who are actively working. The exception to the knee MSD among female is assumed to be due to the high prevalence of knee osteoarthritis in female Korean farmers and fishers,⁴ and they can perform sedentary tasks despite the knee osteoarthritis.

By calculating PAR for multifactorial diseases, it is possible to know which factor contribute more to the prevalence of the disease. The proportion of harmful factors contributing to a disease is the PAF, which means that MSD can be prevented that the extent of that percentage, by improving the occurrence of harmful factors. In this study, significant and modifiable factors for MSDs included ergonomic risk, cold exposure, and mental diseases. Although the contribution of ergonomic risk and mental diseases to MSDs differed according to sex, the contribution of ergonomic risk to MSDs was greater in both sexes. This means that MSD can be prevented by improving ergonomic harmful factors. In particular, intervention to prevent such factors among female fishers are needed to prevent the development of mental diseases and cold exposure.

This study has several limitations. First, we investigated the association between a one-year prevalence and ergonomic risk using a cross-sectional study design. Therefore, unclear temporality and survival bias can act as systemic error. Second, the reliability and validity of the questionnaire used to obtain information for calculating ergonomic risk has not been previously verified. It may be insufficient for reflecting the atypical ergonomic factors among fishers and affected by recall bias. However, since it was impossible to evaluate the atypical and intermittent tasks of fishers in various regions, we accepted the subjective judgment by fishers as the best possible. Third, there is the possibility that fishers who were more interested in their health status participated in this study. Therefore, the prevalence of MSDs may have been overestimated. Fifth, as potential factors, history of injury, whole-body vibration, and hand-arm vibration may be considered, but they were not included in the analysis. Nevertheless, there are strengths to this study. First, this was a relatively large-scale

study targeting fishers among whom research on occupational diseases were lacking, despite their high-risk occupations. Second, we adjusted for almost all potential variables to evaluate the associations of ergonomic risk, mental diseases, and cold exposure with MSDs. Third, we calculated the integrated ergonomic risk reflecting the elements of ergonomic hazard: strength, time, for each of the 5 body parts.

CONCLUSIONS

In this study, significant modifiable indicators of MSDs among fishers were ergonomic risk, mental diseases, and exposure to cold. Ergonomic risk contributed the most to the prevalence of MSDs. These results suggest that MSD can be reduced through improvement of ergonomic risk. Since mental diseases and cold exposure contributed more to MSDs among female than in male, sex differences should be considered in preventive interventions. We believe that the results of this study may provide the basis for the prevention of occupational diseases among fishers.

ACKNOWLEDGEMENTS

The authors thank the Ministry of Oceans and Fisheries Affairs of the Republic of Korea.

SUPPLEMENTARY MATERIALS

Supplementary Data 1

Questionnaire to assess cold, heat, and noise exposure

[Click here to view](#)

Supplementary Table 1

Face-to-face questionnaire for the assessment of ergonomic risk

[Click here to view](#)

Supplementary Table 2

The association between variables and musculoskeletal disease of 5 body sites among male

[Click here to view](#)

Supplementary Table 3

The association between variables and musculoskeletal disease of 5 body sites among female

[Click here to view](#)

Supplementary Table 4

Adjusted OR of variable by multiple logistic regression among male

[Click here to view](#)

Supplementary Table 5

Adjusted OR of variable by multiple logistic regression among female

[Click here to view](#)**REFERENCES**

1. Kim YE, Park H, Jo MW, Oh IH, Go DS, Jung J, et al. Trends and patterns of burden of disease and injuries in Korea using disability-adjusted life years. *J Korean Med Sci* 2019;34(Suppl 1):e75.
[PUBMED](#) | [CROSSREF](#)
2. Oh IH, Yoon SJ, Seo HY, Kim EJ, Kim YA. The economic burden of musculoskeletal disease in Korea: a cross sectional study. *BMC Musculoskelet Disord* 2011;12(1):157.
[PUBMED](#) | [CROSSREF](#)
3. Park J, Kim Y, Han B. Work sectors with high risk for work-related musculoskeletal disorders in Korean men and women. *Saf Health Work* 2018;9(1):75-8.
[PUBMED](#) | [CROSSREF](#)
4. Seok H, Choi SJ, Yoon JH, Song GG, Won JU, Kim JH, et al. the association between osteoarthritis and occupational clusters in the Korean population: a nationwide study. *PLoS One* 2017;12(1):e0170229.
[PUBMED](#) | [CROSSREF](#)
5. Berg-Beckhoff G, Østergaard H, Jepsen JR. Prevalence and predictors of musculoskeletal pain among Danish fishermen - results from a cross-sectional survey. *J Occup Med Toxicol* 2016;11(1):51.
[PUBMED](#) | [CROSSREF](#)
6. Kaerlev L, Jensen A, Hannerz H. Surveillance of hospital contacts among Danish seafarers and fishermen with focus on skin and infectious diseases-a population-based cohort study. *Int J Environ Res Public Health* 2014;11(11):11931-49.
[PUBMED](#) | [CROSSREF](#)
7. Hong C, Lee CG, Song H. Characteristics of lumbar disc degeneration and risk factors for collapsed lumbar disc in Korean farmers and fishers. *Ann Occup Environ Med* 2021;33:e16.
[PUBMED](#) | [CROSSREF](#)
8. Nambiema A, Bertrais S, Bodin J, Fouquet N, Aublet-Cuvelier A, Evanoff B, et al. Proportion of upper extremity musculoskeletal disorders attributable to personal and occupational factors: results from the French Pays de la Loire study. *BMC Public Health* 2020;20(1):456.
[PUBMED](#) | [CROSSREF](#)
9. Punnett L, Prüss-Utün A, Nelson DI, Fingerhut MA, Leigh J, Tak S, et al. Estimating the global burden of low back pain attributable to combined occupational exposures. *Am J Ind Med* 2005;48(6):459-69.
[PUBMED](#) | [CROSSREF](#)
10. Korea Occupational Safety and Health Agency. Guidelines for risk factor survey on tasks with musculoskeletal burden of the KOSHA (H-9-2018). https://kosha.or.kr/kosha/data/musculoskeletalPreventionData_G.do?mode=download&articleNo=296739&attachNo=230707. Updated 2018. Accessed April 4, 2022.
11. European Working Conditions Surveys. EWCTS 2021 – Questionnaire. <https://www.eurofound.europa.eu/surveys/european-working-conditions-surveys/european-working-conditions-survey-2021/ewcs-2021-questionnaire>. Updated 2021. Accessed September 6, 2022.
12. Moriarty AS, Gilbody S, McMillan D, Manea L. Screening and case finding for major depressive disorder using the Patient Health Questionnaire (PHQ-9): a meta-analysis. *Gen Hosp Psychiatry* 2015;37(6):567-76.
[PUBMED](#) | [CROSSREF](#)
13. Spitzer RL, Kroenke K, Williams JB, Löwe B. A brief measure for assessing generalized anxiety disorder: the GAD-7. *Arch Intern Med* 2006;166(10):1092-7.
[PUBMED](#) | [CROSSREF](#)
14. Newson R. REGPAR: Stata module to compute population attributable risks from binary regression models. <https://econpapers.repec.org/scripts/search.pf?ft=REGPAR>. Updated 2015. Accessed September 6, 2022.
15. Nelson DI, Concha-Barrientos M, Driscoll T, Steenland K, Fingerhut M, Punnett L, et al. The global burden of selected occupational diseases and injury risks: methodology and summary. *Am J Ind Med* 2005;48(6):400-18.
[PUBMED](#) | [CROSSREF](#)

16. Haukka E, Leino-Arjas P, Ojajärvi A, Takala EP, Viikari-Juntura E, Riihimäki H. Mental stress and psychosocial factors at work in relation to multiple-site musculoskeletal pain: a longitudinal study of kitchen workers. *Eur J Pain* 2011;15(4):432-8.
[PUBMED](#) | [CROSSREF](#)
17. Kroenke K, Wu J, Bair MJ, Krebs EE, Damush TM, Tu W. Reciprocal relationship between pain and depression: a 12-month longitudinal analysis in primary care. *J Pain* 2011;12(9):964-73.
[PUBMED](#) | [CROSSREF](#)
18. Sheng J, Liu S, Wang Y, Cui R, Zhang X. The link between depression and chronic pain: neural mechanisms in the brain. *Neural Plast* 2017;2017:9724371.
[PUBMED](#) | [CROSSREF](#)
19. Linton SJ, Bergbom S. Understanding the link between depression and pain. *Scand J Pain* 2011;2(2):47-54.
[PUBMED](#) | [CROSSREF](#)
20. Treaster DE, Burr D. Gender differences in prevalence of upper extremity musculoskeletal disorders. *Ergonomics* 2004;47(5):495-526.
[PUBMED](#) | [CROSSREF](#)
21. Mogil JS. Sources of individual differences in pain. *Annu Rev Neurosci* 2021;44(1):1-25.
[PUBMED](#) | [CROSSREF](#)
22. Zhang H, Bi Y, Hou X, Lu X, Tu Y, Hu L. The role of negative emotions in sex differences in pain sensitivity. *Neuroimage* 2021;245:118685.
[PUBMED](#) | [CROSSREF](#)
23. Sandsund M, Øren A, Thorvaldsen T, Holmen I, Sønvisen S, Heidelberg CT, et al. Musculoskeletal symptoms among workers in the commercial fishing fleet of Norway. *Int Marit Health* 2019;70(2):100-6.
[PUBMED](#) | [CROSSREF](#)
24. Pienimäki T. Cold exposure and musculoskeletal disorders and diseases. A review. *Int J Circumpolar Health* 2002;61(2):173-82.
[PUBMED](#) | [CROSSREF](#)
25. Oksa J, Ducharme MB, Rintamäki H. Combined effect of repetitive work and cold on muscle function and fatigue. *J Appl Physiol (1985)* 2002;92(1):354-61.
[PUBMED](#) | [CROSSREF](#)
26. Chiang HC, Chen SS, Yu HS, Ko YC. The occurrence of carpal tunnel syndrome in frozen food factory employees. *Gaoxiong Yi Xue Ke Xue Za Zhi* 1990;6(2):73-80.
[PUBMED](#)
27. Kaikaew K, van den Beukel JC, Neggers SJ, Themmen AP, Visser JA, Grefhorst A. Sex difference in cold perception and shivering onset upon gradual cold exposure. *J Therm Biol* 2018;77:137-44.
[PUBMED](#) | [CROSSREF](#)