

Review Article

The Effect of Periodontitis on Expression of Interleukin-21: A Systematic Review

Archana Mootha, Sankari Malaiappan, N. D. Jayakumar, Sheeja S. Varghese, and Julie Toby Thomas

Department of Periodontics, Saveetha Dental College and Hospitals, No. 162, Poonamallee High Road, Velappanchavadi, Chennai 600077, India

Correspondence should be addressed to Archana Mootha; archanamootha@gmail.com

Received 21 September 2015; Revised 24 December 2015; Accepted 11 January 2016

Academic Editor: Alexander J. Steven

Copyright © 2016 Archana Mootha et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Purpose. Inflammation and tissue breakdown are led by an array of inflammatory destructive mediators associated with initiation and progression of inflammatory diseases like periodontitis. Current evidence shows that these inflammatory mediators have a definitive role in the pathogenesis of various systemic diseases with an inflammatory component. Interleukin-21 (IL-21) has been associated with systemic diseases like rheumatoid arthritis and Crohn's disease that follow a chronic inflammatory cascade. Similarly recent studies have associated Interleukin-21 levels with periodontitis. This systematic review was aimed to assess the levels of IL-21 in subjects with periodontitis. *Methods.* A complete literature search was done in PubMed, Medline, Science Direct, and Cochrane databases and Google Scholar based on the inclusion/exclusion criteria. Six relevant articles were procured. Full text was read individually by two reviewers and data extraction was done based on STROBE statement. *Results.* After data extraction five observational and one interventional study were obtained. All the studies showed an increased expression of IL-21 in periodontitis and the interventional study showed reduction in IL-21 levels after nonsurgical periodontal therapy (NSP). *Conclusion.* Interleukin-21 levels are higher in periodontitis than controls. With this limited evidence further longitudinal studies are required to consider this as a definitive inflammatory marker.

1. Introduction

Periodontitis is a chronic inflammatory condition which was initiated by gram-negative organisms present in the tooth supporting structures [1]. Disease progression occurs as a result of host-immune response to bacteria, leading to destruction of connective tissue and alveolar bone [2]. The pathogens present in the subgingival flora produce various endotoxins which are a prerequisite for periodontal disease [3]. These endotoxins in turn activate a host-immune response to the bacterial challenge by stimulating immune cells like polymorphonuclear neutrophils (PMNs), monocytes, B cells, T cells, and fibroblasts [2] to produce various inflammatory mediators such as cytokines [4, 5], acute phase proteins [6], and proteolytic enzymes [7, 8] that mediate tissue destruction. Progression of the periodontal disease can further trigger the adaptive immune mechanism for release

of inflammatory mediators resulting in further periodontal breakdown.

Cytokines are important in expression of the characteristics of the immune response to bacterial endotoxins [9]. Interleukins comprise a large group of cytokines that are naturally occurring glycoproteins produced by the body [10]. They help in recruitment of neutrophils and macrophages to participate and amplify the inflammatory immune reaction [11].

The role of IL-1, IL-6, IL-8, and IL-12 has previously been established in periodontitis [12–15] proving their definitive role in periodontal destruction. Interleukin-21 (IL-21) is an inflammatory cytokine [16] mainly expressed by activated Th1 and Th17 cells which are distinct proinflammatory lineages, and not by Th2 cells in humans pointing out it to be a proinflammatory cytokine [17]. It functions via the receptor IL-21R which is a type I cytokine receptor. Recently

TABLE 3: Data summation table.
(a)

Sl. number	Author and journal	Study design	Groups	Parameters	Sample and Methodology	Statistical Analysis	Results
1	Dutzan et al. [29] Journal of Periodontology, 2011 "Levels of Interleukin-21 in Patients with Untreated Chronic Periodontitis"	Case control study	Group 1: controls, n = 19 Group 2: CP, n = 15	PI, PD, CAL, PI, BOP, and presence of IL-21	Gingival and GCF IL-21 (healthy-extraction site, CP- periodontal lesions) were analysed by IHC and Western blot (for presence of IL-21) and quantification was done by ELISA	Unpaired <i>t</i> -test and Mann-Whitney <i>U</i> test Chi square test	Gingival IL-21 levels were significantly higher in CP group (3.35 pg/mg) than in controls (0.98 pg/mg) (<i>p</i> < 0.05) Frequency of GCF IL-21 was greater in CP patients (8/10) than in controls (2/10) (<i>p</i> = 0.007) The Western blot and IHC staining confirmed the presence of IL-21 in periodontal tissues and GCF

(b)

Sl. number	Author and journal	Study design	Groups	Parameters	Methodology	Statistical analysis	Results
2	Zhao et al. [30] Journal of Clinical Periodontology, 2011 "Effect of Non-Surgical Periodontal Therapy on the Levels of Th17/Th1/Th2 Cytokines and Their Transcription Factors in Chinese Chronic Periodontitis Patients"	Interventi-onal study	30 participants with CP, n = 30	IL-21	GCF (before and after treatment) IL-21 was analysed by ELISA	Linear correlation Paired <i>t</i> -test	Downregulation of IL-21 (80.3360 ± 39.4188 versus 31.7741 ± 13.2558 pg/mL) (<i>p</i> < 0.05) The quantity of Th17 cells in peripheral blood was decreased especially in IL-17 IFN-γ1 subgroup (<i>p</i> < 0.05)

(c)

Sl. number	Author and journal	Study design	Groups	Parameters	Methodology	Statistical analysis	Results
3	Napimoga et al. [32] Scandinavian Journal of Immunology 2011 "Possible Involvement of IL-21 and IL-10 on Salivary IgA Levels in Chronic Periodontitis Subjects"	Case control study	Group 1: healthy controls, n = 15 Group2: CP, n = 15	IL-21	Gingival IL-21 (healthy-gingivoplasty sites, CP-extraction sites due to advanced periodontitis) using RT-PCR	Student's <i>t</i> -test	The mRNA levels for IL-21 was higher in the CP (1.5-fold) when compared to the healthy group (0.4-fold) (<i>p</i> < 0.05)

(d)

Sl. number	Author and journal	Study design	Groups	Parameters	Methodology	Statistical analysis	Results
4	Dutzan et al. [31] Journal of Periodontology, 2012 "Interleukin-21 Expression and Its Association with Proinflammatory Cytokines in Untreated Chronic Periodontitis subjects"	Case control study	Group 1: controls, $n = 8$ Group 2: CP, $n = 10$	IL-21 mRNA level	Gingival IL-21 (healthy-extraction site, CP- periodontal lesions) was analysed using RT-PCR	The RQ expressed as fold change for each studied cytokine Spearman rank correlation test	A significant overexpression of IL-21 in CP affected tissues compared to healthy gingival tissues (120-fold) Significant positive correlations of IL-21 with PD (0.71) ($p = 0.002$) and CAL (0.60) ($p = 0.01$)

(e)

Sl. number	Author and journal	Study design	Groups	Parameters	Methodology	Statistical analysis	Results
5	Isaza-Guzmán et al., 2015 [33] Journal of Archives of Oral Biology "Association Study between Salivary Levels of Interferon (IFN)-Gamma, Interleukin (IL)-17, IL-21, and IL-22 with Chronic Periodontitis"	Cross-sectional study	Group 1: CP, $n = 105$ Group 2: healthy controls, $n = 44$	IL-21 and IL-21R	Salivary IL-21 and levels were analysed using RT-PCR	Pearson's chi square test	In healthy controls IL-21 (0.00–4.54) than CP (0.00 pg/mL) ($p = 0.470$) Low detection frequency of Th17 cytokines in saliva No significant domination in periodontitis

(f)

Sl. number	Author and journal	Study design	Groups	Parameters	Sample and methodology	Statistical analysis	Results
6	Nizam et al. [28] Journal of Periodontology, 2014 "Salivary Cytokines and the Association between Obstructive Sleep Apnea Syndrome and Periodontal Disease"	Case-control study	Group 1: controls (non-OSAS), $n = 13$ Group 2: mild/moderate OSAS, $n = 17$ Group 3: severe OSAS, $n = 22$	PI, PD, CAL, BOP, and salivary IL-21	Salivary IL-21 was assessed by ELISA	Spearman r rank correlation analysis	Significant correlation was between CAL and IL-21 ($r = -0.347$) ($p = 0.02$) Borderline significance between PI and IL-21 ($r = -0.287$) ($p = 0.05$) IL-21 levels in healthy controls, moderate OSAS, and severe OSAS (90 pg/mL, 85 pg/mL, and 75 pg/mL)

IL: interleukin, OSAS: obstructive sleep apnea syndrome, GCF: gingival crevicular fluid, GI: Gingival Index, PPD: probing pocket depth, PD: probing depth, CAL: clinical attachment level, BOP: bleeding on probing, PCR: polymerase chain reaction, RT-PCR: real time polymerase chain reaction, IHC: immunohistochemistry, and ELISA: enzyme linked immunosorbant assay.

Next study [29] performed qualitative analysis of gingival tissue samples IL-21 using immunohistochemistry and quantitative analysis of salivary and gingival tissue IL-21 using western blot. Gingival tissue IL-21 (3.35 pg/mL) and GCF IL-21 (8/10) were higher in CP (chronic periodontitis) compared to controls.

Another study showed increased expression of IL-21 in gingival tissue samples using PCR, which were 1.5 times more in CP than controls (0.3) [32]. Another study [31] compared IL-21 levels in gingival tissue between CP and healthy controls using RT-PCR. CP group showed 120-fold increase of IL-21 level and showed a direct positive correlation with probing depth ($p = 0.02$) and CAL ($p = 0.02$).

One study showed reduced expression of salivary IL-21 in periodontitis using ELISA [33]. Salivary estimation of IL-21 using ELISA in CP (smokers were included) showed no significant domination of this cytokine (0.00–4.5 pg/mL) and it was concluded that the detection of IL-21 has no predictive value in health/disease.

In the interventional study, the GCF IL-21 levels were increased in CP group and there was a reduction in IL-21 levels after nonsurgical periodontal therapy when estimated using ELISA [30].

Due to variations in the study population, sample source and the methodology of estimation of IL-21 the data obtained were heterogeneous. One study was conducted in Brazilian [32], one study in Turkish [28], and one study in Chinese population [30], whereas three studies were conducted in the American [29, 31, 33] population. Varied sources were used for analysis of IL-21. Two studies used GCF [29, 30], two studies used gingival tissue [29, 31, 32], and two studies used saliva [28, 33]. All the samples showed increased IL-21 levels in periodontitis, among which GCF [30] showed maximum detection of IL-21 (80.336 pg/mL). Several methodologies were used for qualitative and quantitative analysis of IL-21. Three studies detected IL-21 using RT-PCR [31, 32], one study used western blot and immunohistochemistry [29], and three studies used ELISA [28–30, 33] for analysis of IL-21. All methodologies showed increased IL-21 levels in periodontitis, among which ELISA showed maximum detection of IL-21 [30]. Due to the heterogeneity of the data it was not feasible to pool the data to perform a meta-analysis. Only two studies showed significant correlation of IL-21 level with periodontal disease severity [31, 32]. They found a significant correlation of IL-21 with clinical attachment loss (CAL) and borderline significance between plaque index and IL-21. As CAL is the most reliable indicator of periodontal tissue destruction, this correlation indicates better that IL-21 levels increase with the severity of periodontitis.

4. Discussion

Periodontitis is a chronic inflammatory disease that affects the supporting structures of the teeth and is considered as one of the most common reasons for tooth loss [34]. It is one of the most common pathologies of bone and an important modifying factor of several chronic inflammatory systemic diseases like cardiovascular disease and obesity [35]. Evidence shows association of IL-21 with various systemic

conditions such as coronary artery disease [36], rheumatoid arthritis [18], and inflammatory bowel disease [19]. Much about IL-21 has been studied in rheumatoid arthritis (RA), and researchers have reported elevated serum and synovial tissues levels of this cytokine in RA as the disease severity increased [37].

Periodontitis is a chronic inflammatory condition which shows increased local and systemic levels of inflammatory mediators and markers of inflammation like IL-21, and a direct link has been established between periodontitis and rheumatoid arthritis. In this regard, this systematic review aimed to evaluate the expression of IL-21 in periodontitis.

Clinically, periodontal disease is characterized by increase in probing depth and a decrease in clinical attachment level. The ultimate determinant of disease progression and clinical outcome is the immune response of host, which involves the generation of cytokines, activation of osteoclasts, and the recruitment of inflammatory cells like lymphocytes, PMNs, and antigen presenting cells to the site [2]. The recruitment and activation of these host cells lead to overproduction of proinflammatory cytokines like IL-1, IL-6, IL-12, and IL-21 along with other host-destructive mediators that enhance host destruction and activate resident cells to produce proteases which further amplify the host mediated periodontal destruction.

A limited number of eligible studies were identified in this systematic review including five observational [23, 24, 28, 29, 31–33] and one interventional studies [30]. The results of this systematic review show that five [23, 24, 28, 29, 31–33] out of six studies show an increase in IL-21 levels in chronic periodontitis subjects and one study [33] showed no increase in periodontitis. The interventional study [30] showed a reduction in IL-21 levels following nonsurgical therapy.

Nizam et al. [28] hypothesized that OSAS may predispose patients to periodontal disease and associated the salivary cytokines with OSAS severity. He found that salivary IL-6 and IL-33 were similar in OSAS groups but significantly higher than control groups, whereas IL-1 β , IL-21, and PTX were similar in all groups with strong and weak positive correlation of IL-21 with CAL and PI, respectively. The authors concluded, it is likely that elevated IL-6 reflects the degree of subclinical inflammation in periodontal tissues, which can be a link between periodontal disease and OSAS, and the presence or severity of OSAS does not affect the level of IL-1 β and IL-21 in either plasma or saliva.

Dutzan et al. [31] showed an increase in IL-21 levels in chronic periodontitis, a positive correlation of IL-21 with PD, CAL, IL-1 β , IL-6, and Th17 cytokines like IL-17 and IL-23, and a negative correlation with anti-inflammatory cytokines like IL-10 and TGF- β 1 [31].

IL-21 influences the actions of various immune cells like Natural Killer (NK) cells, PMNs, and macrophages. NK cells are a part of the innate immune response and serve as a first line of defence against invading pathogens. The development and activation of NK cells are intrinsically dependent on the activity of class I cytokine receptors, and cells treated with IL-21 displayed enhanced effector functions with greater cytolytic activity. IL-21 increased PMN migration to the local

inflammatory site and caused greater PMN mediated tissue injury. IL-21 also complemented the phagocytic ability of macrophages. In addition to its action in the innate immune response, it also parades the action of the adaptive immune response. IL-21 drives terminal differentiation of B cells into plasma cells, favours the effector functions of T cell, and accelerates memory cell formation for rapid action on encounter of a pathogen for the second time. Furthermore, IL-6 and IL-21 can stimulate Th17 cells itself to produce increased IL-21. Altogether, IL-21 exaggerates the host-immune response and intensifies the local inflammatory actions, and these data support an important role of IL-21 in the pathogenesis of periodontal disease.

Salivary levels of IgA were increased in periodontitis subjects along with gingival levels of IL-21. As it was previously discussed, IL-21 not only controls Th17 activity, but also controls B cell proliferation and maturation into plasma cells; the elevated IgA levels in periodontitis may be explained, as IgA is predominantly produced by B cells and both IgA and IL-21 are increased in periodontitis compared to healthy controls. IgA is the most predominant Ig secreted by oral mucosal sites and is considered as the most important protein against microbial defence. This indicated the role of IL-21 in Ig-isotype switching in chronic periodontitis and its influence on the adaptive immune response and the immunomodulation of oral mucosa under the challenge of periodontal pathogens [32]. In contrast to these findings, Isaza et al. detected low frequency of Th17 cytokines in saliva samples and concluded that its determination is useless for the detection of disease presence and/or its severity.

After nonsurgical periodontal therapy, Zhao et al. found that Th17 cells might participate in the development of periodontitis by upregulating the expression of cytokines IL-17 and IL-21; meanwhile, Th1 cells increased the expression of IFN- γ , but Th2 cells inhibited the expression of IL-4. He concluded that nonsurgical treatment might decrease the expression of Th17-related cytokines, IL-17 and IL-21, and Th1-related cytokine IFN- γ , while increasing the expression of Th2 related cytokine IL-4, thus contributing to the relief of periodontal inflammation [30].

While all the cross-sectional studies showed elevated IL-21 levels in periodontitis, one recent study [33] showed contradictory results by reporting no significant domination of Th17 cytokines in chronic periodontitis. An interesting finding is that there is no positive correlation of smokers with IL-21 levels. No scientific rationale explaining the results is mentioned in the article.

In addition, studies conducted in nonperiodontal oral lesions and animal studies also showed an increase in IL-21 levels in disease state compared to controls. On estimation of IL-21 levels in periapical granulomas, it was demonstrated that IL-21 along with IL-17A, TNF- α (tumour necrosis factor- α), and IFN- γ were higher in active periapical granulomas, while IL-4, IL-9, IL-10, and IL-22 were higher in inactive periapical granulomas. In the secondary lymphoid organs CD4+ T cell subset found in the B cell follicles was described as a major contributor to B cell-mediated antibody responses and an important source of IL-21. Since B cells are predominant in chronic lesions like periapical granulomas, it would be

possible to speculate that IL-21 contributes to a Tfh-B cell response axis in periapical areas, similar to that of tertiary lymphoid tissues associated with chronic infection sites. In view of B cells being potential sources of RANKL, Tfh-B cell axis can directly drive the activity of the lesion via RANKL production [34].

Animal study conducted in rhesus monkeys by Eber-sole et al. [4] reported that disease initiation/progression was characterized by overexpression of Th17/Treg cytokine genes (IL-1 β , IL-6, TGF β , and IL-21) and downregulation of Th1/Th2 cytokine genes (IL-18 and IL-25). They found that several Th17/Treg cytokine genes positively correlated with tissue destruction genes (TDGs), whereas most Th1/Th2 genes exhibited a negative correlation. Overall their observations were consistent with a cytokine profile driving a Th17 type of response early in the initiation process of periodontitis, followed by a persistence with disease progression.

Isaza-Guzmán et al. [33] showed that rats fed with *Actinobacillus actinomycetemcomitans* containing mash were compared to control; both B and T cells were increased and activated with enhanced IgG switched Ig in 2 weeks. Bone resorption was evident at 12 weeks with early increase in the levels of IL-21 along with IL-2, macrophage-inhibiting factor, IL-19, TNF- α , and CD40 ligand, CD70, while IL-16, TNF- α , lymphotoxin- β , IFN- α 1, and IL-1RN were increased from T cells indicating that adaptive immunity appears crucial for bone resorption.

The results of this review conclude that (1) IL-21 was elevated in subjects with chronic periodontitis compared to controls, (2) their levels were correlated with clinical severity of periodontitis (only in 2 studies, and values were given only in one study), and (3) nonsurgical periodontal therapy downregulated IL-21 levels. There are no studies based on comparison of IL-21 levels among types of periodontitis. The present review considers only published data for the interpretation of results and the raw data was not procured from the authors of unpublished studies.

This systematic review included cross-sectional studies (level III) [38] and thus indicates a low level of evidence to prove the definitive role of IL-21 in periodontitis. Future longitudinal studies are required to prove the definitive role of IL-21 in periodontitis and compare IL-21 levels among types of periodontitis. Further research in the field of Interleukin-21 could throw light on a better understanding of its role in the pathogenesis of tissue destruction in periodontitis.

Conflict of Interests

The authors declare that there is no conflict of interests regarding the publication of this paper.

References

- [1] H. Loe, "The role of bacteria in periodontal diseases," *Bulletin of the World Health Organization*, vol. 59, no. 6, pp. 821–825, 1981.
- [2] G. J. Seymour, "Importance of the host response in the periodontium," *Journal of Clinical Periodontology*, vol. 18, no. 6, pp. 421–426, 1991.

- [3] L. Kesic, J. Milasin, M. Igetic, and R. Obradovic, "Microbial etiology of periodontal disease," *Facta Universitatis Series: Medicine and Biology*, vol. 15, no. 1, pp. 1–6, 2008.
- [4] J. L. Ebersole, S. Kirakodu, M. J. Novak et al., "Cytokine gene expression profiles during initiation, progression and resolution of periodontitis," *Journal of Clinical Periodontology*, vol. 41, no. 9, pp. 853–861, 2014.
- [5] D. T. Graves, "The potential role of chemokines and inflammatory cytokines in periodontal disease progression," *Clinical Infectious Diseases*, vol. 28, no. 3, pp. 482–490, 1999.
- [6] R. C. Page, "The role of inflammatory mediators in the pathogenesis of periodontal disease," *Journal of Periodontal Research*, vol. 26, no. 3, pp. 230–242, 1991.
- [7] F. Trindade, F. G. Oppenheim, E. J. Helmerhorst, F. Amado, P. S. Gomes, and R. Vitorino, "Uncovering the molecular networks in periodontitis," *Proteomics—Clinical Applications*, vol. 8, no. 9–10, pp. 748–761, 2014.
- [8] H. M. Fullmer, "Collagenase and periodontal diseases: a review," *Journal of Dental Research*, vol. 50, no. 2, pp. 288–291, 1971.
- [9] S. Stenger and M. Rölinghoff, "Role of cytokines in the innate immune response to intracellular pathogens," *Annals of the Rheumatic Diseases*, vol. 60, no. 3, pp. iii43–iii46, 2001.
- [10] R. E. Elmslie, S. W. Dow, and G. K. Ogilvie, "Interleukins: biological properties and therapeutic potential," *Journal of Veterinary Internal Medicine*, vol. 5, no. 5, pp. 283–293, 1991.
- [11] M. Akdis, S. Burgler, R. Cramer et al., "Interleukins, from 1 to 37, and interferon- γ : receptors, functions, and roles in diseases," *The Journal of Allergy and Clinical Immunology*, vol. 127, no. 3, pp. 701.e70–721.e70, 2011.
- [12] S. P. Engebretson, J. T. Grbic, R. Singer, and I. B. Lamster, "GCF IL-1 β profiles in periodontal disease," *Journal of Clinical Periodontology*, vol. 29, no. 1, pp. 48–53, 2002.
- [13] C. R. Irwin and T. T. Myrillas, "The role of IL-6 in the pathogenesis of periodontal disease," *Oral Diseases*, vol. 4, no. 1, pp. 43–47, 1998.
- [14] A. Harada, N. Sekido, T. Akahoshi, T. Wada, N. Mukaida, and K. Matsushima, "Essential involvement of interleukin-8 [IL-8] in acute inflammation," *Journal of Leukocyte Biology*, vol. 56, pp. 559–564, 1994.
- [15] I.-S. Tsai, C.-C. Tsai, Y.-P. Ho, K.-Y. Ho, Y.-M. Wu, and C.-C. Hung, "Interleukin-12 and interleukin-16 in periodontal disease," *Cytokine*, vol. 31, no. 1, pp. 34–40, 2005.
- [16] C. de Rham, S. Ferrari-Lacraz, S. Jendly, G. Schneiter, J.-M. Dayer, and J. Villard, "The proinflammatory cytokines IL-2, IL-15 and IL-21 modulate the repertoire of mature human natural killer cell receptors," *Arthritis Research and Therapy*, vol. 9, no. 6, article R125, 15 pages, 2007.
- [17] M. Sarra, F. Pallone, and G. Monteleone, "Interleukin-21 in chronic inflammatory diseases," *BioFactors*, vol. 39, no. 4, pp. 368–373, 2013.
- [18] T. K. Rasmussen, T. Andersen, M. Hvid et al., "Increased interleukin 21 (IL-21) and IL-23 are associated with increased disease activity and with radiographic status in patients with early rheumatoid arthritis," *The Journal of Rheumatology*, vol. 37, no. 10, pp. 2014–2020, 2010.
- [19] K. Gerlach, C. Daniel, H. A. Lehr et al., "Transcription factor NFATc2 controls the emergence of colon cancer associated with IL-6-dependent colitis," *Cancer Research*, vol. 72, no. 17, pp. 4340–4350, 2012.
- [20] C. Abraham and J. Cho, "Interleukin-23/Th17 pathways and inflammatory bowel disease," *Inflammatory Bowel Diseases*, vol. 15, no. 7, pp. 1090–1100, 2009.
- [21] R. Nurieva, X. O. Yang, G. Martinez et al., "Essential autocrine regulation by IL-21 in the generation of inflammatory T cells," *Nature*, vol. 448, no. 7152, pp. 480–483, 2007.
- [22] T. Onoda, M. Rahman, H. Nara et al., "Human CD4⁺ central and effector memory T cells produce IL-21: effect on cytokine-driven proliferation of CD4⁺ T cell subsets," *International Immunology*, vol. 19, no. 10, pp. 1191–1199, 2007.
- [23] F.-L. Yuan, W. Hu, W.-G. Lu et al., "Targeting interleukin-21 in rheumatoid arthritis," *Molecular Biology Reports*, vol. 38, no. 3, pp. 1717–1721, 2011.
- [24] F. B. Mercado, R. I. Marshall, A. C. Klestov, and P. M. Bartold, "Relationship between rheumatoid arthritis and periodontitis," *Journal of Periodontology*, vol. 15, pp. 1090–1100, 2001.
- [25] A. C. Araujo-Pires, C. F. Francisoni, C. C. Bigueti et al., "Simultaneous analysis of T helper subsets (Th1, Th2, Th9, Th17, Th22, Tfh, Tr1 and Tregs) markers expression in periapical lesions reveals multiple cytokine clusters accountable for lesions activity and inactivity status," *Journal of Applied Oral Science*, vol. 22, no. 4, pp. 336–346, 2014.
- [26] Y. Li, C. Messina, M. Bendaoud, D. H. Fine, H. Schreiner, and V. K. Tsiagbe, "Adaptive immune response in osteoclastic bone resorption induced by orally administered *Aggregatibacter actinomycetemcomitans* in a rat model of periodontal disease," *Molecular Oral Microbiology*, vol. 25, no. 4, pp. 275–292, 2010.
- [27] Z. M. Wang, J. Zhang, B. S. Pang, L. Q. Zhang, and C. Wang, "Detection of serum interleukin-21 associated with chronic periodontitis," in *Proceedings of the 86th General Session & Exhibition of the IADR*, Toronto, Canada, July 2008.
- [28] N. Nizam, O. K. Basoglu, M. S. Tasbakan, A. Nalbantsoy, and N. Buduneli, "Salivary cytokines and the association between obstructive sleep apnea syndrome and periodontal disease," *Journal of Periodontology*, vol. 85, no. 7, pp. e251–e258, 2014.
- [29] N. Dutzan, C. Rivas, J. García-Sesnich et al., "Levels of interleukin-21 in patients with untreated chronic periodontitis," *Journal of Periodontology*, vol. 82, no. 10, pp. 1483–1489, 2011.
- [30] L. Zhao, Y. Zhou, Y. Xu, Y. Sun, L. Li, and W. Chen, "Effect of non-surgical periodontal therapy on the levels of Th17/Th1/Th2 cytokines and their transcription factors in Chinese chronic periodontitis patients," *Journal of Clinical Periodontology*, vol. 38, no. 6, pp. 509–516, 2011.
- [31] N. Dutzan, R. Vernal, J. P. Vaque et al., "Interleukin-21 expression and its association with proinflammatory cytokines in untreated chronic periodontitis subjects," *Journal of Periodontology*, vol. 82, pp. 1483–1489, 2012.
- [32] M. H. Napimoga, L. H. A. C. Nunes, A. A. B. Maciel et al., "Possible involvement of IL-21 and IL-10 on salivary IgA levels in chronic periodontitis subjects," *Scandinavian Journal of Immunology*, vol. 74, no. 6, pp. 596–602, 2011.
- [33] D. M. Isaza-Guzmán, N. Cardona-Vélez, D. E. Gaviria-Correa, M. C. Martínez-Pabón, M. C. Castaño-Granada, and S. I. Tobón-Arroyave, "Association study between salivary levels of interferon (IFN)-gamma, interleukin (IL)-17, IL-21, and IL-22 with chronic periodontitis," *Archives of Oral Biology*, vol. 60, no. 1, pp. 91–99, 2015.
- [34] H. Löe, A. Anerud, H. Boysen, and M. Smith, "The natural history of periodontal disease in man: the rate of periodontal destruction before 40 years of age," *Journal of Periodontology*, vol. 49, no. 12, pp. 607–620, 1978.
- [35] F. A. Scannapieco and R. J. Genco, "Association of periodontal infections with atherosclerotic and pulmonary diseases," *Journal of Periodontal Research*, vol. 34, no. 7, pp. 340–345, 1999.

- [36] R. Ding, W. Gao, Z. He et al., "Effect of serum interleukin 21 on the development of coronary artery disease," *APMIS*, vol. 122, no. 9, pp. 842–847, 2014.
- [37] J. Kim, S. Kang, J. Kim, G. Kwon, and S. Koo, "Elevated levels of T helper 17 cells are associated with disease activity in patients with rheumatoid arthritis," *Annals of Laboratory Medicine*, vol. 33, no. 1, pp. 52–59, 2013.
- [38] D. L. Sackett, "Rules of evidence and clinical recommendations on the use of antithrombotic agents," *Chest*, vol. 95, no. 2, pp. 2–4, 1989.