Dowling, P.; et al. Characterization of Contractile Proteins from Skeletal Muscle Using Gel-Based Top-Down Proteomics. *Proteomes* 2019, 7, 25

Paul Dowling [1,2], Margit Zweyer [3], Dieter Swandulla [3] and Kay Ohlendieck [1,2,*]

[1]   Department of Biology, Maynooth University, National University of Ireland, Maynooth, W23F2H6 Co. Kildare, Ireland
[2]   MU Human Health Research Institute, Maynooth University, Maynooth, W23F2H6 Co. Kildare, Ireland
[3]   Institute of Physiology II, University of Bonn, D-53115 Bonn, Germany
*   Correspondence: kay.ohlendieck@mu.ie; Tel.: +353-1-708-3842

check for updates

The authors wish to make the following correction to their paper [1]:
The Figures 2 and 5 have some irregular typesetting. The corrected Figures 2 and 5 should be:


Figure 2. Overview of the bioanalytical advantages versus potential technical limitations of gel-based top-down proteomics, especially in relation to two-dimensional gel electrophoresis (2DGE) approaches.

Figure 5. Overview of fluorescence two-dimensional difference in-gel electrophoresis (2D-DIGE) techniques for the comparative analysis of proteoforms. The upper panel shows the two main DIGE approaches for the systematic pre-electrophoretic labelling of protein fractions using fluorescent CyDyes. These methods utilise minimal sub-stochiometric labelling of assessable lysines in proteins or saturation labelling of assessable cysteines in proteins. The lower panel outlines three-dye (Cy2, Cy3, Cy5) versus two-dye (Cy3, Cy5) DIGE labelling methodology.

Reference

1.   Dowling, P.; Zweyer, M.; Swandulla, D.; Ohlendieck, K. Characterization of Contractile Proteins from Skeletal Muscle Using Gel-Based Top-Down Proteomics. *Proteomes* **2019**, *7*, 25. [CrossRef] [PubMed]