



# Initial Development of User-Based Quality Evaluation Questionnaire of Smartwatch Technology for Applying to Healthcare

*Soo-Kyung Lee<sup>1</sup>, Gi Yon Kim<sup>2</sup>, Eun Ji Seo<sup>3</sup>, \*Youn-Jung Son<sup>4</sup>*

1. College of Nursing, Keimyung University, Daegu, 42601, Korea

2. Yonsei University Wonju College of Nursing, Wonju 220751, Korea

3. College of Nursing, Research Institute of Nursing Science, Ajou University, Suwon 16499, Korea

4. Red Cross College of Nursing, Chung-Ang University, 84 Heukseok-ro Dongjak-Gu, Seoul 06974, Korea

\*Corresponding Author: Email: yjson@cau.ac.kr

(Received 11 Nov 2021; accepted 19 Feb 2022)

## Abstract

**Background:** Smartwatches are a consumer wearable device offering a potential, practical, and affordable method to collect personal health data in healthy adults. For patients with chronic diseases, this would enable symptom monitoring and aid clinical decision making. Therefore, providing customized checklists to recommend smartwatches is beneficial. However, few studies have evaluated the practical functions of smartwatches and their influence on user acceptance. We aimed at developing a reliable tool to assess the quality of smartwatches from the users' perspective.

**Methods:** To develop the smartwatch rating scale (SWRS), we conducted a comprehensive literature review as well as reviewed relevant websites. The SWRS includes 22 items for the usability (usability, functionality, safety, material, and display) and five items for the acceptance and adoption domain (satisfaction and intention). We measured the scale's internal consistency and inter-rater reliability by evaluating seven smartwatches.

**Results:** The overall scale demonstrated an excellent level of internal consistency (Cronbach's alpha = 0.91), with each subscale's internal consistency above good level (0.74 ~ 0.92). Inter-rater reliability using intraclass correlation coefficients (ICC) was at good level (2-way random ICC = 0.82, 95% CI 0.09 – 0.97).

**Conclusions:** The SWRS is reliable, which can meet the need for assessment of smartwatch technology for utilizing in personal healthcare. Accounting for users' perspectives will help make the most of technology without impairing the human aspects of care, this study can help consumers choose a smartwatch based on their preferences and provide guidelines for developing user-friendly wearable devices aimed at health behavior changes.

**Keywords:** Smartwatches; Mobile health; Wearable technology; Healthcare; Questionnaires

## Introduction

In recent years, the development of wearable technology is increasing (1,2). Their usage has also increased, and wearable technology now plays a pivotal role in the lives of users (3,4). The worldwide market for wearable devices grew 82.3% in the

fourth quarter of 2019 according to International Data Corporation (5). Wearable technology for the wrist is at the center of the fast-expanding wearable market (6,7). Smartwatches that are the



Copyright © 2023 Lee et al. Published by Tehran University of Medical Sciences.

This work is licensed under a Creative Commons Attribution-NonCommercial 4.0 International license.

(<https://creativecommons.org/licenses/by-nc/4.0/>). Non-commercial uses of the work are permitted, provided the original work is properly cited

electronic watch and computer with smartphone applications installed.

Smartwatches are highly promoted in the information and communications technology industry for their multiple functions that interest users, such as notifications that are synchronized with smartphones and other applications; smartwatches can also offer continuous data monitoring functions, such as step-counting, heart rate tracking, energy consumption as well as physical activity monitoring, which can promote health (7–9). Recent studies have confirmed the positive effects of smartwatches on preventive healthcare and self-management for chronic diseases; they can also function as a tool offering real-time health information to healthcare professionals (10–13). For example, by automatic monitoring of smoking episodes using the smartwatch, motivated smokers were able to facilitate smoking reduction (14). Smartwatches providing real-time feedback also improved the quality of cardio-pulmonary resuscitation performed by healthcare providers (15).

However, despite the usefulness and effectiveness, the effort to assess the user-based quality evaluation of smartwatch has lagged when compared to smartphones and tablets, and few studies have been conducted on the factors of smartwatch adoption (16,17). One of the main reasons for the low user acceptance of smartwatch could be the excessive number of choices available, given little accurate information or knowledge about the devices (18). Previous studies have reported that navigate through the large number of available brands, such as Fitbit, Xiaomi, Apple, Garmin, and Samsung, is a burden for consumers (1,19–21). With such difficulties, therefore, it is no surprise that consumers heavily take into account the brand image and price of the product rather than their functional features (4). In addition, some studies have found that perceived uselessness, perceived price, perceived novelty, and self-efficacy of smartwatches had an impact on consumer resistance to using them (22,23).

Meanwhile, other electronic devices, such as smartphones, tablets, or laptops, have huge amounts of information available to consumers (24). In the case of mobile phone applications,

consumers can readily view comprehensive consumer star ratings and reviews (22,25). For example, the user version of the Mobile Application Rating Scale (MARS), a simple and objective quality evaluation tool for consumers, serves as a credible and reliable guideline for choosing the best health mobile apps available, but no such tool exists for smartwatches (26,27). Thus, there is need to develop a comprehensive rating scale that is parallel to the MARS for smartwatches. However, previous studies regarding smartphones have mostly been focused on the extension of the technology acceptance model or unified theory of acceptance and use of technology and studied their implication on the factors that are associated with usage perception of users (28,29). Such approaches may help understanding the motivation of purchasing a smartwatch but tell little about what practical functions potential users are seeking. Even though Dehghani's (2016) work categorized the adoption factors a more detailed level, the study still does not discuss quality assessment for adopting smartwatches (30).

Thus, we aimed to develop a smartwatch rating scale based on user aspects and to test its reliability for applying to personal healthcare based on user preferences.

## Materials and Methods

### *Development process outline*

This was a methodological study for developing a smartwatch rating scale (SWRS). To create the rating scale, we reviewed a number of published articles about acceptance, adoption, and usability of smartwatch and websites about rating smartwatches. Based on the literature review, keywords for smartwatch rating scales were extracted. By using the extracted keywords, we initially developed the SWRS for assessing the user-based quality evaluation.

### *Smartwatch rating scale (SWRS) development Extracting keywords from literature review and related websites*

We conducted a comprehensive literature search on the factors that influence smartwatch adoption, such as the price, aesthetics, and functionalities, in PubMed, ACM digital library, IEEE-Xplore, and Google scholar. The search keywords were “smartwatch” AND “criteria” OR “assess” OR “evaluate”. Employing our inclusion criteria, we included articles that were focused on the selection criteria of wearable technology.

A total of 56 articles were found. After initial review, we removed 48 articles that did not explore the adoption of smartwatch technology and were left with 8 articles. However, the articles provided broad categories, such as aesthetics and daily use. As an alternative method, we searched readily available websites that provide comprehensive reviews and comparisons of smartwatches with same keywords adding “comparison” OR “chart” for a more explicit criterion. Out of the search results, we selected the websites that provide an explicit and unified rating criterion. Of the smartwatch reviewing websites, 16 met the inclusion criteria. We extracted a total of 141 keywords about the criteria of smartwatch adoption considering the user aspect from 8 articles and 16 websites. The irrelevant keywords, duplicates, and keywords with low frequency (appearing less than twice) were removed; 48 keywords remained (Supplementary Fig. 1- Not published. Readers may contact the authors if needed).

### ***Creating a draft of the SWRS***

An expert panel with extensive experience in medical informatics and ubiquitous health categorized the 48 keywords into the usability domain. Then, each keyword was converted into a corresponding question, utilizing a 5-point Likert scale for a quantitative survey. The usability domain of the scale consisted of 22 items about smartwatch quality, including whether the smartwatch is comfortable for daily use, does it have the functions that consumers demand, is it safe and strong enough for daily use, is its material appropriate, and is the smartwatch’s display clear and readable. The 22 items of assessment criteria identified in previous research were extracted. There is no criteria or evidence

driven framework for smartwatch quality evaluation, we used relevant questions of a reliable tool called MARS which is most popular scale that evaluate the quality of health-related apps (26). The usability domain was divided into usability (five items), functionality (six items), safety (three items), material (four items), and display (four items). A smartwatch could be considered a computer, but at the same time, its aesthetic features significantly influence consumer adoption (4). This implies that potential consumers take into consideration numerous subjective factors when choosing a smartwatch. A rating scale to measure the user-based quality of smartwatches should incorporate this subjective feature. Therefore, we added the acceptance and adoption domain which includes five items to evaluate the satisfaction and intention of user based on the Mobile Application Rating Scale (26). Finally, the SWRS includes two domains with 27 items.

### ***Testing the internal consistency and inter-rater reliability of the final SWRS***

To test the internal consistency and inter-rater reliability of the SWRS, we checked the market ranking data of current mainstream market-leading wrist-wearable devices (3). The seven smartwatches were selected for the following reasons. First, we selected the internationally renowned smart bracelets, according to market research data by NPD and Canalys (31, 32), Fitbit had the largest market share and also exhibited a good market performance. So, Fitbit Surge (Fitbit Inc) was chosen to represent the foreign-made smart bracelets. Second, the shipment of the Mi Band in the health tracking devices market has been second to that of Fitbit. Thus, the Mi Band (Mi, China) was chosen to represent the Chinese brand bracelets (33). Third, when choosing smart watches, given that the functions of smart watches and mobile phones were close and that mobile phones from Samsung and Apple were leading products in the market, Samsung Gear S (Samsung Inc) and Apple Watch (Apple Inc) were chosen to represent the smart watches (34). Then, we selected the seven smartwatches as representative; Fitbit Charge HR, Fitbit

Blaze (Fitbit Inc., San Francisco, CA, USA), Samsung Gear-fit 2, Samsung Gear S2, Samsung Gear S3 (Samsung Electronics Co., Ltd, Suwon, South Korea), Apple Watch 2 (Apple Inc., Cupertino, CA, USA), Xiaomi Mi-Band 2 (Xiaomi Inc., Beijing, China) (Supplementary Table 1- Not published. Readers may contact the authors if needed). After using each device for a minimum of three days, two researchers completed the survey for quality evaluation.

The internal consistency of the SWRS was calculated using Cronbach’s alpha. This indicates the degree (correlations) to which items measuring the same general construct produce similar scores. Inter-rater reliability of the SWRS was determined by the intra-class correlation coefficient (ICC) reflecting the accuracy that the rating process was based on. We performed ICC with a two-way random effect, average measures model with absolute agreement that regards evaluators as random effects, because the developed scale will be used by several evaluators (35,36). Confidence intervals

(CIs) for all ICCs were calculated to assess whether they differed from each other.

## Results

### *Final version of the SWRS*

As shown in Table 1, the final form of questionnaire for assessing user-based quality evaluation of smartwatch composed of usability (five questions), functionality (six questions), safety (three questions), material (four questions), and display (four questions) for the usability domain and five questions for the acceptance and adoption domain, with a total of 27 questions. The self-reported scale for the usability questions was rated on a 5-point Likert scale, from 1-point (“Very bad”) to 5-point (“Very good”). Additionally, responses to the acceptance and adoption domain were categorized into the total score, which was calculated by summing the 27 items. Scores ranged from 27 to 135, and higher scores indicated better overall quality of a smartwatch.

**Table 1:** The final version of the Smartwatch Rating Scale

<i>Items</i>	<i>Rating scale</i>				
Usability and acceptance domain					
Usability	Very bad	Bad	Moderate	Good	Very good
1. Is the size of the smartwatch appropriate/comfortable for daily use?					
2. Is the weight of the smartwatch appropriate/comfortable for daily use?					
3. Does the battery of the smartwatch last long enough?					
4. Is the charging method of the smartwatch convenient?					
5. Is the outer appearance of the smartwatch attractive?					
Functionality	Very bad	Bad	Moderate	Good	Very good
6. Is it convenient to access the main functions of the smartwatch?					
7. Is the gestural movement of the smartwatch (touch screen, scroll, etc.) smooth?					
8. Does the smartwatch respond quickly enough?					
9. Is the smartwatch compatible with user's phone?					
10. Is the memory of the smartwatch large enough?					
11. Does the smartwatch accurately measure what it should be measuring					

(Steps, calorie consumption, sleep, etc.)?						
Safety		Very bad	Bad	Moderate	Good	Very good
12. Does the smartwatch have water resistance?						
13. Does the smartwatch have dust resistance?						
14. Is the smartwatch safe from external shocks or scratch?						
Material		Very bad	Bad	Moderate	Good	Very good
15. Is the material of the watch appropriate (in terms of design, texture, strength, etc.)?						
16. Is the case material of the watch appropriate (in terms of design, texture, strength, etc.)?						
17. Is the band material of the watch appropriate (in terms of design, texture, strength, etc.)?						
18. Is the band lock of the smartwatch comfortable to use?						
Display		Very bad	Bad	Moderate	Good	Very good
19. Is the size of the display screen appropriate for daily use?						
20. Is the screen resolution appropriate?						
21. Is the layout and design of the screen appropriate?						
22. Is the readability of the smartwatch (under sun, in dark, etc.) appropriate?						
Adoption domain	Rating scale					
1. Are you satisfied with the functions of the smartwatch?	Very dissatisfied	No	Somewhat	Yes	Very satisfied	
2. Are you satisfied with the price of the smartwatch?	Very dissatisfied	No	Somewhat	Yes	Very satisfied	
3. Do you intend to recommend the smartwatch to your family members/friends/acquaintances?	Not at all	Maybe	Possibly	Yes	Definitely	
4. How many times do you intend to use the smartwatch over a week?	Not at all	1-2 times	3-4 times	5-6 times	Everyday	
5. What is your overall rating for the smartwatch?	1	2	3	4	5	

### ***Internal consistency and inter-rater reliability of the SWRS***

As shown in Table 2, we confirmed the internal consistency and inter-rater reliability using ICC. The overall score of SWRS for the selected seven smartwatches demonstrated an excellent level of internal consistency (Cronbach's alpha = 0.91). Each subscale's internal consistency examined was

above good level (0.74 ~ 0.92). The SWRS ratings of the seven smartwatches had a significantly good inter-rater reliability by the CI (2-way random ICC = 0.82, 95% CI = 0.09 – 0.97), meaning that the SWRS had an acceptable inter-rater reliability (35,36). However, the ICC of safety and material subscales were not significant.

**Table 2:** Internal consistency and inter-rater reliability of the usability and acceptance domain (N=7)

<i>Item</i>	<i>Subscale</i>	<i>Mean</i>	<i>SD</i>	<i>Internal consistency</i>	<i>Inter-rater reliability (95% CI) / p value</i>
<u>Usability</u>					
1	Size	3.71	0.95	0.91	0.96 (0.79 ~ 0.99)/ <i>P</i> = 0.001
2	Weight	4.14	0.90		
3	Battery	3.14	0.69		
4	Charging	3.71	0.76		
5	Appearance	3.86	1.07		
<u>Functionality</u>					
6	Accessibility	4.14	0.38	0.92	0.85 (0.04 ~ 0.98)/ <i>P</i> = 0.023
7	Gesture	4.59	0.76		
8	Speed	4.14	0.69		
9	Memory	3.43	0.79		
10	Measurement accuracy	3.71	0.76		
11	Phone compatibility	4.29	0.95		
<u>Safety</u>					
12	Water resistance	3.00	0.82	0.76	0.56 (-1.57 ~ 0.93)/ <i>P</i> = 0.177
13	External shock	3.00	0.58		
14	Dust resistance	3.43	0.54		
<u>Material</u>					
15	Material	4.00	0.58	0.74	0.24 (-3.30 ~ 0.87)/ <i>P</i> = 0.375
16	Case	3.86	0.69		
17	Strap	4.29	0.76		
18	Lock	3.14	0.90		
<u>Display</u>					
19	Screen size	3.57	0.98	0.90	0.92 (0.57 ~ 0.99)/ <i>P</i> = 0.003
20	Resolution	3.86	0.90		
21	Layout	3.86	0.69		
22	Readability	4.29	0.49		
	Total	3.77	0.45	0.91	0.82 (0.09 ~ 0.97)/ <i>P</i> = 0.027

*Note*, Internal consistency: Cronbach's alpha / Inter-rater reliability: Intraclass correlation coefficient (ICC)

## Discussion

The SWRS developed in this study provides a function chart which could be used as a checklist when searching for a smartwatch which has a specific function. The SWRS consisted of 27 items to evaluate usability, functionality, safety, materials, comprising the usability domain, and to evaluate user's satisfaction as the acceptance and adoption domain, which is thought to reflect the result of a previous study that revealed that users' intention of a smartwatch is affected by its usefulness, enjoyment, cost, and satisfaction (18). In addition, user-based perspective of the SWRS may help

consumers choose a smartwatch that has functions tailored to their needs and help doctors recommend a wearable device to their patients for personalized symptom monitoring to allow for better disease management. Smartwatches are used in various interventions because they can collect individual information of users in real-time and identify physical information according to the situation (2). It is essential to select the most appropriate smartwatch for each intervention, and the SWRS can be used as an obvious criterion to make this selection. Conducting usability testing on the smartwatch prior to intervention could also increase patients' compliance (8).

Data reliability is a critical issue, especially when data are used to develop the tool or create a program (37). Inter-rater reliability was assessed for the quantitative measurements scored by both raters in the composite scorecard using weighted kappa with a range of zero (no agreement) to one (perfect agreement) (38). In the present study, ICC was used to examine whether there was inter-rater reliability. The standard of ICC is 0.0 to 0.20 for slight, 0.21 to 0.40 for fair, 0.41 to 0.60 for moderate, 0.61 to 0.80 for substantial, and 0.81 to 1.00 for almost perfect (39). Our findings showed that the total ICC for the SWRS has high reliability within and between the assessors. However, the safety and material among the five subscales of the usability domain did not have significance for ICC. Less reliable and accurate ratings may result when data are rated in the middle of the rating scale, are ambivalent, hard to rate, or rated with low confidence. Consequently, it is especially important to the SWRS to review those items, and to carefully consider item length and vocabulary usage. Furthermore, prior training of raters would be helpful as it would standardize the information provided in these components. Future research is required to determine how to best evaluate the safety and material of smartwatch technology. The issue of measurement reliability is an important and necessary first step in determining a tool's utility (36,37). However, further research into other psychometric properties, like test-retest reliability, which would compare the entire assessment process over two time points and exhibit concurrent validity with advanced camera systems, is warranted. One interesting finding is that the number of functions contained within a smartwatch did not necessarily mean higher objective quality. These results seem to be related to the lack of consumer's knowledge of the smartwatch function or lack of interest in function operation. Therefore, it is thought that smartwatches that reflect the common needs of users, regardless of their age or experience in use of smartwatches, will be more acceptable to them. There are some limitations. First, this study included keywords which were extracted from non-scholarly articles and multiple sources, which can

be the most frequently appearing or newly emerging keywords. Accordingly, this approach can draw a selection bias. Second, only a few well-established brands were assessed to test the inter-rater reliability. Third, only two raters assessed the reliability for SWRS, thereby affecting the accuracy of the inter-rater reliability measured. Last over the course of time it took to complete this research, much has changed with technology, market share and the number of related studies has changed. The smartwatches can monitor heart rate and respiration rate with and provide important information regarding arrhythmia.

## **Conclusion**

The SWRS is reliable, which can meet the need for assessment of smartwatch technology. Accordingly, the SWRS could provide evidence-based information to support the need for smartwatches among customers and healthcare providers. Moreover, the SWRS could be used to identify problems between smartwatch types and their intended user populations so that developers can make responsive design changes. We suggest that raters should be sufficiently and appropriated trained. Future research is required to determine the suitability and reliability of the SWRS using larger samples.

## **Journalism Ethics considerations**

Ethical issues (Including plagiarism, informed consent, misconduct, data fabrication and/or falsification, double publication and/or submission, redundancy, etc.) have been completely observed by the authors.

## **Acknowledgments**

This research was supported by the Basic Science Research Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Education (2015R1C1A1A01055753; 2018R1D1A3B07050652).

## Conflicts of Interest

The authors declare no conflict of interest.

## References

- Baba NA, Baharudin AS, Alomari A (2019). Determinants of users' intention to use smartwatch. *J Theor Appl Inf Technol*, 97(18):4738-50.
- Guk K, Han G, Lim J, et al. (2019). Evolution of wearable devices with real-time disease monitoring for personalized healthcare. *Nanomaterials (Basel)*, 9(6):813.
- Canalys, Worldwide wearable band market grew 65% in Q3 2019 while Asia Pacific doubled in volume. <https://www.canalys.com/newsroom/canalys-worldwide-wearable-market-q3-2019>
- Jung Y, Kim S, Choi B (2016). Consumer valuation of the wearables: The case of smartwatches. *Comput. Human Behav*, 63:899-905.
- IDC. Shipments of Wearable Devices Reach 118.9 Million Units in the Fourth Quarter and 336.5 Million for 2019, According to IDC. <https://www.telecomtv.com/content/idc/shipments-of-wearable-devices-reach-118-9-million-units-in-the-fourth-quarter-and-336-5-million-for-2019-according-to-idc-37984/>
- Chun JS. World smartwatch sales in Q2 12.3 million units 44% High Growth. Yonhap News Agency. 2019. <https://www.yna.co.kr/view/AKR20190906024000017>
- Kheirkhahan M, Nair S, Davoudi A, et al (2019). A smartwatch-based framework for real-time and online assessment and mobility monitoring. *J Biomed Inform*, 89:29-40.
- Lu TC, Fu CM, Ma HM, Fang CC, Tumer AM (2016). Healthcare applications of smart watches: A systematic review. *Appl Clin Inform*, 7(3):850-69.
- Piwiek L, Ellis DA, Andrews S, Joinson A (2016). The rise of consumer health wearables: Promises and barriers. *PLoS Med*, 13(2): e1001953.
- King CE, Sarrafzadeh M (2018). A survey of smartwatches in remote health monitoring. *J Healthc Inform Res*, 2(1-2):1-24.
- Jia Y, Wang W, Wen D, Liang L, Gao L, Lei J (2018). Perceived user preferences and usability evaluation of mainstream wearable devices for health monitoring. *PeerJ*, 6:e5350.
- van Helmond N, Freeman CG, Hahnen C, et al (2019). The accuracy of blood pressure measurement by a smartwatch and a portable health device. *Hosp Pract (1995)*, 47(4):211-5.
- Wasserlauf J, You C, Patel R, Valys A, Albert D, Passman R (2019). Smartwatch performance for the detection and quantification of atrial fibrillation. *Circ Arrhythm Electrophysiol*, 12(6): e006834.
- Dar R (2018). Effect of real-time monitoring and notification of smoking episodes on smoking reduction: A pilot study of a novel smoking cessation app. *Nicotine Tob Res*, 20(12): 1515-8.
- Lu TC, Chang YT, Ho TW, et al (2019). Using a smartwatch with real-time feedback improves the delivery of high-quality cardiopulmonary resuscitation by healthcare professionals. *Resuscitation*, 140: 16-22.
- Rauschnabel PA, Krey N, Chuah S, Nguyen B, Lade S, Ramayah T (2016). Exploring the adoption of smartwatches. *Digital Enterprise Computing*, 39-48.
- Strain T, Wijndaele K, Brage S (2019). Physical activity surveillance through smartphone apps and wearable trackers: Examining the UK potential for nationally representative sampling. *JMIR Mhealth Uhealth*, 7(1): e11898.
- Kim KJ (2016). Round or Square? How Screen Shape Affects Utilitarian and Hedonic Motivations for Smartwatch Adoption. *Cyberpsychol Behav Soc Netw*, 19(12): 733-9.
- Anggraini N, Kaburuan ER, Wang G, Jayadi R (2019). Usability study and users' perception of smartwatch: Study on Indonesian customer. *Procedia Comput Sci*, 161: 1266-74.
- Cheung ML, Chau KY, Lam MHS, Tse G, Ho KY, Flint SW (2019). Examining consumers' adoption of wearable healthcare technology: The role of health attributes. *Int J Environ Res Public Health*, 16(13): 2257.
- Gualtieri L, Rosenbluth S, Phillips J (2016). Can a free wearable activity tracker change behavior? The impact of trackers on adults in a physician-led wellness group. *JMIR Res Protoc*, 5(4): e237.
- Mani Z, Chouk I (2017). Drivers of consumers? Resistance to smart products. *J Market Manag*, 33(1-2): 76-97.
- Ha T, Beijnon B, Kim S, Lee S, Kim JH (2017).



- Examining user perceptions of smartwatch through dynamic topic modeling. *Telematics and Informatics*, 34(7): 1262–73.
24. Cecchinato ME, Cox AL, Bird J (2015). Smartwatches: The good, the bad and the ugly? Proceedings of the 33rd annual ACM conference extended abstracts on human factors in computing systems. *A.C.M.*, 2133-8.
  25. Nwosu AC, Quinn C, Samuels J, Mason S, Payne TR (2018). Wearable smartwatch technology to monitor symptoms in advanced illness. *BMJ Support Palliat Care*, 8(2): 237.
  26. Stoyanov SR, Hides L, Kavanagh DJ, Zelenko O, Tjondronegoro D, Mani M (2015). Mobile app rating scale: A new tool for assessing the quality of health mobile apps. *JMIR Mhealth Uhealth*, 3(1): e27.
  27. Foster KR, Torous J (2019). The opportunity and obstacles for smartwatches and wearable sensors. *IEEE Pulse*, 10(1): 22-5.
  28. Chen CC, Shih HS (2014). A study of the acceptance of wearable technology for consumers: An analytical network process perspective. *International Journal of the Analytic Hierarchy Process*, 1-5.
  29. Venkatesh V, Morris MG, Davis GB, Davis FD (2003). User acceptance of information technology: Toward a unified view. *MIS Quarterly*, 27(3): 425-478.
  30. Dehghani M (2016). An assessment towards adoption and diffusion of smart wearable technologies by consumers: The cases of smart watch and fitness wristband products. *27th ACM Conference on Hypertext and Social Media*, CEUR Workshop Proceedings, 1628.
  31. Canals, Fitbit accounted for nearly half of global wearable band shipments in Q1 2014. Available at <https://www.canals.com/newsroom/fitbit-accounted-nearly-half-global-wearable-band-shipments-q1-2014> Accessed 11/20/2018.
  32. Canals, Fitbit accounted for nearly half of global wearable band shipments in Q1. Available at <https://www.canals.com/newsroom/medialert-fitbit-maintains-leadership-share-wearable-band-market-apple-watch-entrance> Accessed 11/20/2018.
  33. Mobihealthnews, Fitbit, Jawbone, Nike had 97 percent of fitness tracker retail sales. Available at <http://www.mobihealthnews.com/28825/fitbit-jawbone-nike-had-97-percent-of-fitness-tracker-retail-sales-in-2013/> Accessed 11/23/2018.
  34. Wen D, Zhang X, Liu X, Lei J (2017). Evaluating the consistency of current mainstream wearable devices in health monitoring: a comparison under free-living conditions. *J Med Internet Res*, 19(3): e68.
  35. Gwet KL (2008). Computing inter-rater reliability and its variance in the presence of high agreement. *Br J Math Stat Psychol*, 61(1): 29-48.
  36. Liljequist D, Elfving B, Skavberg Roaldsen K (2019). Intraclass correlation: A discussion and demonstration of basic features. *PLoS One*, 14(7): e0219854.
  37. Zapf A, Castell S, Morawietz L, Karch A (2016). Measuring inter-rater reliability for nominal data - which coefficients and confidence intervals are appropriate?. *BMC Med Res Methodol*, 16: 1-10.
  38. Sim J, Wright CC (2005). The Kappa Statistic in Reliability Studies: Use, Interpretation, and Sample Size Requirements. *Physl Ther*, 85(3): 257–68.
  39. McHugh ML (2012). Interrater reliability: the kappa statistic. *Biochem Med (Zagreb)*, 22(3): 276-282.