

REVIEW ARTICLE

Genome triplication drove the diversification of *Brassica* plants

Feng Cheng, Jian Wu and Xiaowu Wang

The genus *Brassica* belongs to the plant family Brassicaceae, which includes many important crop species that are used as oilseed, condiments, or vegetables throughout the world. *Brassica* plants comprise many diverse species, and each species contains rich morphotypes showing extreme traits. *Brassica* species experienced an extra whole genome triplication (WGT) event compared with the model plant *Arabidopsis thaliana*. Whole genome sequencing of the *Brassica* species *Brassica rapa*, *Brassica oleracea* and others demonstrated that WGT plays an important role in the speciation and morphotype diversification of *Brassica* plants. Comparative genomic analysis based on the genome sequences of *B. rapa* and *A. thaliana* clearly identified the WGT event and further demonstrated that the translocated Proto-Calepine Karyotype (tPCK, $n=7$) was the diploid ancestor of the three subgenomes in *B. rapa*. Following WGT, subsequent extensive genome fractionation, block reshuffling and chromosome reduction accompanied by paleocentromere descent from the three tPCK subgenomes during the rediploidization process produced stable diploid species. Genomic rearrangement of the diploid species and their hybridization then contributed to *Brassica* speciation. The subgenome dominance effect and biased gene retention, such as the over-retention of auxin-related genes after WGT, promoted functional gene evolution and thus propelled the expansion of rich morphotypes in the *Brassica* species. In conclusion, the WGT event initiated subsequent genomic and gene-level evolution, which further drove *Brassica* speciation and created rich morphotypes in each species.

Horticulture Research (2014) 1, 14024; doi:10.1038/hortres.2014.24; published online: 21 May 2014

GENUS *BRASSICA* IN BRASSICACEAE

Plants of the genus *Brassica* are grouped into the tribe Brassiceae, which belongs to the plant family Brassicaceae. Brassicaceae comprises a large family of plants that exhibit common and distinct features in their flowers. The flowers have cruciform petals and six stamens, two of which are short outer stamens. In total, Brassicaceae is composed of 3709 species and 338 genera,¹ with 308 of the 338 genera further assigned to 44 tribes.² Among the abundant Brassicaceae species, the genus *Brassica* is important because it contains many economically valuable crops that are used as oilseeds, condiments, and culinary vegetables. *Brassica* species share an additional common feature in that they all experienced an extra whole genome triplication (WGT) event, which occurred approximately 9–15 million years ago^{3,4} or even approximately 28 million years ago.^{5–7}

THE U'S TRIANGLE MODEL DESCRIBES THE RELATIONSHIP AMONG *BRASSICA* CROPS

Six species of the genus *Brassica* are used widely throughout the world as oilseed, condiments, fodder or vegetable crops. Three of these species are diploid (*Brassica rapa*, $n=10$; *B. nigra*, $n=8$; and *B. oleracea*, $n=9$), whereas the other three are allotetraploids (*B. juncea*, $n=18$; *B. napus*, $n=19$; *B. carinata*, $n=17$) derived from each pair of the three diploid species. The genetic relationships of these species were identified and confirmed by extensive experimental crosses between tetraploid and/or diploid plants as well as karyotyping or microscopic inspection at the synapsis stage of meiosis in these crosses.⁸ For example, when crosses were performed between *B. napus* and *B. rapa*, F1 plants with a chromosome number of $n=29$ were generated. When there were two sets of chromosomes from the *B. rapa* genome, only nine normal pairs of synaptic chromosomes at the synapsis stage in meiotic cells of the F1 plants were observed by microscopic inspection.⁹ The other 10

chromosomes of *B. napus* that can never form normal pairs of synaptic chromosomes in the F1 were affected by the chromosomes of the A genome, *B. rapa*. This supports the theory that these 10 chromosomes of *B. napus* are homologous to the 10 chromosomes in *B. rapa*.⁹ Similar experiments using crosses of *B. napus* and *B. oleracea* showed that the other nine chromosomes of *B. napus* are homologous with the nine chromosomes of *B. oleracea*. Taken together, these results lead us to conclude that *B. napus* is a tetraploid of *B. rapa* and *B. oleracea*.^{8,9} Based on experimental evidence, the relationships of the six species were simply described by the U's triangle model,⁸ in which the three diploid species *B. rapa*, *B. nigra* and *B. oleracea* are considered to be the basic genomes A, B and C, respectively, and are placed at the three vertices of the triangle. The three allotetraploids, *B. juncea*, *B. napus* and *B. carinata*, which are hybrids of AB, AC and BC, respectively, are placed in the middle of the three edges of the triangle. U's triangle has been successfully applied to aid in understanding the relationships among *Brassica* crops and has fostered the genetic study of these species.

THE RICH DIVERSITY OF *BRASSICA* PLANTS

Brassica plants have rich diversity with respect to both speciation and the abundant morphotypes in each *Brassica* species. *Brassica* crops described by U's triangle are close relatives, and many traits are shared but developed independently and in parallel, such as heading leaves and enlarged roots (Figure 1). One of the important vegetables of *B. rapa* is Chinese cabbage, which has the distinct feature of a leafy head; this feature is also observed in *B. oleracea* and *B. juncea*. Turnip, another morphotype of *B. rapa*, develops enlarged roots as storage organs, and this feature is also found in *B. juncea* and *B. napus*. Furthermore, each *Brassica* species has evolved multiple morphotypes, including leafy heads and enlarged roots, other enlarged organs of stems and inflorescences, oilseeds, sarsons and even ornamental features.¹⁰ Different morphotypes




Figure 1. Rich morphotypes of *Brassica* plants. (a) Morphotypes of *B. rapa*; top two lines from left to right: pak choi, heading *B. rapa*, turnip, oilseed, purple pak choi, caixin, mizuna, purple caitai and takucai; the third line shows additional morphotypes or varieties of the previous morphotypes. (b) Morphotypes of *B. oleracea*; top two lines from left to right: heading cabbage, Brussels sprouts, broccoli, cauliflower, purple cabbage, purple cauliflower, collard; the third line shows additional morphotypes or varieties. Some of the pictures were collected from the Internet.

have different usages. In *B. rapa* (Figure 1a), heading Chinese cabbage and pak choi are consumed as leafy vegetables. Chinese cabbage is distinct for its large leafy head, whereas pak choi has relatively smaller leaves and does not develop heading leaves. Turnip has an enlarged root that is eaten and occasionally used as fodder. Caixin and purple caitai bolt rapidly and generate long, tender stems used as food. Morphotypes of oilseed *B. rapa* produce large, full seeds for oil extraction and sarsons produce seed pods that are eaten in India. Some morphotypes of *B. rapa* develop beautiful leaf patterns and colors, thus are used as ornamental plants. *B. oleracea* also has an abundance of morphotypes, as shown in Figure 1b. Heading *B. oleracea* is consumed as a leaf vegetable, whereas oilseed *B. oleracea* produces edible oil. Cauliflower and broccoli, special morphotypes of *B. oleracea*, have developed enlarged inflorescences that are eaten as vegetables. Other *Brassica* crops, such as *B. juncea*, have even greater morphotype richness than *B. rapa* and *B. oleracea*. In addition to these cultivated crops, there are many wild relatives of the species in U's triangle that have greatly diversified phenotypes, further extending the diversity of *Brassica* plants.

The WGT event was important to the speciation and the expansion of rich morphotypes in the genus *Brassica*. The subsequent genomic rearrangement and gene evolution initiated by WGT promoted the appearance of a variety of *Brassica* plants.

CHROMOSOME EVOLUTION AFTER WGT PROMOTED *BRASSICA* SPECIATION

Genomic blocks (GBs) are collinear chromosome fragments conserved among different genomes. The genomes of Brassicaceae species are composed of 24 GBs labeled A to X. These blocks were defined by a comparative genomic analysis among the genomes of many Brassicaceae species, such as *B. napus*, *Arabidopsis thaliana*, *A. lyrata* and *Capsella rubella*.^{11,12} These GBs formed the basic units in ancestral chromosome reshuffling that generated the present-day species. Parkin *et al.*¹¹ constructed a high-density linkage map of *B. napus*, which is the allotetraploid of *B. rapa* and *B. oleracea*. Based on this map, they defined 21 GBs shared between *B. napus* and *A. thaliana*. Subsequently, Schranz and co-workers¹² combined the *B. napus* linkage map with comparative mapping results from

A. lyrata and *C. rubella*, both of which have eight chromosomes and are defined as the ancestral common karyotype (ACK or AK) of Brassicaceae, resulting in the definition of 24 GBs (A–X) as the basic units of all Brassicaceae genomes. Various combinations of these 24 blocks, occasionally accompanied by whole genome duplication, compose all of the genomes of Brassicaceae species. Genomes that contain only one set of the 24 GBs are considered diploid species. There are many such genomes in Brassicaceae; the main karyotypes include ACK ($n=8$), Proto-Calepine karyotype (PCK, $n=7$), translocated PCK (tPCK, $n=7$) and *A. thaliana* ($n=5$).^{12,13} Previous studies based on phylogenetics and comparative chromosome painting showed that ACK was the ancestral karyotype of Brassicaceae.^{12–15} ACK has eight chromosomes with GBs ordered from A to X across chromosomes one to eight. Examples of extant ACK species include *A. lyrata* and *C. rubella*.^{14,16,17} Both PCK and tPCK have seven chromosomes and differ in one inter-chromosomal translocation.¹³ *Conringia orientalis* has a PCK chromosome order,¹³ whereas *Schrenkiella parvula* has a tPCK genome.¹⁸ Genomes having more than one set of the 24 GBs in Brassicaceae are considered paleopolyploid species, including all *Brassica* crop species, which experienced a WGT event.

The genomic structure of the triplicated-genome species in the *Brassica* genus as well as their ancestral genome evolution were first studied in detail after the whole genome sequencing of the *Brassica* A genome *B. rapa*³ followed by the C genome *B. oleracea*.¹⁹ The sequencing of other *Brassica* species is now underway. Genome datasets of the *Brassica* species are maintained and continuously updated within the *Brassica* database (<http://brassicadb.org>).²⁰ For *B. rapa*, the genome size was estimated to be 485 Mb based on K-mer analysis, and gene prediction suggested 41 020 protein-coding gene models. For *B. oleracea*, the genome size was estimated to be 630 Mb with approximately 45 758 gene models.

Comparative genomic analysis between *B. rapa* and *A. thaliana* clearly indicated the WGT event experienced by *B. rapa*.³ Syntenic gene analysis between the triplicated genome of *B. rapa* and the diploid genome of *A. thaliana* using tool SynOrths showed that most genes inherited from their nearest common diploid ancestor were shared by both species (80.2% and 73.8% for *B. rapa* or *A. thaliana*, respectively).^{21,22} After WGT, the genomic fragments were reshuffled and fractionated. However, the local gene order was conserved, and syntenic genomic fragments can be clearly observed in both *B. rapa* and *A. thaliana*. Furthermore, for each GB in *A. thaliana*, three corresponding syntenic GBs in *B. rapa* were detected, which were generated by the WGT event.^{3,21} Genomic synteny analysis between *B. oleracea* and *A. thaliana* as well as between *B. rapa* and *B. oleracea* showed that *B. oleracea* has good genomic collinearity with genomes of *A. thaliana* and *B. rapa* (Figure 2). It was found that as observed in *B. rapa*, *B. oleracea* has three copies of each GB found in *A. thaliana*, thus confirming that the whole genome sequence level that *B. oleracea* also experienced the extra WGT event.¹⁹ A previous comparative genomic study of *B. juncea* and other *Brassicaceae* based on genetic maps showed that the genome of *B. nigra* also shared the WGT event.²³ Based on the comparison of GB distribution in the *Brassica* A, B and C genomes, we provided a framework for the comparative genomic study for *Brassica* species.

The diploid ancestor of *B. rapa* before WGT had seven chromosomes, which resembles the block arrangement of tPCK.²⁴ In *B. rapa*, there should be three sets of the 24 GBs, ideally 72 GBs in total, that resulted from WGT. Using *A. thaliana* as a reference, the genomic fragments of the three copies of each GB were clearly identified (with the exception of one copy of block G) in the genome of *B. rapa*.²⁴ For a certain GB, the three corresponding copies in *B. rapa* are not always all associated with a same GB. Some block




Figure 2. Framework for the comparative genomic analysis of *Brassica* plants. (a) Chromosomal synteny between *B. rapa* and *B. oleracea* determined based on whole genome sequences. Block associations for each chromosome are listed above or below the chromosome bars. The numbers 1, 2 and 3 placed after each block label (A–X) denote subgenomes LF, MF1 and MF2, respectively. (b) Evolutionary relationships between the chromosomes of *B. rapa* and *B. nigra*. Block information for *B. nigra* was extracted from the genetic map of *B. napus*.^{11,23} Block information is shown below each chromosome bar, and the syntenic chromosomes of *B. rapa* to *B. nigra* are listed above the chromosome bars.

associations are found in the ACK genome, such as block A, which associates with block B (block association A/B). However, some block associations do not exist in ACK, suggesting unique block associations for the diploid ancestor of *B. rapa* but not ACK; this lack of associations may result from genomic reshuffling after WGT. Based on this block distribution information, the block association relationships across the 10 chromosomes of *B. rapa* were observed. The breakage and formation of block associations occur independently; thus, the probability that an ancestral block association was broken more than twice after WGT is low, and the probability that a newly derived block association formed more than twice after WGT is also low. Based on this rule, by counting the copy numbers of all block associations in genome of *B. rapa* and comparing these numbers with the extant diploid karyotypes in Brassicaceae, such as ACK, PCK, tPCK and *A. thaliana*, Cheng and co-workers²⁴ found that the diploid ancestor of *B. rapa* had a tPCK-like karyotype. Furthermore, block association analysis of *B. oleracea* or *B. napus* and *R. sativus* based on genetic maps showed that these species also evolved from an ancestor having a tPCK genome.

The distribution patterns of transposable elements (TEs) support the positions of the 21 tPCK paleocentromeres in the genome of *B. rapa*. It is well known that TEs are enriched in the flanking regions of centromeres; this configuration has been observed in the genomes of many species such as *A. thaliana*, maize and soybean.^{25–27} TE sequences continue to show a relatively high density surrounding the positions of the 21 paleocentromeres in *B. rapa* millions of years after rediploidization following WGT. After reconstructing the three subgenomes of *B. rapa* along the seven tPCK ancestral chromosomes, using a method similar to playing a jigsaw puzzle (Supplementary Fig. S1), we plotted the TEs as a function of their density along the 21 reconstructed tPCK chromosomes (Figure 3).

This plot clearly shows that the TE distribution variation reflects the locations of the 10 inherited centromeres in *B. rapa* as well as the 11 inactivated paleocentromeres. The supported locations of the 21 paleocentromeres accurately match the centromere regions of tPCK, which are positioned between the block associations B/C, G/H, I/J, S/T, P/W, M/E and D/V for tPCK chromosomes one to seven, respectively.^{13,24} Other TE-rich regions, such as the distal end of AK2/5/6/8 in subgenome LF (the least fractionated subgenome) (Figure 3), could represent traces of paleocentromeres from more ancient genome duplications. In addition, as shown in Figure 3, gene fractionations or large genomic fragmental deletions are generally more concentrated near the paleocentromere regions compared to the genomic background.

Chromosomal reduction together with paleocentromere descent from the primal hexaploid ancestor (tPCK \times 3, $n=21$) is important for the speciation of *Brassica* plants. After WGT, extensive chromosome reshuffling during rediploidization contributed to the origin of closely related species in *Brassica*. As mentioned above, in the cross between *B. napus* and *B. rapa*, more than two copies of homologous chromosomes in the synapsis stage of meiosis will result in abnormal synaptonemal complexes, thereby decreasing the fertility of gametes. Logically, natural selection drives the rediploidization process with chromosomal rearrangement that removes the extra homologous chromosomes. Further rounds of genomic reshuffling of the rediploid ancestor at different evolutionary timepoints then created the different species in *Brassica*. In the *B. rapa* genome, the number of chromosomes and paleocentromeres was reduced from 21 to 10. The chromosomes were reduced by multi-chromosome translocation, fusion, and inter-/intrachromosomal recombination. Taking chromosomes A03 and A08 as examples,^{28,29} A03 evolution involved six chromosomes of tPCK (Figure 4a), AK2/5,




Figure 3. Distribution of TEs supporting the positions of the 21 paleocentromeres in the three tPCK subgenomes of the *B. rapa* genome. The color of each bin represents the ratio of TE sequences in the flanking region of a given gene used to reconstruct the tPCK subgenomes. The x-axis shows the position of the reconstructed chromosomes; the y-axis shows the three copies of the seven chromosomes in tPCK, which are the three subgenomes of *B. rapa*.




Figure 4. Chromosomal rearrangement of A03 and A08 in *B. rapa*. (a) The genomic block orders in the seven chromosomes of tPCK; block colors follow the labeling scheme of a previous report.¹² (b) The chromosome evolution of A03 involved six chromosomes of tPCK: AK2/5, AK7, AK2/5/6/8, AK3, AK6/8 and AK4. Red, green, and blue colors denote the subgenomes LF, MF1 and MF2. (c) The process by which two tPCK chromosomes, AK7 and AK1, were reshuffled into chromosome A03 of *B. rapa*.

AK7, AK2/5/6/8, AK3, AK6/8 and AK4 (a proposed chromosomal rearrangement process is shown in Figure 4b), whereas A08 was generated from several rounds of interchromosomal translocation of two tPCK chromosomes, AK1 and AK7 (Figure 4c). The circle model for block associations of M/N/T/U/D and V/K/L/Q/X (Figure 4b) or T/U (Figure 4c) has been used in previous reports.^{28,29} However, the chromosomal rearrangement explained by the circle model (Supplementary Fig. S2a) can also be achieved through an alternative process of chromosome translocation and fusion (Supplementary Fig. S2b).

GENE EVOLUTION AFTER WGT PROPELLED THE EXPANSION OF RICH MORPHOTYPES FOR BRASSICA SPECIES

Subgenome dominance has been detected among the three tPCK subgenomes in *B. rapa*.^{21,30} The subgenome dominance effect resulted in the differentiation of paralogous genes and featured the following characteristics: (i) one subgenome retained more genes than the other two through gene fractionation after WGT; (ii) genes located in the subgenome with high gene density are always expressed at higher levels than their paralogs in the other two subgenomes; and (iii) genes in the dominant subgenome accumulated fewer non-synonymous mutations than did the other

subgenomes.²¹ Gene density differentiation is clearly observed when counting the number of genes within the reconstructed tPCK subgenomes: the subgenome LF has approximately 1.6 times more genes than the other two subgenomes MF1 and MF2 (the more fractionated subgenomes one and two).^{3,21} Using mRNA-Seq data generated for different organs of *B. rapa*, a comparison of paralogous gene pairs showed that a greater number of genes located in subgenome LF are expressed at a higher level (i.e., either those showing at least two-fold greater expression or 'horseshoe' winners) than their paralogs in the MF subgenomes.²¹ The resequencing of different morphotypes of *B. rapa*, such as L144 and a turnip, showed that genes located in LF accumulated fewer functional mutations (non-synonymous single-nucleotide polymorphisms and frame-shift InDels) than those located in the MF subgenomes.²¹ This subgenome dominance effect has also been observed in the genome of maize.³¹

The three aspects of the dominance effect among the subgenomes of *B. rapa* are united by the rule of improving the fitness of the plant. Under this rule, genes that are expressed at higher levels than their paralogs should be more important for the biological function of the plant. Thus, functional mutations of these dominantly expressed genes would be more significant in reducing the plant's fitness than mutations of their syntenic paralogs.

Therefore, natural selection drives the conservation of the dominantly expressed genes against functional mutations, whereas their paralogs accumulate more mutations and eventually become fractionated, resulting in a higher gene density in the dominant subgenome and lower gene density in the dominated subgenomes. This explanation was first suggested following an analysis of the maize genome and subsequently for the genome of *B. rapa*.^{21,31,32}

Short homologous sequence-mediated deletion regulates gene fractionation in *B. rapa*. By investigating the fractionated genes in the *B. rapa* genome, it was found that genes were lost individually rather than via the simultaneous deletion of many genes located in a large fragment. Short repeated sequence-mediated individual gene fractionation has been observed in maize.³³ First, a pair of small direct repeats appear near the gene coding region before fractionation. The small repeated sequences then form a loop for intrachromosome recombination, and the gene sequence located in the middle of the two homologous repeat sequences is deleted. This mechanism was also found to function in the process of gene fractionation in *B. rapa*.³⁰

The 24-bp small RNA-targeted TE methylation that suppressed the expression of nearby genes as well as its biased distribution among the subgenomes of *B. rapa* led to subsequent subgenome dominance.³⁴ Small RNA-Seq data analysis showed that dominantly expressed genes in *B. rapa* always have fewer 24-bp RNA-targeted TEs in their 1-kb flanking regions compared with their paralogs. Previous reports on *A. thaliana* showed that small RNA-targeted TEs were subjected to methylation,^{35,36} and the methylated TEs then suppressed nearby gene expression. All of these observations suggest that the biased distribution of small RNA-targeted TEs played an important role in the formation of the subgenome dominance effect.

WGT provided a bulk of genes that served as both the raw materials and a buffer pool for multicopy genes to evolve disparate or new functions (subfunctionalization or neofunctionalization), whereas the subgenome dominance effect facilitated this process by differentiating the multicopy genes. These newly evolved functions further promoted the evolution of rich morphotypes in *Brassica*. In *A. thaliana*, after several rounds of whole genome duplication (α , β and γ polyploidization), many duplicated genes were subfunctionalized and/or neofunctionalized. For example, in *A. thaliana*, some genes from extra duplications have subfunctionalized compared with those in *Carica papaya*, such as the enzymes

CYP79A and *CYP79B* that catalyze the first step of glucosinolate synthesis.³⁷ Some genes have neofunctionalized to develop extra biosynthetic pathways for indole and methionine-derived aliphatic glucosinolates in *A. thaliana*, which are not detected in *C. papaya*. Analysis of glucosinolate genes in the genome of *B. rapa* showed that for most of these genes, multiple copies were retained after WGT.³⁸ These over-retained genes would be undersubfunctionalization or neofunctionalization to develop new biological functions related to glucosinolate metabolism in *B. rapa*, as in *A. thaliana*. It is expected that there are many more such examples of other over-retained genes in *B. rapa*. The subgenome dominance effect may aid in this evolutionary process by conserving one copy of the dominant gene and letting the other copies differentiate or develop new roles. Finally, these differentiated genes will contribute to the different traits of *B. rapa*.

Biased gene retention after WGT promoted the morphotype diversification of *Brassica* plants. Phytohormones, especially auxin, play important roles in plant morphogenesis.³⁹ The genes involved in plant hormone signaling pathways are thus important for divergent morphotype formation.^{39,40} By comparing the gene contents in *A. thaliana* and other sequenced genomes, such as *Carica papaya* and *Vitis vinifera*, it was found that auxin-related genes were expanded in the *B. rapa* genome.³ Furthermore, by comparing the number of gene categories that retained only one or multiple copies, genes involved in the response to phytohormone signaling were found to be significantly over-retained via gene fractionation following WGT in the genomes of *B. rapa*³ and *B. oleracea*.

TWO-STEP THEORY TO ILLUSTRATE THE WGT PROCESS IN BRASSICA

From a genome evolution perspective, a two-step theory of polyploidization was suggested to illustrate the process of WGT in *Brassica* plants.^{3,21} Based on the results of comparative subgenome analysis in *Brassica*, mainly in *B. rapa* as summarized above, we proposed that the WGT event occurred as two genome duplication steps (Figure 5). In the first step, the two tPCK genomes MF1 and MF2 were merged together. Subsequently, a round of genomic reshuffling and gene fractionation resulted in a new diploid. No significant genome dominance is observed between the MF1 and MF2 subgenomes of *B. rapa* now and thus autotetraploidization cannot be excluded as a possible process for the first duplication. However, based on the observation that there were a greater




Figure 5. Two-step polyplodization theory for the WGT event experienced by *Brassica* plants.

number recent small deletions within the exons of MF1 than those of MF2,³⁰ the first duplication is likely to have been an allotetraploidization. In the second step, the third tPCK genome LF was merged with the MFs (MF1 and MF2). A second round of genomic reshuffling and gene fractionation then resulted in the mesohexaploid ancestor of *B. rapa*. In the second step, the 'two' merged genomes (LF and MFs) had different karyotypes, which produced an allopolyploid, and subsequently resulted in biased genome fractionation and the dominant gene expression phenomenon.

B. rapa genes belonging to different gene families or having important biological functions have been systematically analyzed. Based on the genome sequencing and comparative genomic study of *B. rapa*, the evolution of many gene families or categories such as circadian clock genes,⁴¹ resistance genes,⁴² stress response genes,^{43–45} glucosinolate genes,³⁸ anthocyanin biosynthesis genes,⁴⁶ phytohormone-related genes,⁴⁷ certain transcript factor families⁴⁸ and other genes,^{49–54} were accurately determined individually and studied in detail. Moreover, some functionally important genes related to self-incompatibility,^{55,56} male sterility,⁵⁷ flowering regulation,^{58–60} leaf heading⁶¹ or color⁶² have been identified or cloned, and functional studies have been conducted for some of these genes. These follow-up studies in *B. rapa* helped to further elucidate the evolution of specific genes after the WGT event.

CONCLUSIONS AND DISCUSSION

WGT promoted the diversification of *Brassica* plants with respect to both the speciation and expansion of rich morphotypes for each species. First, WGT promoted genomic reshuffling, i.e., rediploidization, to stabilize the genome and the meiosis process. Genomic reshuffling accompanied by chromosome reduction contributed the speciation of diploid *Brassica* plants, such as *B. rapa*, *B. nigra* and *B. oleracea*. Genomic differentiation of the three basic genomes in U's triangle then generated the stable allotetraploid species *B. carinata*, *B. napus* and *B. juncea*. Second, subgenome differentiation, biased gene retention through gene fractionation after WGT, and further multicopy gene subfunctionalization or neofunctionalization promoted the parallel evolution of many different morphotypes in each *Brassica* species. Therefore, WGT with subsequent genomic and gene-level evolution drove *Brassica* speciation and generated an abundance of rich morphotypes of the *Brassica* species.

In the future, additional research should be conducted to investigate the morphotype evolution of *Brassica* plants. Previous studies based on *de novo* genome sequencing determined the genome- and gene-level evolution of only one accession of *B. rapa*. Subsequently, additional *B. rapa* accessions or other *Brassica* species should be extensively studied to address the following aspects regarding the *Brassica* population: (i) the origins and phylogenetic relationship of different morphotypes in *Brassica* species; (ii) the mechanism of the parallel evolution of similar traits that developed independently in different *Brassica* species (e.g., the leafy head in *B. rapa* and *B. oleracea*); and (iii) the genes involved in the development of different morphotypes or genes that regulate the agronomic important traits of *Brassica* crops. This knowledge will increase our understanding of *Brassica* morphotype diversification and ultimately leverage the benefits of genomic studies for the genetic improvement of *Brassica* crops.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

ACKNOWLEDGMENTS

This work was funded by the 973 program (2012CB113900 and 2013CB127000), the 863 program (2012AA100101) and a National Natural Science Foundation of China NSFC grant (31301771). Research was conducted at the Key Laboratory of Biology

and Genetic Improvement of Horticultural Crops, Ministry of Agriculture, P. R. China, and the Sino-Dutch Joint Lab of Horticultural Genomics Technology.

REFERENCES

- Warwick SI, Francis A, Al-Shehbaz IA. Brassicaceae: species checklist and database on CD-Rom. *Plant Syst Evol* 2006; **259**: 249–258.
- Warwick SI, Mummenhoff K, Sauder CA *et al*. Closing the gaps: phylogenetic relationships in the Brassicaceae based on DNA sequence data of nuclear ribosomal ITS region. *Plant Syst Evol* 2010; **285**: 209–232.
- Wang X, Wang H, Wang J *et al*. The genome of the mesopolyploid crop species *Brassica rapa*. *Nat Genet* 2011; **43**: 1035–1039.
- Beilstein MA, Nagalingum NS, Clements MD *et al*. Dated molecular phylogenies indicate a Miocene origin for *Arabidopsis thaliana*. *Proc Natl Acad Sci USA* 2010; **107**: 18724–18728.
- Lukens LN, Quijada PA, Udall J *et al*. Genome redundancy and plasticity within ancient and recent *Brassica* crop species. *Biol J Linnean Soc* 2004; **82**: 665–674.
- Lysak MA, Koch MA, Pecinka A *et al*. Chromosome triplication found across the tribe Brassicaceae. *Genome Res* 2005; **15**: 516–525.
- Arias T, Beilstein MA, Tang M *et al*. Diversification times among *Brassica* (Brassicaceae) crops suggest hybrid formation after 20 million years of divergence. *Am J Bot* 2014; **101**: 86–91.
- Nagaharu U. Genome analysis in *Brassica* with special reference to the experimental formation of *B. napus* and peculiar mode of fertilization. *Jpn J Bot* 1935; **7**: 389–452.
- Morinaga T. Interspecific hybridization in *Brassica* I. The cytology of F1 hybrids of *B. napella* and various other species with 10 chromosomes. *Cytologia* 1929; **1**: 16–27.
- Zhao J, Wang X, Deng B *et al*. Genetic relationships within *Brassica rapa* as inferred from AFLP fingerprints. *Theor Appl Genet* 2005; **110**: 1301–1314.
- Parkin IA, Gulden SM, Sharpe AG *et al*. Segmental structure of the *Brassica napus* genome based on comparative analysis with *Arabidopsis thaliana*. *Genetics* 2005; **171**: 765–781.
- Schranz ME, Lysak MA, Mitchell-Olds T. The ABC's of comparative genomics in the Brassicaceae: building blocks of crucifer genomes. *Trends Plant Sci* 2006; **11**: 535–542.
- Mandakova T, Lysak MA. Chromosomal phylogeny and karyotype evolution in $x = 7$ crucifer species (Brassicaceae). *Plant Cell* 2008; **20**: 2559–2570.
- Lysak MA, Berr A, Pecinka A *et al*. Mechanisms of chromosome number reduction in *Arabidopsis thaliana* and related Brassicaceae species. *Proc Natl Acad Sci USA* 2006; **103**: 5224–5229.
- Koch MA, Kiefer M. Genome evolution among cruciferous plants: a lecture from the comparison of the genetic maps of three diploid species—*Capsella rubella*, *Arabidopsis lyrata* subsp. *petraea*, and *A. thaliana*. *Am J Bot* 2005; **92**: 761–767.
- Hu TT, Pattyn P, Bakker EG *et al*. The *Arabidopsis lyrata* genome sequence and the basis of rapid genome size change. *Nat Genet* 2011; **43**: 476–481.
- Slotte T, Hazzouri KM, Agren JA *et al*. The *Capsella rubella* genome and the genomic consequences of rapid mating system evolution. *Nat Genet* 2013; **45**: 831–835.
- Dassanayake M, Oh DH, Haas JS *et al*. The genome of the extremophile crucifer *Thellungiella parvula*. *Nat Genet* 2011; **43**: 913–918.
- Liu S, Liu Y, Yang X *et al*. The *Brassica oleracea* genome reveals the asymmetrical evolution of polyploid genomes. *Nature Communication* 2014; in press.
- Cheng F, Liu S, Wu J *et al*. BRAD, the genetics and genomics database for *Brassica* plants. *BMC Plant Biol* 2011; **11**: 136.
- Cheng F, Wu J, Fang L *et al*. Biased gene fractionation and dominant gene expression among the subgenomes of *Brassica rapa*. *PLoS ONE* 2012; **7**: e36442.
- Cheng F, Wu J, Fang L *et al*. Syntenic gene analysis between *Brassica rapa* and other Brassicaceae species. *Front Plant Sci* 2012; **3**: 198.
- Panjabi P, Jagannath A, Bisht NC *et al*. Comparative mapping of *Brassica juncea* and *Arabidopsis thaliana* using Intron Polymorphism (IP) markers: homoeologous relationships, diversification and evolution of the A, B and C *Brassica* genomes. *BMC Genomics* 2008; **9**: 113.
- Cheng F, Mandakova T, Wu J *et al*. Deciphering the diploid ancestral genome of the Mesohexaploid *Brassica rapa*. *Plant Cell* 2013; **25**: 1541–1554.
- Initiative AG. Analysis of the genome sequence of the flowering plant *Arabidopsis thaliana*. *Nature* 2000; **408**: 796–815.
- Schnable PS, Ware D, Fulton RS *et al*. The B73 maize genome: complexity, diversity, and dynamics. *Science* 2009; **326**: 1112–1115.
- Schmutz J, Cannon SB, Schlueter J *et al*. Genome sequence of the palaeopolyploid soybean. *Nature* 2010; **463**: 178–183.
- Mun JH, Kwon SJ, Seol YJ *et al*. Sequence and structure of *Brassica rapa* chromosome A3. *Genome Biol* 2010; **11**: R94.
- Trick M, Kwon SJ, Choi SR *et al*. Complexity of genome evolution by segmental rearrangement in *Brassica rapa* revealed by sequence-level analysis. *BMC Genomics* 2009; **10**: 539.

- 30 Tang H, Woodhouse MR, Cheng F *et al*. Altered patterns of fractionation and exon deletions in *Brassica rapa* support a two-step model of paleohexaploidy. *Genetics* 2012; **190**: 1563–1574.
- 31 Schnable JC, Springer NM, Freeling M. Differentiation of the maize subgenomes by genome dominance and both ancient and ongoing gene loss. *Proc Natl Acad Sci USA* 2011; **108**: 4069–4074.
- 32 Schnable JC, Wang X, Pires JC *et al*. Escape from preferential retention following repeated whole genome duplications in plants. *Front Plant Sci* 2012; **3**: 94.
- 33 Woodhouse MR, Schnable JC, Pedersen BS *et al*. Following tetraploidy in maize, a short deletion mechanism removed genes preferentially from one of the two homologs. *PLoS Biol* 2010; **8**: e1000409.
- 34 Woodhouse MR, Cheng F, Pires JC, Lisch D, Freeling M, Wang X. Origin, inheritance, and gene regulatory consequences of genome dominance in polyploids. *Proc Natl Acad Sci USA* 2014 Apr 8; **111**(14): 5283–5288.
- 35 Hollister JD, Gaut BS. Epigenetic silencing of transposable elements: a trade-off between reduced transposition and deleterious effects on neighboring gene expression. *Genome Res* 2009; **19**: 1419–1428.
- 36 Hollister JD, Smith LM, Guo YL *et al*. Transposable elements and small RNAs contribute to gene expression divergence between *Arabidopsis thaliana* and *Arabidopsis lyrata*. *Proc Natl Acad Sci USA* 2011; **108**: 2322–2327.
- 37 Bekaert M, Edger PP, Hudson CM, Pires JC *et al*. Metabolic and evolutionary costs of herbivory defense: systems biology of glucosinolate synthesis. *New Phytol* 2012; **196**: 596–605.
- 38 Wang H, Wu J, Sun S *et al*. Glucosinolate biosynthetic genes in *Brassica rapa*. *Gene* 2011; **487**: 135–142.
- 39 Santner A, Estelle M. Recent advances and emerging trends in plant hormone signalling. *Nature* 2009; **459**: 1071–1078.
- 40 Gazzarrini S, McCourt P. Cross-talk in plant hormone signalling: what *Arabidopsis* mutants are telling us. *Ann Bot* 2003; **91**: 605–612.
- 41 Lou P, Wu J, Cheng F *et al*. Preferential retention of circadian clock genes during diploidization following whole genome triplication in *Brassica rapa*. *Plant Cell* 2011; **24**: 2415–2426.
- 42 Wu P, Shao ZQ, Wu XZ *et al*. Loss/retention and evolution of NBS-encoding genes upon whole genome triplication of *Brassica rapa*. *Gene* 2014; **540**: 54–61.
- 43 Song X, Li Y, Hou X. Genome-wide analysis of the AP2/ERF transcription factor superfamily in Chinese cabbage (*Brassica rapa* ssp. *pekinensis*). *BMC Genomics* 2013; **14**: 573.
- 44 Lee SC, Lim MH, Yu JG *et al*. Genome-wide characterization of the CBF/DREB1 gene family in *Brassica rapa*. *Plant Physiol Biochem* 2012; **61**: 142–152.
- 45 Ma J, Wang F, Li M *et al*. Genome wide analysis of the NAC transcription factor family in Chinese cabbage to elucidate responses to temperature stress. *Sci Hort* 2014; **165**: 82–90.
- 46 Zhou B, Wang Y, Zhan Y *et al*. Chalcone synthase family genes have redundant roles in anthocyanin biosynthesis and in response to blue/UV-A light in turnip (*Brassica rapa*; Brassicaceae). *Am J Bot* 2013; **100**: 2458–2467.
- 47 Liu Z, Kong L, Zhang M *et al*. Genome-wide identification, phylogeny, evolution and expression patterns of AP2/ERF genes and cytokinin response factors in *Brassica rapa* ssp. *pekinensis*. *PLoS ONE* 2014; **8**: e83444.
- 48 Peng FY, Weselake RJ. Genome-wide identification and analysis of the B3 superfamily of transcription factors in Brassicaceae and major crop plants. *Theor Appl Genet* 2013; **126**: 1305–1319.
- 49 Song XM, Liu TK, Duan WK *et al*. Genome-wide analysis of the GRAS gene family in Chinese cabbage (*Brassica rapa* ssp. *pekinensis*). *Genomics* 2013; **103**: 135–146.
- 50 Sampath P, Lee SC, Lee J *et al*. Characterization of a new high copy Stowaway family MITE, BRAMI-1 in *Brassica* genome. *BMC Plant Biol* 2013; **13**: 56.
- 51 Liu Z, Zhang M, Kong L *et al*. Genome-wide identification, phylogeny, duplication, and expression analyses of two-component system genes in Chinese cabbage (*Brassica rapa* ssp. *pekinensis*). *DNA Res*; e-pub ahead of print 27 February 2014; doi:10.1093/dnares/dsu004.
- 52 Song X, Liu G, Duan W *et al*. Genome-wide identification, classification and expression analysis of the heat shock transcription factor family in Chinese cabbage. *Mol Genet Genomics* e-pub ahead of print 8 March 2014; doi:10.1007/s00438-014-0833-5.
- 53 Hu L, Li S, Gao W. Expression, divergence and evolution of the caleosin gene family in *Brassica rapa*. *Arch Biol Sci* 2013; **65**: 863–876.
- 54 Hu L, Yin W, Chen Y *et al*. Functional divergence and evolutionary dynamics of the putative AAAP gene family in *Brassica rapa*. *Plant Mol Biol Rep* 2014; **32**: 517–530.
- 55 Indriolo E, Tharmapalan P, Wright SI *et al*. The ARC1 E3 ligase gene is frequently deleted in self-compatible Brassicaceae species and has a conserved role in *Arabidopsis lyrata* self-pollen rejection. *Plant Cell* 2012; **24**: 4607–4620.
- 56 Lao X, Suwabe K, Niikura S *et al*. Physiological and genetic analysis of CO₂-induced breakdown of self-incompatibility in *Brassica rapa*. *J Exp Bot* 2014; **65**: 939–951.
- 57 Xu X, Sun X, Zhang J *et al*. Identification of candidate genes associated with male sterility in CMS7311 of heading Chinese cabbage (*Brassica campestris* L. ssp. *pekinensis*). *Acta Physiol Plant* 2013; **35**: 3265–3270.
- 58 Immink RG, Pose D, Ferrario S *et al*. Characterization of SOC1's central role in flowering by the identification of its upstream and downstream regulators. *Plant Physiol* 2012; **160**: 433–449.
- 59 Xiao D, Zhao JJ, Hou XL *et al*. The *Brassica rapa* FLC homologue FLC2 is a key regulator of flowering time, identified through transcriptional co-expression networks. *J Exp Bot* 2013; **64**: 4503–4516.
- 60 Wu J, Wei K, Cheng F *et al*. A naturally occurring InDel variation in BraA.FLC.b (BrFLC2) associated with flowering time variation in *Brassica rapa*. *BMC Plant Biol* 2012; **12**: 151.
- 61 Liu Z, Jia L, Wang H *et al*. HYL1 regulates the balance between adaxial and abaxial identity for leaf flattening via miRNA-mediated pathways. *J Exp Bot* 2011; **62**: 4367–4381.
- 62 Lee S, Lee SC, Byun DH *et al*. Association of molecular markers derived from the BrCRISTO1 gene with polycopene-enriched orange-colored leaves in *Brassica rapa*. *Theor Appl Genet* 2014; **127**: 179–191.



This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivs 3.0 Unported License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by-nc-nd/3.0/>

Supplemental Information for this article can be found on the *Horticulture Research* website (<http://www.nature.com/hortres>).