



Review

Bibliometric Analysis of the Influencing Factors, Derivation, and Application of Heavy Metal Thresholds in Soil

Zhaolin Du ¹, Dasong Lin ¹, Haifeng Li ², Yang Li ¹, Hongan Chen ¹, Weiqiang Dou ³, Li Qin ^{1,*}  and Yi An ^{1,*}

¹ Agro-Environmental Protection Institute, Ministry of Agriculture and Rural Affairs, Tianjin 300191, China; carlcarl1988@163.com (Z.D.); lindasong608@126.com (D.L.); liy0128@163.com (Y.L.); 18702229081@163.com (H.C.)

² Beijing Municipal Key Laboratory of Agriculture Environment Monitoring, Beijing 100097, China; haifli@126.com

³ College of Land Science and Technology, China Agricultural University, Beijing 100193, China; dwq18513262537@126.com

* Correspondence: ql-tj@163.com (L.Q.); simon8601@126.com (Y.A.)

Abstract: The study of threshold levels of heavy metals in soil is essential for the assessment and management of soil environmental quality. This study reviewed the influencing factors, the derivation, and application aspects of heavy metals' threshold values comprehensively by a combination of bibliometric analysis and scientific knowledge mapping. A total of 1106 related studies were comprehensively extracted from the Web of Science database during the period from 2001 to 2020. The results showed that the publication output has been growing strongly. An analysis on the subject, journal, country, and institution was carried out to demonstrate the development and evolution of this research branch during the two decades. According to high-frequency keywords analysis, external factors (e.g., soil physicochemical properties) and internal factors (e.g., crop genotype) can affect heavy metal threshold values in the soil–crop system. The current methods mainly include the Point model (e.g., evaluation factor method), the Probability model (e.g., species sensitivity distribution method), and the Empirical model (e.g., ecological environment effect method). A threshold study can be applicable to the risk assessment for soil heavy metal contamination in order to determinate the soil pollution degree and its spatial and temporal distribution characteristics. Moreover, challenges and prospects of the study of heavy metal threshold values are proposed, indicating that research should focus on the relationships between human health risks and the established threshold values of heavy metals in the soil, long-term field trials and bioavailability of heavy metals for the derivation of the thresholds, and the establishment of more scientific and rational soil environmental benchmarks.

Keywords: soil; crop; heavy metal; threshold; risk assessment



Citation: Du, Z.; Lin, D.; Li, H.; Li, Y.; Chen, H.; Dou, W.; Qin, L.; An, Y. Bibliometric Analysis of the Influencing Factors, Derivation, and Application of Heavy Metal Thresholds in Soil. *Int. J. Environ. Res. Public Health* **2022**, *19*, 6561. <https://doi.org/10.3390/ijerph19116561>

Academic Editors: Jinling Liu and Jun Zhou

Received: 19 April 2022

Accepted: 25 May 2022

Published: 27 May 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Soil contamination has attracted widespread attention globally [1–3]. With the continuous advancement of urbanization and industrialization, heavy metals are constantly emitted into the terrestrial environment and accumulate in the soil, posing great threats to the quality of soil, groundwater, and crops [4–6]. Once heavy metals were in the food chain, the health of humans and other organisms could be affected [7–10]. Owing to the potential toxicity and difficult biodegradation of heavy metals, their content in soil has become of great concern [9,11,12].

Threshold levels of heavy metals in soil are the maximum heavy metal concentrations that the soil can possess without destroying ecosystems [13]. The derivation of heavy metal thresholds in the soil is essential for effectively protecting soil quality, guaranteeing crop yields and quality, and ensuring human health. It also provides a scientific basis and an important foundation for ecological risk assessment and the formulation of soil environmental quality standards. Hence, it plays an indispensable role in soil management.

The academic community has conducted much research on heavy metal thresholds in the soil. Some studies focused on various types of heavy metals, including certain common potentially toxic elements (e.g., cadmium (Cd)) and emerging “high-tech” critical elements (e.g., lanthanides) [13]. Some studies focused on the migration and transformation of heavy metals in soil–crop systems rather than on a simple soil system. Previous research mainly focused on protection for a sustainable use of soil [14], and current research has begun to examine the quality and safety of agricultural products as well as human health risks [15,16]. Thus, research on the thresholds of heavy metals has been extensive and diverse. However, most previous research was limited to specific situations for soil risk assessment or to qualitative analyses of small amounts of literature data. A systematic review based on big data analysis to demonstrate the development of heavy metal threshold studies is still lacking.

Bibliometric analysis is a method to summarize historical achievements and development trends comprehensively, systematically, and objectively. It can also explore possible future research directions of the research field with the help of mathematics and statistics, computer analysis and information visualization technology, combined with the analysis of the literature [17,18]. Scientific knowledge mapping, which is a scientific and effective literature research tool for large amounts of literature data to study a certain knowledge domain, is becoming increasingly popular [19]. It can vividly reveal the dynamic development of a certain research field by visualizing its development, research hotspots, and frontier knowledge [20,21].

In this study, a literature review was combined with a bibliometric analysis and scientific knowledge mapping to analyze studies on heavy metal threshold levels in soil published in the last two decades. The influence factors, derivation, and application aspects of the threshold levels were summarized through high-frequency keywords analysis. Moreover, the challenges and promising research frontiers of research on heavy metal thresholds are discussed.

2. Materials and Methods

2.1. Data Collection

Publications were obtained from the online version of the database of Science Citation Index-Expanded (SCI-E) of the Web of Science (WoS) core collection. Records retrieval started with a topic search for “soil,” “heavy metal,” and “threshold” in the publication year range from 2001 to 2020. Some repeated or irrelevant records were removed after analyzing each record. Finally, a total of 1106 records which met the study requirements were obtained.

2.2. Methodology of Bibliometric Analysis

CiteSpace, regarded as a useful, efficient scientific visualization tool, was used to analyze the related literature and construct scientific knowledge maps [22]. It was jointly developed by Dr. Chaomei Chen and the WISE Laboratory of Dalian University of Technology and is provided to researchers free of charge. CiteSpace V Software was used in the study for visualizing and analyzing trends and patterns in the scientific literature. Its primary source of input data was ISI WoS. It could vividly provide key information which included the publication output and journal information, country and institution cooperation, subject and research hotspot evolution through analysis functions of co-citation, collaboration and co-occurrence, respectively.

3. Results

3.1. Publication Output Analysis

As shown in Supplementary material Figure S1, the annual number of studies increased steadily from 13 in 2001 to 140 in 2020, demonstrating an almost nine-fold increase over 20 years. The predominant publication type in this field was Article, accounting

for 97.92% of the studies (1083 records) [23], followed by other types such as Review (22 records) and Letter (1 record) [23,24].

3.2. Subject Analysis

Publications in the field of research of heavy metal thresholds in soil were distributed in 55 unique subject categories. The most productive discipline in this field was Environmental Sciences and Ecology, accounting for 67.18% of the studies (743 records), followed by Agriculture, Engineering, Water Resources, Toxicology, Geology, Public and Environmental Occupational Health, Chemistry, Plant Sciences, Science Technology, Other Topics, Geochemistry Geophysics, and Biodiversity Conservation (Table S1).

Figure 1 shows that Environmental Science and Ecology, Environmental Science, Agriculture, Soil Science, Water Resources, Agronomy, Plant Science, Chemistry and other disciplines are basic disciplines in the research on heavy metal thresholds in soil. They have laid a solid theoretical foundation for research and development in this field. Since 2005, several disciplines including Engineering, Environmental Engineering, Geology and Geosciences, gradually became the supporting disciplines of research in this field, indicating that the research was no longer limited to theoretical research but also began to consider practical applications [25]. In 2011, the subject of “Public, Environmental and Occupational Health” appeared, which revealed that many studies began to attach importance to the relationships between the thresholds of heavy metals in soil and human health risks [26]. After 2015, certain disciplines such as “Biodiversity Conservation” and “Green and Sustainable Science and Technology” gradually became important research disciplines in this research field, demonstrating that the derivation of heavy metal thresholds in soil no longer only considered the effects on a single species but had acquired a macro view, considering the effects on whole ecosystems [27].




Figure 1. Evolution mapping of disciplines development during 2001–2020. Note: Each node represents a discipline, and the larger the node indicates the higher the number of articles related to this discipline. A pivotal point with high betweenness centrality is highlighted with a purple ring, showing this discipline has a great influence in the cooperation between disciplines.

3.3. Journal, Country, and Institution Analysis

Science of the Total Environment was the journal with the highest number of citations, followed by *Environmental Pollution*, *Chemosphere*, and *Journal of Hazardous Materials* (Figure S2). China, the United States, and Italy were the three most productive countries. Collaborative relationships between countries were highlighted by colored lines, and a close cooperation meant that research on heavy metal thresholds in soil was still one of the most important

the Point model (e.g., evaluation factor method), the Probability model (e.g., SSD method), and the Empirical model (e.g., ecological environment effect method). The advantages and disadvantages of each method are shown in Table 1.

Table 1. Analysis on the advantages, disadvantages, and uncertainty of the methods for determining the threshold levels of heavy metals in agricultural land.

Model	Representative Method	Advantages and Disadvantages	References
Point model	Evaluation factor method	Advantages: simple and easy to operate, taking account of species sensitivity, independent of any theoretical model; quantitative result description Disadvantages: selection of evaluation factors depending on experiences and national policies, single type and quantity of sensitive species without representativity	[57,58]
Probability model	Species sensitivity distribution (SSD)	Advantages: taking account of differences in species sensitivity, soil physical and chemical properties, bioavailability and sources of pollutants, and different pollution risk levels; quantitative result description Disadvantages: rarely taking account of the interrelationship of food chains between species, unable to provide potential recovery information for the environment	[58–60]
Empirical model	Ecological environment effect method	Advantages: taking account of the effects of different soil physical and chemical properties; quantitative result description Disadvantages: rarely taking account of species sensitivity	[59,61–63]

4.3. Application of Heavy Metal Thresholds in Soil

Based on threshold studies, risk assessment can determine a soil pollution degree and spatial and temporal distribution characteristics [64]. It could effectively allow locating areas containing toxic elements through these risk assessment procedures: setting different threshold levels for common heavy metals in soil (taking China as an example, four threshold levels of Cd were established for major edible agricultural products' safety (e.g., rice) in consideration of soil physical and chemical properties, i.e., 0.3 mg/kg for $\text{pH} \leq 5.5$, 0.4 mg/kg for $5.5 < \text{pH} \leq 6.5$, 0.6 mg/kg for $6.5 < \text{pH} \leq 7.5$, and 0.8 mg/kg for $\text{pH} > 7.5$), conducting more targeted investigations on heavy metal pollution characteristics, selecting appropriate methods, and defining a soil pollution classification. Hence, this analysis could play an important role in setting soil environmental benchmarks and guiding follow-up risk control measures. The common soil heavy metal pollution risk assessment methods can be divided into two categories: index and model methods [65,66]. Index methods mainly included the single pollution index method [67], the Nemerow index method [65], the pollution load index method, the environmental risk index method, and the geoaccumulation index method [68]. Model methods mainly include the enrichment factor method and the fuzzy comprehensive evaluation method. Another method is the GIS-based geostatistical legal evaluation method [69]. Index methods are relatively easy to operate, and model methods are applicable to solving complicated problems. Research mainly uses certain indicators such as enrichment factor, geoaccumulation index, Nemerow index, and average enrichment ratio to characterize the pollution degree of an area [53,70]. The content of heavy metals in crops grown on contaminated soil could also be experimentally and quantitatively evaluated with the help of the bioaccumulation factor to identify heavily polluted areas [52,55]. However, these methods could hardly determine quantitative sources of potentially toxic elements in soil management.

4.4. Challenges and Prospects

Frequency bursts can indicate that the scientific community has paid or is paying special attention to the contribution of specific factors, and burst detection can identify burst keywords as indicators of emerging trends [71]. As shown in Figure 4, the discussion of the factors influencing heavy metal thresholds in soil will continue in the future, especially concerning the three aspects of “health risk assessment”, “ecological risk assessment”, and “apportionment and identification of pollution sources” [72]. The topics of “health risk” and “health risk assessment” have emerged as research frontiers, with the strongest strength of frequency bursts corresponding to 10.4624 and 6.8364, respectively, indicating that this research will be of interest to scientists in the future. Performing a comprehensive ecological risk assessment plays a remarkable role in formulating scientific, reasonable heavy metal thresholds in soil, but specific adaptation and accuracy of ecological risk assessment remain a challenge. Methods and tools for identifying and analyzing sources of heavy metal pollution still need further research. The current highly complex and changeable environmental situation and the considerable effects on soil pollution of human activities make heavy metal pollution sources difficult to identify. This may also be one of the reasons why the terms “source apportionment” and “source identification” have become burst keywords.














Top 13 Keywords with the Strongest Citation Bursts				
Keywords	Strength	Begin	End	2015–2020
health risk	10.4624	2018	2020	
health risk assessment	6.8364	2018	2020	
standards	6.6933	2018	2020	
source apportionment	6.5342	2018	2020	
ecological risk	5.9887	2017	2020	
threshold	5.2999	2018	2020	
food crop	5.275	2018	2020	
bioavailability	4.2179	2017	2020	
street dust	4.0823	2017	2020	
potentially toxic elements	4.0205	2018	2020	
source identification	3.881	2018	2020	
uncertainty	3.869	2018	2020	
compost	3.804	2018	2020	

Figure 4. Top 13 keywords with the strongest citation bursts. Note: the duration of the citation burst of a keyword is represented by the red bar.

4.4.1. Relationships between Human Health Risks and Heavy Metal Thresholds in Soil

Heavy metals in soil can enter the food chain after enrichment in crops, which can pose a severe threat to human health. Moreover, human beings accumulate heavy metals also through inhalation of particles in soil and by dermal contact [15]. Under the combined effects of various exposure pathways, the derivation of heavy metal thresholds in soil requires multidisciplinary joint research. Therefore, it is worthy to explore how to establish heavy metal thresholds in soil, ensuring that these contaminants do not cause risks to human health in the future.

4.4.2. Derivation of Heavy Metal Thresholds Based on Long-Term Field Trials

The current studies mostly use short-term pot experiments or field trials to understand quantitative relationships regarding heavy metal concentrations in soil–crop systems. Large spatiotemporal differences in the experimental conditions between pot experiments and field trials often lead to inconsistent experimental results. Consequently, heavy metal thresholds in soil often failed to meet the requirements of soil pollution risk assessment, with relatively poor reliability. Determining quantitative relationships and studying the

dynamic changes of heavy metal thresholds in long-term field trials may become future research topics.

4.4.3. Derivation of Heavy Metal Thresholds Based on Their Bioavailability in Soil

At present, methods for determining heavy metal thresholds in farmland soil mainly consider the total amount of heavy metals as evaluation standards. However, the total amount of heavy metals cannot fully represent the actual toxic effects on crops and humans. The pH, organic matter content, and other factors should be considered for the derivation of heavy metal thresholds, and this requires research innovations. Therefore, the construction of new derivation methods for thresholds based on the bioavailable state of heavy metals in the soil may be an important development direction.

4.4.4. Establishment of Soil Environmental Benchmarks

Soil environmental benchmarks based on heavy metal thresholds can scientifically reflect the risk of heavy metal contamination in soil on related receptors. They can guide the formulation of soil environmental quality standards and subsequent pollution risk assessment. However, the current soil environmental benchmarks in many countries are hard to meet; therefore, there is a need to scientifically formulate soil environmental quality standards. For example, heavy metal threshold values based on different pH levels and land-use types were established in the China's current farmland soil environmental quality standard (GB15618-2018). This standard can be beneficial to determine different farmland soil pollution levels preliminarily. However, this standard did not take into account other factors influencing heavy metal thresholds, such as soil types, soil physical and chemical properties (such as redox potential, soil organic matter, and ionic strength), and crop varieties. As a result, it may cause risk assessment results to deviate from reality, which can lead to "overprotection" or to "inadequate protection" in relation to soil environmental quality safety. Therefore, establishing more scientific and rational soil environmental benchmarks regarding heavy metal thresholds in soil may be important research direction.

5. Conclusions

Based on a big data analysis of 1106 studies, the following results on heavy metal thresholds in soils were obtained: (1) Environmental Sciences and Ecology, Agriculture, and Engineering were the three main disciplines concerned with heavy metal threshold research. (2) Journals, countries and institutions with the highest number of articles published were identified, and the development and evolution of these studies during the two decades were demonstrated. (3) Systematical analysis of external influencing factors (such as heavy metal levels in soil, soil pH, redox potential, soil organic matter, and ionic strength) and internal factors (e.g., crop genotype), of derivation methods (such as point model, probability model, and empirical model), and of application aspects of heavy metal threshold studies from 2001 to 2020 was carried out. (4) Challenges and future prospects of heavy metal threshold studies were proposed, indicating that research should focus on the relationships between human health risks and the established thresholds of heavy metals in soil, the derivation of thresholds based on the bioavailability of heavy metals and long-term field trials, and the establishment of more scientific and rational soil environmental benchmarks.

Supplementary Materials: The following supporting information can be downloaded at: <https://www.mdpi.com/article/10.3390/ijerph19116561/s1>, Figure S1: Annual number of various types of publications (2001–2020). Note: the bar chart shows the annual number of various types of publications, and the line chart shows cumulative frequency; Figure S2: Most cited journals. Note: each node represents a journal, and a brighter node indicates this journal is highly cited; Figure S3: Development and evolution of the studies in the most productive countries. Note: each node represents a country, and a larger node indicates a higher number of articles published by this country. The pivotal point with high betweenness centrality is highlighted with a purple ring, showing that country has a great influence in the cooperation relationships between countries; Table S1: Twelve

most productive disciplines; Table S2: Six most productive journals; Table S3: Fifteen most productive institutions.

Author Contributions: Conceptualization, Z.D.; methodology, Z.D.; formal analysis, Y.A.; investigation, L.Q.; resources, H.L.; data curation, W.D. and D.L.; writing—original draft preparation, Z.D., Y.L. and H.C.; writing—review and editing, Z.D., L.Q., Y.A. and H.L. All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by the Open Project Foundation in Beijing Research Center for Agricultural Standards and Testing (ZBZX202104) and the National Natural Science Foundation of China (No. 42107081).

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: Not applicable.

Conflicts of Interest: The authors declare no conflict of interest. The funders had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript, or in the decision to publish the results.

References

- Toth, G.; Hermann, T.; Szatmari, G.; Pasztor, L. Maps of heavy metals in the soils of the European Union and proposed priority areas for detailed assessment. *Sci. Total Environ.* **2016**, *565*, 1054–1062. [[CrossRef](#)] [[PubMed](#)]
- Hu, B.; Shao, S.; Ni, H.; Fu, Z.; Shi, Z. Current status, spatial features, health risks, and potential driving factors of soil heavy metal pollution in China at province level. *Environ. Pollut.* **2020**, *266 Pt 3*, 114961. [[CrossRef](#)] [[PubMed](#)]
- Zhang, H.; Zhang, F.; Song, J.; Tan, M.L.; Kung, H.T.; Johnson, V.C. Pollutant source, ecological and human health risks assessment of heavy metals in soils from coal mining areas in Xinjiang, China. *Environ. Eng. Res.* **2021**, *202*, 111702. [[CrossRef](#)] [[PubMed](#)]
- Varol, M.; Gunduz, K.; Sunbul, M.R. Pollution status, potential sources and health risk assessment of arsenic and trace metals in agricultural soils: A case study in Malatya province, Turkey. *Environ. Eng. Res.* **2021**, *202*, 111806. [[CrossRef](#)]
- Qu, J.; Wei, S.; Liu, Y.; Zhang, X.; Zhang, Y. Effective lead passivation in soil by bone char/CMC-stabilized FeS composite loading with phosphate-solubilizing bacteria. *J. Hazard. Mater.* **2021**, *423*, 127043. [[CrossRef](#)]
- Negahban, S.; Mokarram, M.; Pourghasemi, H.R.; Zhang, H. Ecological risk potential assessment of heavy metal contaminated soils in Ophiolitic formations. *Environ. Eng. Res.* **2021**, *192*, 110305. [[CrossRef](#)]
- Huang, J.; Wang, Y.; Xu, L.; Xu, N.; Fan, W. Cumulative risk assessment for an urban agglomeration based on a relative risk model and multi-media fugacity model: A case study in Guangdong province, China. *Environ. Impact Assess. Rev.* **2021**, *91*, 106637. [[CrossRef](#)]
- Taati, A.; Salehi, M.H.; Mohammadi, J.; Mohajer, R.; Diez, S. Pollution assessment and spatial distribution of trace elements in soils of Arak industrial area, Iran: Implications for human health. *Environ. Eng. Res.* **2020**, *187*, 109577. [[CrossRef](#)]
- Du, Z.L.; Chen, H.A.; Guo, X.Y.; Qin, L.; Lin, D.S.; Huo, L.L.; Yao, Y.P.; Zhang, Z.H. Mechanism and industrial application feasibility analysis on microwave-assisted rapid synthesis of amino-carboxyl functionalized cellulose for enhanced heavy metal removal. *Chemosphere* **2021**, *268*, 11. [[CrossRef](#)]
- Aschner, M.; Paoliello, M.M.B.; Tsatsakis, A.; Bowman, A.B.; Dorea, J.G.; Hartung, T.; Domingo, J.L.; Barbosa, F. Social injustice in environmental health: A call for fortitude. *Environ. Eng. Res.* **2021**, *194*, 110675. [[CrossRef](#)]
- Qu, J.; Meng, Q.; Lin, X.; Han, W.; Jiang, Q.; Wang, L.; Hu, Q.; Zhang, L.; Zhang, Y. Microwave-assisted synthesis of beta-cyclodextrin functionalized celluloses for enhanced removal of Pb(II) from water: Adsorptive performance and mechanism exploration. *Sci. Total Environ.* **2021**, *752*, 141854. [[CrossRef](#)] [[PubMed](#)]
- Qu, J.H.; Liu, Y.; Cheng, L.; Jiang, Z.; Zhang, G.S.; Deng, F.X.; Wang, L.; Han, W.; Zhang, Y. Green synthesis of hydrophilic activated carbon supported sulfide nZVI for enhanced Pb(II) scavenging from water: Characterization, kinetics, isotherms and mechanisms. *J. Hazard. Mater.* **2021**, *403*, 11. [[CrossRef](#)] [[PubMed](#)]
- Reimann, C.; Fabian, K.; Birke, M.; Filzmoser, P.; Demetriades, A.; Negrel, P.; Oorts, K.; Matschullat, J.; de Caritat, P.; Team, G.P. GEMAS: Establishing geochemical background and threshold for 53 chemical elements in European agricultural soil. *Appl. Geochem.* **2018**, *88*, 302–318. [[CrossRef](#)]
- Keller, A.; Schulin, R. Modelling regional-scale mass balances of phosphorus, cadmium and zinc fluxes on arable and dairy farms. *Eur. J. Agron* **2003**, *20*, 181–198. [[CrossRef](#)]
- Hu, W.Y.; Huang, B.; Tian, K.; Holm, P.E.; Zhang, Y.X. Heavy metals in intensive greenhouse vegetable production systems along Yellow Sea of China: Levels, transfer and health risk. *Chemosphere* **2017**, *167*, 82–90. [[CrossRef](#)] [[PubMed](#)]
- Xiao, R.; Guo, D.; Ali, A.; Mi, S.S.; Liu, T.; Ren, C.Y.; Li, R.H.; Zhang, Z.Q. Accumulation, ecological-health risks assessment, and source apportionment of heavy metals in paddy soils: A case study in Hanzhong, Shaanxi, China. *Environ. Pollut.* **2019**, *248*, 349–357. [[CrossRef](#)]

17. He, K.; Zhang, J.; Zeng, Y. Knowledge domain and emerging trends of agricultural waste management in the field of social science: A scientometric review. *Sci. Total Environ.* **2019**, *670*, 236–244. [[CrossRef](#)]
18. Bezak, N.; Mikos, M.; Borrelli, P.; Alewell, C.; Alvarez, P.; Anache, J.A.A.; Baartman, J.; Ballabio, C.; Biddoccu, M.; Cerda, A.; et al. Soil erosion modelling: A bibliometric analysis. *Environ. Eng. Res.* **2021**, *197*, 111087. [[CrossRef](#)]
19. Li, X.; Li, Y.; Li, G. A scientometric review of the research on the impacts of climate change on water quality during 1998–2018. *Environ. Sci. Pollut. Res.* **2020**, *27*, 14322–14341. [[CrossRef](#)]
20. Li, X.; Nan, R. A bibliometric analysis of eutrophication literatures: An expanding and shifting focus. *Environ. Sci. Pollut. Res.* **2017**, *24*, 17103–17115. [[CrossRef](#)]
21. Li, C.; Wu, K.; Wu, J. A bibliometric analysis of research on haze during 2000–2016. *Environ. Sci. Pollut. Res.* **2017**, *24*, 24733–24742. [[CrossRef](#)] [[PubMed](#)]
22. Ouyang, W.; Wang, Y.D.; Lin, C.Y.; He, M.C.; Hao, F.H.; Liu, H.B.; Zhu, W.H. Heavy metal loss from agricultural watershed to aquatic system: A scientometrics review. *Sci. Total Environ.* **2018**, *637*, 208–220. [[CrossRef](#)] [[PubMed](#)]
23. Galhardi, J.A.; Mello, J.; Wilkinson, K.J. Bioaccumulation of potentially toxic elements from the soils surrounding a legacy uranium mine in Brazil. *Chemosphere* **2020**, *261*, 127679. [[CrossRef](#)] [[PubMed](#)]
24. Grzebisz, W.; Diatta, J.B.; Ciesla, L. Spatial Distribution of Copper in Arable Soils and in Non-Consumable Crops (Flax, Oil-Seed Rape) Cultivated Near a Copper Smelter. *Pol. J. Environ. Stud.* **2001**, *10*, 269–274.
25. Liu, X.; Wu, J.; Xu, J. Characterizing the risk assessment of heavy metals and sampling uncertainty analysis in paddy field by geostatistics and GIS. *Environ. Pollut.* **2006**, *141*, 257–264. [[CrossRef](#)]
26. Vilavert, L.; Sisteré, C.; Schuhmacher, M.; Nadal, M.; Domingo, J.L. Environmental Concentrations of Metals in the Catalan Stretch of the Ebro River, Spain: Assessment of Temporal Trends. *Biol. Trace Elem. Res.* **2015**, *163*, 48–57. [[CrossRef](#)]
27. Kumar, V.; Sharma, A.; Pandita, S.; Bhardwaj, R.; Thukral, A.K.; Cerda, A. A review of ecological risk assessment and associated health risks with heavy metals in sediment from India. *Int. J. Sediment Res.* **2020**, *35*, 516–526. [[CrossRef](#)]
28. Cao, H.C.; Luan, Z.Q.; Wang, J.D.; Zhang, X.L. Potential ecological risk of cadmium, lead and arsenic in agricultural black soil in Jilin Province, China. *Stoch. Environ. Res. Risk Assess.* **2009**, *23*, 57–64. [[CrossRef](#)]
29. Li, P.Y.; Qian, H.; Howard, K.W.F.; Wu, J.H.; Lyu, X.S. Anthropogenic pollution and variability of manganese in alluvial sediments of the Yellow River, Ningxia, northwest China. *Environ. Monit. Assess.* **2014**, *186*, 1385–1398. [[CrossRef](#)]
30. Gillet, S.; Ponge, J.F. Humus forms and metal pollution in soil. *Eur. J. Soil Sci.* **2002**, *53*, 529–539. [[CrossRef](#)]
31. Queiroz, H.M.; Nóbrega, G.; Ferreira, T.O.; Almeida, L.S.; Romero, T.B.; Santaella, S.T.; Bernardino, A.F.; Otero, X.L. The Samarco mine tailing disaster: A possible time-bomb for heavy metals contamination? *Sci. Total Environ.* **2018**, *637–638*, 498–506. [[CrossRef](#)] [[PubMed](#)]
32. Jain, P.; Kim, H.; Townsend, T.G. Heavy metal content in soil reclaimed from a municipal solid waste landfill. *Waste Manag.* **2005**, *25*, 25–35. [[CrossRef](#)] [[PubMed](#)]
33. Bolan, N.S.; Adriano, D.C.; Mahimairaja, S. Distribution and bioavailability of trace elements in livestock and poultry manure by-products. *Crit. Rev. Environ. Sci. Technol.* **2004**, *34*, 291–338. [[CrossRef](#)]
34. Li, Y.; Gou, X.; Wang, G.; Zhang, Q.; Su, Q.; Xiao, G.J. Heavy metal contamination and source in and agricultural soil in central Gansu Province, China. *J. Environ. Sci.* **2008**, *20*, 607–612. [[CrossRef](#)]
35. Hani, A.; Pazira, E. Heavy metals assessment and identification of their sources in agricultural soils of Southern Tehran, Iran. *Environ. Monit. Assess.* **2011**, *176*, 677–691. [[CrossRef](#)] [[PubMed](#)]
36. Bigalke, M.; Ulrich, A.; Rehmus, A.; Keller, A. Accumulation of cadmium and uranium in arable soils in Switzerland. *Environ. Pollut.* **2017**, *221*, 85–93. [[CrossRef](#)]
37. Pedrot, M.; Dia, A.; Davranche, M.; Bouhnik-Le Coz, M.; Henin, O.; Gruau, G. Insights into colloid-mediated trace element release at the soil/water interface. *J. Colloid Interface Sci.* **2008**, *325*, 187–197. [[CrossRef](#)]
38. Relic, D.; Dordevic, R.; Popovic, R.; Jadranin, R.; Polic, R. Fractionation and potential mobility of trace metals in Danube alluvial aquifer within an industrialized zone. *Environ. Monit. Assess.* **2010**, *171*, 229–248. [[CrossRef](#)]
39. Ispas, G.; Roba, C.; Blc, R.; Coad, M.; Gligor, D. The quality of water and soil in Hane mining area, Alba Country, Romania. *Carpathian J. Earth Environ. Sci.* **2018**, *13*, 505–514. [[CrossRef](#)]
40. Rivera-Hernández, J.R.; Green-Ruiz, C. Strong acid mixture and sequential geochemical arsenic extractions in surface sediments from the Santa Maria La Reforma Coastal Lagoon, Mexico: A bioavailability assessment. *Arch. Environ. Contam. Toxicol.* **2016**, *70*, 348–360. [[CrossRef](#)]
41. Hao, L.; Chen, L.P.; Ai, Y.W.; Xia, Y.; Yu, Y.H.; Zuo, Y.B.; Fu, G.Y. Heavy metal contamination in soil alongside mountain railway in Sichuan, China. *Environ. Monit. Assess.* **2009**, *152*, 25–33.
42. Song, W.E.; Chen, S.B.; Liu, J.F.; Chen, L.; Song, N.N.; Ning, L.I.; Liu, B. Variation of Cd concentration in various rice cultivars and derivation of cadmium toxicity thresholds for paddy soil by species-sensitivity distribution. *J. Integr. Agric.* **2015**, *14*, 1845–1854. [[CrossRef](#)]
43. Oorts, K.; Ghesquiere, U.; Swinnen, K.; Smolders, E. Soil properties affecting the toxicity of CuCl₂ and NiCl₂ for soil microbial processes in freshly spiked soils. *Environ. Toxicol. Chem.* **2006**, *25*, 836–844. [[CrossRef](#)] [[PubMed](#)]
44. Irem, S.; Islam, E.; Maathuis, F.; Niazi, N.K.; Li, T. Assessment of potential dietary toxicity and arsenic accumulation in two contrasting rice genotypes: Effect of soil amendments. *Chemosphere* **2019**, *225*, 104–114. [[CrossRef](#)]

45. Paschke, M.W.; Redente, E. Copper toxicity thresholds for important restoration grass species of the western United States. *Environ. Toxicol. Chem.* **2002**, *21*, 2692–2697. [[CrossRef](#)]
46. Liu, H.Y.; Probst, A.; Liao, B.H. Metal contamination of soils and crops affected by the Chenzhou lead/zinc mine spill (Hunan, China). *Sci. Total Environ.* **2005**, *339*, 153–166. [[CrossRef](#)]
47. Smolders, E.; Oorts, K.; van Sprang, P.; Schoeters, I.; Janssen, C.R.; McGrath, S.P.; McLaughlin, M.J. Toxicity of trace metals in soil as affected by soil type and aging after contamination: Using calibrated bioavailability models to set ecological soil standards. *Environ. Toxicol. Chem.* **2009**, *28*, 1633–1642. [[CrossRef](#)]
48. Guala, S.D.; Vega, F.A.; Covelo, E.F. The dynamics of heavy metals in plant-soil interactions. *Ecol. Model.* **2010**, *221*, 1148–1152. [[CrossRef](#)]
49. Ordóñez, A.; Álvarez, R.; Charlesworth, S.; Miguel, E.D.; Loredó, J. Risk assessment of soils contaminated by mercury mining, Northern Spain. *J. Environ. Monit.* **2011**, *13*, 128–136. [[CrossRef](#)]
50. Harmanescu, M.; Alda, L.M.; Bordean, D.M.; Gogoasa, I.; Gergen, I. Heavy metals health risk assessment for population via consumption of vegetables grown in old mining area; a case study: Banat County, Romania. *Chem. Cent. J.* **2011**, *5*, 64–73. [[CrossRef](#)]
51. Dung, T.T.T.; Cappuyns, V.; Swennen, R.; Phung, N.K. From geochemical background determination to pollution assessment of heavy metals in sediments and soils. *Rev. Environ. Sci. Biotechnol.* **2013**, *12*, 335–353. [[CrossRef](#)]
52. Zhang, S.; Song, J.; Gao, H.; Zhang, Q.; Lv, M.-C.; Wang, S.; Liu, G.; Pan, Y.Y.; Christie, P.; Sun, W. Improving prediction of metal uptake by Chinese cabbage (*Brassica pekinensis* L.) based on a soil-plant stepwise analysis. *Sci. Total Environ.* **2016**, *569–570*, 1595–1605. [[CrossRef](#)] [[PubMed](#)]
53. Doabi, S.A.; Karami, M.; Afyuni, M.; Yeganeh, M. Pollution and health risk assessment of heavy metals in agricultural soil, atmospheric dust and major food crops in Kermanshah province, Iran. *Ecotoxicol. Environ. Saf.* **2018**, *163*, 153–164. [[CrossRef](#)] [[PubMed](#)]
54. Antoniadis, V.; Golia, E.E.; Liu, Y.T.; Wang, S.L.; Shaheen, S.M.; Rinklebe, J. Soil and maize contamination by trace elements and associated health risk assessment in the industrial area of Volos, Greece. *Environ. Int.* **2019**, *124*, 79–88. [[CrossRef](#)] [[PubMed](#)]
55. Rutigliano, F.A.; Marzaioli, R.; De Crescenzo, S.; Trifuoggi, M. Human health risk from consumption of two common crops grown in polluted soils. *Sci. Total Environ.* **2019**, *691*, 195–204. [[CrossRef](#)]
56. Li, K.; Cao, C.L.; Ma, Y.B.; Su, D.C.; Li, J.M. Identification of cadmium bioaccumulation in rice (*Oryza sativa* L.) by the soil-plant transfer model and species sensitivity distribution. *Sci. Total Environ.* **2019**, *692*, 1022–1028. [[CrossRef](#)]
57. Zabel, T.F.; Cole, S.M. The Derivation of Environmental Quality Standards for the Protection of Aquatic Life in the UK. *Water Environ. J.* **2010**, *13*, 436–440. [[CrossRef](#)]
58. Wheeler, J.R.; Grist, E.M.; Leung, K.; Morrith, D.; Crane, M. Species sensitivity distributions: Data and model choice. *Mar. Pollut. Bull.* **2020**, *45*, 192–202. [[CrossRef](#)]
59. Fang, F.S.; Fu, H.W.; Xu, W.; Wu, H.; Xiang, X.L. Soil threshold values of total and available cadmium for vegetable growing based on field data in Guangdong province, South China. *J. Sci. Food Agric.* **2013**, *93*, 1967–1973.
60. Shi, Y.; Pan, Y.; Wang, J.; Cai, Y. Perfluorinated Chemicals Related Environmental Problems. *Prog. Chem.* **2009**, *21*, 369–376.
61. Ding, C.; Li, X.; Zhang, T.; Ma, Y.; Wang, X. Phytotoxicity and accumulation of chromium in carrot plants and the derivation of soil thresholds for Chinese soils. *Ecotoxicol. Environ. Saf.* **2014**, *108*, 179–186. [[CrossRef](#)]
62. Ding, C.; Ma, Y.; Li, X.; Zhang, T.; Wang, X. Determination and validation of soil thresholds for cadmium based on food quality standard and health risk assessment. *Sci. Total Environ.* **2017**, *619*, 700–706. [[CrossRef](#)]
63. Wang, Z.; Meng, B.; Zhang, W.; Bai, J.; Ma, Y.; Liu, M. Multi-target risk assessment of potentially toxic elements in farmland soil based on the environment-ecological-health effect. *Int. J. Environ. Res. Public Health* **2018**, *15*, 1101. [[CrossRef](#)]
64. Yotova, G.; Padareva, M.; Hristova, M.; Astel, A.; Georgieva, M.; Dinev, N.; Tsakovski, S. Establishment of geochemical background and threshold values for 8 potential toxic elements in the Bulgarian soil quality monitoring network. *Sci. Total Environ.* **2018**, *643*, 1297–1303. [[CrossRef](#)]
65. Birch, G.F. Determination of sediment metal background concentrations and enrichment in marine environments—A critical review. *Sci. Total Environ.* **2017**, *580*, 813–831. [[CrossRef](#)]
66. Naz, A.; Khan, S.; Muhammad, S.; Ahmad, R.; Rahman, Z.U. Risk assessment of hazardous elements in wastewater irrigated soil and cultivated vegetables in Pakistan. *Arab. J. Geosci.* **2020**, *13*, 1201–1210. [[CrossRef](#)]
67. Hu, B.; Zhao, R.; Chen, S.; Yue, Z.; Jin, B.; Li, Y.; Shi, Z. Heavy metal pollution delineation based on uncertainty in a coastal industrial city in the Yangtze River Delta, China. *Int. J. Environ. Res. Public Health* **2018**, *15*, 710. [[CrossRef](#)]
68. Ibezim-Ezeani, M.; Dibofori-Orji, A.; Udo, K.; Ihunwo, O. Assessment of metal contamination in sediment from Woji Creek, Niger Delta Region of Nigeria. *Int. J. Environ. Anal. Chem.* **2020**, *3*, 1–12. [[CrossRef](#)]
69. Ztürk, E.; Dengiz, O. Assessment and selection of suitable microbasins for organic agriculture under subhumid ecosystem conditions: A case study from Trabzon Province, Turkey. *Arab. J. Geosci.* **2020**, *13*, 1222–1234. [[CrossRef](#)]
70. Jia ZM, Li SY, Wang L: Assessment of soil heavy metals for eco-environment and human health in a rapidly urbanization area of the upper Yangtze Basin. *Sci. Rep.* **2018**, *8*, 3256–3269. [[CrossRef](#)]

71. Chen, C.M.; Dubin, R.; Kim, M.C. Emerging trends and new developments in regenerative medicine: A scientometric update (2000–2014). *Expert Opin. Biol. Ther.* **2014**, *14*, 1295–1317. [[CrossRef](#)]
72. Puthusseri, R.M.; Nair, H.P.; Johnny, T.K.; Bhat, S.G. Insights into the response of mangrove sediment microbiomes to heavy metal pollution: Ecological risk assessment and metagenomics perspectives. *J. Environ. Manag.* **2021**, *298*, 113492. [[CrossRef](#)]