

Meeting abstract

Open Access

**Epigenetic therapy with hydralazine and valproate associated to cisplatin chemoradiation in FIGO stage IIIB. A phase II study**

Myrna Candelaria<sup>1</sup>, Lucely Cetina<sup>1</sup>, Alicia Garcia<sup>1</sup>, Talia Wegman-Ostrosky<sup>1</sup>, Elizabeth Robles<sup>1</sup>, Aurora González-Fierro<sup>1</sup>, Carlos López-Graniel<sup>2</sup>, Aaron González<sup>2</sup>, David Cantú<sup>2</sup>, Lesbia Ribera<sup>3</sup>, Lucia Taja-Chayeb<sup>1</sup>, Catalina Trejo-Becerril<sup>1</sup>, Enrique Pérez-Cárdenas<sup>1</sup>, Carlos Pérez-Plasencia<sup>1</sup>, Alma Chavez-Blanco<sup>1</sup> and Alfonso Dueñas-González\*<sup>1,4</sup>

Address: <sup>1</sup>Division of Clinical Research. INCAN, Mexico, <sup>2</sup>Department of Gynecology-Oncology. INCAN, Mexico, <sup>3</sup>Division of Radiation Oncology. INCAN, Mexico and <sup>4</sup>Unit of Biom Res on Cancer, IIB UNAM/INCAN, Mexico

Email: Alfonso Dueñas-González\* - [alfonso\\_duenasg@yahoo.com](mailto:alfonso_duenasg@yahoo.com)

\* Corresponding author

from 24<sup>th</sup> Annual Meeting of the National Cancer Institute of Mexico  
Mexico City, Mexico. 14–17 February 2007

Published: 5 February 2007

BMC Cancer 2007, 7(Suppl 1):A28 doi:10.1186/1471-2407-7-S1-A28

This article is available from: <http://www.biomedcentral.com/1471-2407/7/S1/A28>

© 2007 Candelaria et al; licensee BioMed Central Ltd.

**Background**

Standard chemoradiation with cisplatin has improved the results of treatment, however, newer therapeutical modalities are needed. DNA demethylating and histone deacetylase inhibitors have shown radiosensitizing effects as well as potentiation of chemotherapy drugs. This study was set to evaluate hydralazine and valproate as epigenetic therapy added to chemoradiation with cisplatin in cervical cancer.

**Materials and methods**

Eligibility criteria included untreated disease, histological diagnosis and adequate bone marrow, hepatic and liver function. Patients were typed for acetylator phenotype and then oral hydralazine at 182 mg/day or 83 mg/day for rapid and slow-acetylators respectively. Oral magnesium valproate was used at 30 mg/Kg/day. Treatment started 7 days before commencing chemoradiation. Chemoradiation consisted on EBRT 50 Gy in 2G fractions plus cisplatin at 40 mg/m<sup>2</sup> weekly × 6. Brachytherapy was scheduled after EBRT. Response, toxicity and survival were evaluated as well as the biological effects of hydralazine and valproate in blood and primary tumor samples.

**Results**

Twenty one patients were included. Mean age was 50.4 years (range: 28–69 y). Overall, 85% of patients completed both phases, external beam and intracavitary therapy. Three patients did not receive brachytherapy: two abandoned treatment during EBRT and one died after the first cycle of chemotherapy. Mean + SD dose administered to point A was 84.6 + 2.2 (range: 79.6 – 87 for point A). Median number of cisplatin cycles administered was five (range 1–6). The 18 patientes evaluable for response achieved complete clinical response after EBRT. The most frequent grade 3/4 hematological toxicity was neutropenia (45%), followed by thrombocytopenia (10%) and anemia (10%). Non-hematological toxicity was mild in most of cases. Three out of 18 cases have relapsed (16.6%): one local, one had local/systemic relapse and one systemic, at a median follow-up of 11 months (range: 1 – 13.4 m). The hydralazine/valproate-induced changes in the expression of genes analyzed by microarrays will be presented in the meeting.

**Conclusion**

Epigenetic therapy with hydralazine and valproate associated to chemoradiation in cervical cancer is well-tolerated

and seems effective as all evaluable patients achieved clinical complete response during the external radiation.

Publish with **BioMed Central** and every scientist can read your work free of charge

*"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."*

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:  
[http://www.biomedcentral.com/info/publishing\\_adv.asp](http://www.biomedcentral.com/info/publishing_adv.asp)

