

# Transcriptional regulation of oil biosynthesis in seed plants: Current understanding, applications, and perspectives

Yuzhou Yang<sup>1,6</sup>, Que Kong<sup>1,6</sup>, Audrey R.Q. Lim<sup>1</sup>, Shaoping Lu<sup>2,3</sup>, Hu Zhao<sup>2,3</sup>, Liang Guo<sup>2,3,\*</sup>, Ling Yuan<sup>4,5,\*</sup> and Wei Ma<sup>1,\*</sup>

<sup>1</sup>School of Biological Sciences, Nanyang Technological University, Singapore 637551, Singapore

<sup>2</sup>National Key Laboratory of Crop Genetic Improvement, Huazhong Agricultural University, Wuhan 430070, China

<sup>3</sup>Hubei Hongshan Laboratory, Wuhan 430070, China

<sup>4</sup>Department of Plant and Soil Sciences, Kentucky Tobacco Research and Development Center, University of Kentucky, Lexington, KY 40546, USA

<sup>5</sup>Guangdong Provincial Key Laboratory of Applied Botany, South China Botanical Garden, Chinese Academy of Sciences, Guangzhou 510650, China

<sup>6</sup>These authors contributed equally to this article.

\*Correspondence: Liang Guo ([guoliang@mail.hzau.edu.cn](mailto:guoliang@mail.hzau.edu.cn)), Ling Yuan ([lyuan3@uky.edu](mailto:lyuan3@uky.edu)), Wei Ma ([weima@ntu.edu.sg](mailto:weima@ntu.edu.sg))

<https://doi.org/10.1016/j.xplc.2022.100328>

## ABSTRACT

Plants produce and accumulate triacylglycerol (TAG) in their seeds as an energy reservoir to support the processes of seed germination and seedling development. Plant seed oils are vital not only for the human diet but also as renewable feedstocks for industrial use. TAG biosynthesis consists of two major steps: *de novo* fatty acid biosynthesis in the plastids and TAG assembly in the endoplasmic reticulum. The latest advances in unraveling transcriptional regulation have shed light on the molecular mechanisms of plant oil biosynthesis. We summarize recent progress in understanding the regulatory mechanisms of well-characterized and newly discovered transcription factors and other types of regulators that control plant fatty acid biosynthesis. The emerging picture shows that plant oil biosynthesis responds to developmental and environmental cues that stimulate a network of interacting transcriptional activators and repressors, which in turn fine-tune the spatiotemporal regulation of the pathway genes.

**Keywords:** plant oil biosynthesis, oil accumulation, seed development, environmental and developmental signals, transcription factor, transcriptional regulation


Yang Y., Kong Q., Lim A.R.Q., Lu S., Zhao H., Guo L., Yuan L., and Ma W. (2022). Transcriptional regulation of oil biosynthesis in seed plants: Current understanding, applications, and perspectives. *Plant Comm.* **3**, 100328.

## INTRODUCTION

Plant oils (often referred to as vegetable oils) are generated and stored mainly in the form of triacylglycerol (TAG) in seeds. For plants, TAG serves as a major carbon and energy source to support seed germination and seedling development. For humans, plant oils are not only essential as a component of the diet, providing about one-fourth of dietary calories in developed countries, but also important as a source of carbon-neutral fuels and raw materials for industrial products, such as lubricants, detergents, and nylon (Chapman and Ohlrogge, 2012; Covas et al., 2015; Kong et al., 2020a; Song et al., 2021b). Global demand for vegetable oils is increasing rapidly and is projected to double by 2030 (Chapman and Ohlrogge, 2012), incentivizing a tremendous amount of research effort to enhance plant oil biosynthesis.

The mechanism of fatty acid (FA) biosynthesis has been extensively studied and reviewed (Chapman and Ohlrogge, 2012; Bates et al., 2013; Li-Beisson et al., 2013; Allen, 2016; Bates, 2016; Xu and Shanklin, 2016; Kong et al., 2020a; Song et al., 2021b; Miray et al., 2021). In brief, FA synthesis starts with the provision of carbon from glycolysis. After glycolysis, pyruvate dehydrogenase converts pyruvate into acetyl-CoA, the initial substrate for *de novo* FA biosynthesis. Acetyl-CoA carboxylase (ACCase) catalyzes the first committed step, converting acetyl-CoA and bicarbonate into malonyl-CoA. A key co-factor, acyl-carrier protein (ACP), then accepts the malonyl group to form

Published by the Plant Communications Shanghai Editorial Office in association with Cell Press, an imprint of Elsevier Inc., on behalf of CSPB and CEMPS, CAS.



**Figure 1. Environmental and developmental cues mediate gene expression and transcriptional activity during seed development.**

Environmental signals affect embryogenesis as well as phytohormones and sugar levels, which also control embryogenesis. Developmental signals significantly affect the LAFL network (which includes *LEC1*, *LEC2*, *L1L*, *ABI3*, and *FUS3*), *WRI1* and its interacting partners, and other transcription factors, leading to dynamic transcriptional regulation of plant oil biosynthetic pathways. Epigenetic regulators such as *CURLY LEAF* (CLF) and *PICKLE* (PKL) also determine the transcription levels of several key regulators in the LAFL network and *WRI1*. The combined transcriptional regulation leads to responsive metabolic readout, resulting in alteration of oil content and composition.

malonyl-ACP. After the first cycle of FA synthesis produces an acetoacetyl-ACP, additional cycles, facilitated by the fatty acid synthase complex, elongate the FA chain in a two-carbon increment, whereby acetyl units provided from malonyl-CoA are added successively to a lengthening acyl chain esterified to an ACP. During this process, 3-ketoacyl-ACP synthase (KAS) catalyzes the condensation reaction to connect new carbons to the growing acyl chain. KAS III initiates FA biosynthesis by combining acetyl-CoA and malonyl-ACP. KAS I subsequently extends the acyl chain to C12–C16. Eventually, KAS II completes the biosynthesis to C18. The acyl chains are released from acyl-ACPs by hydrolysis reactions catalyzed by thioesterases. The products of thioesterases are activated to acyl-CoA molecules prior to export to the endoplasmic reticulum. Currently, there are still some ambiguities regarding the specific steps and the sequence of events that occur during fatty acyl activation and export out of plastid envelopes (Bates et al., 2013; Bates, 2016). The pathway to TAG assembly from glycerol-3-phosphate and acyl-CoAs consists of multiple enzymatic steps (Chapman and Ohlrogge, 2012; Li-Beisson et al., 2013; Allen, 2016; Bates, 2016). First, glycerol-3-phosphate acyltransferase generates lysophosphatidic acid by transfer of an FA from acyl-CoA to the *sn*-1 hydroxyl of glycerol-3-phosphate. Then lysophosphatidic acid acyltransferase (LPAAT) esterifies a second FA to the *sn*-2 hydroxyl of lysophosphatidic acid to produce phosphatidic acid. Third, phosphatidic

acid phosphatase mediates the removal of the *sn*-3 phosphate to produce *de novo* diacylglycerol (DAG). The *de novo* DAG pool is vital, as it can be used for biosynthesis of membrane lipids, such as phosphatidylcholine (PC) and phosphatidylethanolamine, or for TAG production. The DAG for TAG biosynthesis can also be obtained from PC. Acyl flux of the *de novo* DAG into PC and the ensuing acyl flux of PC-derived DAG out of PC that generates the DAG substrate for producing TAG may occur through a phosphocholine headgroup exchange mechanism. In the final stage, diacylglycerol acyltransferase (DGAT) or phospholipid:diacylglycerol acyltransferase (PDAT) catalyzes the conversion of DAG to TAG (Zou et al., 1999; Dahlqvist et al., 2000; Zhang et al., 2009). TAG is then coated with oleosins and stored as lipid droplets (Chapman et al., 2012).

TAG biosynthesis is a complex biological process that involves three main metabolic pathways: glycolysis, FA *de novo* synthesis and elongation, and TAG assembly. Glycolysis provides carbons that can be esterified with glycerol to produce TAG. Hence, seed oil accumulation requires the balanced transcriptional regulation of multiple branch pathways (Chapman and Ohlrogge, 2012; Lu et al., 2018; Kong et al., 2019). Transcription factors are proteins that recognize and bind, directly or indirectly, to specific DNA sequences in gene promoters to control gene expression (Lambert et al., 2018; Zhu et al., 2018). Numerous transcription factors have been shown to participate in regulating the expression of TAG biosynthetic genes (Cernac and Benning, 2004; Baud et al., 2007; Mu et al., 2008; Yamamoto et al., 2010; Ma et al., 2013; Li et al., 2017; Pelletier et al., 2017; Kong and Ma, 2018b; Tian et al., 2020). Among them, the *APETALA2* transcription factor *WRINKLED1* (*WRI1*) directly regulates the expression of various genes involved in glycolysis and FA biosynthesis, functioning to “push” FA *de novo* synthesis from upstream (Vanhercke et al., 2014, 2019). Other transcriptional regulators also participate in regulation of the FA biosynthetic pathway by mediating the expression of *WRI1* (Baud et al., 2007; Yamamoto et al., 2010; Li et al., 2017; Kong et al., 2020a; Tian et al., 2020). Unlike animals, plants are sessile organisms that are incapable of escaping from various environmental hazards. Plants have therefore evolved sophisticated systems to translate environmental cues into metabolic modulation to mitigate environmental impacts. FA biosynthesis is part of seed development, which responds to various environmental signals, including light, humidity, and temperature. Transcriptional and other regulatory networks integrate the balanced regulation of environmental signal perception/transduction, seed development, and FA biosynthesis (Figure 1). In this review, we focus on understanding some of the key regulators in these networks.

### The essential roles of key regulators of seed development in transcriptional regulation of seed oil accumulation

Numerous studies suggest that four transcriptional regulators, *LEAFY COTYLEDON1* (*LEC1*), *LEAFY COTYLEDON2* (*LEC2*), *FUSCA3* (*FUS3*), and *ABSCISIC ACID INSENSITIVE3* (*ABI3*), play central roles in the process of seed development and maturation (Giraudat et al., 1992; Parcy et al., 1997; Lotan et al., 1998; Luerssen et al., 1998; Stone et al., 2001; Verdier and Thompson, 2008; Yamamoto et al., 2010; Pelletier et al., 2017; Jo et al., 2019;

Song et al., 2021a). Recent evidence shows that LEC1, LEC2, FUS3, and ABI3 are also involved in the transcriptional regulation of seed oil production (Baud et al., 2007; Mu et al., 2008; Shen et al., 2010; Tan et al., 2011; Zhang et al., 2016; Manan et al., 2017; Tian et al., 2020; Yang et al., 2021). Genes encoding these regulators are expressed predominantly during seed development and have a considerable effect on seed oil accumulation.

LEC1 encodes a HAP3 subunit of the CCAAT-binding transcription factor. Inducible transgenic lines of *Arabidopsis* overexpressing *AtLEC1* (*AtLEC1-OXI*) showed enhanced FA production (Mu et al., 2008). Global transcriptomic analysis of *AtLEC1-OXI* lines revealed that LEC1 regulates the expression of genes involved in FA metabolism, including key reactions of condensation, chain elongation, and desaturation of FA biosynthesis (Mu et al., 2008). Elevation of *LEC1* enhanced seed oil content. Overexpression of *AtLEC1*, *BnLEC1* (*Brassica napus* *LEC1*), and *ZmLEC1* (maize *LEC1*) increased oil content in transgenic plants (Mu et al., 2008; Shen et al., 2010; Tan et al., 2011). LEC1-LIKE (L1L), a close homolog of LEC1, displays a similar function as LEC1 in the regulation of FA biosynthesis in *Arabidopsis* and *B. napus* (Mu et al., 2008; Tan et al., 2011). Notably, a recent study illustrated the significance of the endosperm-produced *AtLEC1* in facilitating the seed developmental process, including seed maturation (Song et al., 2021a). *LEC1* expression occurs in the endosperm, followed by LEC1 trafficking to the embryo to regulate seed maturation, indicating the complex roles of LEC1 in mediating seed development and controlling metabolic activities, such as FA production (Song et al., 2021a).

LEC2 is a B3 domain transcription factor vital for embryo development (Stone et al., 2001). In addition to its essential function in controlling the developmental process, LEC2 is involved in transcriptional regulation of FA biosynthesis and has therefore been used in engineering plants to enhance oil production (Santos Mendoza et al., 2005; Baud et al., 2007; Vanhercke et al., 2014, 2019; Kim et al., 2015). Transgenic *Arabidopsis* seedlings expressing an inducible *LEC2* showed oil accumulation in leaves upon induction, accompanied by increased expression of genes encoding key transcription factors, such as LEC1, ABI3, and FUS3 (Santos Mendoza et al., 2005).

In the transcriptional network, LEC1 and LEC2 positively regulate the expression of *WRI1* (Baud et al., 2007; Mu et al., 2008; Pelletier et al., 2017; Kong et al., 2019). *WRI1* expression is elevated in an *Arabidopsis* gain-of-function mutant of *LEC1*, *tnp*, as well as inducible *LEC1* overexpression lines (Casson and Lindsey, 2006; Mu et al., 2008). Inducible *LEC2* overexpression after cycloheximide treatment (to block translation) showed the activation of *AtWRI1* without the participation of other intermediate proteins, and *AtWRI1* expression is decreased in the *lec2* mutant compared with that in the wild-type (WT) control (Baud et al., 2007). Chromatin immunoprecipitation (ChIP) experiments confirmed *WRI1* as a direct target of LEC1 (Pelletier et al., 2017). Effects of LEC1 and LEC2 on *WRI1* expression are conserved across other plant species, such as maize and soybean (Shen et al., 2010; Manan et al., 2017).

The B3 domain transcription factor FUSCA3 (FUS3) triggers the expression of various FA biosynthetic genes, such as *FATTY ACID DESATURASE3* (*FAD3*), *KAS I*, and *FATTY ACID ELONGATION1* (*FAE1*) (Yamamoto et al., 2010; Wang and Perry, 2013). FUS3 also binds directly to the promoters of *LEC1*, *L1L*, and *ABI3* to mediate their expression (Wang and Perry, 2013). Transcriptomic analysis revealed that the expression of *WRI1* is decreased in the *fus3* mutant compared with the WT (Yamamoto et al., 2010). *WRI1* was identified as a direct target of FUS3 in a genome-wide ChIP-chip analysis (Wang and Perry, 2013). The *fus3* mutant displays fewer oil droplets in the cotyledons of developing seeds, as well as reduced seed oil content compared with the WT (Meinke et al., 1994; Tiedemann et al., 2008; Roscoe et al., 2015). Transgenic overexpression of *FUS3* in *Arabidopsis* significantly triggers TAG accumulation in vegetative tissues, suggesting a useful role for FUS3 in the bioengineering of vegetable oil production (Zhang et al., 2016).

The B3 domain transcription factor ABI3 plays an important role in the abscisic acid (ABA) signaling pathway that mediates physiological processes such as seed maturation and dormancy (Rohde et al., 2000; Monke et al., 2012; Boulard et al., 2017; Tian et al., 2020). Recent ChIP-chip experiments identified direct targets of ABI3; they included *WRI1* and other genes encoding FA biosynthetic genes, including *FAD3* and *FATTY ACID BIOSYNTHESIS 2* (*FAB2*) (Tian et al., 2020). The combined results of ChIP-chip and comparative transcriptomic analysis of developing *Arabidopsis* seeds of *abi3* and WT showed that 317 genes, including *WRI1*, were directly activated by ABI3 (Tian et al., 2020). Furthermore, a recent study indicated that transgenic tobacco BY2 cell lines or transgenic *Arabidopsis* seedlings expressing an inducible *ABI3* showed activation of oil production, further supporting a role for ABI3 in the regulation of oil biosynthesis. In the *fus3* mutant, induction of *ABI3* triggers strong TAG production in transgenic seedlings, suggesting that ABI3 activates oil biosynthesis in a FUS3-independent manner. Further transcriptomic analysis showed that *LIPID DROPLET PROTEIN* (*LDP*) genes and *WRI1* are activated by ABI3 in transgenic *fus3* plants (Yang et al., 2021).

AGAMOUS-Like15 (AGL15) is a well-characterized MADS transcription factor (Thakare et al., 2008; Zheng et al., 2009; Serivichyaswat et al., 2015; Cosio et al., 2017; Joshi et al., 2021). ChIP-chip analysis detected various direct targets of AGL15, including *LEC2*, *FUS3*, and *ABI3* (Zheng et al., 2009). Recent ChIP analysis revealed that HIGH-LEVEL EXPRESSION OF SUGAR INDUCIBLE GENE2 (*HSI2*)/VP1/ABI3-LIKE1 (*VAL1*) directly binds to the *AGL15* promoter via the RY element to repress the expression of *AGL15*. In the *hsi2-2* mutant, the expression of *AGL15* (as well as *LEC1*, *LEC2*, *ABI3*, and *FUS3*) was significantly increased compared with the WT (Chen et al., 2018b). *BABY BOOM* (*BBM*), encoding an APETALA2 transcription factor, is essential for transcriptional regulation of plant cell totipotency (Horstman et al., 2014, 2017; Lutz et al., 2015). Functional analysis revealed that BBM binds directly to the promoters of *LEC1*, *LEC2*, and *ABI3*. In BBM-overexpressing transgenic plants, expression of *LEC1*, *LEC2*, *FUS3*, and *ABI3* was upregulated, suggesting a positive role for BBM in the activation of these FA-biosynthesis-related genes in



*planta* (Horstman et al., 2017). The direct involvement of AGL15 or BBM in oil production remains to be explored.

### The pivotal role of WRI1 in transcriptional control of plant oil biosynthesis


The *AtWRI1* loss-of-function mutant *wri1-1* displays a near 80% reduction in seed oil content compared with the WT (Focks and Benning, 1998). Transcriptomic analysis of developing *wri1-1* seeds showed that most genes with decreased expression encoded key enzymes in late glycolysis and the FA pathways (Ruuska et al., 2002). Subsequent studies confirmed that *AtWRI1* directly regulates many of these genes (Baud et al., 2007; Maeo et al., 2009), including *pyruvate kinase beta subunit 1* (*PKP-β1*), *biotin carboxyl carrier protein* (*BCCP2*; encoding a subunit of the ACCase), *ACP1*, and *KAS I*. *WRI1* binds directly to the promoter region and activates the expression of the *GLB1* gene, which encodes a PII protein, a key integrator of central metabolism (Baud et al., 2010). PII displays inhibitory effect on ACCase activity and is involved in fine-tuning FA biosynthesis in developing seeds (Baud et al., 2010). Several *BIOTIN ATTACHMENT DOMAIN-CONTAINING* (*BADC*) genes that repress FA biosynthesis are activated by *AtWRI1* (Liu et al., 2019), suggesting another level of mediation of *WRI1*-regulated oil biosynthesis. Furthermore, a recent discovery showed that *carboxyltransferase interactor* (*CTI*), encoding an envelope membrane regulator of ACCase, is also positively regulated by *WRI1* (Ye et al., 2020). In contrast to the positive regulation by *LEC1/LEC2/FUS3*, the *Arabidopsis* *R2R3-MYB* transcription factor *MYB89* represses *AtWRI1* expression (Li et al., 2017) (Figure 2A). It should be noted that alternative transcriptional regulators may mediate *WRI1* expression in other plant species. In oil palm mesocarp (where most palm oil is generated), *EgWRI1* was highly repressed during fruit ripening. However, no transcripts of orthologs of known *WRI1*-regulator genes (e.g., *LEC1*, *LEC2*, *ABI3*, *FUS3*) were detected, suggesting that additional upstream transcriptional regulators (possibly fruit-specific factors) control the expression of *EgWRI1* (Bourgis et al., 2011). Transcriptional regulators, including *EgNF-YA3*, *EgNF-YC2*, and *EgABI5*, participate in the transcriptional control of *EgWRI1* by binding to the *EgWRI1* promoter (Yeap et al., 2017).

To prevent hyperactivation of essential biological processes, which might result in metabolic imbalance, the activities of transcriptional regulators are frequently calibrated through interactions with other regulators (Hafner et al., 2019; Memelink and Gantet, 2007; Mouchiroud et al., 2014; Pireyre and Burow, 2015; Sui et al., 2018). As the protein–protein interaction network of *WRI1* has been unfolded, various interacting partners of *AtWRI1* have been found to modulate its activity (summarized in Figure 2A). The CULLIN3 (*CUL3*)-based E3 ligase adaptor BTB/POZMATH (*BPM*) interacts with *AtWRI1* to bridge *AtWRI1* to *CUL3*, which mediates *AtWRI1* degradation by the 26S proteasome (Chen et al., 2013). A recent study showed that *BPM* mRNA stability is mediated by *Arabidopsis* *PUMILIO PROTEIN24* (*APUM24*, encoding a Pumilio homology domain-containing protein that is involved in rRNA processing and mRNA degradation), subsequently fine-tuning the function of the *BPM-WRI1* module (Huang et al., 2021). The 14-3-3 proteins, a family of phosphopeptide-binding proteins, physically interact with *AtWRI1* in yeast two-hybrid and bimolecular fluorescence complementation assays (Ma et al., 2016). Transient

overproduction of 14-3-3 with *AtWRI1* results in increased oil biosynthesis in *Nicotiana benthamiana* leaves by enhancing the protein stability and transcriptional activity of *AtWRI1* (Ma et al., 2016). *BPM1* and 14-3-3 independently interact with *AtWRI1* protein in an overlapping region, leading to the hypothesis that 14-3-3 interacts with *AtWRI1* at the expense of the *AtWRI1-BPM* interaction (Ma et al., 2016; Kong and Ma, 2018a). The SNF1-related protein kinase *KIN10* is another *AtWRI1*-interacting partner that phosphorylates *AtWRI1*, resulting in the degradation of *AtWRI1* protein (Zhai et al., 2017). In addition, phosphorylation-deficient mutagenesis at amino acid residues T70 and S166 of *AtWRI1* abolished *KIN10*-mediated phosphorylation and improved *AtWRI1* stability (Zhai et al., 2017). Trehalose 6-phosphate is involved in stabilizing *AtWRI1* and boosting FA biosynthesis by suppressing *KIN10* activity (Zhai et al., 2018).

Furthermore, interacting partners may modulate the assembly of the *WRI1* transcriptional complex. In eukaryotic cells, recruitment of mediator (*MED*) subunits by transcription factors to initiate transcription is a conserved regulatory mechanism (Taates, 2010). The *MED15* subunit of the *Arabidopsis* mediator complex has been shown to physically interact with *WRI1* both *in vitro* and *in vivo* (Kim et al., 2016). Transgenic *Arabidopsis* overexpressing *MED15* displays upregulation of the FA biosynthetic genes targeted by *AtWRI1*. ChIP experiments also revealed that *MED15* binds to the promoters of the *AtWRI1* target genes. However, the transgenic *wri1* mutant that overproduces *MED15* also displays enhanced expression of *AtWRI1* target genes, suggesting that alternative transcription factors interact with *MED15* to modulate the expression of the oil biosynthetic genes (Kim et al., 2016).

Intra- and inter-family interactions of transcriptional regulators significantly increase the complexity of combinatorial transcriptional regulation. In plants, protein complexes formed by members of a transcription factor family are well documented; nonetheless, the importance of cross-family interactions of transcriptional regulators in gene regulation is increasingly recognized (Bemer et al., 2017). Members of a transcription factor family are generally a group of *cis*-element-specific DNA-binding factors with overlapping or divergent functions. Different transcription factor families vary in their DNA binding properties, transactivation activities (e.g., functions as repressors or activators), and responses to environmental/developmental stimuli. As such, cross-family transcription factor interactions are a key molecular mechanism for fine-tuning gene expression in response to diverse environmental and developmental cues. Screening of an *Arabidopsis* transcription factor library led to the identification of *TCP4* as a cross-family interacting regulator of *AtWRI1* (Kong et al., 2020c). Functional analyses revealed that *TCP4* reduces *AtWRI1*-regulated oil accumulation by repressing *AtWRI1* transactivation activity. The *tcp4* loss-of-function mutants exhibit elevated seed oil content compared with the WT (Kong et al., 2020c). Until recently, little was known about the transcriptional repressors of *WRI1* activity. This work uncovered the potential function of *TCP4* as a transcriptional repressor of *AtWRI1*-regulated oil biosynthesis. However, the molecular mechanisms of the *WRI1-TCP4* regulatory module remain to be elucidated. For instance, phosphorylation of *WRI1* or *TCP4* individually affects their activities and functions (Ma et al., 2015; Kubota et al., 2017; Zhai et al., 2017;



**Figure 2. The complex and multi-level regulatory mechanisms for the control of plant seed oil accumulation.**

**(A)** LEC1, LEC2, FUS3, and ABI3 are the major upstream transcriptional regulators that mediate embryo development, seed maturation, and metabolism. Together with HSI2/VAL1, AGL15, BBM, TT8, MYB89, they form a network that tightly controls the expression of *WRI1*. These upstream regulators also mutually regulate one another. The upstream regulators are affected by epigenetic factors, including CLF and PKL. *WRI1* physically interacts with several transcription factors, such as BPMs, 14-3-3s, MED15, and TCP4, as well as post-translational modifiers, such as KIN10 kinase. *WRI1* is subject to post-translational modifications, such as phosphorylation (circled P) and ubiquitination (Ub), that result in protein degradation. The targets of *WRI1* include genes involved in the FA biosynthetic pathway and late glycolysis. The positive (arrows) and negative (T-bars) regulatory mechanisms, together with post-translational modifications, form a network that fine-tunes oil biosynthesis.

**(B)** Other transcriptional regulators that participate in the regulation of plant oil biosynthesis by upregulating oil pathway genes. MYB96, DOF4, DOF11, TZF, WRKY10, WRKY43, DREB2C, and bZIP67 positively regulate FA accumulation in seeds. Other factors, including TT2, TTG1, GL2, MYB76, MYB118, and WRKY6, negatively affect seed oil accumulation. The crosstalk among the positive and negative regulatory modules enables balanced and fine-tuned responses to environmental and developmental signals during seed oil accumulation. The green arrows indicate positive regulation. The red T-bars represent negative regulation. Black lines indicate the formation of protein–protein interacting complexes. Genes in light blue shaded circles are downstream targets of the transcription factors.

Kong et al., 2020b, 2020c); however, it is unclear how phosphorylation affects the DNA binding, subcellular localization, protein stability, protein complex formation, and transcriptional activity of the *WRI1*–TCP4 module. In addition, TCPs are a plant-specific transcription factor family with diverse functions, and it will be interesting to investigate the interactions between *WRI1* and other TCP members and their roles in the regulation of FA biosynthesis.

Although the *WRI1* binding sequence AW-box has been established as a key *cis*-regulatory element (Maeo et al., 2009),

*AtWRI1* is capable of binding to the promoter of *GH3.3* (involved in auxin degradation) at a non-AW-box element (Kong et al., 2017). Because *AtWRI1* recognizes and binds to the AW-box in the promoters of the auxin carrier-protein *PIN* genes, a role for *AtWRI1* in auxin homeostasis has been suggested (Kong et al., 2017). This hypothesis is supported in soybean, as overexpression of *GmWRI1* elevates, and knockdown of *GmWRI1* decreases, the expression of auxin-related genes (Chen et al., 2020). How *WRI1* modulates alternative targets to affect plant growth and development is currently being explored. Recent interesting work showed that *AtWRI1* itself

## Plant Communications

neither activates nor suppresses the activity of the *GH3.3* promoter (Kong et al., 2022). Yeast two-hybrid screening identified TCP20 as an interacting regulator of AtWRI1 (Kong et al., 2022). TCP20 was found to regulate the expression of *GH3.3* by binding to a *cis*-acting element. Intriguingly, AtWRI1 physically interacts with TCP20 to mitigate *GH3.3* expression, highlighting the role of a WRI1–TCP20 regulatory module in fine-tuning auxin homeostasis (Kong et al., 2022).

### The complex transcriptional network involved in controlling seed oil accumulation

In addition to LEC1, LEC2, FUS3, ABI3, and WRI1, significant progress has been made in understanding other transcriptional regulators that control seed oil biosynthesis (Figure 2B). The transcription factor bZIP67 binds to the G-box of the *FAD3* promoter and controls the biosynthesis of  $\alpha$ -linolenic acid in *Arabidopsis* seeds in a L1L- and NF-YC2-dependent manner (Mendes et al., 2013). In addition, overproduction of a soybean bZIP transcription factor (GmbZIP123) elevates seed oil content in transgenic *Arabidopsis* plants (Song et al., 2013). In this case, the expression of genes involved in *de novo* FA biosynthesis is not affected, but rather several sucrose transport-related genes are upregulated, suggesting that GmbZIP123 mediates FA biosynthesis by regulating sucrose metabolism (Song et al., 2013).

In addition to MYB89, other MYB transcription factors regulate seed oil biosynthesis. MYB96 regulates seed oil accumulation by controlling the expression of *DGAT1* and *PDAT1* (Lee et al., 2018) and controls the biosynthesis of very long chain fatty acids by regulating the expression of *FAE1* by directly binding to its promoter (Lee et al., 2015). MYB76, MYB118, and MYB123 (TRANSPARENT TESTA2 [TT2]) also affect seed oil biosynthesis (Chen et al., 2012a; Barthole et al., 2014; Duan et al., 2017). However, their functions seem to be distinct from that of MYB96. For example, MYB118 acts as a repressor of FA biosynthesis in the endosperm by repressing maturation-associated genes (Barthole et al., 2014). TT2, known for regulating proanthocyanidin production (Nesi et al., 2001), also modulates seed oil content. The *tt2* mutant displays increased seed oil content and increased expression of genes involved in FA biosynthesis, such as *FUS3*, *FAE1*, and *KAS II*, suggesting that TT2 negatively regulates seed oil accumulation (Chen et al., 2012a). Moreover, TT8 (bHLH42) and TRANSPARENT TESTA GLABRA1 (TTG1), known to mediate the biosynthesis of proanthocyanidin and anthocyanin, respectively (Baudry et al., 2004), play negative roles in regulating seed FA biosynthesis. TT8 represses seed FA production by binding to the promoters of *LEC1*, *LEC2*, and *FUS3* (Chen et al., 2014). TTG1, encoding a WD40 protein, is involved in various physiological processes, including anthocyanin biosynthesis, embryogenesis, seed coat mucilage production, and formation of trichomes and root hairs in *Arabidopsis* (Walker et al., 1999; Debeaujon et al., 2000; Western et al., 2001; Tsuchiya et al., 2004; Ramsay and Glover, 2005; Tominaga-Wada et al., 2011). The seeds of the *ttg1* mutant contain higher total proteins, starch, and fatty acids (Chen et al., 2015). Further analysis revealed that TTG1 indirectly represses the expression of various genes involved in seed developmental processes and FA biosynthesis (Chen et al., 2015). Overproduction of soybean GmMYB73 resulted in increased lipid

## Transcriptional regulation of plant oil biosynthesis

accumulation in seeds of *Arabidopsis* and *Lotus*, possibly through downregulation of *GLABRA2* (*GL2*) expression (Liu et al., 2014).

Other transcriptional regulators, including GL2, WRKY, Dof (DNA binding one finger), and tandem CCCH zinc finger (TZF) proteins, are also involved in controlling seed oil accumulation. *GL2* encodes a homeobox protein that plays key roles in root development, trichome formation, and seed coat mucilage biosynthesis (Cheng et al., 2014; Hung et al., 1998; Masucci et al., 1996; Rerie et al., 1994; Western et al., 2001). The *gl2* loss-of-function mutant displays an 8% increase in seed oil content compared with the WT (Shen et al., 2006). Further study revealed that *GL2* is expressed in both embryo and seed coat, and its target gene *MUCILAGE MODIFIED 4* (*MUM4*) is involved in mucilage biosynthesis. The *mum4* mutant displays a similar increased seed oil phenotype as the *gl2* mutant, possibly because the *gl2* mutant generates more oil with elevated carbon allocation to the embryo at the expense of seed coat mucilage production (Shi et al., 2012). WRKY6 was previously reported to play diverse roles in leaf senescence and responses to low phosphate stress (Chen et al., 2009; Ye et al., 2018b; Zhang et al., 2018). A recent study revealed a new role of WRKY6 in mediating plant seed oil accumulation. The *wrky6* loss-of-function mutant displays enhanced seed oil content compared with the WT (Song et al., 2020). The expression levels of *FUS3* and *DGAT1* are upregulated in the seeds of *wrky6*; however, how WRKY6 controls the expression of the oil biosynthetic genes remains unclear (Song et al., 2020). Mutation of the *MINISEED3* gene, encoding WRKY10, which controls seed development in *Arabidopsis*, leads to reduced total FA content, as well as decreased seed size and seed weight (Fatihi et al., 2013). The disruption of *WRKY43* in *Arabidopsis* did not change total FA content in seeds but significantly altered the ratio of polyunsaturated fatty acids (including linoleic acid and linolenic acid), suggesting a potential role for WRKY43 in modulating seed FA desaturation (Geilen et al., 2017). Dofs are plant-specific transcription factors that regulate diverse plant physiological processes, such as light response, seed germination, plant defense, vascular development, and phytohormone signaling (Yanagisawa, 2002; Le Hir and Bellini, 2013; Ruta et al., 2020). Overexpression of two soybean Dofs (*GmDof4* and *GmDof11*) in *Arabidopsis* increased seed oil content (Wang et al., 2007). The two GmDofs activate the expression of *ACCase* and *long-chain-acyl CoA synthetase* (*LACS*) genes while repressing *CRA1* (encoding a seed storage protein) by binding to the Dof *cis*-element in their promoters (Wang et al., 2007). Plant TZF proteins participate in a variety of phytohormone signaling events and stress responses to regulate gene expression (Kim et al., 2008; Lin et al., 2011; Bogamuwa and Jang, 2013). A recent work revealed that a soybean TZF protein, GmZF392, functions as a transcriptional activator in a seed-specific manner (Lu et al., 2021). Overexpression of *GmZF392* enhanced seed oil accumulation in transgenic *Arabidopsis* and soybean plants. GmZF392 controls the expression of FA biosynthetic genes by binding to the bipartite elements in their promoters (Lu et al., 2021).

### Epigenetic regulators involved in mediating seed oil accumulation

Epigenetic regulators affect transcriptional regulation by altering the regional state of chromatin. Recent evidence suggests that



Environmental and developmental cues	Effect on seed oil biosynthesis	Species	Reference(s)
High light intensity	Increases seed oil content Increases oleic acid content	<i>Brassica napus</i> , <i>Arabidopsis thaliana</i> , <i>Zea mays</i> , <i>Helianthus annuus</i>	Goffman et al., 2005; Ruuska et al., 2004; Schwender et al., 2004; Karki and Bates, 2018; Li et al., 2006; Andrade and Ferreiro, 1996; de la Vega et al., 2001; Singer et al., 2016; Seiler, 1986
High temperature	Reduces seed oil content Reduces unsaturated fatty acid content	<i>Glycine max</i> , <i>Helianthus annuus</i> , <i>Brassica napus</i> , <i>Linum usitatissimum</i> , <i>Glycine max</i>	Canvin, 1965; Carrera et al., 2009; Harris et al., 1978; Singer et al., 2016; Schulte et al., 2013; Wolf et al., 1982
Drought	Reduces seed oil content Increases oleic acid content, reduces linoleic acid content	<i>Helianthus annuus</i> , <i>Glycine max</i> , <i>Brassica napus</i>	Singer et al., 2016
Salinity	Reduces seed oil content Increases oleic acid content, reduces linoleic acid and/or linolenic acid content	<i>Helianthus annuus</i> , <i>Carthamus tinctorius</i> L., <i>Gossypium hirsutum</i>	Singer et al., 2016
Gibberellins	Alter seed oil content Modify fatty acid composition	<i>Arabidopsis thaliana</i>	Cao et al., 2006; Chen et al., 2012b; Zentella et al., 2007
Auxin and jasmonates	Alter seed oil content Modify fatty acid composition	<i>Brassica napus</i>	Niu et al., 2009

**Table 1. Effects of environmental and developmental cues on seed oil biosynthesis.**

epigenetic regulators participate in controlling seed oil accumulation (Figure 2A). CURLY LEAF (CLF), a histone methyltransferase of Polycomb Repressive Complex 2 (PRC2), which modulates the trimethylation of histone H3 Lys 27 (H3K27me3), functions as a key regulator in repressing gene expression during embryonic development (Liu et al., 2016). The *clf* mutant produces larger seeds with increased seed mass, greater seed oil content, and modified long-chain FA composition compared with the WT (Liu et al., 2016). Genes encoding transcriptional regulators involved in controlling FA biosynthesis, such as LEC1, LEC2, FUS3, WRI1, as well as various FA biosynthetic genes, are upregulated in developing seeds of the *clf* mutant, suggesting a role for CLF in the transcriptional repression of FA biosynthesis (Liu et al., 2016). PICKLE (PKL) is a nuclear-localized ATP-dependent chromatin remodeler from the chromodomain helicase DNA-binding domain (CHD3) family (Ogas et al., 1999; Dean Rider et al., 2003; Zhang et al., 2012; Park et al., 2017; Zha et al., 2020). PKL represses the expression of *LEC1*, *LEC2*, and *FUS3*. Expression levels of *LEC1/LEC2/FUS3* significantly increased in roots of the *pk1* mutant compared with the WT (Ogas et al., 1999; Dean Rider et al., 2003), correlated with the upregulation of storage lipid biosynthesis (Ogas et al., 1997, 1999).

### Environmental and developmental cues that affect seed oil biosynthesis

Light potentially affects seed oil accumulation (Table 1). Most Brassicaceae family members, including *Arabidopsis*, produce green seeds with photosynthetic activity (Eastmond et al., 1996; Eastmond and Rawsthorne, 1998). The light passing through the silique wall is an important driving force for FA production (Ruuska et al., 2004; Schwender et al., 2004;

Goffman et al., 2005). Analysis of *Arabidopsis* plants grown under different light conditions revealed a positive correlation between seed oil content (as well as seed yield and weight) and light intensity (Li et al., 2006; Karki and Bates, 2018). Other studies have also indicated that reduced sunlight during seed filling leads to decreased oil content in oil crops (Andrade and Ferreiro, 1996; de la Vega et al., 2001; Singer et al., 2016). Increased light intensity also alters seed FA composition, increasing the oleic acid to linoleic acid ratio (Seiler, 1986). Although the molecular mechanism by which light affects seed lipid metabolism remains to be elucidated, it is rational to hypothesize that photosynthesis affected by light conditions modifies carbon supply, which eventually influences FA biosynthesis.

Temperature affects seed oil accumulation and composition. The influence of higher global temperatures due to climate change on FA biosynthesis may change crop oil yield and/or FA composition. For example, when Canada experienced a season of abnormally warm temperatures in 2012, *B. napus* seed oil yield was significantly reduced from the normal average of 44.4% to 43.5% (Singer et al., 2016). For certain *B. napus* varieties, cooler growing temperatures, particularly during the seed filling period, tend to result in greater oil contents (Harris et al., 1978; Carrera et al., 2009; Singer et al., 2016). Similar to seed oil content, seed FA composition is also affected by temperature. The FA unsaturation level tends to be inversely correlated with increasing environmental temperature. Plant species such as *B. napus* and sunflower show reduced levels of linoleic acid and  $\alpha$ -linolenic acid when grown at higher temperatures, and the decrease in these polyunsaturated FAs is compensated by an increase in oleic acid (Canvin, 1965; Wolf et al., 1982; Schulte et al., 2013; Singer et al., 2016). The molecular

mechanism that underlies temperature effects on seed oil yield and composition remains unclear. Nonetheless, it has been proposed that higher temperature increases nitrogen bioavailability, which leads to a competition for the carbon backbone required for storage lipid production, consistent with the inverse correlation between FA and protein yields under heat stress (Canvin, 1965; Singer et al., 2016). Lower FA desaturation at elevated temperatures can also be due to lower oxygen solubility, as has been demonstrated in *Acanthamoeba* (Jones et al., 1991; Rutter et al., 2002), although such a mechanism has not been reported in higher plants.

Drought generally leads to a substantial reduction in seed oil content and modification of FA composition in plants (Singer et al., 2016) (Table 1). It is proposed that drought leads to reduced seed filling but faster embryo development, which affects FA biosynthesis (Singer et al., 2016). In addition, salinity stress causes reductions in seed oil and changes in FA composition (Singer et al., 2016). Salinity generates reactive oxygen species, which damage cellular redox homeostasis and trigger oxidative stress, negatively affecting metabolic enzymes, including those involved in FA biosynthesis. Interestingly, overproduction of DGAT1 in *B. napus* reduces the adverse effects of drought stress on seed oil production (Weselake et al., 2008). Transgenic *Camelina* co-producing CsMYB96A and CsDGAT1C also shows increased resistance to drought stress with improved seed oil content (Kim et al., 2019). Furthermore, the expression of DGAT1 is upregulated under stresses such as salinity (Lu and Hills, 2002; Kong et al., 2013), suggesting a wider role for DGAT1 in response to environmental stresses. Transgenic plants overexpressing DGAT1 are ideal materials for gaining an in-depth understanding of the role of DGAT1 under various stresses.

FA biosynthesis is associated with phytohormone signaling. Gibberellins (GAs) are important phytohormones involved in regulating numerous plant developmental processes, including stem elongation, germination, dormancy, flowering, flower development, and leaf and fruit senescence (Sasaki et al., 2003; Fleet and Sun, 2005; Swain and Singh, 2005). GAs potentially affect FA metabolism in plant cells. The DELLA proteins are essential players in the complex GA signal transduction cascade (Thomas and Sun, 2004; Jiang and Fu, 2007; Zentella et al., 2007). A transcriptomic analysis showed that DELLAs mediate the down-regulation of various GDSL-type lipase genes in seeds and flower buds, indicating a possible function of DELLAs in affecting seed FA metabolism (Cao et al., 2006). Further study showed that exogenous GA treatment or DELLA mutations caused alterations in total seed FA content, modification of FA composition, and various morphological changes in seeds (Chen et al., 2012b). A global transcriptomic analysis revealed that GA3 treatment of *ga1-3* (a severe GA-deficient mutant) significantly changes the expression of *WRI1* (Zentella et al., 2007). As essential phytohormones, auxin and jasmonate (JA) participate in mediating various developmental processes (Woodward and Bartel, 2005; Zhao, 2010; Lavy and Estelle, 2016; Huang et al., 2017; Guo et al., 2018). Transcriptomic analysis of developing *B. napus* seeds revealed that multiple auxin *de novo* biosynthesis-associated genes and a JA-inducible gene are co-expressed with FA biosynthetic genes, implying a potential role for auxin and JA in FA biosynthesis in *B. napus* seed development

(Niu et al., 2009) (Table 1). These studies shed light on the relationships among FA biosynthesis, phytohormone signaling, and seed development; however, the detailed molecular mechanisms by which phytohormone signaling modulates FA metabolism during embryogenesis remain to be fully elucidated.

Earlier studies provided evidence for the involvement of WRI1 in ABA and sugar signaling, particularly in developing seedlings (Masaki et al., 2005; Cernac et al., 2006). The interaction between ABA and sugar signaling pathways fine-tunes gene expression during plant developmental processes (Finkelstein and Gibson, 2002; Dekkers et al., 2008; Sakr et al., 2018). ABA broadly participates in various stress-related responses and developmental regulation. Unraveling the position of WRI1 in phytohormone crosstalk, particularly related to seed filling under stress conditions, is of agricultural importance. Plant-fungi interactions have been crucial for initial land plant colonization and vascular plant evolution. The arbuscular mycorrhizal (AM) symbiosis is a close evolutionary relationship between land plants and Glomeromycotina fungi (Remy et al., 1994; Bonfante and Genre, 2010). Recent studies demonstrated that WRI-like proteins, including CTCF MOTIF-BINDING TRANSCRIPTION FACTOR1 (CBX1) and WRI5a (from *Lotus japonicus* and *Medicago truncatula*, respectively), play essential roles in FA production and phosphate uptake that are necessary for the AM symbiosis (Jiang et al., 2018; Xue et al., 2018). These findings shed light on the molecular mechanism underlying bidirectional nutrient exchanges during the AM symbiosis, suggesting that, evolutionarily, WRI-like proteins are ancient transcriptional regulators. A recent study showed a conserved transcriptional change in land plants during the AM symbiosis. A WRI-like transcriptional regulator identified from *Marchantia paleacea* functions in the transcriptional regulation of FA biosynthesis and transport, which are vital for mutualistic interactions in *M. paleacea* symbiosis (Rich et al., 2021).

### Translational applications of plant transcriptional regulators for bioengineering plant oil accumulation

The function of WRI1 seems to be conserved across various angiosperms, both dicots and monocots. The orthologs of WRI1 have been characterized in a variety of plant species. Transgenic plants overproducing AtWRI1 or WRI1 orthologs display substantially enhanced seed oil content (An and Suh, 2015; Cernac and Benning, 2004; Chen et al., 2020; Chen et al., 2018a; Liu et al., 2010; Shen et al., 2010; Sun et al., 2017; Yang et al., 2015; Ye et al., 2018a) (Table 2). The shared regulatory network controlling both development and FA biosynthesis should be considered when using these transcriptional regulators to engineer plant oil production. Choosing appropriate promoters to drive *WRI1* expression is crucial for oil bioengineering. For instance, transgenic maize showed considerably elevated oil accumulation when *ZmWRI1* expression was driven by the embryo-preferred OLEOSIN (OLE) promoter but not by the starch endosperm-specific 19 KD ZEIN promoter (Shen et al., 2010). Another study showed that use of the *FUS3* promoter to drive AtWRI1 expression, aiming to extend the oil accumulation period during the mid-phase of seed development, is an efficient strategy for improving seed oil accumulation (Kanai et al., 2016) (Table 2).

Transient overexpression of AtWRI1 or other WRI1 orthologs effectively activates TAG production in tobacco leaves



Promoter	Gene	Host species	Phenotype	Reference(s)
<b>AtWRI1 and WRI1 orthologs</b>				
CaMV 35S	<i>AtWRI1</i>	<i>Arabidopsis thaliana</i>	Increases seed oil content	Cernac and Benning, 2004
SiW6 promoter	<i>AtWRI1</i>	<i>Camelina sativa</i>	Enhances seed oil content and seed mass	An and Suh, 2015
CaMV 35S	<i>GmWRI1a/GmWRI1b</i>	<i>Arabidopsis thaliana</i>	Enhance total TAG content	Chen et al., 2020
Napin promoter	<i>GmWRI1a</i>	<i>Glycine max</i>	Increases the content of total oil and total fatty acids in seeds	Chen et al., 2018a
CaMV 35S	<i>BnWRI1</i>	<i>Arabidopsis thaliana</i>	Increases seed oil content, seed weight, and size	Liu et al., 2010
OLE promoter	<i>ZmWRI1</i>	<i>Zea mays</i>	Increases seed oil content with no significant difference in embryo size	Shen et al., 2010
EnP2 promoter	<i>CoWRI1</i>	<i>Oryza sativa</i>	Increases seed oil content and alters lipid composition	Sun et al., 2017
EnP2 promoter	<i>CoWRI1</i>	<i>Arabidopsis thaliana</i>	Increases seed oil content and alters lipid composition	Sun et al., 2017
ZmUBI1 promoter	<i>BdWRI1</i>	<i>Brachypodium distachyon</i>	Increases grain dry weight and leads to higher TAG content in endosperm and leaf blades	Yang et al., 2015
Glycinin promoter	<i>HaWRI1</i>	<i>Arabidopsis thaliana</i>	Increases seed oil content, seed weight, and size	Lim et al., 2022
CaMV 35S	<i>JcWRI1</i>	<i>Jatropha curcas</i>	Elevates seed lipid content and increases seed mass	Ye et al., 2018a
CaMV 35S	<i>StWRI1/PtWRI1/AsWRI1/CeWRI1</i>	<i>Nicotiana benthamiana</i> leaves	Cause significant increases in oil accumulation	Grimberg et al., 2015
FUS3 promoter	<i>AtWRI1</i>	<i>Arabidopsis thaliana</i>	Prolonged expression of <i>AtWRI1</i> during seed development leads to increased oil content in seeds	Kanai et al., 2016
<b>AtWRI1 variants</b>				
CaMV 35S	<i>AtWRI1</i> <sup>1-397</sup>	<i>Nicotiana benthamiana</i> leaves	Deletion of the IDR-PEST motif of <i>AtWRI1</i> leads to increased <i>AtWRI1</i> protein stability and higher TAG content	Ma et al., 2015
CaMV 35S	<i>AtWRI1</i> <sup>PEST/AA</sup>	<i>Nicotiana benthamiana</i> leaves, <i>Arabidopsis thaliana</i> ( <i>wri1-1</i> )	Mutation of multiple putative phosphorylation sites in PEST motif of <i>AtWRI1</i> ( <i>AtWRI1</i> <sup>S398A/S401A/S402A/S407A/S415A/S416A/T420A/T421A/T422A/S423A</sup> ) leads to increased stability of <i>AtWRI1</i> and higher TAG content	Ma et al., 2015
CaMV 35S	<i>AtWRI1</i> <sup>4SA</sup>	<i>Nicotiana benthamiana</i> leaves	Mutation of four residues ( <i>AtWRI1</i> <sup>S398A/S401A/S402A/S407A</sup> ) in PEST motif of <i>AtWRI1</i> leads to increased stability and higher TAG content	Ma et al., 2015
CaMV 35S	<i>AtWRI1</i> <sup>RR</sup>	<i>Nicotiana benthamiana</i> leaves	The lysine to arginine mutation in the dilysine motif ( <i>AtWRI1</i> <sup>K2R/K3R</sup> ) in <i>AtWRI1</i> results in increased stability and TAG accumulation	Zhai et al., 2017

Table 2. Translational applications of plant transcriptional regulators in bioengineering plant oil.

(Continued on next page)

Promoter	Gene	Host species	Phenotype	Reference(s)
<b>WRI1-interacting partners</b>				
<i>CaMV 35S</i>	<i>14-3-3<math>\kappa</math></i> or <i>14-3-3<math>\lambda</math></i>	<i>Nicotiana benthamiana</i> leaves, <i>Arabidopsis thaliana</i>	Enhancement of AtWRI1 stability and transcriptional activity leads to increased oil production	Ma et al., 2016
<i>CaMV 35S</i>	<i>MED15</i>	<i>Arabidopsis thaliana</i>	Increases total fatty acid content in seeds	Kim et al., 2016
<i>CaMV 35S</i>	<i>KIN10</i>	<i>Nicotiana benthamiana</i> leaves	Significantly reduces TAG production; KIN10 negatively regulates AtWRI1 by triggering phosphorylation of sites in the AP2 DNA binding domains, leading to its degradation	Zhai et al., 2017
<i>CaMV 35S</i>	<i>TCP4</i>	<i>Nicotiana benthamiana</i> leaves	TCP4 negatively regulates the activity of AtWRI1, leading to reduced oil biosynthesis	Kong et al., 2020b
<i>CaMV 35S</i>	<i>BPM-microRNA</i>	<i>Arabidopsis thaliana</i>	Increases total fatty acid content in seeds; AtWRI1 interacts with E3 ligase adaptor BPMs for degradation through the 26S proteasome	Chen et al., 2013
<b>Combination of WRI1 and other genes</b>				
<i>CaMV 35S</i>	<i>AtWRI1</i> , <i>DGAT1</i>	<i>Nicotiana benthamiana</i> leaves	Increase oil accumulation due to the upregulation of late glycolysis and fatty acid biosynthesis by WRI1 and TAG synthesis by DGAT1	Vanhercke et al., 2013
<i>CaMV 35S</i>	<i>AtWRI1</i>	<i>Arabidopsis thaliana</i>	Lead to significant accumulation of oil in vegetative tissues due to the downregulation of starch formation by APS1 and directing carbon to fatty acid synthesis mediated by AtWRI1	Sanjaya et al., 2011
<i>Patatin B33 promoter</i>	<i>AtAPS1-RNAi</i>			
<i>Rubi3 promoter</i>	<i>AtWRI1</i>	Sugarcane ( <i>Saccharum</i> spp. hybrids)	Elevate TAG accumulation in leaves and stems	Zale et al., 2016
<i>Mubi promoter</i>	<i>ZmDGAT1-2</i>			
<i>CaMV 35S</i>	<i>AtOLE1</i>			
<i>CaMV 35S</i>	<i>AGPase-RNAi</i>			
<i>CaMV 35S</i>	<i>PXA1-RNAi</i>			

Table 2. Continued

(Grimberg et al., 2015; Ma et al., 2015; Vanhercke et al., 2013) (Table 2). Transient co-expression of *AtWRI1* and *DGAT1* in tobacco leaves leads to markedly elevated oil production compared with the expression of *AtWRI1* alone, indicating a synergistic effect of the two genes (Vanhercke et al., 2013). Transient overproduction of the more stable *AtWRI1* variants, or transient co-expression of *AtWRI1* with *AtWRI1*-interacting regulators that stabilize *AtWRI1*, results in enhanced oil production compared with expression of the native *WRI1* (Ma et al., 2015, 2016; Zhai et al., 2017). In an effort to improve oil biosynthesis in vegetative tissues, transgenic *Arabidopsis* were designed to overexpress *AtWRI1* and repress *AGPase* (the small subunit of ADP-glucose pyrophosphorylase) by RNAi, and they showed increased TAG accumulation in the leaves (Sanjaya et al., 2011). In sugarcane, co-expression of *WRI1*, *DGAT1*, and *oleosin 1* (*OLE1*), together with the repression of *AGPase* and *PXA1* (en-

coding a subunit of the peroxisomal ABC transporter) by RNAi, considerably increased TAG generation in vegetative tissues (Zale et al., 2016) (Table 2). The engineering scheme of “push, pull, package, and protect” involves concurrently augmenting (pushing) *de novo* FA biosynthesis by activating a key transcriptional regulator, pulling the precursors into the final products by rate-limiting enzymes, packaging TAG molecules into oil droplets, and shielding the TAG from decay by deactivating lipases (Vanhercke et al., 2014, 2019).

Enhancement of seed oil accumulation can also be achieved by the manipulation of other transcriptional regulators that mediate the expression of genes involved in alternative metabolic pathways (e.g., seed storage and carbohydrate metabolism). Transgenic *Arabidopsis* overproducing the soybean transcription factors GmbZIP123, GmDof4, and GmDof7 showed increased

seed oil content, possibly because of the upregulation of genes involved in sucrose metabolism and seed storage (Wang et al., 2007; Song et al., 2013).

Plants produce diverse unusual fatty acids, such as hydroxy fatty acids (HFAs), that are highly valuable for industrial applications because of their unique physical and chemical characteristics. Ectopic expression of a hydroxylase gene in *Arabidopsis*, while producing a small amount of HFAs, significantly decreases seed oil content (Bates and Browse, 2011; Bates et al., 2014), perhaps because of feedback inhibition of *de novo* FA synthesis in the plastids (Bates et al., 2014). Using an improved strategy in which *AtWRI1* and *RcFAH12* (a castor fatty acid hydroxylase) are co-expressed, both HFA and seed oil content are considerably increased, indicating that *AtWRI1* efficiently overcomes the feedback inhibition by the FA hydroxylase (Adhikari et al., 2016).

### Concluding remarks and perspectives

Plants are sessile organisms that consistently face environmental challenges. Plants have developed elegant systems to respond to environmental signals and modify their metabolic activities to overcome these stresses. The available evidence clearly shows that an integrated regulatory network balances environmental responses, seed development, and FA biosynthesis. Although it has been recognized that various environmental cues affect seed oil biosynthesis and FA composition (Li et al., 2006; Singer et al., 2016; Karki and Bates, 2018), the molecular mechanisms that orchestrate these responses require further investigation. For instance, although many individual transcription factors in the regulatory network have been characterized, how they work in concert to perceive, transduce, and respond to environmental and developmental signals is poorly understood.

Manipulation of transcriptional regulators is a proven strategy for improving vegetable oil yield compared with the single-enzyme approach (Broun, 2004; Grotewold, 2008; Shen et al., 2010). Although various degrees of success have been achieved, a deeper understanding of the transcriptional regulation of plant FA biosynthesis is required to overcome the remaining challenges and obstacles. Since the identification of *AtWRI1* more than 15 years ago, a wealth of information suggests that *WRI1* plays an essential role in the transcriptional control of plant oil biosynthesis. However, how the key seed development regulators (LEC1, LEC2, FUS3, and ABI3) mediate *WRI1* expression remains to be fully elucidated. Whether additional upstream transcriptional regulators modulate *WRI1* expression, particularly in response to different environmental and developmental cues, is an important question to be addressed in the future. The recently discovered non-seed-specific transcription factors in oil palm that mediate *EgWRI1* expression beg the question of whether similar regulators exist in oil-producing tissues of other plant species (Yeap et al., 2017). To tackle this question, it is necessary to conduct more thorough analyses of *WRI1* promoters from diverse plant species; the resulting identification of transcription factor binding sites may shed light on additional regulators of *WRI1*. Recent functional and structural analyses of different *WRI1* orthologs and variants have concluded that the phosphorylation

residues, IDR3-PEST motif, interacting motif of BPMs, and 14-3-3s are pertinent to the stability and transcriptional activity of *WRI1* that affect plant oil accumulation (Ma et al., 2013, 2015, 2016; Zhai et al., 2017). The missing “VYL” motif in *AsWRI1c*, *RcWRI1-B*, and *OsWRI1-1* casts doubt on the functional significance of “VYL” (Ji et al., 2018; Mano et al., 2019; Yang et al., 2019). Therefore, solving the three-dimensional structure, together with computational modeling and simulation, will facilitate our understanding of the structure/function relationship of *WRI1*.

Recent advances suggest that the crosstalk between *WRI1*-interacting partners is regulated at the post-translational level, underlining the collaborative nature of the *WRI1* transcriptional apparatus. Although dual roles of phosphorylation in modulating the activity of *WRI1* have been proposed (Ma et al., 2016; Kong and Ma, 2018a), future work will focus on illustrating the molecular mechanisms by which *WRI1* activity is fine-tuned through phosphorylation and dephosphorylation by diverse kinases and phosphatases in response to a variety of developmental and environmental stimuli. The recently established *WRI1* interactome has contributed to the discovery of novel *WRI1*-interacting regulators (Figure 2A), opening new doors to the investigation of the molecular action of the regulatory network.

Alternative targets of *WRI1* beyond the oil biosynthetic pathway might be factors to consider in translational application of *WRI1* for oil bioengineering in crops. Nevertheless, numerous studies have demonstrated that *WRI1* is one of the most effective transcriptional regulators for the metabolic engineering of vegetable oil production. The use of protein engineering and genome editing approaches to improve *WRI1* binding specificity to genes in the oil biosynthetic pathway should be considered in future work on oil bioengineering.

The *AtWRI* gene family remains to be explored. *AtWRI3* and *AtWRI4* are *WRI1*-like genes capable of complementing the phenotype of the *wri1* mutant (To et al., 2012). The functions of *AtWRI3* and *AtWRI4* in transcriptional control of seed oil accumulation are not certain, as their genes are preferentially expressed in non-seed tissues (e.g., stems and flowers), despite their overlapping roles with *AtWRI1* in supplying acyl chains for cutin production (To et al., 2012). No obvious changes in plant growth were observed in the *wri1 wri3 wri4* triple mutant, and the expression of FA biosynthetic genes remained at a basal level in the triple mutant, leading to speculation that alternative transcriptional regulators may control FA biosynthetic genes in vegetative tissues (To et al., 2012; Marchive et al., 2014). Dissecting the distinct and overlapping roles of the *WRI1*-like proteins will contribute to our understanding of the complex transcriptional regulation of FA biosynthesis, particularly in vegetative tissues.

Seed oil biosynthesis involves hundreds of genes in multiple regulatory and metabolic pathways. How this dynamic process is regulated during seed development remains to be explored at the whole-genome and transcriptome levels. The development of population transcriptome analysis in recent years has led to the exploration of genes that regulate lipid metabolism. Using population transcriptome data from maize, expression



quantitative trait loci analysis revealed an intergenic locus that affected three oil-related genes, *long-chain fatty alcohol dehydrogenase (FADD)*, *glycosylphosphatidylinositol (GPI)*, and *glycolipid transfer protein 1 (GLTP1)*, resulting in variation in kernel oil concentration (Liu et al., 2017). Combining expression quantitative trait locus with co-expression networks helped to identify some key regulatory genes of lipid metabolism. The transcription factor *BrMD-2* was identified using transcriptomic data from a *B. rapa* doubled haploid population (Basnet et al., 2016). Recently, transcriptome-wide association studies were used to dissect oil biosynthesis in *B. napus*, and 690 genes, including *LPAAT*, *TT4*, and *C4H*, were identified as being associated with seed oil content (Tang et al., 2021). Thus, bioinformatics analysis based on transcriptome data from multiple tissues and multiple time series at the population level may provide more insights into the coordination of seed oil biosynthesis.

In summary, identification and characterization of essential regulatory factors that play roles in improving oil yield and agricultural traits are vital missions in vegetable oil research. To reach this goal, we require a profound understanding of the regulatory mechanisms that impact critical steps in the FA biosynthetic pathway and how these regulatory mechanisms are connected to environmental and developmental signaling. The knowledge gained in such an endeavor will guide our design to enhance vegetable oil yield in crops with integrated approaches that include genome editing, protein engineering, and metabolic engineering.

## FUNDING

This work was supported by Ministry of Education (MOE) of Singapore Tier 1 to W.M. (RG29/20), MOE of Singapore Tier 2 to W.M. (MOE-T2EP30220-0011), the National Key R&D Program of China to L.Y. (2019YFC1711100), the Hubei Hongshan Laboratory Research Fund to L.G. (2021HSZD004), and the HZAU-AGIS Cooperation Fund to L.G. (SZYJY2021004).

## AUTHOR CONTRIBUTIONS

Y.Y., Q.K., A.R.Q.L., L.G., L.Y., and W.M. conceived the ideas, wrote the first draft, and prepared the figures and tables. Q.K., S.L., H.Z., L.G., L.Y., and W.M. reviewed and edited the manuscript, figures, and tables. All authors have read and approved the final version of the manuscript.

## ACKNOWLEDGMENTS

We apologize to the authors whose important works are not cited because of space limitations. The authors declare no competing interests.

Received: November 5, 2021

Revised: February 28, 2022

Accepted: April 14, 2022

Published: April 20, 2022

## REFERENCES

- Adhikari, N.D., Bates, P.D., and Browse, J. (2016). WRINKLED1 rescues feedback inhibition of fatty acid synthesis in hydroxylase-expressing seeds. *Plant Physiol.* **171**:179–191. <https://doi.org/10.1104/pp.15.01906>.
- Allen, D.K. (2016). Assessing compartmentalized flux in lipid metabolism with isotopes. *Biochim. Biophys. Acta* **1861**:1226–1242. <https://doi.org/10.1016/j.bbalip.2016.03.017>.
- An, D., and Suh, M.C. (2015). Overexpression of Arabidopsis WRI1 enhanced seed mass and storage oil content in *Camelina sativa*. *Plant Biotechnol. Rep.* **9**:137–148. <https://doi.org/10.1007/s11816-015-0351-x>.

- Andrade, F.H., and Ferreiro, M.A. (1996). Reproductive growth of maize, sunflower and soybean at different source levels during grain filling. *Field Crops Res.* **48**:155–165. [https://doi.org/10.1016/s0378-4290\(96\)01017-9](https://doi.org/10.1016/s0378-4290(96)01017-9).
- Barthole, G., To, A., Marchive, C., Brunaud, V., Soubigou-Taconnat, L., Berger, N., Dubreucq, B., Lepiniec, L., and Baud, S. (2014). MYB118 represses endosperm maturation in seeds of Arabidopsis. *Plant Cell* **26**:3519–3537. <https://doi.org/10.1105/tpc.114.130021>.
- Basnet, R.K., Del Carpio, D.P., Xiao, D., Bucher, J., Jin, M., Boyle, K., Fobert, P., Visser, R.G., Maliepaard, C., and Bonnema, G. (2016). A systems genetics approach identifies gene regulatory networks associated with fatty acid composition in *Brassica rapa* seed. *Plant Physiol.* **170**:568–585. <https://doi.org/10.1104/pp.15.00853>.
- Bates, P.D. (2016). Understanding the control of acyl flux through the lipid metabolic network of plant oil biosynthesis. *Biochim. Biophys. Acta* **1861**:1214–1225. <https://doi.org/10.1016/j.bbalip.2016.03.021>.
- Bates, P.D., and Browse, J. (2011). The pathway of triacylglycerol synthesis through phosphatidylcholine in Arabidopsis produces a bottleneck for the accumulation of unusual fatty acids in transgenic seeds. *Plant J.* **68**:387–399. <https://doi.org/10.1111/j.1365-313X.2011.04693.x>.
- Bates, P.D., Stymne, S., and Ohlrogge, J. (2013). Biochemical pathways in seed oil synthesis. *Curr. Opin. Plant Biol.* **16**:358–364. <https://doi.org/10.1016/j.pbi.2013.02.015>.
- Bates, P.D., Johnson, S.R., Cao, X., Li, J., Nam, J.W., Jaworski, J.G., Ohlrogge, J.B., and Browse, J. (2014). Fatty acid synthesis is inhibited by inefficient utilization of unusual fatty acids for glycerolipid assembly. *Proc. Natl. Acad. Sci. U S A* **111**:1204–1209. <https://doi.org/10.1073/pnas.1318511111>.
- Baud, S., Mendoza, M.S., To, A., Harscoet, E., Lepiniec, L., and Dubreucq, B. (2007). WRINKLED1 specifies the regulatory action of LEAFY COTYLEDON2 towards fatty acid metabolism during seed maturation in Arabidopsis. *Plant J.* **50**:825–838. <https://doi.org/10.1111/j.1365-313X.2007.03092.x>.
- Baud, S., Feria Bourrellier, A.B., Azzopardi, M., Berger, A., Dechorgnat, J., Daniel-Vedele, F., Lepiniec, L., Miquel, M., Rochat, C., Hodges, M., et al. (2010). PII is induced by WRINKLED1 and fine-tunes fatty acid composition in seeds of Arabidopsis thaliana. *Plant J.* **64**:291–303. <https://doi.org/10.1111/j.1365-313X.2010.04332.x>.
- Baudry, A., Heim, M.A., Dubreucq, B., Caboche, M., Weisshaar, B., and Lepiniec, L. (2004). TT2, TT8, and TTG1 synergistically specify the expression of BANYULS and proanthocyanidin biosynthesis in Arabidopsis thaliana. *Plant J.* **39**:366–380. <https://doi.org/10.1111/j.1365-313X.2004.02138.x>.
- Bemer, M., van Dijk, A.D.J., Immink, R.G.H., and Angenent, G.C. (2017). Cross-family transcription factor interactions: an additional layer of gene regulation. *Trends Plant Sci.* **22**:66–80. <https://doi.org/10.1016/j.tplants.2016.10.007>.
- Bogamuwa, S., and Jang, J.C. (2013). The Arabidopsis tandem CCH zinc finger proteins AtTZF4, 5 and 6 are involved in light-abscisic acid- and gibberellic acid-mediated regulation of seed germination. *Plant Cell Environ.* **36**:1507–1519. <https://doi.org/10.1111/pce.12084>.
- Bonfante, P., and Genre, A. (2010). Mechanisms underlying beneficial plant-fungus interactions in mycorrhizal symbiosis. *Nat. Commun.* **1**:48. <https://doi.org/10.1038/ncomms1046>.
- Boulard, C., Fatihi, A., Lepiniec, L., and Dubreucq, B. (2017). Regulation and evolution of the interaction of the seed B3 transcription factors with NF-Y subunits. *Biochim. Biophys. Acta Gene Regul. Mech.* **1860**:1069–1078. <https://doi.org/10.1016/j.bbagrm.2017.08.008>.
- Bourgis, F., Kilaru, A., Cao, X., Ngando-Ebongue, G.F., Drira, N., Ohlrogge, J.B., and Arondel, V. (2011). Comparative transcriptome

- and metabolite analysis of oil palm and date palm mesocarp that differ dramatically in carbon partitioning. *Proc. Natl. Acad. Sci. U S A* **108**:12527–12532. <https://doi.org/10.1073/pnas.1106502108>.
- Broun, P.** (2004). Transcription factors as tools for metabolic engineering in plants. *Curr. Opin. Plant Biol.* **7**:202–209. <https://doi.org/10.1016/j.pbi.2004.01.013>.
- Carvin, D.T.** (1965). The effect of temperature on the oil content and fatty acid composition of oils from several oil seed crops. *Can. J. Bot.* **43**:63–69. <https://doi.org/10.1139/b65-008>.
- Cao, D., Cheng, H., Wu, W., Soo, H.M., and Peng, J.** (2006). Gibberellin mobilizes distinct DELLA-dependent transcriptomes to regulate seed germination and floral development in Arabidopsis. *Plant Physiol.* **142**:509–525. <https://doi.org/10.1104/pp.106.082289>.
- Carrera, C., Martínez, M.J., Dardanelli, J., and Mónica Balzarini, M.** (2009). Water deficit effect on the relationship between temperature during the seed fill period and soybean seed oil and protein concentrations. *Crop Sci.* **49**:990–998. <https://doi.org/10.2135/cropsci2008.06.0361>.
- Casson, S.A., and Lindsey, K.** (2006). The turnip mutant of Arabidopsis reveals that LEAFY COTYLEDON1 expression mediates the effects of auxin and sugars to promote embryonic cell identity. *Plant Physiol.* **142**:526–541. <https://doi.org/10.1104/pp.106.080895>.
- Cernac, A., and Benning, C.** (2004). WRINKLED1 encodes an AP2/EREB domain protein involved in the control of storage compound biosynthesis in Arabidopsis. *Plant J.* **40**:575–585. <https://doi.org/10.1111/j.1365-3113X.2004.02235.x>.
- Cernac, A., Andre, C., Hoffmann-Benning, S., and Benning, C.** (2006). WRI1 is required for seed germination and seedling establishment. *Plant Physiol.* **141**:745–757. <https://doi.org/10.1104/pp.106.079574>.
- Chapman, K.D., and Ohlrogge, J.B.** (2012). Compartmentation of triacylglycerol accumulation in plants. *J. Biol. Chem.* **287**:2288–2294. <https://doi.org/10.1074/jbc.R111.290072>.
- Chapman, K.D., Dyer, J.M., and Mullen, R.T.** (2012). Biogenesis and functions of lipid droplets in plants. *J. Lipid Res.* **53**:215–226. <https://doi.org/10.1194/jlr.R021436>.
- Chen, B., Zhang, G., Li, P., Yang, J., Guo, L., Benning, C., Wang, X., and Zhao, J.** (2020). Multiple GmWRI1s are redundantly involved in seed filling and modulation by regulating plastidic glycolysis, lipid biosynthesis and hormone signalling in soybean (*Glycine max*). *Plant Biotechnol. J.* **18**:155–171. <https://doi.org/10.1111/pbi.13183>.
- Chen, L., Lee, J.H., Weber, H., Tohge, T., Witt, S., Roje, S., Fernie, A.R., and Hellmann, H.** (2013). Arabidopsis BPM proteins function as substrate adaptors to a cullin3-based E3 ligase to affect fatty acid metabolism in plants. *Plant Cell* **25**:2253–2264. <https://doi.org/10.1105/tpc.112.107292>.
- Chen, L., Zheng, Y., Dong, Z., Meng, F., Sun, X., Fan, X., Zhang, Y., Wang, M., and Wang, S.** (2018a). Soybean (*Glycine max*) WRINKLED1 transcription factor, GmWRI1a, positively regulates seed oil accumulation. *Mol. Genet. Genomics* **293**:401–415. <https://doi.org/10.1007/s00438-017-1393-2>.
- Chen, M., Zhang, B., Li, C., Kulaveerasingam, H., Chew, F.T., and Yu, H.** (2015). TRANSPARENT TESTA GLABRA1 regulates the accumulation of seed storage reserves in arabidopsis. *Plant Physiol.* **169**:391–402. <https://doi.org/10.1104/pp.15.00943>.
- Chen, M., Wang, Z., Zhu, Y., Li, Z., Hussain, N., Xuan, L., Guo, W., Zhang, G., and Jiang, L.** (2012a). The effect of transparent TESTA2 on seed fatty acid biosynthesis and tolerance to environmental stresses during young seedling establishment in Arabidopsis. *Plant Physiol.* **160**:1023–1036. <https://doi.org/10.1104/pp.112.202945>.
- Chen, M., Xuan, L., Wang, Z., Zhou, L., Li, Z., Du, X., Ali, E., Zhang, G., and Jiang, L.** (2014). TRANSPARENT TESTA8 inhibits seed fatty acid accumulation by targeting several seed development regulators in arabidopsis. *Plant Physiol.* **165**:905–916. <https://doi.org/10.1104/pp.114.235507>.
- Chen, M., Du, X., Zhu, Y., Wang, Z., Hua, S., Li, Z., Guo, W., Zhang, G., Peng, J., and Jiang, L.** (2012b). Seed Fatty Acid Reducer acts downstream of gibberellin signalling pathway to lower seed fatty acid storage in Arabidopsis. *Plant Cell Environ.* **35**:2155–2169. <https://doi.org/10.1111/j.1365-3040.2012.02546.x>.
- Chen, N., Veerappan, V., Abdelmageed, H., Kang, M., and Allen, R.D.** (2018b). HSI2/VAL1 silences AGL15 to regulate the developmental transition from seed maturation to vegetative growth in arabidopsis. *Plant Cell* **30**:600–619. <https://doi.org/10.1105/tpc.17.00655>.
- Chen, Y.F., Li, L.Q., Xu, Q., Kong, Y.H., Wang, H., and Wu, W.H.** (2009). The WRKY6 transcription factor modulates PHOSPHATE1 expression in response to low Pi stress in Arabidopsis. *Plant Cell* **21**:3554–3566. <https://doi.org/10.1105/tpc.108.064980>.
- Cheng, Y., Zhu, W., Chen, Y., Ito, S., Asami, T., and Wang, X.** (2014). Brassinosteroids control root epidermal cell fate via direct regulation of a MYB-bHLH-WD40 complex by GSK3-like kinases. *Elife* **3**:e02525. <https://doi.org/10.7554/eLife.02525>.
- Cosio, C., Ranocha, P., Francoz, E., Burlat, V., Zheng, Y., Perry, S.E., Ripoll, J.J., Yanofsky, M., and Dunand, C.** (2017). The class III peroxidase PRX17 is a direct target of the MADS-box transcription factor AGAMOUS-LIKE15 (AGL15) and participates in lignified tissue formation. *New Phytol.* **213**:250–263. <https://doi.org/10.1111/nph.14127>.
- Covas, M.I., de la Torre, R., and Fito, M.** (2015). Virgin olive oil: a key food for cardiovascular risk protection. *Br. J. Nutr.* **113** (Suppl 2):S19–S28. <https://doi.org/10.1017/S0007114515000136>.
- Dahlqvist, A., Stahl, U., Lenman, M., Banas, A., Lee, M., Sandager, L., Ronne, H., and Stymne, S.** (2000). Phospholipid:diacylglycerol acyltransferase: an enzyme that catalyzes the acyl-CoA-independent formation of triacylglycerol in yeast and plants. *Proc. Natl. Acad. Sci. U S A* **97**:6487–6492. <https://doi.org/10.1073/pnas.120067297>.
- de la Vega, A.J., Chapman, S.C., and Hall, A.J.** (2001). Genotype by environment interaction and indirect selection for yield in sunflower. *Field Crops Res.* **72**:17–38. [https://doi.org/10.1016/s0378-4290\(01\)00162-9](https://doi.org/10.1016/s0378-4290(01)00162-9).
- Dean Rider, S., Jr., Henderson, J.T., Jerome, R.E., Edenberg, H.J., Romero-Severson, J., and Ogas, J.** (2003). Coordinate repression of regulators of embryonic identity by PICKLE during germination in Arabidopsis. *Plant J.* **35**:33–43. <https://doi.org/10.1046/j.1365-313x.2003.01783.x>.
- Debeaujon, I., Leon-Kloosterziel, K.M., and Koornneef, M.** (2000). Influence of the testa on seed dormancy, germination, and longevity in Arabidopsis. *Plant Physiol.* **122**:403–414. <https://doi.org/10.1104/pp.122.2.403>.
- Dekkers, B.J.W., Schuurmans, J.A.M.J., and Smeekens, S.C.M.** (2008). Interaction between sugar and abscisic acid signalling during early seedling development in Arabidopsis. *Plant Mol. Biol.* **67**:151–167. <https://doi.org/10.1007/s11103-008-9308-6>.
- Duan, S., Jin, C., Li, D., Gao, C., Qi, S., Liu, K., Hai, J., Ma, H., and Chen, M.** (2017). MYB76 inhibits seed fatty acid accumulation in arabidopsis. *Front. Plant Sci.* **8**:226. <https://doi.org/10.3389/fpls.2017.00226>.
- Eastmond, P., Koláček, L., and Rawsthorne, S.** (1996). Photosynthesis by developing embryos of oilseed rape (*Brassica napus* L.). *J. Exp. Bot.* **47**:1763–1769. <https://doi.org/10.1093/jxb/47.11.1763>.
- Eastmond, P.J., and Rawsthorne, S.** (1998). Comparison of the metabolic properties of plastids isolated from developing leaves or embryos of *Brassica napus* L. *J. Exp. Bot.* **49**:1105–1111. <https://doi.org/10.1093/jxb/49.324.1105>.
- Fathi, A., Bzierszak, A.M., and Dormann, P.** (2013). Alterations in seed development gene expression affect size and oil content of

- Arabidopsis seeds. *Plant Physiol.* **163**:973–985. <https://doi.org/10.1104/pp.113.226761>.
- Finkelstein, R.R., and Gibson, S.I. (2002). ABA and sugar interactions regulating development: cross-talk or voices in a crowd? *Curr. Opin. Plant Biol.* **5**:26–32. [https://doi.org/10.1016/s1369-5266\(01\)00225-4](https://doi.org/10.1016/s1369-5266(01)00225-4).
- Fleet, C.M., and Sun, T.P. (2005). A DELLAcate balance: the role of gibberellin in plant morphogenesis. *Curr. Opin. Plant Biol.* **8**:77–85. <https://doi.org/10.1016/j.pbi.2004.11.015>.
- Focks, N., and Benning, C. (1998). *wrinkled1*: a novel, low-seed-oil mutant of arabidopsis with a deficiency in the seed-specific regulation of carbohydrate Metabolism1. *Plant Physiol.* **118**:91–101. <https://doi.org/10.1104/pp.118.1.91>.
- Geilen, K., Heilmann, M., Hillmer, S., and Bohmer, M. (2017). WRKY43 regulates polyunsaturated fatty acid content and seed germination under unfavourable growth conditions. *Sci. Rep.* **7**:14235. <https://doi.org/10.1038/s41598-017-14695-0>.
- Giraudat, J., Hauge, B.M., Valon, C., Smalle, J., Parcy, F., and Goodman, H.M. (1992). Isolation of the Arabidopsis ABI3 gene by positional cloning. *Plant Cell* **4**:1251–1261. <https://doi.org/10.1105/tpc.4.10.1251>.
- Goffman, F.D., Alonso, A.P., Schwender, J., Shachar-Hill, Y., and Ohlrogge, J.B. (2005). Light enables a very high efficiency of carbon storage in developing embryos of rapeseed. *Plant Physiol.* **138**:2269–2279. <https://doi.org/10.1104/pp.105.063628>.
- Grimberg, A., Carlsson, A.S., Marttila, S., Bhalerao, R., and Hofvander, P. (2015). Transcriptional transitions in Nicotiana benthamiana leaves upon induction of oil synthesis by WRINKLED1 homologs from diverse species and tissues. *BMC Plant Biol.* **15**:192. <https://doi.org/10.1186/s12870-015-0579-1>.
- Grotewold, E. (2008). Transcription factors for predictive plant metabolic engineering: are we there yet? *Curr. Opin. Biotechnol.* **19**:138–144. <https://doi.org/10.1016/j.copbio.2008.02.002>.
- Guo, Q., Major, I.T., and Howe, G.A. (2018). Resolution of growth-defense conflict: mechanistic insights from jasmonate signaling. *Curr. Opin. Plant Biol.* **44**:72–81. <https://doi.org/10.1016/j.pbi.2018.02.009>.
- Hafner, A., Bulyk, M.L., Jambhekar, A., and Lahav, G. (2019). The multiple mechanisms that regulate p53 activity and cell fate. *Nat. Rev. Mol. Cell Biol.* **20**:199–210. <https://doi.org/10.1038/s41580-019-0110-x>.
- Harris, H.C., McWilliam, J.R., and Mason, W.K. (1978). Influence of temperature on oil content and composition of sunflower seed. *Aust. J. Agric. Res.* **29**:1203–1212. <https://doi.org/10.1071/ar9781203>.
- Horstman, A., Willemsen, V., Boutilier, K., and Heidstra, R. (2014). AINTEGUMENTA-LIKE proteins: hubs in a plethora of networks. *Trends Plant Sci.* **19**:146–157. <https://doi.org/10.1016/j.tplants.2013.10.010>.
- Horstman, A., Li, M., Heidmann, I., Weemen, M., Chen, B., Muino, J.M., Angenent, G.C., and Boutilier, K. (2017). The BABY BOOM transcription factor Activates the LEC1-ABI3-FUS3-LEC2 network to induce somatic embryogenesis. *Plant Physiol.* **175**:848–857. <https://doi.org/10.1104/pp.17.00232>.
- Huang, H., Liu, B., Liu, L., and Song, S. (2017). Jasmonate action in plant growth and development. *J. Exp. Bot.* **68**:1349–1359. <https://doi.org/10.1093/jxb/erw495>.
- Huang, R., Liu, M., Gong, G., Wu, P., Patra, B., Yuan, L., Qin, H., Wang, X., Wang, G., Liao, H., et al. (2021). The Pumilio RNA-binding protein APUM24 regulates seed maturation by fine-tuning the BPM-WRI1 module in Arabidopsis. *J. Integr. Plant Biol.* **63**:1240–1259. <https://doi.org/10.1111/jipb.13092>.
- Hung, C.Y., Lin, Y., Zhang, M., Pollock, S., David Marks, M., and Schiefelbein, J. (1998). A common position-dependent mechanism controls cell-type patterning and GLABRA2 regulation in the root and hypocotyl epidermis of Arabidopsis1. *Plant Physiol.* **117**:73–84. <https://doi.org/10.1104/pp.117.1.73>.
- Ji, X.J., Mao, X., Hao, Q.T., Liu, B.L., Xue, J.A., and Li, R.Z. (2018). Splice variants of the Castor WRI1 gene upregulate fatty acid and oil biosynthesis when expressed in tobacco leaves. *Int. J. Mol. Sci.* **19**:146. <https://doi.org/10.3390/ijms19010146>.
- Jiang, C., and Fu, X. (2007). GA action: turning on de-DELLA repressing signaling. *Curr. Opin. Plant Biol.* **10**:461–465. <https://doi.org/10.1016/j.pbi.2007.08.011>.
- Jiang, Y., Xie, Q., Wang, W., Yang, J., Zhang, X., Yu, N., Zhou, Y., and Wang, E. (2018). Medicago AP2-domain transcription factor WRI5a is a master regulator of lipid biosynthesis and transfer during mycorrhizal symbiosis. *Mol. Plant* **11**:1344–1359. <https://doi.org/10.1016/j.molp.2018.09.006>.
- Jo, L., Pelletier, J.M., and Harada, J.J. (2019). Central role of the LEAFY COTYLEDON1 transcription factor in seed development. *J. Integr. Plant Biol.* **61**:564–580. <https://doi.org/10.1111/jipb.12806>.
- Jones, A.L., Pruitt, N.L., Lloyd, D., and Harwood, J.L. (1991). Temperature-induced changes in the synthesis of unsaturated fatty acids by Acanthamoeba castellanii. *J. Protozoology* **38**:532–536. <https://doi.org/10.1111/j.1550-7408.1991.tb06076.x>.
- Joshi, S., Keller, C., and Perry, S.E. (2021). The EAR motif in the arabidopsis MADS transcription factor AGAMOUS-like 15 is not necessary to promote somatic embryogenesis. *Plants (Basel)* **10**:758. <https://doi.org/10.3390/plants10040758>.
- Kanai, M., Mano, S., Kondo, M., Hayashi, M., and Nishimura, M. (2016). Extension of oil biosynthesis during the mid-phase of seed development enhances oil content in Arabidopsis seeds. *Plant Biotechnol. J.* **14**:1241–1250. <https://doi.org/10.1111/pbi.12489>.
- Karki, N., and Bates, P.D. (2018). The effect of light conditions on interpreting oil composition engineering in Arabidopsis seeds. *Plant Direct.* **2**:e00067. <https://doi.org/10.1002/pld3.67>.
- Kim, D.H., Yamaguchi, S., Lim, S., Oh, E., Park, J., Hanada, A., Kamiya, Y., and Choi, G. (2008). SOMNUS, a CCCH-type zinc finger protein in Arabidopsis, negatively regulates light-dependent seed germination downstream of PIL5. *Plant Cell* **20**:1260–1277. <https://doi.org/10.1105/tpc.108.058859>.
- Kim, H.U., Lee, K.R., Jung, S.J., Shin, H.A., Go, Y.S., Suh, M.C., and Kim, J.B. (2015). Senescence-inducible LEC2 enhances triacylglycerol accumulation in leaves without negatively affecting plant growth. *Plant Biotechnol. J.* **13**:1346–1359. <https://doi.org/10.1111/pbi.12354>.
- Kim, M.J., Jang, I.C., and Chua, N.H. (2016). The mediator complex MED15 subunit mediates activation of downstream lipid-related genes by the WRINKLED1 transcription factor. *Plant Physiol.* **171**:1951–1964. <https://doi.org/10.1104/pp.16.00664>.
- Kim, R.J., Kim, H.U., and Suh, M.C. (2019). Development of camelina enhanced with drought stress resistance and seed oil production by co-overexpression of MYB96A and DGAT1C. *Ind. Crops Prod.* **138**:111475. <https://doi.org/10.1016/j.indcrop.2019.111475>.
- Kong, Q., and Ma, W. (2018a). WRINKLED1 as a novel 14-3-3 client: function of 14-3-3 proteins in plant lipid metabolism. *Plant Signal. Behav.* **13**:e1482176. <https://doi.org/10.1080/15592324.2018.1482176>.
- Kong, Q., and Ma, W. (2018b). WRINKLED1 transcription factor: how much do we know about its regulatory mechanism? *Plant Sci.* **272**:153–156. <https://doi.org/10.1016/j.plantsci.2018.04.013>.
- Kong, Q., Yuan, L., and Ma, W. (2019). WRINKLED1, a "master regulator" in transcriptional control of plant oil biosynthesis. *Plants (Basel)* **8**:238. <https://doi.org/10.3390/plants8070238>.



- Kong, Q., Yang, Y., Guo, L., Yuan, L., and Ma, W. (2020a). Molecular basis of plant oil biosynthesis: insights gained from studying the WRINKLED1 transcription factor. *Front. Plant Sci.* **11**:24. <https://doi.org/10.3389/fpls.2020.00024>.
- Kong, Q., Ma, W., Yang, H., Ma, G., Mantyla, J.J., and Benning, C. (2017). The Arabidopsis WRINKLED1 transcription factor affects auxin homeostasis in roots. *J. Exp. Bot.* **68**:4627–4634. <https://doi.org/10.1093/jxb/erx275>.
- Kong, Q., Yang, Y., Low, P.M., Guo, L., Yuan, L., and Ma, W. (2020b). The function of the WRI1-TCP4 regulatory module in lipid biosynthesis. *Plant Signal. Behav.* **15**:1812878. <https://doi.org/10.1080/15592324.2020.1812878>.
- Kong, Q., Low, P.M., Lim, A.R.Q., Yang, Y., Yuan, L., and Ma, W. (2022). Functional antagonism of WRI1 and TCP20 modulates GH3.3 expression to maintain auxin homeostasis in roots. *Plants (Basel)* **11**:454. <https://doi.org/10.3390/plants11030454>.
- Kong, Q., Singh, S.K., Mantyla, J.J., Pattanaik, S., Guo, L., Yuan, L., Benning, C., and Ma, W. (2020c). TEOSINTE BRANCHED1/CYCLOIDEA/PROLIFERATING CELL FACTOR4 interacts with WRINKLED1 to mediate seed oil biosynthesis. *Plant Physiol.* **184**:658–665. <https://doi.org/10.1104/pp.20.00547>.
- Kong, Y., Chen, S., Yang, Y., and An, C. (2013). ABA-insensitive (ABI) 4 and ABI5 synergistically regulate DGAT1 expression in Arabidopsis seedlings under stress. *FEBS Lett.* **587**:3076–3082. <https://doi.org/10.1016/j.febslet.2013.07.045>.
- Kubota, A., Ito, S., Shim, J.S., Johnson, R.S., Song, Y.H., Breton, G., Goraloglia, G.S., Kwon, M.S., Laboy Cintron, D., Koyama, T., et al. (2017). TCP4-dependent induction of CONSTANS transcription requires GIGANTEA in photoperiodic flowering in Arabidopsis. *Plos Genet.* **13**:e1006856. <https://doi.org/10.1371/journal.pgen.1006856>.
- Lambert, S.A., Jolma, A., Campitelli, L.F., Das, P.K., Yin, Y., Albu, M., Chen, X., Taipale, J., Hughes, T.R., and Weirauch, M.T. (2018). The human transcription factors. *Cell* **172**:650–665. <https://doi.org/10.1016/j.cell.2018.01.029>.
- Lavy, M., and Estelle, M. (2016). Mechanisms of auxin signaling. *Development* **143**:3226–3229. <https://doi.org/10.1242/dev.131870>.
- Le Hir, R., and Bellini, C. (2013). The plant-specific dof transcription factors family: new players involved in vascular system development and functioning in Arabidopsis. *Front. Plant Sci.* **4**:164. <https://doi.org/10.3389/fpls.2013.00164>.
- Lee, H.G., Park, B.Y., Kim, H.U., and Seo, P.J. (2015). MYB96 stimulates C18 fatty acid elongation in Arabidopsis seeds. *Plant Biotechnol. Rep.* **9**:161–166. <https://doi.org/10.1007/s11816-015-0352-9>.
- Lee, H.G., Kim, H., Suh, M.C., Kim, H.U., and Seo, P.J. (2018). The MYB96 transcription factor regulates triacylglycerol accumulation by activating DGAT1 and PDAT1 expression in arabidopsis seeds. *Plant Cell Physiol.* **59**:1432–1442. <https://doi.org/10.1093/pcp/pcy073>.
- Li-Beisson, Y., Shorrosh, B., Beisson, F., Andersson, M.X., Arondel, V., Bates, P.D., Baud, S., Bird, D., Debono, A., Durrett, T.P., et al. (2013). Acyl-lipid metabolism. *Arabidopsis Book* **11**:e0161. <https://doi.org/10.1199/tab.0161>.
- Li, D., Jin, C., Duan, S., Zhu, Y., Qi, S., Liu, K., Gao, C., Ma, H., Zhang, M., Liao, Y., et al. (2017). MYB89 transcription factor represses seed oil accumulation. *Plant Physiol.* **173**:1211–1225. <https://doi.org/10.1104/pp.16.01634>.
- Li, Y., Beisson, F., Pollard, M., and Ohlrogge, J. (2006). Oil content of Arabidopsis seeds: the influence of seed anatomy, light and plant-to-plant variation. *Phytochemistry* **67**:904–915. <https://doi.org/10.1016/j.phytochem.2006.02.015>.
- Lim, A.R.Q., Kong, Q., Singh, S.K., Guo, L., Yuan, L., and Ma, W. (2022). Sunflower WRINKLED1 plays a key role in transcriptional regulation of oil biosynthesis. *Int. J. Mol. Sci.* **23**:3054. <https://doi.org/10.3390/ijms23063054>.
- Lin, P.C., Pomeranz, M.C., Jikumaru, Y., Kang, S.G., Hah, C., Fujioka, S., Kamiya, Y., and Jang, J.C. (2011). The Arabidopsis tandem zinc finger protein AtTZF1 affects ABA- and GA-mediated growth, stress and gene expression responses. *Plant J.* **65**:253–268. <https://doi.org/10.1111/j.1365-3113X.2010.04419.x>.
- Liu, H., Zhai, Z., Kuczynski, K., Keereetaweep, J., Schwender, J., and Shanklin, J. (2019). WRINKLED1 regulates BIOTIN ATTACHMENT DOMAIN-CONTAINING proteins that inhibit fatty acid synthesis. *Plant Physiol.* **181**:55–62. <https://doi.org/10.1104/pp.19.00587>.
- Liu, H., Luo, X., Niu, L., Xiao, Y., Chen, L., Liu, J., Wang, X., Jin, M., Li, W., Zhang, Q., et al. (2017). Distant eQTLs and non-coding sequences play critical roles in regulating gene expression and quantitative trait variation in maize. *Mol. Plant* **10**:414–426. <https://doi.org/10.1016/j.molp.2016.06.016>.
- Liu, J., Hua, W., Zhan, G., Wei, F., Wang, X., Liu, G., and Wang, H. (2010). Increasing seed mass and oil content in transgenic Arabidopsis by the overexpression of wri1-like gene from Brassica napus. *Plant Physiol. Biochem.* **48**:9–15. <https://doi.org/10.1016/j.plaphy.2009.09.007>.
- Liu, J., Deng, S., Wang, H., Ye, J., Wu, H.W., Sun, H.X., and Chua, N.H. (2016). CURLY LEAF regulates gene sets coordinating seed size and lipid biosynthesis. *Plant Physiol.* **171**:424–436. <https://doi.org/10.1104/pp.15.01335>.
- Liu, Y.F., Li, Q.T., Lu, X., Song, Q.X., Lam, S.M., Zhang, W.K., Ma, B., Lin, Q., Man, W.Q., Du, W.G., et al. (2014). Soybean GmMYB73 promotes lipid accumulation in transgenic plants. *BMC Plant Biol.* **14**:73. <https://doi.org/10.1186/1471-2229-14-73>.
- Lotan, T., Ohto, M.a., Yee, K.M., West, M.A., Lo, R., Kwong, R.W., Yamagishi, K., Fischer, R.L., Goldberg, R.B., and Harada, J.J. (1998). Arabidopsis LEAFY COTYLEDON1 is sufficient to induce embryo development in vegetative cells. *Cell* **93**:1195–1205. [https://doi.org/10.1016/S0092-8674\(00\)81463-4](https://doi.org/10.1016/S0092-8674(00)81463-4).
- Lu, C., and Hills, M.J. (2002). Arabidopsis mutants deficient in diacylglycerol acyltransferase display increased sensitivity to abscisic acid, sugars, and osmotic stress during germination and seedling development. *Plant Physiol.* **129**:1352–1358. <https://doi.org/10.1104/pp.006122>.
- Lu, L., Wei, W., Li, Q.T., Bian, X.H., Lu, X., Hu, Y., Cheng, T., Wang, Z.Y., Jin, M., Tao, J.J., et al. (2021). A transcriptional regulatory module controls lipid accumulation in soybean. *New Phytol.* **231**:661–678. <https://doi.org/10.1111/nph.17401>.
- Lu, S., Sturtevant, D., Aziz, M., Jin, C., Li, Q., Chapman, K.D., and Guo, L. (2018). Spatial analysis of lipid metabolites and expressed genes reveals tissue-specific heterogeneity of lipid metabolism in high- and low-oil Brassica napus L. seeds. *Plant J.* **94**:915–932. <https://doi.org/10.1111/tbj.13959>.
- Luerßen, H., Kirik, V., Herrmann, P., and Misera, S. (1998). FUSCA3 encodes a protein with a conserved VP1/ABI3-like B3 domain which is of functional importance for the regulation of seed maturation in Arabidopsis thaliana. *Plant J.* **15**:755–764. <https://doi.org/10.1046/j.1365-3113x.1998.00259.x>.
- Lutz, K.A., Martin, C., Khairzada, S., and Maliga, P. (2015). Steroid-inducible BABY BOOM system for development of fertile Arabidopsis thaliana plants after prolonged tissue culture. *Plant Cell Rep.* **34**:1849–1856. <https://doi.org/10.1007/s00299-015-1832-7>.
- Ma, W., Kong, Q., Mantyla, J.J., Yang, Y., Ohlrogge, J.B., and Benning, C. (2016). 14-3-3 protein mediates plant seed oil biosynthesis through interaction with AtWRI1. *Plant J.* **88**:228–235. <https://doi.org/10.1111/tbj.13244>.

- Ma, W., Kong, Q., Grix, M., Mantyla, J.J., Yang, Y., Benning, C., and Ohlrogge, J.B. (2015). Deletion of a C-terminal intrinsically disordered region of WRINKLED1 affects its stability and enhances oil accumulation in Arabidopsis. *Plant J.* **83**:864–874. <https://doi.org/10.1111/tpj.12933>.
- Ma, W., Kong, Q., Arondel, V., Kilaru, A., Bates, P.D., Thrower, N.A., Benning, C., and Ohlrogge, J.B. (2013). Wrinkled1, a ubiquitous regulator in oil accumulating tissues from Arabidopsis embryos to oil palm mesocarp. *PLoS One* **8**:e68887. <https://doi.org/10.1371/journal.pone.0068887>.
- Maeo, K., Tokuda, T., Ayame, A., Mitsui, N., Kawai, T., Tsukagoshi, H., Ishiguro, S., and Nakamura, K. (2009). An AP2-type transcription factor, WRINKLED1, of Arabidopsis thaliana binds to the AW-box sequence conserved among proximal upstream regions of genes involved in fatty acid synthesis. *Plant J.* **60**:476–487. <https://doi.org/10.1111/j.1365-3113X.2009.03967.x>.
- Manan, S., Ahmad, M.Z., Zhang, G., Chen, B., Haq, B.U., Yang, J., and Zhao, J. (2017). Soybean LEC2 regulates subsets of genes involved in controlling the biosynthesis and catabolism of seed storage substances and seed development. *Front. Plant Sci.* **8**:1604. <https://doi.org/10.3389/fpls.2017.01604>.
- Mano, F., Aoyanagi, T., and Kozaki, A. (2019). Atypical splicing accompanied by skipping conserved micro-exons produces unique WRINKLED1, an AP2 domain transcription factor in rice plants. *Plants (Basel)* **8**:207. <https://doi.org/10.3390/plants8070207>.
- Marchise, C., Nikovics, K., To, A., Lepiniec, L., and Baud, S. (2014). Transcriptional regulation of fatty acid production in higher plants: molecular bases and biotechnological outcomes. *Eur. J. Lipid Sci. Technol.* **116**:1332–1343. <https://doi.org/10.1002/ejlt.201400027>.
- Masaki, T., Mitsui, N., Tsukagoshi, H., Nishii, T., Morikami, A., and Nakamura, K. (2005). ACTIVATOR of Spomin::LUC1/WRINKLED1 of Arabidopsis thaliana transactivates sugar-inducible promoters. *Plant Cell Physiol.* **46**:547–556. <https://doi.org/10.1093/pcp/pci072>.
- Masucci, J.D., Rerie, W.G., Foreman, D.R., Zhang, M., Galway, M.E., Marks, M.D., and Schiefelbein, J.W. (1996). The homeobox gene GLABRA2 is required for position-dependent cell differentiation in the root epidermis of Arabidopsis thaliana. *Development* **122**:1253–1260. <https://doi.org/10.1242/dev.122.4.1253>.
- Meinke, D.W., Franzmann, L.H., Nickle, T.C., and Yeung, E.C. (1994). Leafy cotyledon mutants of arabidopsis. *Plant Cell* **6**:1049–1064. <https://doi.org/10.1105/tpc.6.8.1049>.
- Memelink, J., and Gantet, P. (2007). Transcription factors involved in terpenoid indole alkaloid biosynthesis in Catharanthus roseus. *Phytochemistry Rev.* **6**:353–362. <https://doi.org/10.1007/s11101-006-9051-z>.
- Mendes, A., Kelly, A.A., van Erp, H., Shaw, E., Powers, S.J., Kurup, S., and Eastmond, P.J. (2013). bZIP67 regulates the omega-3 fatty acid content of Arabidopsis seed oil by activating fatty acid desaturase3. *Plant Cell* **25**:3104–3116. <https://doi.org/10.1105/tpc.113.116343>.
- Miray, R., Kazaz, S., To, A., and Baud, S. (2021). Molecular control of oil metabolism in the endosperm of seeds. *Int. J. Mol. Sci.* **22**:1621. <https://doi.org/10.3390/ijms22041621>.
- Monke, G., Seifert, M., Keilwagen, J., Mohr, M., Grosse, I., Hahnel, U., Junker, A., Weisshaar, B., Conrad, U., Baumlein, H., et al. (2012). Toward the identification and regulation of the Arabidopsis thaliana ABI3 regulon. *Nucleic Acids Res.* **40**:8240–8254. <https://doi.org/10.1093/nar/gks594>.
- Mouchiroud, L., Eichner, L.J., Shaw, R.J., and Auwerx, J. (2014). Transcriptional coregulators: fine-tuning metabolism. *Cell Metab.* **20**:26–40. <https://doi.org/10.1016/j.cmet.2014.03.027>.
- Mu, J., Tan, H., Zheng, Q., Fu, F., Liang, Y., Zhang, J., Yang, X., Wang, T., Chong, K., Wang, X.J., et al. (2008). LEAFY COTYLEDON1 is a key regulator of fatty acid biosynthesis in Arabidopsis. *Plant Physiol.* **148**:1042–1054. <https://doi.org/10.1104/pp.108.126342>.
- Nesi, N., Jond, C., Debeaujon, I., Caboche, M., and Lepiniec, L. (2001). The arabidopsis TT2 gene encodes and R2R3 MYB domain protein that acts as a key determinant for proanthocyanidin accumulation in developing seed. *Plant Cell* **13**:2099–2114. <https://doi.org/10.2307/3871430>.
- Niu, Y., Wu, G.Z., Ye, R., Lin, W.H., Shi, Q.M., Xue, L.J., Xu, X.D., Li, Y., Du, Y.G., and Xue, H.W. (2009). Global analysis of gene expression profiles in Brassica napus developing seeds reveals a conserved lipid metabolism regulation with Arabidopsis thaliana. *Mol. Plant* **2**:1107–1122. <https://doi.org/10.1093/mp/ssp042>.
- Ogas, J., Cheng, J.C., Sung, Z.R., and Somerville, C. (1997). Cellular differentiation regulated by gibberellin in the Arabidopsis thaliana pickle mutant. *Science* **277**:91–94. <https://doi.org/10.1126/science.277.5322.91>.
- Ogas, J., Kaufmann, S., Henderson, J., and Somerville, C. (1999). PICKLE is a CHD3 chromatin-remodeling factor that regulates the transition from embryonic to vegetative development in Arabidopsis. *Proc. Natl. Acad. Sci. U S A* **96**:13839–13844. <https://doi.org/10.1073/pnas.96.24.13839>.
- Parcy, F., Valon, C., Kohara, A., Misera, S., and Giraudat, J. (1997). The ABSCISIC ACID-INSENSITIVE3, FUSCA3, and LEAFY COTYLEDON1 loci act in concert to control multiple aspects of Arabidopsis seed development. *Plant Cell* **9**:1265–1277. <https://doi.org/10.2307/3870380>.
- Park, J., Oh, D.H., Dassanayake, M., Nguyen, K.T., Ogas, J., Choi, G., and Sun, T.P. (2017). Gibberellin signaling requires chromatin remodeler PICKLE to promote vegetative growth and phase transitions. *Plant Physiol.* **173**:1463–1474. <https://doi.org/10.1104/pp.16.01471>.
- Pelletier, J.M., Kwong, R.W., Park, S., Le, B.H., Baden, R., Cagliari, A., Hashimoto, M., Munoz, M.D., Fischer, R.L., Goldberg, R.B., et al. (2017). LEC1 sequentially regulates the transcription of genes involved in diverse developmental processes during seed development. *Proc. Natl. Acad. Sci. U S A* **114**:E6710–E6719. <https://doi.org/10.1073/pnas.1707957114>.
- Pireyre, M., and Burow, M. (2015). Regulation of MYB and bHLH transcription factors: a glance at the protein level. *Mol. Plant* **8**:378–388. <https://doi.org/10.1016/j.molp.2014.11.022>.
- Ramsay, N.A., and Glover, B.J. (2005). MYB-bHLH-WD40 protein complex and the evolution of cellular diversity. *Trends Plant Sci.* **10**:63–70. <https://doi.org/10.1016/j.tplants.2004.12.011>.
- Remy, W., Taylor, T.N., Hass, H., and Kerp, H. (1994). Four hundred-million-year-old vesicular arbuscular mycorrhizae. *Proc. Natl. Acad. Sci. U S A* **91**:11841–11843. <https://doi.org/10.1073/pnas.91.25.11841>.
- Rerie, W.G., Feldmann, K.A., and Marks, M.D. (1994). The GLABRA2 gene encodes a homeo domain protein required for normal trichome development in Arabidopsis. *Genes Dev.* **8**:1388–1399. <https://doi.org/10.1101/gad.8.12.1388>.
- Rich, M.K., Vigneron, N., Libourel, C., Keller, J., Xue, L., Hajheidari, M., Radhakrishnan, G.V., Le Ru, A., Diop, S.I., Potente, G., et al. (2021). Lipid exchanges drove the evolution of mutualism during plant terrestrialization. *Science* **372**:864–868. <https://doi.org/10.1126/science.abg0929>.
- Rohde, A., Kurup, S., and Holdsworth, M. (2000). ABI3 emerges from the seed. *Trends Plant Sci.* **5**:418–419. [https://doi.org/10.1016/s1360-1385\(00\)01736-2](https://doi.org/10.1016/s1360-1385(00)01736-2).
- Roscoe, T.T., Guilleminot, J., Bessoule, J.J., Berger, F., and Devic, M. (2015). Complementation of seed maturation phenotypes by ectopic expression of ABSCISIC ACID INSENSITIVE3, FUSCA3 and LEAFY

- COTYLEDON2 in arabidopsis. *Plant Cell Physiol.* **56**:1215–1228. <https://doi.org/10.1093/pcp/pcv049>.
- Ruta, V., Longo, C., Lepri, A., De Angelis, V., Occhigrossi, S., Costantino, P., and Vittorioso, P. (2020). The DOF transcription factors in seed and seedling development. *Plants (Basel)* **9**:218. <https://doi.org/10.3390/plants9020218>.
- Rutter, A.J., Thomas, K.L., Herbert, D., Henderson, R.J., Lloyd, D., and Harwood, J.L. (2002). Oxygen induction of a novel fatty acid n-6 desaturase in the soil protozoan, *Acanthamoeba castellanii*. *Biochem. J.* **368**:57–67. <https://doi.org/10.1042/BJ20020189>.
- Ruuska, S.A., Schwender, J., and Ohlrogge, J.B. (2004). The capacity of green oilseeds to utilize photosynthesis to drive biosynthetic processes. *Plant Physiol.* **136**:2700–2709. <https://doi.org/10.1104/pp.104.047977>.
- Ruuska, S.A., Girke, T., Benning, C., and Ohlrogge, J.B. (2002). Contrapuntal networks of gene expression during arabidopsis seed filling[W]. *Plant Cell* **14**:1191–1206. <https://doi.org/10.1105/tpc.000877>.
- Sakr, S., Wang, M., Dedaldechamp, F., Perez-Garcia, M.D., Oge, L., Hamama, L., and Atanassova, R. (2018). The sugar-signaling hub: overview of regulators and interaction with the hormonal and metabolic network. *Int. J. Mol. Sci.* **19**:2506. <https://doi.org/10.3390/ijms19092506>.
- Sanjaya, Durrett, T.P., Weise, S.E., and Benning, C. (2011). Increasing the energy density of vegetative tissues by diverting carbon from starch to oil biosynthesis in transgenic Arabidopsis. *Plant Biotechnol. J.* **9**:874–883. <https://doi.org/10.1111/j.1467-7652.2011.00599.x>.
- Santos Mendoza, M., Dubreucq, B., Miquel, M., Caboche, M., and Lepiniec, L. (2005). LEAFY COTYLEDON 2 activation is sufficient to trigger the accumulation of oil and seed specific mRNAs in Arabidopsis leaves. *FEBS Lett.* **579**:4666–4670. <https://doi.org/10.1016/j.febslet.2005.07.037>.
- Sasaki, A., Itoh, H., Gomi, K., Ueguchi-Tanaka, M., Ishiyama, K., Kobayashi, M., Jeong, D.H., An, G., Kitano, H., Ashikari, M., et al. (2003). Accumulation of phosphorylated repressor for gibberellin signaling in an F-box mutant. *Science* **299**:1896–1898. <https://doi.org/10.1126/science.1081077>.
- Schulte, L.R., Ballard, T., Samarakoon, T., Yao, L., Vadlani, P., Staggenborg, S., and Rezac, M. (2013). Increased growing temperature reduces content of polyunsaturated fatty acids in four oilseed crops. *Ind. Crops Prod.* **51**:212–219. <https://doi.org/10.1016/j.indcrop.2013.08.075>.
- Schwender, J., Goffman, F., Ohlrogge, J.B., and Shachar-Hill, Y. (2004). Rubisco without the Calvin cycle improves the carbon efficiency of developing green seeds. *Nature* **432**:779–782. <https://doi.org/10.1038/nature03145>.
- Seiler, G.J. (1986). Analysis of the relationships of environmental factors with seed oil and fatty acid concentrations of wild annual sunflower. *Field Crops Res.* **15**:57–72. [https://doi.org/10.1016/0378-4290\(86\)90101-2](https://doi.org/10.1016/0378-4290(86)90101-2).
- Serivichyaswat, P., Ryu, H.S., Kim, W., Kim, S., Chung, K.S., Kim, J.J., and Ahn, J.H. (2015). Expression of the floral repressor miRNA156 is positively regulated by the AGAMOUS-like proteins AGL15 and AGL18. *Mol. Cells* **38**:259–266. <https://doi.org/10.14348/molcells.2015.2311>.
- Shen, B., Sinkevicius, K.W., Selinger, D.A., and Tarczynski, M.C. (2006). The homeobox gene GLABRA2 affects seed oil content in Arabidopsis. *Plant Mol. Biol.* **60**:377–387. <https://doi.org/10.1007/s11103-005-4110-1>.
- Shen, B., Allen, W.B., Zheng, P., Li, C., Glassman, K., Ranch, J., Nubel, D., and Tarczynski, M.C. (2010). Expression of ZmLEC1 and ZmWRI1 increases seed oil production in maize. *Plant Physiol.* **153**:980–987. <https://doi.org/10.1104/pp.110.157537>.
- Shi, L., Katavic, V., Yu, Y., Kunst, L., and Haughn, G. (2012). Arabidopsis glabra2 mutant seeds deficient in mucilage biosynthesis produce more oil. *Plant J.* **69**:37–46. <https://doi.org/10.1111/j.1365-3113.2011.04768.x>.
- Singer, S.D., Zou, J., and Weselake, R.J. (2016). Abiotic factors influence plant storage lipid accumulation and composition. *Plant Sci.* **243**:1–9. <https://doi.org/10.1016/j.plantsci.2015.11.003>.
- Song, G., Li, X., Munir, R., Khan, A.R., Azhar, W., Yasin, M.U., Jiang, Q., Bancroft, I., and Gan, Y. (2020). The WRKY6 transcription factor affects seed oil accumulation and alters fatty acid compositions in Arabidopsis thaliana. *Physiol. Plant* **169**:612–624. <https://doi.org/10.1111/ppl.13082>.
- Song, J., Xie, X., Chen, C., Shu, J., Thapa, R.K., Nguyen, V., Bian, S., Kohalmi, S.E., Marsolais, F., Zou, J., et al. (2021a). LEAFY COTYLEDON1 expression in the endosperm enables embryo maturation in Arabidopsis. *Nat. Commun.* **12**:3963. <https://doi.org/10.1038/s41467-021-24234-1>.
- Song, J.M., Zhang, Y., Zhou, Z.W., Lu, S., Ma, W., Lu, C., Chen, L.L., and Guo, L. (2021b). Oil plant genomes: current state of the science. *J. Exp. Bot.* **10**:1093. <https://doi.org/10.1093/jxb/erab472>.
- Song, Q.X., Li, Q.T., Liu, Y.F., Zhang, F.X., Ma, B., Zhang, W.K., Man, W.Q., Du, W.G., Wang, G.D., Chen, S.Y., et al. (2013). Soybean GmbZIP123 gene enhances lipid content in the seeds of transgenic Arabidopsis plants. *J. Exp. Bot.* **64**:4329–4341. <https://doi.org/10.1093/jxb/ert238>.
- Stone, S.L., Kwong, L.W., Yee, K.M., Pelletier, J., Lepiniec, L., Fischer, R.L., Goldberg, R.B., and Harada, J.J. (2001). LEAFY COTYLEDON2 encodes a B3 domain transcription factor that induces embryo development. *Proc. Natl. Acad. Sci. U S A* **98**:11806–11811. <https://doi.org/10.1073/pnas.201413498>.
- Sui, X., Singh, S.K., Patra, B., Schluttenhofer, C., Guo, W., Pattanaik, S., and Yuan, L. (2018). Cross-family transcription factor interaction between MYC2 and GBFs modulates terpenoid indole alkaloid biosynthesis. *J. Exp. Bot.* **69**:4267–4281. <https://doi.org/10.1093/jxb/ery229>.
- Sun, R., Ye, R., Gao, L., Zhang, L., Wang, R., Mao, T., Zheng, Y., Li, D., and Lin, Y. (2017). Characterization and ectopic expression of CoWRI1, an AP2/EREBP domain-containing transcription factor from coconut (*Cocos nucifera* L.) endosperm, changes the seeds oil content in transgenic Arabidopsis thaliana and rice (*oryza sativa* L.). *Front. Plant Sci.* **8**:63. <https://doi.org/10.3389/fpls.2017.00063>.
- Swain, S.M., and Singh, D.P. (2005). Tall tales from sly dwarves: novel functions of gibberellins in plant development. *Trends Plant Sci.* **10**:123–129. <https://doi.org/10.1016/j.tplants.2005.01.007>.
- Taatjes, D.J. (2010). The human Mediator complex: a versatile, genome-wide regulator of transcription. *Trends Biochem. Sci.* **35**:315–322. <https://doi.org/10.1016/j.tibs.2010.02.004>.
- Tan, H., Yang, X., Zhang, F., Zheng, X., Qu, C., Mu, J., Fu, F., Li, J., Guan, R., Zhang, H., et al. (2011). Enhanced seed oil production in canola by conditional expression of Brassica napus LEAFY COTYLEDON1 and LEC1-LIKE in developing seeds. *Plant Physiol.* **156**:1577–1588. <https://doi.org/10.1104/pp.111.175000>.
- Tang, S., Zhao, H., Lu, S., Yu, L., Zhang, G., Zhang, Y., Yang, Q.Y., Zhou, Y., Wang, X., Ma, W., et al. (2021). Genome- and transcriptome-wide association studies provide insights into the genetic basis of natural variation of seed oil content in Brassica napus. *Mol. Plant* **14**:470–487. <https://doi.org/10.1016/j.molp.2020.12.003>.
- Thakare, D., Tang, W., Hill, K., and Perry, S.E. (2008). The MADS-domain transcriptional regulator AGAMOUS-LIKE15 promotes



- somatic embryo development in Arabidopsis and soybean. *Plant Physiol.* **146**:1663–1672. <https://doi.org/10.1104/pp.108.115832>.
- Thomas, S.G., and Sun, T.P. (2004). Update on gibberellin signaling. A tale of the tall and the short. *Plant Physiol.* **135**:668–676. <https://doi.org/10.1104/pp.104.040279>.
- Tian, R., Wang, F., Zheng, Q., Niza, V.M.A.G.E., Downie, A.B., and Perry, S.E. (2020). Direct and indirect targets of the arabidopsis seed transcription factor ABCISIC ACID INSENSITIVE3. *Plant J.* **103**:1679–1694. <https://doi.org/10.1111/tpj.14854>.
- Tiedemann, J., Rutten, T., Monke, G., Vorwieger, A., Rolletschek, H., Meissner, D., Milkowski, C., Petereck, S., Mock, H.P., Zank, T., et al. (2008). Dissection of a complex seed phenotype: novel insights of FUSCA3 regulated developmental processes. *Dev. Biol.* **317**:1–12. <https://doi.org/10.1016/j.ydbio.2008.01.034>.
- To, A., Joubes, J., Barthole, G., Lecureuil, A., Scagnelli, A., Jasinski, S., Lepiniec, L., and Baud, S. (2012). WRINKLED transcription factors orchestrate tissue-specific regulation of fatty acid biosynthesis in arabidopsis. *Plant Cell* **24**:5007–5023. <https://doi.org/10.1105/tpc.112.106120>.
- Tominaga-Wada, R., Ishida, T., and Wada, T. (2011). New insights into the mechanism of development of Arabidopsis root hairs and trichomes. *Int. Rev. Cell Mol. Biol.* **286**:67–106. <https://doi.org/10.1016/B978-0-12-385859-7.00002-1>.
- Tsuchiya, Y., Nambara, E., Naito, S., and McCourt, P. (2004). The FUS3 transcription factor functions through the epidermal regulator TTG1 during embryogenesis in Arabidopsis. *Plant J.* **37**:73–81. <https://doi.org/10.1046/j.1365-3113x.2003.01939.x>.
- Vanhercke, T., El Tahchy, A., Shrestha, P., Zhou, X.R., Singh, S.P., and Petrie, J.R. (2013). Synergistic effect of WRI1 and DGAT1 coexpression on triacylglycerol biosynthesis in plants. *FEBS Lett.* **587**:364–369. <https://doi.org/10.1016/j.febslet.2012.12.018>.
- Vanhercke, T., Dyer, J.M., Mullen, R.T., Kilaru, A., Rahman, M.M., Petrie, J.R., Green, A.G., Yurchenko, O., and Singh, S.P. (2019). Metabolic engineering for enhanced oil in biomass. *Prog. Lipid Res.* **74**:103–129. <https://doi.org/10.1016/j.plipres.2019.02.002>.
- Vanhercke, T., El Tahchy, A., Liu, Q., Zhou, X.R., Shrestha, P., Divi, U.K., Ral, J.P., Mansour, M.P., Nichols, P.D., James, C.N., et al. (2014). Metabolic engineering of biomass for high energy density: oilseed-like triacylglycerol yields from plant leaves. *Plant Biotechnol. J.* **12**:231–239. <https://doi.org/10.1111/pbi.12131>.
- Verdier, J., and Thompson, R.D. (2008). Transcriptional regulation of storage protein synthesis during dicotyledon seed filling. *Plant Cell Physiol.* **49**:1263–1271. <https://doi.org/10.1093/pcp/pcn116>.
- Walker, A.R., Davison, P.A., Bolognesi-Winfield, A.C., James, C.M., Srinivasan, N., Blundell, T.L., Esch, J.J., Marks, M.D., and Gray, J.C. (1999). The TRANSPARENT TESTA GLABRA1 locus, which regulates trichome differentiation and anthocyanin biosynthesis in Arabidopsis, encodes a WD40 repeat protein. *Plant Cell* **11**:1337–1350. <https://doi.org/10.2307/3870753>.
- Wang, F., and Perry, S.E. (2013). Identification of direct targets of FUSCA3, a key regulator of Arabidopsis seed development. *Plant Physiol.* **161**:1251–1264. <https://doi.org/10.1104/pp.112.212282>.
- Wang, H.W., Zhang, B., Hao, Y.J., Huang, J., Tian, A.G., Liao, Y., Zhang, J.S., and Chen, S.Y. (2007). The soybean Dof-type transcription factor genes, GmDof4 and GmDof11, enhance lipid content in the seeds of transgenic Arabidopsis plants. *Plant J.* **52**:716–729. <https://doi.org/10.1111/j.1365-3113x.2007.03268.x>.
- Weselake, R.J., Shah, S., Tang, M., Quant, P.A., Snyder, C.L., Furukawa-Stoffer, T.L., Zhu, W., Taylor, D.C., Zou, J., Kumar, A., et al. (2008). Metabolic control analysis is helpful for informed genetic manipulation of oilseed rape (*Brassica napus*) to increase seed oil content. *J. Exp. Bot.* **59**:3543–3549. <https://doi.org/10.1093/jxb/ern206>.
- Western, T.L., Burn, J., Tan, W.L., Skinner, D.J., Martin-McCaffrey, L., Moffatt, B.A., and Haughn, G.W. (2001). Isolation and characterization of mutants defective in seed coat mucilage secretory cell development in Arabidopsis. *Plant Physiol.* **127**:998–1011. <https://doi.org/10.1104/pp.010410>.
- Wolf, R.B., Cavins, J.F., Kleiman, R., and Black, L.T. (1982). Effect of temperature on soybean seed constituents: oil, protein, moisture, fatty acids, amino acids and sugars. *J. Am. Oil Chemists' Soc.* **59**:230–232. <https://doi.org/10.1007/bf02582182>.
- Woodward, A.W., and Bartel, B. (2005). Auxin: regulation, action, and interaction. *Ann. Bot.* **95**:707–735. <https://doi.org/10.1093/aob/mci083>.
- Xu, C., and Shanklin, J. (2016). Triacylglycerol metabolism, function, and accumulation in plant vegetative tissues. *Annu. Rev. Plant Biol.* **67**:179–206. <https://doi.org/10.1146/annurev-arplant-043015-111641>.
- Xue, L., Klinnawee, L., Zhou, Y., Saridis, G., Vijayakumar, V., Brands, M., Dormann, P., Gigolashvili, T., Turck, F., and Bucher, M. (2018). AP2 transcription factor CBX1 with a specific function in symbiotic exchange of nutrients in mycorrhizal *Lotus japonicus*. *Proc. Natl. Acad. Sci. U S A* **115**:E9239–E9246. <https://doi.org/10.1073/pnas.1812275115>.
- Yamamoto, A., Kagaya, Y., Usui, H., Hobo, T., Takeda, S., and Hattori, T. (2010). Diverse roles and mechanisms of gene regulation by the Arabidopsis seed maturation master regulator FUS3 revealed by microarray analysis. *Plant Cell Physiol.* **51**:2031–2046. <https://doi.org/10.1093/pcp/pcq162>.
- Yanagisawa, S. (2002). The Dof family of plant transcription factors. *Trends Plant Sci.* **7**:555–560. [https://doi.org/10.1016/s1360-1385\(02\)02362-2](https://doi.org/10.1016/s1360-1385(02)02362-2).
- Yang, Y., Munz, J., Cass, C., Zienkiewicz, A., Kong, Q., Ma, W., Sedbrook, J., and Benning, C. (2015). Ectopic expression of WRINKLED1 affects fatty acid homeostasis in *Brachypodium distachyon* vegetative tissues. *Plant Physiol.* **169**:1836–1847. <https://doi.org/10.1104/pp.15.01236>.
- Yang, Z., Liu, X., Wang, K., Li, Z., Jia, Q., Zhao, C., and Zhang, M. (2021). ABI3 with or without FUS3 highly up-regulates lipid droplet proteins and activates oil accumulation. *J. Exp. Bot.* **73**:2077–2092. <https://doi.org/10.1093/jxb/erab524>.
- Yang, Z., Liu, X., Li, N., Du, C., Wang, K., Zhao, C., Wang, Z., Hu, Y., and Zhang, M. (2019). WRINKLED1 homologs highly and functionally express in oil-rich endosperms of oat and castor. *Plant Sci.* **287**:110193. <https://doi.org/10.1016/j.plantsci.2019.110193>.
- Ye, J., Wang, C., Sun, Y., Qu, J., Mao, H., and Chua, N.H. (2018a). Overexpression of a transcription factor increases lipid content in a Woody perennial *Jatropha curcas*. *Front. Plant Sci.* **9**:1479. <https://doi.org/10.3389/fpls.2018.01479>.
- Ye, Q., Wang, H., Su, T., Wu, W.H., and Chen, Y.F. (2018b). The ubiquitin E3 ligase PRU1 regulates WRKY6 degradation to modulate phosphate homeostasis in response to low-pi stress in arabidopsis. *Plant Cell* **30**:1062–1076. <https://doi.org/10.1105/tpc.17.00845>.
- Ye, Y., Nikovics, K., To, A., Lepiniec, L., Fedosejevs, E.T., Van Doren, S.R., Baud, S., and Thelen, J.J. (2020). Docking of acetyl-CoA carboxylase to the plastid envelope membrane attenuates fatty acid production in plants. *Nat. Commun.* **11**:6191. <https://doi.org/10.1038/s41467-020-20014-5>.
- Yeap, W.C., Lee, F.C., Shabari Shan, D.K., Musa, H., Appleton, D.R., and Kulaveerasingam, H. (2017). WRI1-1, ABI5, NF-YA3 and NF-YC2 increase oil biosynthesis in coordination with hormonal signaling

- during fruit development in oil palm. *Plant J.* **91**:97–113. <https://doi.org/10.1111/tpj.13549>.
- Zale, J., Jung, J.H., Kim, J.Y., Pathak, B., Karan, R., Liu, H., Chen, X., Wu, H., Candreva, J., Zhai, Z., et al.** (2016). Metabolic engineering of sugarcane to accumulate energy-dense triacylglycerols in vegetative biomass. *Plant Biotechnol. J.* **14**:661–669. <https://doi.org/10.1111/pbi.12411>.
- Zentella, R., Zhang, Z.L., Park, M., Thomas, S.G., Endo, A., Murase, K., Fleet, C.M., Jikumaru, Y., Nambara, E., Kamiya, Y., et al.** (2007). Global analysis of *della* direct targets in early gibberellin signaling in *Arabidopsis*. *Plant Cell* **19**:3037–3057. <https://doi.org/10.1105/tpc.107.054999>.
- Zha, P., Liu, S., Li, Y., Ma, T., Yang, L., Jing, Y., and Lin, R.** (2020). The evening complex and the chromatin-remodeling factor PICKLE coordinately control seed dormancy by directly repressing *DOG1* in *arabidopsis*. *Plant Commun.* **1**:100011. <https://doi.org/10.1016/j.xplc.2019.100011>.
- Zhai, Z., Liu, H., and Shanklin, J.** (2017). Phosphorylation of WRINKLED1 by KIN10 results in its proteasomal degradation, providing a link between energy homeostasis and lipid biosynthesis. *Plant Cell* **29**:871–889. <https://doi.org/10.1105/tpc.17.00019>.
- Zhai, Z., Keereetaweep, J., Liu, H., Feil, R., Lunn, J.E., and Shanklin, J.** (2018). Trehalose 6-phosphate positively regulates fatty acid synthesis by stabilizing WRINKLED1. *Plant Cell* **30**:2616–2627. <https://doi.org/10.1105/tpc.18.00521>.
- Zhang, H., Bishop, B., Ringenberg, W., Muir, W.M., and Ogas, J.** (2012). The CHD3 remodeler PICKLE associates with genes enriched for trimethylation of histone H3 lysine 27. *Plant Physiol.* **159**:418–432. <https://doi.org/10.1104/pp.112.194878>.
- Zhang, M., Fan, J., Taylor, D.C., and Ohlrogge, J.B.** (2009). DGAT1 and PDAT1 acyltransferases have overlapping functions in *Arabidopsis* triacylglycerol biosynthesis and are essential for normal pollen and seed development. *Plant cell* **21**:3885–3901. <https://doi.org/10.1105/tpc.109.071795>.
- Zhang, M., Cao, X., Jia, Q., and Ohlrogge, J.** (2016). FUSCA3 activates triacylglycerol accumulation in *Arabidopsis* seedlings and tobacco BY2 cells. *Plant J.* **88**:95–107. <https://doi.org/10.1111/tpj.13233>.
- Zhang, Y., Liu, Z., Wang, X., Wang, J., Fan, K., Li, Z., and Lin, W.** (2018). DELLA proteins negatively regulate dark-induced senescence and chlorophyll degradation in *Arabidopsis* through interaction with the transcription factor WRKY6. *Plant Cell Rep.* **37**:981–992. <https://doi.org/10.1007/s00299-018-2282-9>.
- Zhao, Y.** (2010). Auxin biosynthesis and its role in plant development. *Annu. Rev. Plant Biol.* **61**:49–64. <https://doi.org/10.1146/annurev-arplant-042809-112308>.
- Zheng, Y., Ren, N., Wang, H., Stromberg, A.J., and Perry, S.E.** (2009). Global identification of targets of the *Arabidopsis* MADS domain protein AGAMOUS-Like15. *Plant Cell* **21**:2563–2577. <https://doi.org/10.1105/tpc.109.068890>.
- Zhu, F., Farnung, L., Kaasinen, E., Sahu, B., Yin, Y., Wei, B., Dodonova, S.O., Nitta, K.R., Morgunova, E., Taipale, M., et al.** (2018). The interaction landscape between transcription factors and the nucleosome. *Nature* **562**:76–81. <https://doi.org/10.1038/s41586-018-0549-5>.
- Zou, J., Wei, Y., Jako, C., Kumar, A., Selvaraj, G., and Taylor, D.C.** (1999). The *Arabidopsis thaliana* TAG1 mutant has a mutation in a diacylglycerol acyltransferase gene. *Plant J.* **19**:645–653. <https://doi.org/10.1046/j.1365-313x.1999.00555.x>.