

# Global Transcriptome Profiling of the Pine Shoot Beetle, *Tomicus yunnanensis* (Coleoptera: Scolytinae)

Jia-Ying Zhu<sup>1\*</sup>, Ning Zhao<sup>2</sup>, Bin Yang<sup>1\*</sup>

**1** Key Laboratory of Forest Disaster Warning and Control of Yunnan Province, College of Forestry, Southwest Forestry University, Kunming, China, **2** College of Life Sciences, Southwest Forestry University, Kunming, China

## Abstract

**Background:** The pine shoot beetle *Tomicus yunnanensis* (Coleoptera: Scolytinae) is an economically important pest of *Pinus yunnanensis* in southwestern China. Developed resistance to insecticides due to chemical pesticides being used for a long time is a factor involved in its serious damage, which poses a challenge for management. In addition, highly efficient adaptation to divergent environmental ecologies results in this pest posing great potential threat to pine forests. However, the molecular mechanisms remain unknown as only limited nucleotide sequence data for this species is available.

**Methodology/Principal Findings:** In this study, we applied next generation sequencing (Illumina sequencing) to sequence the adult transcriptome of *T. yunnanensis*. A total of 51,822,230 reads were obtained. They were assembled into 140,702 scaffolds, and 60,031 unigenes. The unigenes were further functionally annotated with gene descriptions, Gene Ontology (GO), Clusters of Orthologous Groups (COG), and Kyoto Encyclopedia of Genes and Genome (KEGG). In total, 80,932 unigenes were classified into GO, 13,599 unigenes were assigned to COG, and 33,875 unigenes were found in KO categories. A biochemical pathway database containing 219 predicted pathways was also created based on the annotations. In depth analysis of the data revealed a large number of genes related to insecticides resistance and heat shock protein genes associated with environmental stress.

**Conclusions/Significance:** The results facilitate the investigations of molecular resistance mechanisms to insecticides and environmental stress. This study lays the foundation for future functional genomics studies of important biological questions of this pest.

**Citation:** Zhu J-Y, Zhao N, Yang B (2012) Global Transcriptome Profiling of the Pine Shoot Beetle, *Tomicus yunnanensis* (Coleoptera: Scolytinae). PLoS ONE 7(2): e32291. doi:10.1371/journal.pone.0032291

**Editor:** Christian Schönbach, Kyushu Institute of Technology, Japan

**Received:** December 15, 2011; **Accepted:** January 25, 2012; **Published:** February 23, 2012

**Copyright:** © 2012 Zhu et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Funding:** This work was supported by the Special Fund for Forest Scientific Research in the Public Interest of China (201004067), Science Foundation for the Excellent Youth Academic and Technological Scholars of Yunnan Province of China (2007YP01-18), Science Foundation of Southwest Forestry University (111128), and the Key Subject of Forest Protection of Yunnan Province of China (XKZ200905). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

**Competing Interests:** The authors have declared that no competing interests exist.

\* E-mail: jyzhu001@gmail.com (J-YZ); yangbin48053@yahoo.com.cn (BY)

## Introduction

The bark beetle genus *Tomicus* (Coleoptera: Curculionidae: Scolytinae) comprises seven described species [1]. Among them, *T. piniperda* and *T. minor* have a Palearctic distribution from Europe to Japan and China and repeatedly introduced to North America, *T. destruens* is found only around the Mediterranean Basin, while *T. yunnanensis*, *T. brevipilosus*, *T. pilifer* and *T. puellus* are restricted to southwestern, eastern, and central China [2]. *Tomicus* damages various pine species mainly due to its shoot feeding behavior and trunk stem attacking, which is among the most damaging pests of pine forests in the outbreak countries. Especially, some species of *Tomicus* have evolved towards different ecological strategies of host use, wider range distribution, and high levels of resistance to major classes of insecticides owing to biological and genetic factors. They are as exotic pests frequently invaded into new areas. *Tomicus* imposes a potentially serious damage upon pine forest.

Having a globally economical importance and ecological threat, it has been studied well from general biology and ecological viewpoints [3–5]. But it has remained poorly investigated at the

molecular level. Only few studies have been developed, leading to elucidate their population genetic and species evolution [2,6]. Up to September 26, 2011, there are only 719 nucleotide sequences deposited on National Center for Biotechnology Information (NCBI) for *Tomicus*. Due to insufficient genetic background, the molecular mechanisms of *Tomicus* in forest ecological systems are poorly understood.

The next generation sequencing (high-throughput deep sequencing) technology recently enables efficient approach on large-scale and genome-wide for gene discovery and expression profiling, and studies in functional, comparative and evolutionary genomics in non-model organisms with little or no previous genomic information exists [7,8]. In the past few years, several studies based on this technology have, indeed, allowed the efficient, massive and successful molecular mechanisms investigation of some insect species lacking genome information, such as *Bemisia tabaci*, *Nilaparvata lugens*, and *Trialeurodes vaporariorum* [9–11].

*T. yunnanensis* is a newly described pine shoot beetle only found in Yunnan, Sichuan, and Guizhou Provinces in southwestern China, and has affected 200,000 ha. of *Pinus yunnanensis* forests

over the past 30 years [12]. Control of this pest until now depended largely on chemical insecticides. It has evolved high level of resistance to insecticides. This might be due to the increasing of metabolic capability of detoxificative systems and/or reducing target site sensitivity. In addition, this pest has been found distributed in divergent environmental ecologies. The distributional areas of *T. yunnanensis* are with high mounts and deep valley as geographical barriers, and its ecological environments are in large difference. It suggested that *T. yunnanensis* adapted efficiently to environmental stress for surviving in diversifiable ecologies during the past years. Even resistance of *T. yunnanensis* to insecticides and severe environmental stress is an ongoing challenge for pest management, there is no information available for uncovering the molecular mechanisms under these.

In the present work, we characterized the first global transcriptome using Illumina technology of that *Tomicus* species *T. yunnanensis*. A systematic bioinformatics strategy was engaged to functional annotation of the transcriptome data. Additionally, important homologues of genes involved in insecticide resistance and heat shock protein (HSP) genes associated with environmental stress were identified. This study dramatically provides a foundation and increases the significant promise for further functional genomics studies of *T. yunnanensis* and other *Tomicus* species.

**Materials and Methods**

**Ethics statement**

Regarding the field study, no specific permits were required. The location is not privately-owned or protected in any way. The field studies did not involve endangered or protected species.

**Insects**

Adult beetles identified based on morphological characters [1] as *T. yunnanensis* by their trunk attacking phase on *P. yunnanensis* were collected from Qujing city, Yunnan province, China. The samples were frozen at  $-80^{\circ}\text{C}$  until use.

**cDNA library and Illumina sequencing**

Total RNA was extracted from each 20 female and male adult beetles using TRIzol<sup>®</sup> Reagent (Invitrogen) according to the manufacturer’s instructions. RNA quality and yield were assessed by 2100 Bioanalyzer (Agilent Technologies) with a minimum RNA integrated number value of 8. The cDNA library was prepared using Illumina’s kit following manufacturer’s recommendations. Briefly, messenger RNA was isolated from 20  $\mu\text{g}$  total RNA (pooled RNA of female and male adults) using oligo(dT) magnetic beads and fragmented into short sequences using divalent cations under elevated temperature. First and second strand cDNA were synthesized from cleaved RNA fragments. After the end repair and ligation of adaptors, the products were cleaned up with a QIAquick PCR Purification Kit to create the final cDNA library. The library was sequenced on the Illumina sequencing platform (GAII) to obtain short sequences from both ends.

**Bioinformatic analysis**

The raw reads from the images and quality value calculation were performed by the Illumina data processing pipeline (version 1.6). Before the assembly, the raw reads were cleaned by adaptor sequences, empty reads and low quality sequences (reads with unknown sequences ‘N’) to obtain the high-quality clean reads. Raw reads were then assembled into sequence contigs, scaffolds, and unigenes using SOAPdenovo software [13], and clustered using TGI Clustering tools [14]. All Illumina assembled unigenes

were searched against nr database in NCBI, Swiss-Prot, Kyoto Encyclopedia of Genes and Genome (KEGG), and Cluster of Orthologous Groups (COG) with the BLASTX algorithm. The E-value cut-off was set at  $10^{-5}$ . Genes were tentatively identified according to the best hits against known sequences. Blast2GO [15] was used to predict the functions of the sequences, assign Gene Ontology (GO) terms, and predict the metabolic pathways in COG and KEGG databases. Amino acid sequence comparisons were conducted with Clustal X (v1.83) program [16]. Phylogenetic tree was constructed by MEGA 5.0 package [17] using the neighbour-joining method with the Poisson correction model after sequence alignment performed by Clustalx. Bootstrap analysis of 1,000 replication trees was performed in order to evaluate the branch strength of each tree.

**Data availability**

The Illumina reads of *T. yunnanensis* have been submitted to NCBI Short Read Archive under the accession number of SRA047283.

**Results and Discussion**


**Sequencing and *de novo* assembly**

Illumina sequencing resulted in 51,822,230 raw reads, corresponding to an accumulated length of 4,664,000,700 bp (Table 1). The average raw reads length is 90 bp, which is consistent with the Illumina sequencing capacity. Using SOAPdenovo software, the raw reads were assembled into contigs after adaptor sequences, empty reads and low quality sequences filtered out. The raw reads were assembled into 642,521 contigs with a mean length of 117 bp. The range length of contigs is from 50 to 2953 bp. Although the majority of the contigs are between 50 to 400 bp, we obtained 9,331 contigs which were greater than 400 bp in length. The size distribution of these contigs is shown in Figure S1. Using paired end-joining and gap-filling, these contigs were further assembled into 140,702 scaffolds with a mean length of 233 bp, and range length of 100–2953 bp. 2,293 scaffolds were longer than 800 bp (Figure S2). Using TGI software, scaffold sequences were assembled into clusters. We obtained 60,031 unigenes with a mean length of 355 bp. The lengths of the 8,953 and 1,055 unigenes were  $\geq 500$  bp and 1000 bp, respectively, revealing that 85% of them fell between 100 and 500 bp in length (Figure 1). The result demonstrated that the scaffold and unigene length distribution followed the contig length distribution closely, with the majority being shorter sequences with relatively little redundancy, which was in a similar to other insect transcriptome projects using this technology [9–11,18,19]. The majority of scaffolds and unigenes after assembly were still less than 500 bp, which might be due to the short length sequencing capacity of Illumina sequencing and/or the low coverage of the transcriptome represented in this dataset [20]. The assembled abundant sequence data provided a rich source of information for further

**Table 1.** Sequence statistics of the Illumina sequencing assembly.

	Reads	Contigs	scaffolds	Unigenes
Number of sequences	51,822,230	642,521	140,702	60,031
Mean length (bp)	90	117	233	355
Total length (bp)	4,664,000,700	74,951,788	32,749,102	21,338,135

doi:10.1371/journal.pone.0032291.t001



**Figure 1. Length distribution of unigenes.** The number of y-axis has been transfer into logarithmic scale. doi:10.1371/journal.pone.0032291.g001

investigation, thus allowing for rapid characterization of a large portion of the transcriptome and better reference of interesting genes.


**Annotation of predicted proteins**

The assembled unigenes were used as a query for Blastx searches in the NCBI nr protein database with a cut-off E-value of


$10^{-5}$ . The search produced 34,702 hits, which comprised 57.81% of all the unigenes (Table S1). A large proportion of them (about 40%) apparently have no significant match in any of the existing databases, indicating many of them may contain novel sequences and a high number of Coleoptera or species-specific transcripts or transcript parts (orphan UTRs). This is expected, as there is very little sequence information from closely related species. The E-value distribution of the top hits in the nr database showed that 17% of the mapped sequences have strong homology (smaller than  $1.0E^{-49}$ ), whereas 83% of the homolog sequences ranged between  $1.0E^{-5}$  to  $1.0E^{-49}$  (Figure 2A). The species distribution of the best match result for each sequence showed that the *T. yunnanensis* sequences have 62.48% matches with sequences from the Coleoptera species (*Tribolium castaneum*), while very low proportion (<5%) of them have matches to other insects (Figure 2B). It demonstrated that *T. yunnanensis* have near evolution distance with *T. castaneum*.

**GO assignments**

Gene ontology is widely used to standardize representation of genes across species and provide a controlled vocabulary of terms for describing gene products [21]. In total, 80,932 unigenes were assigned for GO terms based on BlastX matches with sequences whose function is previously known (Figure 3, Table S2). These GO terms were summarized into the 3 main GO categories (biological process, cellular component, and molecular function)



**Figure 2. Characteristics of homology search of Illumina sequences against the nr database.** (A) E-value distribution of BLAST hits for each unique sequence with a cut-off E-value of  $1.0E^{-5}$ . (B) Species distribution of the BLASTX results. We used the first hit of each sequence for analysis. doi:10.1371/journal.pone.0032291.g002



**Figure 3. Distribution of second level GO of *Tomicus yunnanensis* transcriptome.** (A) biological process, (B) cellular component and (C) molecular function. The percentage of a specific category of genes in that main category is shown. doi:10.1371/journal.pone.0032291.g003

according to the standard GO terms and 47 sub-categories. Compared to the GO annotations of *Drosophila melanogaster* genome [22], our sequence data do not contain any notable biases towards particular categories of genes. Biological process made up the majority of the GO annotations (38,578, 47.67% of the total), followed by cellular component (26,119, 32.27%), and molecular

function (16,235, 20.06%). Among biological process category, cellular process (17.97%), and metabolic process (15.12%) were the most dominant subcategories, reflecting that the analyzed tissues were undergoing rapid growth and extensive metabolic activities. The following subcategories were multicellular organismal process (8.54%), developmental process (8.21%), biological

regulation (7.85%), localization (6.88%), and cellular component organization (5.81%). The biological process illustrated all of the major cellular processes from transport and cellular organization to transcription, translation, and metabolism. Under the category of cellular component, cell (29.91%), cell part (29.91%), and organelle (17.75%) were among the most highly represented subcategories. Most of the unigenes annotated with a cellular component are localized to plastids or mitochondria. The molecular function category was mainly comprised of proteins involved in binding (46.54%), predominantly heat shock proteins (Hsp), and catalytic activities (36.91%) including hydrolases, kinases, and transferases, allowing for the identification of genes involved in the secondary metabolite synthesis pathways. Similar observations for metabolic processes were reported in transcriptomic studies of other insects [18]. These GO annotations represent a general gene expression profile signature for *T. yunnanensis* adults, which demonstrates that the expressed genes in this species encode diverse structural, regulatory and stress proteins.

**COG classification**

A total of 13,599 unigenes were assigned to the appropriate COG clusters (Figure 4). These COG classifications were grouped into 25 function categories that correspond to the categories observed in GO analysis. The cluster for general function prediction only (16.10%) represents the largest group. The other three largest categories include: (1) translation, ribosomal structure and biogenesis (9.85%), (2) posttranslational modification, protein turnover, chaperones (8.54%), and (3) replication, recombination and repair (7.54%). The category of secondary metabolites biosynthesis, transport and catabolism was highlighted with


2.86%, because of the importance of secondary metabolites to the insecticides in insects. The most abundant sequences in this category are cytochrome P450 monooxygenases. To some extent, the COG classifications shed light on specific responses and functions involved in the molecular processes of *T. yunnanensis*.

**KEGG analysis**

To identify the biological pathways that are active in *T. yunnanensis*, we mapped the 34,702 annotated sequences to the referential canonical pathways in KEGG. A total of 16,727 unigenes were assigned to 219 KEGG pathways. All the pathways are summarized in Table S3. The top 10 pathways are metabolic pathways (3339 members), spliceosome (665 members), huntington’s disease (663 members), pathways in cancer (586 members), lysosome (527 members), purine metabolism (509), regulation of actin cytoskeleton (472 members), alzheimer’s disease (471 members), ubiquitin mediated proteolysis (467 members), and focal adhesion (466 members). These annotations provide a valuable resource for investigating specific processes, functions and pathways in *T. yunnanensis* research.

**Putative insecticides resistance related genes**

**Cytochrome P450 (P450).** Because of the genetic diversity, broad substrate specificity, and catalytic versatility, P450s can mediate resistance to all classes of insecticides [23]. Approximately a total of 146 P450 related unigenes were identified in the transcriptome. Even 23 *T. yunnanensis* P450 sequences were identified in the dataset as length longer than 600 bp (Table 2), and the majority of them were as short fragments listed in Table S4. In general, insect genomes harbor ~100 different P450s, which can be divided into four clades (CYP2, CYP3, CYP4 and



**Figure 4. COG function classification of the *Tomiscus yunnanensis* transcriptome.** All putative proteins were aligned to the COG database and can be classified functionally into at least 25 molecular families. doi:10.1371/journal.pone.0032291.g004

**Table 2.** Putatively identified P450 genes in *Tomicus yunnanensis*.

Unigene ID	Putative assignment	Length (bp)	First hit	Identity (%)	E_value	Blast annotation/Organism
Unigene12883	CYP3	612	XP_975568	44	2.00E-28	PREDICTED: similar to cytochrome P450 [ <i>Tribolium castaneum</i> ]
Unigene15066	CYP4	819	EFA01330	58	9.00E-59	cytochrome P450-like protein [ <i>Tribolium castaneum</i> ]
Unigene1549	CYP4	679	AAT38513	42	8.00E-43	ubiquitous cytochrome P450 [ <i>Phyllopertha diversa</i> ]
Unigene1697	CYP3	902	EFA04564	39	5.00E-56	cytochrome P450 347A1 [ <i>Tribolium castaneum</i> ]
Unigene21111	CYP3	771	XP_972794	56	9.00E-34	PREDICTED: similar to Probable cytochrome P450 9f2 (CYP1XF2) [ <i>Tribolium castaneum</i> ]
Unigene22371	CYP4	694	XP_001814854	45	1.00E-45	PREDICTED: similar to cytochrome P450 [ <i>Tribolium castaneum</i> ]
Unigene23650	mito	645	XP_974252	60	7.00E-65	PREDICTED: similar to CYP302a1 [ <i>Tribolium castaneum</i> ]
Unigene23993	CYP2	1092	NP_503303	38	2.00E-57	Cytochrome P450 family member (cyp-33C11) [ <i>Caenorhabditis elegans</i> ]
Unigene24457	CYP3	1192	EEZ99338	62	2.00E-63	cytochrome P450 6BQ13 [ <i>Tribolium castaneum</i> ]
Unigene24875	CYP2	1173	XP_968477	53	1.00E-98	PREDICTED: similar to cytochrome P450, partial [ <i>Tribolium castaneum</i> ]
Unigene551	mito	640	NP_001123894	57	2.00E-54	cytochrome P450 CYP314A1 [ <i>Tribolium castaneum</i> ]
Unigene58533	CYP4	603	ABF06546	67	2.00E-44	CYP4BD1 [ <i>Ips paraconfusus</i> ]
Unigene58724	CYP3	627	NP_001164248	50	5.00E-50	cytochrome P450 9Z4 [ <i>Tribolium castaneum</i> ]
Unigene58733	CYP3	628	NP_001164248	47	4.00E-51	cytochrome P450 9Z4 [ <i>Tribolium castaneum</i> ]
Unigene59248	CYP4	720	ABF06544	46	5.00E-59	CYP4AY1 [ <i>Ips paraconfusus</i> ]
Unigene59398	CYP4	760	XP_973153	45	1.00E-44	PREDICTED: similar to cytochrome P450 [ <i>Tribolium castaneum</i> ]
Unigene59490	CYP4	788	XP_001602395	36	7.00E-41	PREDICTED: similar to cytochrome P450 [ <i>Nasonia vitripennis</i> ]
Unigene59572	CYP3	818	EFA02819	45	6.00E-67	cytochrome P450 6BQ5 [ <i>Tribolium castaneum</i> ]
Unigene59684	CYP3	871	NP_496108	42	4.00E-47	Cytochrome P450 family member (cyp-13A1) [ <i>Caenorhabditis elegans</i> ]
Unigene59753	CYP2	916	XP_969587	69	1.00E-115	cytochrome P450 307A1 [ <i>Tribolium castaneum</i> ]
Unigene59997	CYP4	1443	EEZ99364	45	1.00E-100	cytochrome P450-like protein [ <i>Tribolium castaneum</i> ]
Unigene8212	CYP4	650	ABF06550	60	6.00E-71	CYP4BG1 [ <i>Ips paraconfusus</i> ]
Unigene8705	CYP4	686	EEZ99364	32	4.00E-32	cytochrome P450-like protein [ <i>Tribolium castaneum</i> ]

Fragments less than 600 bp listed in Table S4.

doi:10.1371/journal.pone.0032291.t002

mito) [24,25]. The number of unigenes in the *T. yunnanensis* transcriptome belongs to the range of P450s identified in other insect species, while the accurate gene number still remains to be identified by gene cloning based on the fragments obtained here. Of the 23 unigenes which contained longer length, it was possible to differentiate them into four clades by phylogenetic analysis. The majority of these P450s belonged to the CYP3 and CYP4 clades compared with CYP2 and mito clades, which is in agreement with *Tribolium castaneum* and other insect systems [25,26]. Because some functionality of P450s can be assigned to each of the four clades based on known functionalities characterized in other insects, we

can select some interesting candidates for a further investigation according to the phylogenetic results. Until recently, the increased production of P450s in resistant insects has shown to occur almost exclusively through up regulation via changes in *cis*-or *trans*-acting regulatory loci, but gene duplication or amplification of P450s has now been implicated in the resistance of four insect species [27]. From phylogeny, gene duplication events specific to *T. yunnanensis* are apparent, with the best example being the two CYP3-type sequences Unigene58724 and Unigene58733, and two CYP4-type sequences Unigene59997 and Unigene8705, if they were paired in the phylogeny with bootstrap support greater than 70%. However,

**Table 3.** Putatively identified GST genes in *Tomicus yunnanensis*.

Unigene ID	Putative assignment	Length (bp)	First hit	Identity (%)	E_value	Blast annotation/Organism
Unigene22819	Epsilon	521	XP_971136	49	2.00E-25	PREDICTED: similar to Glutathione S transferase E8 CG17533-PA [ <i>Tribolium castaneum</i> ]
Unigene23097	Epsilon	542	ACU09495	41	9.00E-23	glutathione S-transferase 16 [ <i>Helicoverpa armigera</i> ]
Unigene25020	Sigma	693	XP_002633726	35	2.00E-16	C. briggsae CBR-GST-3 protein [ <i>Caenorhabditis briggsae</i> ]
Unigene25058	Epsilon	715	XP_966966	41	1.00E-37	PREDICTED: similar to Glutathione S transferase E6 CG17530-PA [ <i>Tribolium castaneum</i> ]
Unigene28476	Epsilon	209	XP_971389	65	4.00E-15	PREDICTED: similar to Glutathione S transferase E5 CG17527-PA [ <i>Tribolium castaneum</i> ]
Unigene31256	Epsilon	218	XP_966787	42	7.00E-10	PREDICTED: similar to Glutathione S transferase E7 CG17531-PA [ <i>Tribolium castaneum</i> ]
Unigene34958	Epsilon	230	XP_971268	37	2.00E-07	PREDICTED: similar to Glutathione S transferase E7 CG17531-PA [ <i>Tribolium castaneum</i> ]
Unigene35169	Epsilon	231	XP_971449	54	2.00E-11	PREDICTED: similar to Glutathione S transferase E7 CG17531-PA [ <i>Tribolium castaneum</i> ]
Unigene37693	Sigma	241	NP_001165920	52	6.00E-14	glutathione S-transferase S3 [ <i>Nasonia vitripennis</i> ]
Unigene40017	Epsilon	252	NP_611325	56	3.00E-19	glutathione S transferase E3 [ <i>Drosophila melanogaster</i> ]
Unigene44809	Epsilon	279	ADD19697	62	2.00E-27	glutathione S-transferase [ <i>Glossina morsitans morsitans</i> ]
Unigene44817	Epsilon	279	ADD19697	61	1.00E-27	glutathione S-transferase [ <i>Glossina morsitans morsitans</i> ]
Unigene49693	Epsilon	323	XP_967234	43	3.00E-19	PREDICTED: similar to glutathione S-transferase 6A [ <i>Tribolium castaneum</i> ]
Unigene50873	Delta	339	XP_974204	76	4.00E-45	PREDICTED: similar to GST [ <i>Tribolium castaneum</i> ]
Unigene54951	Epsilon	415	BAE80117	45	4.00E-26	glutathione S-transferase [ <i>Plutella xylostella</i> ]
Unigene5052	Epsilon	443	XP_971136	45	2.00E-31	PREDICTED: similar to Glutathione S transferase E8 CG17533-PA [ <i>Tribolium castaneum</i> ]
Unigene8150	Epsilon	416	NP_611328	44	2.00E-14	glutathione S transferase E6 [ <i>Drosophila melanogaster</i> ]

doi:10.1371/journal.pone.0032291.t003

the relevance of duplication resistance of *T. yunnanensis* requires a further investigation.

**Glutathione S-transferase (GST).** GSTs catalyse the glutathione conjugation reaction with reduced glutathione (GSH) to convert them, resulting in less toxic water-soluble products that can eventually be excreted, which are widespread in both prokaryotes and eukaryotes [28]. The increased expression and activity of GSTs, and amplification of the structural GST genes has been documented as a mechanism of insect resistance [27]. Seventeen putative GST unigenes were identified in *T. yunnanensis* transcriptome (Table 3). Based on the closest BLAST hits in the NCBI nr database and, when possible, through applying a phylogenetic analysis 14, 2, and 1 unigenes were assigned to the Epsilon, Sigma, and Delta classes, respectively. In insects, there are two ubiquitously distributed distantly related groups of GSTs, classified according to their location within the cell: microsomal and cytosolic [29]. The microsomal class contains few gene duplicates, while the cytosolic class contains highly diverse larger gene family divided into six major subclasses: Delta, Epsilon, Sigma, Omega, Theta, and Zeta [30]. In regard to microsomal class, it may await discovery due to its absence from the current transcriptomic dataset, but this group has not been implicated in the metabolism of insecticides [29]. In microsomal class, only three subclasses have been identified from the current database, which is similar to some insect species that do not have six subclasses like *Drosophila melanogaster* [31]. For instance, only Sigma, Delta, and Theta type GSTs were found in *Nasonia vitripennis*, and Epsilon, Sigma, and Omega type GSTs were found in *T. castaneum* based on genomic analysis [30]. Although the Delta subclass is absent in *T. castaneum*, one unigene assigned to Delta subclass was identified

in *T. yunnanensis*. In addition, the majority of *T. yunnanensis* GSTs were assigned to Epsilon subclass, which is in accordance with studied insect GSTs with extensive cases of gene expansions in Epsilon and Delta types [32].

**Other candidates.** In addition to the detailed analyses above, further unigenes were identified with a high sequence similarity to important genes related to insecticide metabolism and targets. As shown in Table 4, a number of unigenes annotated as

**Table 4.** Putative genes of interest related to insecticide resistance.

Gene name	Number of unigenes had a hit with nr database
Cytochrome P450	146
Glutathione S-transferase	17
Carboxylesterase	128
Superoxide dismutase	20
Acetyl-CoA carboxylase	25
Acetylcholinesterase	6
$\gamma$ -aminobutyric acid (GABA) receptor	8
Nicotinic acetylcholine receptor	17
Sodium channel	5
Chloride channel	63
Ryanodine receptor	36

doi:10.1371/journal.pone.0032291.t004

```

Unigene52967 : MLFSALRLSALKNFTBFFDRVVVERFAPELKTKGGIMPEKTQGVLEGTVIAVGPYRT : 60
Api          : -----MASVATKFRPLFDRVLVKRLDAVKQSKGGIMLPESASKTREATVIAVGPARN : 54
Gmo         : -----MAAQAIKKIMPLDRIILVQRAESLTTTKGGIVLPEKSOAKMMQGTVVAVGPGARN : 55
Nvi         : ----MAAAAIVKRLIPLFDRVLIQRAEALTKTKGGIVIPEKAQGVLRGTVVAIGPGSRN : 56
Tca         : -----MSKHVVPILLNRVLLKRFDPAAKTKGGVVIPEGWRKKISKGKVVAVGPGTVN : 51

Unigene52967 : ED-GKTIPPAVKTGDKVLLPEYGGSKVVVEDKE-YHIFRESDLIGKFN- : 106
Api          : QD-GKPVFIDVNVGDRVLLPEYGGTAIQLDLDDDSYTIKFESSELLAKVE- : 101
Gmo         : NQTGAHIAPSVKEGDRVLLPEYGGTKVELEDKKEYLLFRESDILAKYE- : 103
Nvi         : DK-GEHIPPSSIKVGDVLLPEYGGTKVELEENKEFHLLFRESDILAKLEA : 104
Tca         : NQ-GKTVPCCLKAGDVLLPEYGGTKVQYDEKQEYLYRENDILAKINE : 99
    
```

**Figure 5. Alignment of the predicted amino acid sequences of *Tomicus yunnanensis* HSP10s with that of other insects.** Identical residues are shaded black, conserved substitutions are shaded grey. Dash (-) indicates insertion or deletion. Api, *Acyrtosiphon pisum* (NP\_001119666); Gmo, *Glossina morsitans morsitans* (ADD19718); Nvi, *Nasonia vitripennis* (XP\_001599992); Tca, *Tribolium castaneum* (XP\_969732). doi:10.1371/journal.pone.0032291.g005

enzymes related to insecticide metabolic resistance, such as carboxylesterase, and superoxide dismutase; and insecticide targets, such as acetyl-CoA carboxylase, acetylcholinesterase, and  $\gamma$ -aminobutyric acid (GABA) receptor, were present in *T. yunnanensis* transcriptome. Although most of these unigenes are not full length, they will nevertheless facilitate a further characterisation of these targets by RACE to retrieve the full length cDNAs. The abundance of these transcripts demonstrates the quality of our sequencing data. It provided new leads for functional studies of dissecting the potential insecticide resistance role each these genes plays.

**Detection of HSP genes**

**Heat shock protein 10 (HSP10).** HSPs are highly conserved molecules that play vital roles in all cells. Among them, HSP10 is a near 10 kDa, highly conserved protein. In eukaryotes, HSP10, originally identified as a mitochondrial chaperone, now is also known to be present in other places such as cytosol, cell surface,

and extracellular space [33]. Here, six unigenes were identified putatively encoded HSP10 (Table S5). Two of them (Unigene4430 and Unigene52967) appeared to be complete. Amino acid alignment revealed a low homology of these four HSP10s to HSP10s from the *Acyrtosiphon pisum* (46%), *Glossina morsitans morsitans* (49%), *Nasonia vitripennis* (51%), and *Tribolium castaneum* (41%) (Figure 5). HSP10 is known as a co-chaperone for heat shock protein 60 (HSP60), and exerts immunosuppressive activity in mammals [34]. To our best knowledge, HSP10 in insects has not been structurally and functionally studied in detail.

**Small heat shock protein (sHSP).** sHSP are a family of molecular chaperones, with molecular mass ranging from 12 kDa to 43 kDa, usually below 30 kDa, which is involved in cellular defense under environmental stress conditions [35,36]. Up to data, knowledge of insect sHsps is much less than of sHsps in plants and vertebrates. Twenty three unigenes were found to be with similarities to sHSP (Table S5). Among of them, five (Unigene24062, Unigene56764, and Unigene56997,

```

Unigene56764 : -----MSIEVTHNWAENWDWP----- : 17
Unigene56997 : -----MSIPVSWDGAAPTWDVW----- : 17
Unigene24062 : -----MSHNFRRSDRWRWDWPSDWPRMDA-----IVPRF : 30
Unigene58448 : -----MAMLLRPSFYSPISSSFYDPISCSPPWSMFDDVDRY-----IKDVE : 40
Unigene59017 : MTAIQPICRTFSRLTLWKNVASTLAVPSTSTRCFSGKSLVPFGGGFYRPSNVIGEFQNRV : 60


Unigene56764 : -----LQH-NDGVVKVHNT--ADSWEVGLDAAFFTPK : 46
Unigene56997 : -----LSKENGVVTSTVT--DNEWSIQLEAGPFKPD : 47
Unigene24062 : SSHLDRFDRNWRDDPFWRDLYPRWAEPKFKEGIDVKTNIVND--RNRFAVDIDTYQFKPE : 88
Unigene58448 : EEMLDTER-----AVEKDLGTRVRVTAEPQDGKFGKLCNLTGFKPE : 81
Unigene59017 : EQMERDEFDRYFGLGGFGFRPFVDFEHPSTSTELFRLKNPIVEEDGVKFKLEFDVRRFRPE : 120

Unigene56764 : EIDVKVLG--DQLVTHCVHEERQDKHGAVKREVNRTYHLPGDVDPSTIRSHLTS-RGILH : 103
Unigene56997 : EIDVTVVH--GVLIVNLKHAVRTDNHGTVRHI TRSYHLPSWIDPGTVKSSLSLSP-RGFLA : 104
Unigene24062 : ELQVKTLD--DTLLIEGRHEDVRDADNYTKMHFVRKYQLPSDVMPQDIVSSIDS-HGRIM : 145
Unigene58448 : EMKCELDG--DHLVLEGEHKE----GNNFRSFHHRIWLPPSVNKDSLKCFDPDKGELA : 134
Unigene59017 : DVKISTDSKAGTITVEAKHSD----EHSKFEYVRKISIPGVKPEELTCTYKG-NGVLS : 174

Unigene56764 : ITAQKKH----- : 110
Unigene56997 : ITAQKKQ----- : 111
Unigene24062 : VEARKRVP AIDGRERSIPIESAAFPPPSSTAIRPGSSSSSNYSRENQNLQHDAGYQSQRQ : 205
Unigene58448 : IEAKATPPAITQQEQQQTVENKSS----- : 158
Unigene59017 : LEAPYVATPEKEPPKDTPIQIKHE----- : 198
    
```

**Figure 6. Alignment of the predicted amino acid sequences of *Tomicus yunnanensis* sHSPs. The conserved  $\alpha$ -crystallin domain is underlined.** Identical residues are shaded black, conserved substitutions are shaded grey. Dash (-) indicates insertion or deletion. doi:10.1371/journal.pone.0032291.g006






**Figure 7. Alignment of the *Tomicus yunnanensis* HSP40 amino acid sequence with other insects HSP40 amino acid sequences.** The conserved J domain is underlined. The Gly/Phe-rich domain is double underlined. The cysteine-rich domain is indicated as dotted line. Identical residues are shaded black, conserved substitutions are shaded grey. Dash (-) indicates insertion or deletion. Tca, *Tribolium castaneum* (EFA11191); Lsa, *Liriomyza sativae* (ABE57132); Lmi, *Locusta migratoria* (ABC84495); Bmo, *Bombyx mori* (BAD90846). doi:10.1371/journal.pone.0032291.g007

Unigene58448 and Unigene59017) appeared to be complete. These five sHSPs showed variable sequence identity, ranged from 29% to 82%, to the first hits under blast search, suggesting that sHSPs are diverse. An alignment of the predicted proteins deduced from the complete unigenes is shown in Figure 6. Interestingly, the alignment indicated that they were with very low similarity between each other, revealing that they do not display a highly conservation. Previous studies have demonstrated that sHSPs are abundant and

ubiquitous in almost all organisms with different numbers, from bacteria to algae single celled and even to the higher organisms including human [37]. Representative sequences of the many demonstrated sHSPs available share a conserved sequence of approximately 90 amino acid residues, termed  $\alpha$ -crystallin domain responsible for dimer formation and form large multimeric complexes that are known to be crucial to their chaperone activities [38–40]. There is  $\alpha$ -crystallin domain in the resided



**Figure 8. Alignment of the partial amino acid sequence of *Tomicus yunnanensis* HSP60 with that of other insects.** (A) the ATP-binding motif. (B) The classical mt-HSP60 signature (underlined) motif. Identical amino acids are shaded black, and conserved residues are shaded grey. Tca, *Tribolium castaneum* (XP\_971630), Ame, *Apis mellifera* (XP\_392899), Csu, *Chilo suppressalis* (ACT52824); Lsa, *Liriomyza sativae* (AAW49251). doi:10.1371/journal.pone.0032291.g008



**Figure 9. Alignment of the partial amino acid sequence of *Tomicus yunnanensis* HSP70s with that from other insects.** (A) Three highly conserved HSP70 family signatures labeled I, II, III. (B) The ATP-GTP binding site. (C) Putative bipartite nuclear localization signals. (D) The nonorganellar consensus motif. (E) The C-terminus. Identical amino acids are shaded black, and conserved residues are shaded grey. Dash (-) indicates insertion or deletion. Mci, *Macrocentrus cingulum* (ACD84944); Pxy, *Plutella xylostella* (BAF95560); Tca, *Tribolium castaneum* (XP\_974442); Lmi, *Locusta migratoria* (AAO21473). doi:10.1371/journal.pone.0032291.g009

C-terminus of *T. yunnanensis* predicted sHSPs. The N-terminal coding sequences and N-terminal ends of sHSPs are more variable than C-terminal with the  $\alpha$ -crystallin domain in different species [41]. As expected, the deduced amino acid sequence of *T. yunnanensis* HSP20s showed high conservation in  $\alpha$ -crystallin domain, and high divergence in N-terminal region and variable N-terminal end. However, there are still some conserved amino acid residues in N-terminal.

**Heat shock protein 40 (HSP40).** HSP40 family is one of the HSPs containing the DNAJ homologous region of *Escherichia coli*, J-domain of DNAJ [42]. HSP40 performs an essential molecular chaperone function in protein translation, folding, unfolding, translocation and degradation, in protein translocation across membranes, and in protecting cells from the effects of heat and other stress factors, primarily acting as cofactors or regulators of

heat shock protein 70 (HSP70) [43]. Thirty seven unigenes with similarity to HSP40 were present in *T. yunnanensis* transcriptome. Among them, Unigene59953 is with full open reading frame. The amino acid identity is about 30% compared with those of *T. castaneum*, *Liriomyza sativae*, *Locusta migratoria*, and *Bombyx mori* (Figure 7). Typically, HSP40 proteins have three distinct domains: (1) the J domain of 70 amino acid residues in length which constitutes the most conserved region of these proteins and interacts with HSP70 and stimulates its ATPase activity; (2) a glycine and phenylalanine-rich region (G/F domain), postulated to act as a flexible hinge needed to activate the substrate binding properties of Hsp70 when it interacts with Hsp40; and (3) a cysteine-rich region (C domain) resembling a zinc-finger like structure, suggested to mediate dimer formation and molecular chaperone-peptide interactions [44–47]. These three domains

were found in the predicted amino acid sequence of *T. yunnanensis* HSP40. However, not all HSP40 necessarily contain all of these 3 domains. Based on the differences in these regions, HSP40 proteins can be categorized into three groups: Type I homologs have all 3 domains (J, G/F, and C), Type II have the J and G/F but not the C domain, and Type III have the J domain alone [48,49]. According to this, *T. yunnanensis* HSP40 obtained here belongs to Type I.

**Heat shock protein 60 (HSP60).** HSP60, a major group of the heat shock proteins, includes stress inducible and constitutively expressed members. It is believed to be predominately mitochondrial, although some are also reported in cytosol and in extracellular compartments [50]. Eighteen unigenes coding for putative HSP60 were identified in the database (Table S5). Compared with orthologues from other organisms, the percentage identity of the deduced amino acid sequence of these unigenes is high, between 45–93%, confirming the remarkable conservation of this family. Among them, only Unigene59944 was with the length above 1 kb, the others were shorter than 600 bp. Alignment of the predicted amino acid sequence of Unigene59944 with that of other insects revealed it contained the signature peptide known as conserved ATP-binding motif [51] (Figure 8). The classical mitochondrial HSP60 signature motif was found in the deduced amino acid sequence [52], suggesting the HSP60 coded by Unigene59944 presented here is a member of the mitochondrial HSP60 chaperone family. As Unigene59944 was 3' and 5'-truncated, a fragment of about 30 amino acids at the N terminus required for import into the mitochondria and a typical GGM repeat motif for HSP60 at the C terminus were not observed.

**Heat shock protein 70 (HSP70).** In the HSP family, the most studied are HSP70s. There are two main forms of these 70 kD proteins, the heat shock cognate (HSC70) which is expressed constitutively and an inducible form (HSP70) which is normally expressed in response to external stimuli [53]. Seventeen unigenes produced the best sequence matches to HSP70 (Table S5). The majority of them showed highly conserved identity above 70% to the classic inducible form of HSP70 from other organisms. All these sequences were not in completeness, and only six sequences (Unigene1, Unigene16402, Unigene20916, Unigene59995, Unigene6391, and Unigene6586) were in length longer than 1 kb. After alignment of these 6 non-redundant sequences to that of other insects (Figure 9), three signature motifs of the HSP70 family [54] were conserved. Putative ATP-GTP binding site [55] and nonorganellar consensus motif were located at the amino acid sequences. Putative bipartite nuclear localization signals, which are needed for the selective translocation of HSC70 into the nucleus [56] were identified. The terminal sequence motif of the HSP70 family identifies their cellular location, with the EEDV sequence motif attesting to its cytoplasmic localisation, whereas K/HDEL and PEAEEYEEAKK characterise members localised to the endoplasmic reticulum and mitochondria respectively [53,57]. With regard to this, the terminal sequence motif of Unigene20916 and Unigene6391 were K/HDEL and EEDV, indicating they belonged to cytoplasmic localization and endoplasmic reticulum member, respectively.

**Heat shock protein 90 (HSP90).** In normal physiological conditions, HSP90 is abundant, accounting for 1% of the total soluble cytosolic protein, which is a highly conserved molecular

chaperone contributing to the folding, maintenance of structural integrity and proper regulation of a subset of cytosolic protein [54,58,59]. Like HSP70, HSP90s has also been extensively studied. Twenty four unigenes with short length less than 800 bp were identified coding for putative HSP90 (Table S5). Except for Unigene20269 and Unigene46492, they showed above 60% amino acid identity to HSP90s from other organisms. Other sequence information was not analyzed here because of only relatively short fragments available.

## Conclusions

In this work, transcriptome database has been produced on a large scale for *T. yunnanensis* using Illumina sequencing. The results provided new insights into the genomics of this forest pest. It contributed significantly to the rapid discovery of a wide diversity candidate gene for this organism that lacks complete genome sequences and other genetic tools and resources. This transcriptome can be used as a reference to provide new leads for comparative studies within the family. Based on this database, the predicted repertoire of genes related to insecticide resistance and environmental stress are constructed, providing valuable information regarding further investigations of the detailed mechanisms. However, as the sequences obtained by Illumina sequencing are short, most of the interesting putative genes discovery by this comprehensive tool is need to rely on RACE PCR in order to obtain full-length sequence data. Furthermore, the database will continue to be an enormous resource for genome-wide association studies of the whole picture of other important biological questions in the future.

## Supporting Information

**Figure S1** Length distribution of contigs. (TIF)

**Figure S2** Length distribution of scaffolds. (TIF)

**Table S1** Top BLAST hits from NCBI nr database. BLAST results against the NCBI nr database for all the distinct sequences with a cut-off E value above  $10^{-5}$  are shown. (XLS)

**Table S2** Gene Ontology of *Tomicus yunnanensis* unigenes. (XLS)

**Table S3** KEGG summary of *Tomicus yunnanensis* transcriptome. (XLS)

**Table S4** Putative P450 genes identified in *Tomicus yunnanensis* transcriptome. (XLS)

**Table S5** Putative heat shock protein genes identified in *Tomicus yunnanensis* transcriptome. (XLS)

## Author Contributions

Conceived and designed the experiments: JYZ BY. Performed the experiments: JYZ NZ. Analyzed the data: JYZ BY. Contributed reagents/materials/analysis tools: NZ. Wrote the paper: JYZ BY.

## References

- Kirkendall LR, Faccoli M, Ye H (2008) Description of the Yunnan shoot borer, *Tomicus yunnanensis* Kirkendall & Faccoli sp. n. (Curculionidae: Scolytinae), an

unusually aggressive pine shoot beetle from southern China, with a key to the species of *Tomicus*. *Zootaxa* 1819: 25–39.

2. Duan Y, Kerdelhué C, Ye H, Lieutier F (2004) Genetic study of the forest pest *Tomicus piniperda* (Col., Scolytinae) in Yunnan province (China) compared to Europe: new insights for the systematics and evolution of the genus *Tomicus*. *Heredity* 93: 416–422.
3. Haack RA, Petrice TR, Poland TM (2000) *Tomicus piniperda* (Coleoptera: Scolytidae) emergence in relation to burial depth of brood logs. *J Econ Entomol* 93: 342–346.
4. Eager TA, Berisford CW, Dalusky MJ, Nielsen DG, Brewer JW, et al. (2004) Suitability of some southern and western pines as hosts for the pine shoot beetle, *Tomicus piniperda* (Coleoptera: Scolytidae). *J Econ Entomol* 97: 460–467.
5. Faccoli M, Anfora G, Tasin M (2008) Responses of the Mediterranean pine shoot beetle *Tomicus destruens* (Wollaston) to pine shoot and bark volatiles. *J Chem Ecol* 34: 1162–1169.
6. Horn A, Stauffer C, Lieutier F, Kerdelhué C (2009) Complex postglacial history of the temperate bark beetle *Tomicus piniperda* L. (Coleoptera, Scolytinae). *Heredity* 103: 238–247.
7. Hegedus Z, Zakrzewska A, Agoston VC, Ordas A, Rácz P, et al. (2009) Deep sequencing of the zebrafish transcriptome response to mycobacterium infection. *Mol Immunol* 46: 2918–2930.
8. Der JP, Barker MS, Wickett NJ, de Pamphilis CW, Wolf PG (2011) *De novo* characterization of the gametophyte transcriptome in bracken fern, *Pteridium aquilinum*. *BMC Genomics* 12: 99.
9. Wang XW, Luan JB, Li JM, Bao YY, Zhang CX, et al. (2010) *De novo* characterization of a whitefly transcriptome and analysis of its gene expression during development. *BMC Genomics* 11: 400.
10. Xue J, Bao YY, Li BL, Cheng YB, Peng ZY, et al. (2010) Transcriptome analysis of the brown planthopper *Nilaparvata lugens*. *PLoS One* 5: e14233.
11. Karatolos N, Pauchet Y, Wilkinson P, Chauhan R, Denholm I, et al. (2011) Pyrosequencing the transcriptome of the greenhouse whitefly, *Trialeurodes vaporariorum* reveals multiple transcripts encoding insecticide targets and detoxifying enzymes. *BMC Genomics* 12: 56.
12. Liu H, Zhang Z, Ye H, Wang H, Clarke SR, et al. (2010) Response of *Tomicus yunnanensis* (Coleoptera: Scolytinae) to infested and uninfested *Pinus yunnanensis* bolts. *J Econ Entomol* 103: 95–100.
13. Li R, Zhu H, Ruan J, Qian W, Fang X, et al. (2010) *De novo* assembly of human genomes with massively parallel short read sequencing. *Genome Res* 20: 265–272.
14. Pertea G, Huang X, Liang F, Antonescu V, Sultana R, et al. (2003) TIGR Gene Indices clustering tools (TGICL): a software system for fast clustering of large EST datasets. *Bioinformatics* 19: 651–652.
15. Aparicio G, Gotz S, Conesa A, Segrelles D, Blanquer I, et al. (2006) Blast2GO goes grid: developing a grid-enabled prototype for functional genomics analysis. *Stud Health Technol Inform* 120: 194–204.
16. Thompson JD, Gibson TJ, Plewniak F, Jeanmougin F, Higgins DG (1997) The CLUSTAL\_X windows interface: flexible strategies for multiple sequence alignment aided by quality analysis tools. *Nucleic Acid Res* 25: 4876–4882.
17. Tamura K, Peterson D, Peterson N, Stecher G, Nei M, et al. (2011) MEGA5: molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Mol Biol Evol* 28: 2731–2739.
18. Bai X, Zhang W, Orantes L, Jun TH, Mittapalli O, et al. (2010) Combining next-generation sequencing strategies for rapid molecular resource development from an invasive aphid species, *Aphis glycines*. *PLoS One* 5: e11370.
19. Crawford JE, Guelbeogo WM, Sanou A, Traoré A, Vermick KD, et al. (2010) *De novo* transcriptome sequencing in *Anopheles funestus* using Illumina RNA-seq technology. *PLoS One* 5: e14202.
20. Yang H, Hu L, Hurek T, Reinhold-Hurek B (2010) Global characterization of the root transcriptome of a wild species of rice, *Oryza longistaminata*, by deep sequencing. *BMC Genomics* 11: 705.
21. Gene Ontology Consortium (2008) The Gene Ontology project in 2008. *Nucleic Acids Res* 36: D440–D444.
22. Ewen-Campen B, Shaner N, Panfilio KA, Suzuki Y, Roth S, et al. (2011) The maternal and early embryonic transcriptome of the milkweed bug *Oncopeltus fasciatus*. *BMC Genomics* 12: 61.
23. Li X, Schuler MA, Berenbaum MR (2007) Molecular mechanisms of metabolic resistance to synthetic and natural xenobiotics. *Annu Rev Entomol* 52: 231–253.
24. Feyereisen R (2005) Insect Cytochrome P450. In *Comprehensive Molecular Insect Science* Volume 4 Gilbert LI, Iatrou K, Gill SS, eds. Elsevier. pp 1–77.
25. Feyereisen R (2006) Evolution of insect P450. *Biochem Soc Trans* 34: 1252–1255.
26. Tribolium Genome Sequencing Consortium (2008) The genome of the model beetle and pest *Tribolium castaneum*. *Nature* 452: 949–955.
27. Bass C, Field LM (2011) Gene amplification and insecticide resistance. *Pest Manag Sci* 67: 886–890.
28. Yamamoto K, Teshiba S, Shigeoka Y, Aso Y, Banno Y, et al. (2011) Characterization of an omega-class glutathione S-transferase in the stress response of the silkworm. *Insect Mol Biol* 20: 379–386.
29. Enayati AA, Ranson H, Hemingway J (2005) Insect glutathione transferases and insecticide resistance. *Insect Mol Biol* 14: 3–8.
30. Friedman R (2011) Genomic organization of the glutathione S-transferase family in insects. *Mol Phylogenet Evol* 61: 924–932.
31. Ding Y, Ortelli F, Rossiter LC, Hemingway J, Ranson H (2003) The *Anopheles gambiae* glutathione transferase supergene family: annotation, phylogeny and expression profiles. *BMC Genomics* 4: 35.
32. Gullipalli D, Arif A, Aparoy P, Svenson GJ, Whiting MF, et al. (2010) Identification of a developmentally and hormonally regulated Delta-Class glutathione S-transferase in rice moth *Coryca cephalonica*. *Comp Biochem Physiol B* 156: 33–39.
33. Jia H, Halilou AI, Hu L, Cai W, Liu J, et al. (2011) Heat shock protein 10 (Hsp10) in immune-related diseases: one coin, two sides. *Int J Biochem Mol Biol* 2: 47–57.
34. Tazir Y, Steisslinger V, Soblik H, Younis AE, Beckmann S, et al. (2009) Molecular and functional characterisation of the heat shock protein 10 of *Strongyloides ratti*. *Mol Biochem Parasitol* 168: 149–157.
35. Kim KK, Kim R, Kim SH (1998) Crystal structure of a small heat-shock protein. *Nature* 394: 595–599.
36. Waters ER, Aebermann BD, Sanders-Reed Z (2008) Comparative analysis of the small heat shock proteins in three angiosperm genomes. *Cell Stress Chaperones* 13: 127–142.
37. Li ZW, Li X, Yu QY, Xiang ZH, Kishino H, et al. (2009) The small heat shock protein (sHSP) genes in the silkworm, *Bombyx mori*, and comparative analysis with other insect sHSP genes. *BMC Evol Biol* 9: 215.
38. van Montfort RLM, Basha E, Friedrich KL, Slingsby C, Vierling E (2001) Crystal structure and assembly of a eukaryotic small heat shock protein. *Nat Struct Biol* 8: 1025–1030.
39. Chen H, Chu Z, Zhang Y, Yang S (2006) Over-expression and characterization of the recombinant small heat shock protein from *Pyrococcus furiosus*. *Biotechnol Lett* 28: 1089–1094.
40. Aebermann BD, Waters ER (2008) A comparative genomic analysis of the small heat shock proteins in *Caenorhabditis elegans* and *briggssae*. *Genetica* 133: 307–319.
41. Sun Y, MacRae TH (2005) Small heat shock proteins: molecular structure and chaperone function. *Cell Mol Life Sci* 62: 2460–2476.
42. Hamajima F, Hasegawa T, Nakashima I, Isobe K (2002) Genomic cloning and promoter analysis of the GAHSP40 gene. *J Cell Biochem* 84: 401–407.
43. Qiu XB, Shao YM, Miao S, Wang L (2006) The diversity of the DnaJ/Hsp40 family, the crucial partners for Hsp70 chaperones. *Cell Mol Life Sci* 63: 2560–2570.
44. Michels AA, Kanon B, Bensaude O, Kampinga HH (1999) Heat shock protein (Hsp) 40 mutants inhibit Hsp70 in mammalian cells. *J Biol Chem* 274: 36757–36763.
45. Hata M, Ohtsuka K (2000) Murine cDNA encoding a novel type I HSP40/DNAJ homolog, mmDjA4<sup>1</sup>. *Biochim Biophys Acta* 1493: 208–210.
46. Wang HS, Wang XH, Zhou CS, Huang LH, Zhang SF, et al. (2007) cDNA cloning of heat shock proteins and their expression in the two phases of the migratory locust. *Insect Mol Biol* 16: 207–219.
47. Huang LH, Kang L (2007) Cloning and interspecific altered expression of heat shock protein genes in two leafminer species in response to thermal stress. *Insect Mol Biol* 16: 491–500.
48. Cheetham ME, Caplan AJ (1998) Structure, function and evolution of DnaJ: conservation and adaptation of chaperone function. *Cell Stress Chaperones* 3: 28–36.
49. Ohtsuka K, Hata M (2000) Mammalian HSP40/DNAJ homologs: cloning of novel cDNAs and a proposal for their classification and nomenclature. *Cell Stress Chaperones* 5: 98–112.
50. Arya R, Lakhota SC (2008) Hsp60D is essential for caspase-mediated induced apoptosis in *Drosophila melanogaster*. *Cell Stress Chaperones* 13: 509–526.
51. Choresch O, Loya Y, Müller WE, Wiedenmann J, Azem A (2004) The mitochondrial 60-kDa heat shock protein in marine invertebrates: biochemical purification and molecular characterization. *Cell Stress Chaperones* 9: 38–48.
52. Zhou J, Wang WN, He WY, Zheng Y, Wang L, et al. (2010) Expression of HSP60 and HSP70 in white shrimp, *Litopenaeus vannamei* in response to bacterial challenge. *J Invertebr Pathol* 103: 170–178.
53. Clark MS, Thorne MA, Toullec JY, Meng Y, Guan LL, et al. (2011) Antarctic krill 454 pyrosequencing reveals chaperone and stress transcriptome. *PLoS One* 6: e15919.
54. Sonoda S, Fukumoto K, Izumi Y, Yoshida H, Tsumuki H (2006) Cloning of heat shock protein genes (hsp90 and hsc70) and their expression during larval diapause and cold tolerance acquisition in the rice stem borer, *Chilo suppressalis* Walker. *Arch Insect Biochem Physiol* 63: 36–47.
55. Saraste M, Sibbald PR, Wittinghofer A (1990) The P-loop: a common motif in ATP- and GTP-binding proteins. *Trends Biochem Sci* 11: 430–434.
56. Knowlton AA, Salfity M (1996) Nuclear localization and the heat shock proteins. *J Biosci* 21: 123–132.
57. Clark MS, Peck LS (2009) HSP70 heat shock proteins and environmental stress in Antarctic marine organisms: A mini-review. *Mar Gen* 2: 11–18.
58. Welch WJ, Feramisco JR (1982) Purification of the major mammalian heat shock proteins. *J Biol Chem* 257: 14949–14959.
59. Picard D (2002) Heat-shock protein 90, a chaperone for folding and regulation. *Cell Mol Life Sci* 59: 1640–1648.