

SCIENTIFIC REPORTS



OPEN

Reduced heterotrophy in the stony coral *Galaxea fascicularis* after life-long exposure to elevated carbon dioxide

Received: 05 January 2016

Accepted: 12 May 2016

Published: 03 June 2016

Joy N. Smith^{1,2,3}, Julia Strahl^{1,4}, Sam H. C. Noonan¹, Gertraud M. Schmidt², Claudio Richter^{2,3} & Katharina E. Fabricius¹

Ocean acidification imposes many physiological, energetic, structural and ecological challenges to stony corals. While some corals may increase autotrophy under ocean acidification, another potential mechanism to alleviate some of the adverse effects on their physiology is to increase heterotrophy. We compared the feeding rates of *Galaxea fascicularis* colonies that have lived their entire lives under ocean acidification conditions at natural carbon dioxide (CO₂) seeps with colonies living under present-day CO₂ conditions. When provided with the same quantity and composition of zooplankton as food, corals acclimatized to high CO₂ showed 2.8 to 4.8 times depressed rates of zooplankton feeding. Results were consistent over four experiments, from two expeditions and both in field and chamber measurements. Unless replenished by other sources, reduced zooplankton uptake in *G. fascicularis* acclimatized to ocean acidification is likely to entail a shortage of vital nutrients, potentially jeopardizing their health and survival in future oceans.

Corals evolved in oligotrophic waters to be mixotrophs, i.e. both auto- and heterotrophs. Autotrophy is the more studied component of the two nutritional modes. However, heterotrophy is just as important, even though its role in coral health is often ignored or underestimated¹. In addition to supplementing the organic carbon supplied by endosymbiotic zooxanthellae living within their tissue², heterotrophy provides corals with essential micro- and macronutrients that are not attained through autotrophy³. These nutrients, including nitrogen and phosphorus, are needed for tissue growth, zooxanthellae regulation, and reproduction^{4–8}. Corals obtain these nutrients by the uptake of dissolved organic matter⁹, detrital particulates suspended in the water column¹⁰, bacteria¹¹, and zooplankton¹². Some species of corals increase their reliance on heterotrophy when under stress due to high turbidity^{10,13}, increased seawater temperatures that lead to the loss of their endosymbionts (coral bleaching)^{1,14}, and short-term exposure to elevated carbon dioxide (CO₂) concentrations^{15,16}. As environmental stressors from anthropogenic causes continue to increase, heterotrophy may become more relevant in the future to maintain coral health. Here, we explore coral heterotrophy with respect to one of the biggest environmental threats of all, ocean acidification. The term ‘ocean acidification’ describes the shift in seawater carbonate chemistry as anthropogenic CO₂ is absorbed by the oceans¹⁷. Under ocean acidification, seawater pH and calcium carbonate saturation states are both reduced. The reduced concentration of carbonate ions increases energy demands to maintain rates of calcification and growth, and triggers other physiological and energetic changes¹⁸.

The number of studies is limited, but some suggest that coral heterotrophy may reduce the impacts caused by ocean acidification^{15,16,19}. Several laboratory experiments show that adult and juvenile corals can maintain calcification rates with heterotrophy under ocean acidification^{15,19,20}. Other studies found that feeding or nutrient loading did not offset the impacts to coral calcification by increased CO₂^{21,22}. For example, calcification rates of *Porites rus* reduced during short-term high CO₂ exposure but were unaffected by the provision of food²³. It also remains unresolved whether coral heterotrophy may be affected by ocean acidification, and any underlying mechanisms explaining those changes. Previous studies that have investigated the effects of elevated CO₂ on coral

¹Australian Institute of Marine Science, Townsville, Queensland, Australia. ²Alfred Wegener Institute Helmholtz Centre for Polar and Marine Research, Bremerhaven, Germany. ³University of Bremen, Germany. ⁴The Carl von Ossietzky University of Oldenburg, Germany. Correspondence and requests for materials should be addressed to J.N.S. (email: JZ.Smith@aims.gov.au)




Figure 1. Rates of heterotrophy in the coral *Galaxea fascicularis* in all experiments from two methods (field and chamber), two expeditions, and two CO₂ levels (control and high CO₂).

heterotrophy have shown mixed results. For example, *Porites lutea* expanded its polyps more in high CO₂ waters, perhaps in an attempt to feed more and ameliorate the negative effects of ocean acidification²⁴. Also, the corals *Acropora cervicornis* and *Porites rus* displayed increased rates of heterotrophy under elevated CO₂^{15,16}, mitigating the adverse effects of elevated CO₂ on calcification, while *Stylophora pistillata* had reduced rates under laboratory conditions²⁵.

In this study, we investigated the impacts of ocean acidification on zooplankton capture rates in a coral species known for its voracity in feeding, *Galaxea fascicularis*. This coral feeds on zooplankton by extending mesenteric filaments through the polyp mouth, capturing particles, and then either ingesting them or digesting them externally, outside the coelenteron^{26,27}. Our study was based on four complementary field and laboratory experiments. They were conducted during two expeditions to fringing reefs in Papua New Guinea where CO₂ seeps create natural pH gradients. We compared the morphology, behavior and feeding rates of *G. fascicularis* colonies grown in seawater with elevated CO₂ (pH_T [total scale]~7.8, pCO₂~760 μatm) against those grown at control CO₂ (pH_T~8.1, pCO₂~420 μatm). We tested the following hypotheses: colonies acclimatized to elevated CO₂ (1) have smaller polyps due to energetic constraints for calcification, (2) expand their polyps further, and (3) have increased rates of heterotrophy. We also tested for (4) food selectivity in *G. fascicularis* as a function of CO₂ levels, and (5) whether the neurotransmitter receptor GABA_A was involved in the observed changes in the feeding ability of *G. fascicularis* under ocean acidification.

GABA (gamma-amino butyric acid) is one of many neurotransmitters within the central nervous system of cnidarians that helps regulate circadian rhythms in corals^{28,29} and modulates feeding responses in the cnidarian *Hydra vulgaris*^{30–32}. Two receptors are associated with the neurotransmitter GABA: GABA_A and GABA_B. There are multiple binding sites on each receptor with various possible agonists (chemicals that activate a biological response) and antagonists (chemicals that block the action of any agonist), which can attach to the binding site. The functioning of the GABA receptor GABA_A is of particular interest within ocean acidification research and has been linked to interference of neurotransmitter functioning in fish, mollusks, and other marine organisms^{33,34}. Sensory and behavioral impairment of these organisms can effectively be reversed with one of the antagonist to the GABA_A receptor, gabazine, although it has never been tested in corals. *G. fascicularis* colonies under ocean acidification were treated with gabazine to determine its possible influences on heterotrophy.

G. fascicularis fragments used in this study have been exposed to high CO₂ conditions their entire life; therefore, all observations of feeding behavior of *G. fascicularis* reflect heterotrophy of corals with life-long acclimation to ocean acidification.

Results

Feeding rates. *G. fascicularis* colonies acclimatized to high CO₂ conditions (average pH_T of 7.8) consumed less zooplankton compared to colonies under control conditions (pH_T 8.1; Fig. 1). This result was consistent for all experiments across methods and expeditions. The difference in the total number of zooplankton consumed per surface area was statistically different between CO₂ levels, but not between methods (i.e. field versus chamber), or

Factors and Interactions	F _(df,df)	P-value
Method	F _(1,61) = 0.46	0.50
Expedition	F _(1,60) = 1.9	0.18
CO ₂	F _(1,62) = 51.9	<0.001*
Method: Expedition	F _(1,57) = 9.4	0.003*
Method: CO ₂	F _(1,59) = 0.39	0.53
Expedition: CO ₂	F _(1,58) = 0.25	0.62
Method: Expedition: CO ₂	F _(1,56) = 0.48	0.49

Table 1. Results of a generalized linear model regression of coral feeding rates in response to method, expedition, CO₂, and their interaction terms.




Figure 2. Community analysis of zooplankton consumed under contrasting CO₂ regimes. Ordination plot from a canonical correlation analysis (CCA).

between expeditions (Table 1). The interaction between method and expedition had a significant influence on the total number of zooplankton consumed, although there was no difference for the main effect variables of method and expedition (Table 1).

Following the observation of reduced feeding rates during the first expedition, in the second expedition we assessed whether the reduced heterotrophy was caused by CO₂-induced impairment of neurotransmitters. The addition of gabazine during the chamber experiment from expedition 2 had no significant impact on the feeding rates (one-way ANOVA: F_(2,22) = 0.51; P = 0.48). Thus, heterotrophy rates under high CO₂ were not restored by the treatment with gabazine, the GABA_A receptor antagonist.

Composition of consumed food and selective feeding. Although the total number of zooplankton consumed was different between CO₂ levels, the types of zooplankton consumed by *G. fascicularis* were not different between CO₂ levels. Taxonomic richness of the zooplankton prey consumed was not different between CO₂ levels (three-way ANOVA: F_(1,13) = 2.74; P = 0.10), although it was higher in the chamber experiments compared to the field experiments (F_(1,15) = 20.2; P < 0.001), and higher in expedition 2 compared to expedition 1 (F_(1,15) = 8.17; P = 0.006). Multivariate community analyses on the prey consumed by corals supported these results and indicated that the zooplankton community consumed was also not different between CO₂ levels (three-way ANOVA: F_(1,56) = 1.45; P = 0.14; Fig. 2; supplementary information Figure S2), although differed between methods (F_(1,56) = 2.86; P = 0.003), expeditions (F_(1,56) = 14.5; P = 0.001), and the interaction across the two variables (F_(1,56) = 2.95; P = 0.005).

The types of prey identified in the coelenteron of dissected corals had much lower taxonomic richness than the plankton available in the water column: corals contained only 11–17 zooplankton taxa of the 26–33 taxa present in the water. Corals preferentially ingested some zooplankton taxa, including *Pontellidae* and *Paracalanidae* copepods, decapods, amphipods, and chaetognaths, whereas *Oithonidae* copepods that are abundant in the water column were scarce in the food consumed (Fig. 3).

Results from logistic regressions that examined the effects of elevated CO₂, expedition, and method, on the probability that each zooplankton taxon may be consumed indicated slight variation in the rates of consumption of the various taxa in response to these factors (Table 2). There was no difference in selectivity between high CO₂ and control corals for the most available and most frequently consumed zooplankton taxa. However, the rare *Acartidae* copepodites, *Harpacticoida*, *Isopoda*, *Ostracoda*, and *Polychaeta* appeared preferentially consumed at the control CO₂ level. These taxa all represent a small proportion of the plankton available and consumed (<2%).




Figure 3. The percent composition of the top available and consumed zooplankton taxa is shown for both expeditions, methods, and between CO_2 levels. Plots for the 16 most commonly consumed zooplankton taxa compare the percent of each taxon consumed by the coral represented in the coelenteron (blue symbols) to the percent of the community that each zooplankton is available in the water column (red symbols). Each zooplankton taxon has two rows, with the top row (circles) representing the control site and the bottom row (triangles) representing the elevated CO_2 site. Each panel represents a separate experiment (two expeditions and two methods). Asterisks indicate a significant difference between the percent consumed and percent available in the water column (t-tests, p -value < 0.05).

Furthermore, consumption rates of several zooplankton taxa differed between expeditions and methods. For example, *Tortanidae* copepods were rarely consumed during the first expedition, and yet during the second expedition they constituted on average 30.2% of the coral diet in the field experiment and 22.6% in the chamber experiment. Similarly, uptake rates of decapods and chaetognaths were relatively high during the second expedition.

Corallite size and polyp expansion between CO_2 levels. No difference was observed in the size of *G. fascicularis* corallites between colonies originating at the seep and control sites (1-way ANOVA: $F_{(1,62)} = 2.7$, $P = 0.11$). Elevated CO_2 also had no effect on polyp expansion of *G. fascicularis* at the seep and control sites, neither in the field nor in the chamber experiments. While coral polyps were not expanded more under elevated CO_2 compared to control CO_2 levels (4-way ANOVA: $F_{(1,124)} = 1.1$; $P = 0.29$), they were expanded significantly more in the field compared to the chamber experiments ($F_{(1,126)} = 22.0$; $P < 0.001$), and in expedition 2 compared to expedition 1 ($F_{(1,125)} = 12.2$; $P < 0.001$; see supplementary Table S1). Furthermore, corals expanded their polyps more at the end of each experiment compared to the beginning ($F_{(1,123)} = 6.3$; $P = 0.013$).

Discussion

The observed effects of ocean acidification on heterotrophy in the stony coral *Galaxea fascicularis* contradicted our initial hypothesis. We expected corals to ingest more zooplankton under high CO_2 . Instead, we found that food consumption rates were reduced under elevated CO_2 , both in the field and in chamber experiments, and during two expeditions. Since the colonies in our high and ambient CO_2 treatments had been subjected to life-long exposure to their respective CO_2 environments, this study presents the first investigation of heterotrophy in corals that were fully acclimatized to elevated CO_2 throughout their entire post-settlement lives.

The taxonomic composition of the zooplankton consumed by *G. fascicularis* was different compared to the zooplankton community available to the corals. Such selectivity is known for corals¹². Selectivity may be indicative of plankton behavior; for example, some zooplankton taxa swim more slowly or clumsily making it easier to capture them, while some taxa have chemical defenses that make them unpalatable to corals³⁵. Whether it is from their own choosing or more from the behavior or chemical defenses of the zooplankton, there was strong

Taxon	CO ₂		Expedition		Method		CO ₂ : Expedition		CO ₂ : Method		Expedition: Method	
	X ²	p-value	X ²	p-value	X ²	p-value	X ²	p-value	X ²	p-value	X ²	p-value
COPEPODS												
Acartiidae (adults)	8.78	0.679	8.06	0.011	7.87	0.196	7.52	0.074	7.29	0.15	6.92	0.071
Acartiidae (copepodites)	5.28	<0.001	2.44	<0.001	2.26	0.003	2.26	0.999	1.26	<0.001	1.26	0.999
Corycaeidae	2.71	0.889	2.15	<0.001	1.58	<0.001	1.58	1.000	1.57	0.611	1.57	0.999
Harpacticoida	0.81	<0.001	0.81	0.907	0.63	0.001	0.63	0.999	0.63	0.999	0.60	0.169
Oithonidae	5.01	0.464	2.98	<0.001	2.54	<0.001	2.27	0.008	2.11	0.034	2.08	0.441
Oncaeidae	6.33	0.151	6.28	0.464	6	0.077	6.00	0.932	6.00	0.943	5.90	0.303
Paracalanidae	10.1	0.585	9.46	0.047	9.29	0.324	8.27	0.142	8.23	0.654	7.93	0.179
Pontellidae	13.9	0.614	13.6	0.199	12.6	0.029	12.4	0.412	11.9	0.110	11.6	0.217
Temoridae	2.62	0.171	2.17	<0.001	2	0.025	2.00	0.999	1.47	<0.001	1.47	0.999
Tortanidae	18.4	0.989	8.70	<0.001	8.37	0.096	8.38	1.000	8.03	0.083	8.03	1.000
NON-COPEPODS												
Amphipoda	2.98	0.922	2.70	0.030	2.61	0.218	2.56	0.385	2.45	0.178	2.45	0.885
Chaetognatha	8.83	0.656	6.23	<0.001	4.94	<0.001	4.94	0.999	4.94	0.873	4.94	0.999
Decapoda	9.24	0.173	7.72	<0.001	7.7	0.695	7.69	0.771	7.46	0.177	7.25	0.196
Isopoda	0.35	<0.001	0.34	0.203	0.34	0.390	0.34	0.999	0.34	0.999	0.26	0.545
Ostracoda	2.23	0.002	2.22	0.574	2.22	0.766	2.06	0.013	1.74	<0.001	1.45	<0.001
Polychaeta	1.41	<0.001	1.41	0.841	1.25	0.042	1.25	0.999	1.25	0.999	1.14	0.082

Table 2. Probability for each of the 16 most common zooplankton taxon to be consumed by *Galaxea fascicularis*, as a function of CO₂ (seep vs. control), expedition (one vs. two), method (field vs. chamber), and the interactions of these parameters (three-way interactions were non-significant for all taxa and are not shown). X² (with df = 1 for all parameters) and p-values from the logistic regression analysis are presented, with bold print indicating significances at p < 0.05.

selection for certain zooplankton, and this selectivity appeared to be largely unaffected by CO₂ treatments, with the exception of only a few uncommon taxa (Table 2).

Selectivity results may be slightly biased towards larger zooplankton taxa since smaller groups digest faster than larger zooplankton³⁶. However, the feeding time in this study was purposely chosen to be one hour so that complete digestion could be avoided. Complete digestion takes hours to days, and even small nauplii are still recognizable after only 60 minutes in the coelenteron^{12,26,36,37}. Furthermore, since the mesh size of the plankton net was 100 μm, the smallest zooplankton types were excluded from the experiment. In fact, most zooplankton consumed were easily identifiable to species level even when partially digested, hence the category 'unidentified consumed zooplankton' represented only 13% of the items retrieved from the coelenteron.

G. fascicularis consumed less zooplankton in the high CO₂ water despite having the same access to food, the same state of polyp expansion, and the same corallite sizes between CO₂ treatments. The reasons for the observed reduction in feeding rates could be many, however our study negated several potential causes. Reduced heterotrophy was not caused by a reduction in corallite size since *G. fascicularis* corallites were the same size between CO₂ levels, even though exposure to elevated CO₂ reduces corallite sizes in some other coral species³⁸. For example, the temperate coral *Oculina patagonica* showed smaller corallites at elevated CO₂ due to high energetic costs for calcification; however, after one month of acidic conditions the skeleton completely dissolved and polyp sizes increased when calcification ceased and the resulting free energy was channeled into somatic growth³⁹. With respect to *G. fascicularis*, it is possible that net calcification rates may change under ocean acidification conditions despite the morphology of the corallites remaining similar for both CO₂ levels.

Reduced heterotrophy was also not caused by a difference in polyp expansion, which remained unaffected by ocean acidification for *G. fascicularis*. In contrast, another study observed that polyps from the coral *P. lutea* extended further under high CO₂ conditions²⁴. During the second expedition, however, *G. fascicularis* polyps were expanded more, which happened to occur during a new moon compared to the first expedition that had a full moon. Corals are known to feed differently with the lunar cycle, coinciding with lunar effects on zooplankton migration patterns^{28,40}. Also, polyps were expanded more in the field experiments compared to the chamber experiment, probably because the corals were undisturbed in the field.

A deficiency in the functioning of GABA_A neurotransmitter receptors in *G. fascicularis* was also not a likely cause for the observed reduction in heterotrophy. Gabazine plays a role in *Hydra vulgaris* feeding response³¹, therefore we expected it to also influence coral feeding behavior of *G. fascicularis* since both of these cnidarians share similar nervous systems. Despite our predictions, the treatment of *G. fascicularis* with gabazine yielded no change in coral heterotrophy. The effect of ocean acidification on coral neurotransmitters cannot be completely excluded, however, because different chemicals besides gabazine may bind to the neurotransmitter receptors (e.g. the agonist muscimol and the antagonist bicuculline)³². To thoroughly understand the effect of ocean acidification on neurotransmitters of *G. fascicularis*, the reactions of other receptor antagonists and agonists to elevated CO₂ need to be evaluated.

Additional experiments are needed to reveal the underlying mechanisms responsible for the reduced feeding rates in *G. fascicularis*. Potential causes or contributors that deserve further study include reduced particle retention, changes in cellular homeostasis of the tentacle cells, reduced nematocyst functioning, altered mucus production, physiological stress that makes them less capable to feed, an increase in autotrophy, and potential changes in plankton behavior, as briefly outlined here. *G. fascicularis* exerted similar effort to capture zooplankton between CO₂ levels by extending their polyps to the same level. That they ingested fewer food particles in ocean acidification conditions may reflect upon the polyps' ability to capture food. Food retention may be reduced if the functionality of their stinging cells (nematocysts) is disrupted^{41,42}. Nematocyst performance may be vulnerable to changes in pH since the acid-base balance in cells corresponds to the intracellular concentration of free H⁺ ions. A study on the jellyfish *Pelagica noctiluca* indicated that the cell homeostasis of nematocysts is profoundly compromised by acidification of the surrounding seawater impairing the cells' discharge capability⁴³. Although cellular homeostasis in nematocysts may vary between jellyfish and corals, nematocyst functioning may be impaired for corals under ocean acidification and merits further investigation.

Another possible cause for the observed reduced feeding rates could be that the polyps themselves lose their ability to retain food particles. Food particles may be stung or killed, but the mucosal or tentacular action of the polyps may not trap the particles, resulting in the loss of prey items²⁶. Mucus enhances coral heterotrophy⁴⁴, therefore heterotrophy will likely be vulnerable to any changes in mucus production, but nothing is known about how ocean acidification may affect coral mucus.

It is perceivable that *G. fascicularis* may also have reduced rates of heterotrophy in response to a reduced energy demand. Elevated CO₂ enhances the photosynthetic-derived energy supply in some coral species, and this energy is available to support critical functions like calcification. Coral calcification is generally considered to decline with elevated CO₂ levels⁴⁵, although some studies report parabolic and even positive calcification responses to ocean acidification conditions^{46,47}. However, corals are more nutrient limited than carbon limited in oligotrophic and shallow (high-light) environments². Furthermore, feeding rates of corals only reach saturation when food concentrations are high, with heterotrophy generally more efficient in oligotrophic habitats⁴⁸. Considering that *G. fascicularis* from the CO₂ seep sites live in a nutrient-poor and high-light environment, it is highly unlikely that feeding becomes saturated and their need for essential nutrients not attained from photosynthesis would still be prevalent. Therefore, *G. fascicularis* would likely continue to feed on zooplankton at the CO₂ seep sites if they were still capable even under an increased carbon supply from photosynthesis.

Regardless of the underlying mechanisms, reduced heterotrophy under elevated CO₂ will have biological impacts on corals. Growth, reproduction, zooxanthellae maintenance⁴⁹, and other metabolic processes depend on nitrogen, phosphorus, and other essential trace elements, which are exclusively attained through heterotrophy^{6,50–52}. We are only starting to understand the long-term impacts of ocean acidification on tissue growth, phototrophy, respiration, heterotrophy, and their energetic interdependencies, in selected species of coral. Many but not all coral species increase their rates of photosynthesis at higher pCO₂ levels⁵³. Reduced heterotrophy may also impact coral lipid content and fatty acid composition, since they are co-determined by zooplankton consumption⁵⁴. Furthermore, lower feeding rates may slow skeletal and tissue growth considering that growth is positively correlated with rates of heterotrophy for several coral species^{6,52}, so lower feeding rates may slow growth. Heterotrophy is certainly beneficial to corals and yet clearly heterotrophy declines for *G. fascicularis* under elevated CO₂. Any potential impact to their basic biology warrants further research.

Despite the remaining knowledge gaps, decreased heterotrophy will have important implications for the health and resilience of corals. As ocean conditions increasingly become unfavorable for many coral species, their ability to react to such stress will become imperative to their survival. Some coral species will persist while others will not, and our data show that some *G. fascicularis* colonies are able to survive under high CO₂ in the field, despite their lifetime exposure to elevated CO₂ conditions and associated reduced zooplankton feeding rates. However, it was beyond the scope of this study to measure their physiology (tissue biomass, lipid content, calcification rates, or other biophysical parameters indicative of their overall health). Such measurements should be conducted to better understand coral long-term survivability under ocean acidification.

Methods

Study site. The feeding experiments were conducted at Upa-Upasina Reef, a fringing reef in Milne Bay Province, Papua New Guinea, where a natural volcanic CO₂ seep provides gradients in seawater pH⁵⁵. A spatial map of the seawater carbonate chemistry, along with a detailed description of the Upa-Upasina high CO₂ and control site can be found in Fabricius *et al.*^{55,56}. *G. fascicularis* colonies were collected near the seep site where seawater approximates 7.8 pH_T (total scale), and from a control site with control CO₂ at ~8.1 pH_T. The chamber feeding experiments were conducted aboard the back deck of the ship while moored near Upa-Upasina Reef, with *G. fascicularis* fragments that were freshly collected from the reef. The field and chamber experiments were conducted during two ship expeditions to the site (12–14 April 2014 and 18–20 November 2014).

Seawater carbonate chemistry. The carbonate chemistry for the field sites varied through time and long-term measurements have been reported in previous literature⁵⁶. Additionally, seawater pH at total scale (pH_T) was recorded at the control and elevated CO₂ sites for several days surrounding the commencement of the feeding experiments using SeaFET pH sensors (Supplementary Information Figure S3). pH_T values had similar ranges compared to previous expeditions^{55,56}. Water samples were also collected, fixed with saturated mercuric chloride solution (HgCl₂), and later analyzed for their dissolved inorganic carbon (DIC: μmol kg⁻¹) and total alkalinity (A_T: μmol kg⁻¹) using the Versatile Instrument for the Determination of Total Inorganic Carbon and Titration Alkalinity (VINDTA 3C).

Carbonate chemistry was also measured for the seawater used for the chamber experiments and water temperature (°C) was recorded on site. Water samples saturated with HgCl₂ were stored and later measured for DIC

Expedition	Treatment	pH _T	Temperature (°C)	A _T (μmol kg ⁻¹)	DIC (μmol kg ⁻¹)	pCO ₂ (μatm)	HCO ₃ ⁻ (μmol kg ⁻¹)	CO ₃ ²⁻ (μmol kg ⁻¹)	CO _{2(aq)} (μmol kg ⁻¹)	Ω _{CA}	Ω _{AR}
1	control	8.05	28.0	2206	1887	381	1652	225	9.7	5.54	3.69
1	elevated-CO ₂	7.70	28.0	2282	2135	1028	1987	121	26.3	2.97	1.98
2	control	8.08	29.5	2270	1938	359	1693	236	9.5	5.78	3.84
2	elevated-CO ₂	7.75	29.5	2336	2171	906	2015	132	24.0	3.25	2.15

Table 3. Seawater carbonate chemistry of the chamber experiments with dissolved inorganic carbon (DIC) and total alkalinity (A_T) measured from water samples fixed with saturated mercuric chloride solution (HgCl₂). DIC and A_T were inputted into the Excel macro CO₂SYS and used to calculate pH at total scale (pH_T), partial pressure of carbon dioxide (pCO₂), bicarbonate (HCO₃⁻), carbonate (CO₃²⁻), aqueous carbon dioxide (CO_{2(aq)}), the saturation state of calcite (Ω_{CA}), and the saturation state of aragonite (Ω_{AR}).

and A_T. The water temperature was 25 °C at the time the samples were analyzed in the laboratory for its carbonate chemistry using the VINDTA 3C. DIC and A_T were used to calculate other seawater parameters (Table 3), including pH at total scale (pH_T), partial pressure of carbon dioxide (pCO₂; μatm), bicarbonate (HCO₃⁻; μmol kg⁻¹), carbonate (CO₃²⁻; μmol kg⁻¹), aqueous carbon dioxide (CO_{2(aq)}; μmol kg⁻¹), the saturation state of calcite (Ω_{CA}), and the saturation state of aragonite (Ω_{AR}), using the Excel macro CO₂SYS⁵⁷ under the constraints set by Dickson and Millero (1987)⁵⁸.

Food collection. Zooplankton were freshly collected via plankton net tows from the control site at approximately 9 pm, i.e. 2–3 h after sunset, and shortly before the start of the field and the chamber experiments. Each net tow was very slow to minimize stress to the zooplankton. Live samples were handled with care and only living zooplankton were used as food for corals (i.e. zooplankton still suspended in the water column and actively swimming. No zooplankton that had settled at the bottom of the collection container were used). Three to six zooplankton samples were preserved in 4% formalin and kept as references to determine variation in the number and taxonomic composition of zooplankton between samples.

Field feeding experiment. Tents of 100 μm plankton mesh and 25 cm base diameter (approximately 8 L volume) were used to contain zooplankton close to corals for the duration of the feeding experiment. Five tents were placed over separate *G. fascicularis* colonies each at the high CO₂ and the control sites. To prevent corals from consuming zooplankton that are naturally in the water column, the tents were deployed during daylight when zooplankton numbers are low and demersal zooplankton have not emerged into the water column yet. At approximately 9 pm, SCUBA divers injected three 60 ml syringes of freshly collected and concentrated zooplankton into each tent. Polyp expansion (25, 50, 75 or 100 percent expanded) was recorded at the beginning and end of the feeding period. After approximately one hour, the tents were removed and a fragment of each colony was extracted with a hammer and chisel and preserved in 4% formalin. The field experiments were conducted once during expedition 1 (three replicate coral colonies per CO₂ level), and twice on two consecutive nights during expedition 2 (five replicate coral colonies per CO₂ level for both nights). The plankton fed to the corals during the second expedition had similar composition and concentration in the two consecutive nights, so the results from both nights were pooled together and considered one experiment.

Chamber feeding experiment. *G. fascicularis* fragments were collected from both the high CO₂ site and the control site. They were placed in flow-through aquaria for four days to recover. The aquaria consisted of two 60 L bins with an outboard pump supplying a constant inflow of fresh seawater. For 12 hours prior to the feeding experiment, 100 μm mesh was placed over the input valve to starve *G. fascicularis*, allowing any previously consumed food to be digested. Three hours prior to the feeding experiment, each coral fragment was transferred onto a raised grid platform in individual cylindrical incubation chambers (89 mm diameter, 106 mm height, 637 ml volume) without exposing them to air. Corals collected from the seeps were placed in the chambers filled with seawater from the seep site, while those from the control site were placed in chambers filled with seawater from the control site (seawater carbonate chemistry for chamber experiments found in Table 3).

Chambers were 80% immersed in a water bath. Airspace in the chamber and a hole in its upper lid facilitated gas exchange. To generate a current within the chamber, a battery driven pulley system activated magnetic stirrer bars underneath the grid⁵³. *G. fascicularis* were fed at around 9 pm. Taking care to supply only living zooplankton, concentrated zooplankton was injected through a hole in the top lid of the chamber with a volumetric pipette. The zooplankton concentration was lower during the second expedition compared to the first, so a larger volume of plankton solution was inserted into the chamber during the second (30 ml) compared to the first expedition (20 ml). An additional three samples of the food were preserved in 4% formalin and kept as references. The feeding experiment was conducted in the dark, although red light was used for a few minutes at the commencement and cessation of the experiment to assess their state of polyp expansion. *G. fascicularis* fed for approximately one hour and then each coral piece was removed and immediately stored in 4% formalin. The chamber experiment was conducted once through an initial pilot study during expedition 1 (7 replicate coral colonies per CO₂ level), and repeated during expedition 2 with additional replicates (12 replicate coral colonies per CO₂ level).

To determine if elevated CO₂ interferes with neurotransmitter receptor functioning, six of the coral fragments per CO₂ treatment were exposed to gabazine (SR-95531, Sigma-Aldrich) at a concentration of 4 mg L⁻¹ seawater for 30 min (chamber experiment, second expedition). Coral fragments were gently washed and transferred into

their chambers filled with gabazine-free seawater. The other six colonies per CO₂ treatment were exposed to the same handling procedure, but their 30 min transfer was into a container without gabazine. Experiments were then conducted as outlined above.

Food samples for corals. Food samples given to corals were compared within and between experiments. Food samples given to each replicate coral fragment were similar in quantity and composition within each experiment, and they were not different between high CO₂ (7.8 pHT) and control treatments (8.1 pHT) and replicates. However, food samples varied in quantity and composition between the four field and chamber experiments. Details about the analysis of food samples are in the supplementary information, including Figure S1.

Laboratory analysis. Coral consumption was measured through coelenteron content analysis. *G. fascicularis* fragments were removed from formalin and placed in freshwater. Every polyp coelenteron was probed using a tungsten needle and dissecting forceps. Extracted zooplankton were identified to their major taxonomic groups. Total corallite number and corallites containing food particles were enumerated. Each coral fragment was photographed and the surface area calculated within the image-processing program, ImageJ. Corallite size was calculated by dividing the surface area of each coral fragment by the number of corallites.

Statistics. All statistical analyses were computed in R, version 3.2.2 (R Development Core Team, 2015). Generalized linear models (GLMs) were used to determine if: (1) the number of zooplankton consumed (standardized by surface area) differed across CO₂ regimes (seep vs. control), expedition (one vs. two), or methods (field vs. chamber), (2) species richness (Shannon-diversity index) of the zooplankton taxa consumed by corals differed between CO₂ regimes, expedition, or methods, (3) gabazine affected coral feeding rates, (4) zooplankton concentration in the food samples was different between each of the experimental runs, (5) corallite sizes were different between corals originating from seep and control sites, and (6) polyp expansion differed across CO₂ levels, methods, expeditions, or from the beginning to the end of the experiment. Appropriate data distributions and link functions were chosen for each GLM. Model assumptions of independence, homogeneity of variance, and normality of error were evaluated through diagnostic tests of leverage, Cook's distance, and dfbetas⁵⁹. Checks for all GLMs indicated that no influential data points or outliers existed in the data and model assumptions were met. ANOVAS (Type II) were used to determine the minimal adequate GLM with the 'Anova' function in the R library 'car' (version 2.1-1)⁶⁰. The effects of the explanatory variables on the response variables were then reported based on these GLMs.

Canonical correspondence analysis (CCA) was used to determine if the zooplankton community composition of the food available to the corals, and the food consumed by the corals, differed in relation to the explanatory variables (CO₂, expedition, method). To account for many zeros in the data where some zooplankton taxonomic groups were rarely present or rarely consumed, the community data was standardized using the Hellinger (square root) method within the decostand function of the vegan package in R⁶¹. A Monte-Carlo permutation test was used to determine the optimal CCA model and to assess the significance of the variation in species composition attributable to the explanatory variables (CO₂, expedition, method).

For each zooplankton taxon, its percent representation in the coral coelenteron content was compared against its percent in the available food using two-tailed t-tests, assuming unequal variances between samples. Logistic regressions were used to model the response of each zooplankton taxon contained in the corals to the explanatory variables of CO₂, expedition, and method. Logistic regressions use binary data of 'successes' and 'failures'. In this example, 'success' equals the probability of each taxon being consumed (p), and 'failure' equals the probability of not being consumed (1-p). Logistic regressions are within the framework of GLMs and use log-odd-ratios, defined by the logit link function, to estimate the (log) odds of each taxon being consumed under each independent variable. GLMs with a binary data distribution and logit link function were checked for overdispersion. Overdispersion (residual deviance greater than the residual degrees of freedom) existed, so the data distribution was changed to quasibinomial. Anovas with a Chi-square test were applied to the results of each GLM for each zooplankton taxon.

References

- Grotto, A. G., Rodrigues, L. J. & Palardy, J. E. Heterotrophic plasticity and resilience in bleached corals. *Nature* **440**, 1186–1189 (2006).
- Muscatine, L. & Porter, J. W. Reef Corals: Mutualistic Symbioses Adapted to Nutrient-Poor Environments. *Bioscience* **27**, 454–460 (1977).
- Houlbrèque, F. & Ferrier-Pagès, C. Heterotrophy in tropical scleractinian corals. *Biol. Rev.* **84**, 1–17 (2009).
- Ferrier-Pagès, C., Witting, J., Tambutté, E. & Sebens, K. P. Effect of natural zooplankton feeding on the tissue and skeletal growth of the scleractinian coral *Stylophora pistillata*. *Coral Reefs* **22**, 229–240 (2003).
- Piniak, G. A. & Lipschultz, F. Effects of nutritional history on nitrogen assimilation in congeneric temperate and tropical scleractinian corals. *Mar. Biol.* **145**, 1085–1096 (2004).
- Rodolfo-Metalpa, R., Peirano, A., Houlbrèque, F., Abbate, M. & Ferrier-Pagès, C. Effects of temperature, light and heterotrophy on the growth rate and budding of the temperate coral *Cladocora caespitosa*. *Coral Reefs* **27**, 17–25 (2008).
- Johannes, R. E., Coles, S. L. & Kuenzel, N. T. The role of zooplankton in the nutrition of some scleractinian corals. *Limnol. Oceanogr.* **15**, 579–586 (1970).
- D'Elia, C. F. The uptake and release of dissolved phosphorus by reef corals. *Limnol. Oceanogr.* **22**, 301–315 (1977).
- Goreau, T. F., Goreau, N. I. & Yonge, C. M. Reef corals: autotrophs or heterotrophs? *Biol. Bull.* **141**, 247–260 (1971).
- Anthony, K. R. N. Coral suspension feeding on fine particulate matter. *J. Exp. Mar. Biol. Ecol.* **232**, 85–106 (1999).
- Ferrier-Pagès, C., Gattuso, J. P., Cauwet, G., Jaubert, J. & Allemand, D. Release of dissolved organic carbon and nitrogen by the zooxanthellate coral *Galaxea fascicularis*. *Mar. Ecol. Prog. Ser.* **172**, 265–274 (1998).
- Sebens, K. P., Vandersall, K. S., Savina, L. A. & Graham, K. R. Zooplankton capture by two scleractinian corals, *Madracis mirabilis* and *Montastrea cavernosa*, in a field enclosure. *Mar. Biol.* **127**, 303–317 (1996).
- Anthony, K. R. N. Enhanced energy status of corals on high-turbidity reefs. *Mar. Ecol. Prog. Ser.* **319**, 111–116 (2005).

14. Palardy, J. E., Rodrigues, L. J. & Grotto, A. G. The importance of zooplankton to the daily metabolic carbon requirements of healthy and bleached corals at two depths. *J. Exp. Mar. Biol. Ecol.* **367**, 180–188 (2008).
15. Edmunds, P. J. Zooplanktivory ameliorates the effects of ocean acidification on the reef coral *Porites* spp. *Limnol. Oceanogr.* **56**, 2402–2410 (2011).
16. Towle, E. K., Enochs, I. C. & Langdon, C. Threatened Caribbean coral is able to mitigate the adverse effects of ocean acidification on calcification by increasing feeding rate. *PLoS One* **e0123394**, doi: 10.1371/journal.pone.0123394 (2015).
17. Doney, S. C., Fabry, V. J., Feely, R. A. & Kleypas, J. A. Ocean acidification: the other CO₂ problem. *Ann. Rev. Mar. Sci.* **1**, 169–192 (2009).
18. Hoegh-Guldberg, O. *et al.* Coral reefs under rapid climate change and ocean acidification. *Science*. **318**, 1737–1742 (2007).
19. Drenkard, E. J. *et al.* Calcification by juvenile corals under heterotrophy and elevated CO₂. *Coral Reefs* **32**, 727–735 (2013).
20. Cohen, A. L., McCorkle, D. C., De Putron, S., Gaetani, G. A. & Rose, K. A. Morphological and compositional changes in the skeletons of new coral recruits reared in acidified seawater: Insights into the biomineralization response to ocean acidification. *Geochemistry, Geophys. Geosystems* **10**, doi: 10.1029/2009GC002411 (2009).
21. Chauvin, A., Denis, V. & Cuet, P. Is the response of coral calcification to seawater acidification related to nutrient loading? *Coral Reefs* **30**, 911–923 (2011).
22. Holcomb, M., Cohen, A. L. & McCorkle, D. C. An investigation of the calcification response of the scleractinian coral *Astrangia poculata* to elevated pCO₂ and the effects of nutrients, zooxanthellae and gender. *Biogeosciences* **9**, 29–39 (2012).
23. Comeau, S., Carpenter, R. C. & Edmunds, P. J. Effects of feeding and light intensity on the response of the coral *Porites rus* to ocean acidification. *Mar. Biol.* **160**, 1127–1134 (2013).
24. Pachterres, C. O., Schmidt, G. M. & Richter, C. Coral growth and bioerosion of *Porites lutea* in response to large amplitude internal waves. *J. Exp. Biol.* **216**, 4365–4374 (2013).
25. Houlbrèque, F. *et al.* Ocean acidification reduces feeding rates in the scleractinian coral *Stylophora pistillata*. *Limnol. Oceanogr.* **60**, 89–99 (2015).
26. Hii, Y. S., Soo, C. L. & Liew, H. C. Feeding of scleractinian coral, *Galaxea fascicularis*, on *Artemia salina* nauplii in captivity. *Aquac. Int.* **17**, 363–376 (2009).
27. Wijgerde, T., Diantari, R., Lewaru, M. W., Verreth, J. A. J. & Osinga, R. Extracoelenteric zooplankton feeding is a key mechanism of nutrient acquisition for the scleractinian coral *Galaxea fascicularis*. *J. Exp. Biol.* **214**, 3351–3357 (2011).
28. Hoadley, K. D., Szmant, A. M. & Pyott, S. J. Circadian clock gene expression in the coral *Favia fragum* over diel and lunar reproductive cycles. *PLoS One* **6**, (2011).
29. Bertucci, A., Forêt, S., Ball, E. E. & Miller, D. J. Transcriptomic differences between day and night in *Acropora millepora* provide new insights into metabolite exchange and light-enhanced calcification in corals. *Mol. Ecol.* **24**, 4489–4504 (2015).
30. Pierobon, P. *et al.* Biochemical and functional identification of GABA receptors in *Hydra vulgaris*. *Life Sci.* **56**, 1485–1497 (1995).
31. Pierobon, P., Tino, A., Minei, R. & Marino, G. Different roles of GABA and glycine in the modulation of chemosensory responses in *Hydra vulgaris* (Cnidaria, Hydrozoa). *Hydrobiologia* **530–531**, 59–66 (2004).
32. Pierobon, P. Coordinated modulation of cellular signaling through ligand-gated ion channels in *Hydra vulgaris* (Cnidaria, Hydrozoa). *Int. J. Dev. Biol.* **56**, 551–565 (2012).
33. Nilsson, G. E. *et al.* Near-future carbon dioxide levels alter fish behaviour by interfering with neurotransmitter function. *Nat. Clim. Change*. **2**, 201–204 (2012).
34. Watson, S.-A. *et al.* Marine mollusc predator-escape behaviour altered by near-future carbon dioxide levels. *Proc. Biol. Sci.* **281**, <http://dx.doi.org/10.1098/rspb.2013.2377> (2014).
35. Lindquist, N. Palatability of invertebrate larvae to corals and sea anemones. *Marine Biology* **126**, 745–755 (1996).
36. Porter, J. W. Zooplankton feeding by the Caribbean reef-building coral *Montastrea cavernosa*. *Proceedings of the Second International Coral Reef Symposium* **1**, 111–125 (1974).
37. Leal, M. C., Nejstgaard, J. C., Calado, R., Thompson, M. E. & Frischer, M. E. Molecular assessment of heterotrophy and prey digestion in zooxanthellate cnidarians. *Mol. Ecol.* **23**, 3838–3848 (2014).
38. Suwa, R. *et al.* Effects of acidified seawater on early life stages of scleractinian corals (Genus *Acropora*). *Fish. Sci.* **76**, 93–99 (2010).
39. Fine, M. & Tchernov, D. Scleractinian coral species survive and recover from decalcification. *Science* **315**, 1811 (2007).
40. Alldredge, A. L. & King, J. M. Effects of moonlight on the vertical migration patterns of demersal zooplankton. *J. Exp. Mar. Biol. Ecol.* **44**, 133–156 (1980).
41. Greenwood, P. G. Acquisition and use of nematocysts by cnidarian predators. *Toxicon* **54**, 1065–1070 (2009).
42. Özbek, S., Balasubramanian, P. G. & Holstein, T. W. Cnidocyst structure and the biomechanics of discharge. *Toxicon* **54**, 1038–1045 (2009).
43. Morabito, R., Marino, A., Lauf, P. K., Adragna, N. C. & La Spada, G. Sea water acidification affects osmotic swelling, regulatory volume decrease and discharge in nematocytes of the jellyfish *Pelagia noctiluca*. *Cell. Physiol. Biochem.* **32**, 77–85 (2013).
44. Bythell, J. C. & Wild, C. Biology and ecology of coral mucus release. *J. Exp. Mar. Biol. Ecol.* **408**, 88–93 (2011).
45. Cohen, A. & Holcomb, M. Why corals care about ocean acidification: uncovering the mechanism. *Oceanography* **22**, 118–127 (2009).
46. Castillo, K. D., Ries, J. B., Bruno, J. F. & Westfield, I. T. The reef-building coral *Siderastrea siderea* exhibits parabolic responses to ocean acidification and warming. *Proc. Biol. Sci.* **281**, <http://dx.doi.org/10.1098/rspb.2014.1856> (2014).
47. Rodolfo-Metalpa, R. *et al.* Coral and mollusc resistance to ocean acidification adversely affected by warming. *Nat. Clim. Change*. **1**, 308–312 (2011).
48. Anthony, K. R. N. Enhanced particle-feeding capacity of corals on turbid reefs (Great Barrier Reef, Australia). *Coral Reefs* **19**, 59–67 (2000).
49. van Os, N. *et al.* Influence of heterotrophic feeding on the survival and tissue growth rates of *Galaxea fascicularis* (Octocorralia: Occulinidae) in aquaria. *Aquaculture* **330–333**, 156–161 (2012).
50. Ferrier-Pagès, C., Hoogenboom, M. & Houlbrèque, F. *The role of plankton in coral trophodynamics*. *Coral Reefs: An Ecosystem in Transition* (Springer Science, 2011), doi: 10.1007/978-94-007-0114-4.
51. Sérè, M. G., Massé, L. M., Perissinotto, R. & Schleyer, M. H. Influence of heterotrophic feeding on the sexual reproduction of *Pocillopora verrucosa* in aquaria. *J. Exp. Mar. Biol. Ecol.* **395**, 63–71 (2010).
52. Houlbrèque, F., Tambutté, E. & Ferrier-Pagès, C. Effect of zooplankton availability on the rates of photosynthesis, and tissue and skeletal growth in the scleractinian coral *Stylophora pistillata*. *J. Exp. Mar. Biol. Ecol.* **296**, 145–166 (2003).
53. Strahl, J. *et al.* Physiological and ecological performance differs in four coral taxa at a volcanic carbon dioxide seep. *Comp. Biochem. Physiol. Part A* **184**, 179–186 (2015).
54. Al-Moghrabi, S., Allemand, D., Couret, J. M. & Jaubert, J. Fatty acids of the scleractinian coral *Galaxea fascicularis*: effect of light and feeding. *J. Comp. Physiol. B* **165**, 183–192 (1995).
55. Fabricius, K. E. *et al.* Losers and winners in coral reefs acclimatized to elevated carbon dioxide concentrations. *Nat. Clim. Change*. **1**, 165–169 (2011).
56. Fabricius, K. E., Kluebenschedl, A., Harrington, L., Noonan, S. & De'ath, G. *In situ* changes of tropical crustose coralline algae along carbon dioxide gradients. *Sci. Rep.* **5**, 9537 (2015).
57. Lewis, E. & Wallace, D. Program developed for CO₂ system calculations. ORNL/CDIAC-105. *Carbon Dioxide Information Analysis Center*, Oak Ridge National Laboratory, U.S. Department of Energy, Oak Ridge, Tennessee (1998).

58. Dickson, A. G. & Millero, F. J. A comparison of the equilibrium constants for the dissociation of carbonic acid in seawater media. *Deep Sea Res. Part A, Oceanogr. Res. Pap.* **34**, 1733–1743 (1987).
59. Cohen, Y. & Cohen, J. Y. *Statistics and Data with R: An Applied Approach Through Examples*. (John Wiley & Sons Ltd., 2008), doi: 10.1111/j.1751-5823.2010.00109_8.x.
60. Fox, J. & Weisberg, S. *An R Companion to Applied Regression*. (Sage, 2011), at <http://socserv.socsci.mcmaster.ca/jfox/Books/Companion>.
61. Legendre, P. & Gallagher, E. D. Ecologically meaningful transformations for ordination of species data. *Oecologia* **129**, 271–280 (2001).

Acknowledgements

We thank the community at Upa-Upasina for their permission to study the corals on their reef, as well as the crew of the M.V. Chertan, especially Obedi Daniel and Robin ('Lei') Luke, for their support during the fieldwork. We additionally thank Johanna Arndt and Meliha Bademci for helping hands in the laboratory, and Glenn De'ath and Karin Boos for statistical advice. This project was funded in part by the Erasmus Mundus funded joint doctoral program MARES (FPA 2011-0016), the Great Barrier Reef Foundation's 'Resilient Coral Reefs Successfully Adapting to Climate Change' Program in collaboration with the Australian Government, the BIOACID Phase II Programme of the German Science Ministry BMBF (Grant 03F0655B), and the Australian Institute of Marine Science. The authors also acknowledge the contribution by J. Ries and an anonymous referee, whose helpful reviews improved the manuscript.

Author Contributions

J.N.S., J.S., G.M.S., C.R. and K.E.F. designed the experiment. J.N.S., J.S., S.H.C.N. and K.E.F. conducted the fieldwork. J.N.S. did the laboratory work and data analysis. J.N.S. and K.E.F. wrote the manuscript. All authors reviewed and edited the manuscript.

Additional Information

Supplementary information accompanies this paper at <http://www.nature.com/srep>

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Smith, J. N. *et al.* Reduced heterotrophy in the stony coral *Galaxea fascicularis* after life-long exposure to elevated carbon dioxide. *Sci. Rep.* **6**, 27019; doi: 10.1038/srep27019 (2016).



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>