



Draft Genome Sequence of the Chitin-Degrading Psychrotolerant Bacterium *Pedobacter jejuensis* TN23, Isolated from Antarctic Soil

Ahnna Cho,^a Yong-Joon Cho,^{b,c}  Ok-Sun Kim^a

^aDivision of Polar Life Sciences, Korea Polar Research Institute, Incheon, Republic of Korea

^bDepartment of Biological Sciences, Seoul National University, Seoul, Republic of Korea

^cResearch Institute of Basic Sciences, Seoul National University, Seoul, Republic of Korea

ABSTRACT *Pedobacter jejuensis* TN23 was isolated from soil from Terra Nova Bay, Victoria Land, Antarctica. The assembled draft genome size is 4,795,808 bp, and it contains a total of 4,095 genes with 3,970 coding sequences, including genes putatively involved in the degradation of chitin.

Pedobacter jejuensis TN23, a Gram-negative strict aerobe (1), was isolated from soil from Terra Nova Bay, Victoria Land, Antarctica (74°37'26.0"S, 164°13'49.2"E). A soil sample was collected on 3 February 2011, suspended (0.5 g/2 ml) in phosphate-buffered saline (PBS), and spread onto a Reasoner's 2A (R2A) agar plate. The plate was incubated at 10°C for 2 weeks in an aerobic state, and then a single colony was transferred to a new R2A agar plate for pure culture. This strain showed 98.01% similarity with the type strain *Pedobacter jejuensis* THG-DR3 (2) by 16S rRNA gene typing using universal 27F-1492R primers and the EzTaxon-e database (3).

The genomic DNA of *P. jejuensis* TN23 was extracted from a culture in R2A liquid medium using a PowerSoil DNA isolation kit (Qiagen, USA). The concentration and purity were determined using a Qubit 2.0 fluorometer (Invitrogen, USA), and 1 µg of genomic DNA was processed to the next step. A Nextera DNA Flex library prep kit (catalog number 20018704; Illumina, USA) was used to prepare the DNA library for sequencing following the manufacturer's protocol. The sequencing was performed using the Illumina MiSeq platform and generated 1,567,670 reads (2 × 250-bp paired-end format).

The raw data were cleaned using Trimmomatic v0.36 (4) with default parameters for adapter removal and quality trimming. The trimmed reads were assembled using SPAdes v3.1.2 (5) in standard mode with read error and mismatch correction. The final assembly yielded 45 contigs with a length of 4,795,808 bp, an N_{50} value of 316,667 bp, a GC content of 35.8%, and a genome coverage of 196.13×. Gene prediction and annotation were carried out using the NCBI Prokaryotic Genome Annotation Pipeline (6). Finally, a total of 4,095 genes were identified, including 3,970 coding genes, 3 rRNAs, 49 tRNAs, 3 noncoding RNAs (ncRNAs), and 70 pseudogenes.

The closest strain in the public genome databases was identified as *Pedobacter* sp. strain RP-3-11 by average nucleotide identity comparison (7), with a value of 80.09%. Several species of *Pedobacter* have been reported to survive in the Arctic and Antarctic environments (8, 9). They are known to have the ability to degrade chitin and cellulose in the soil (8, 10). The genome of *P. jejuensis* TN23 also has one chitinase (D7004_05180) and one chitin disaccharide deacetylase (D7004_02630) related to chitin degradation. While the optimal temperature of most chitin deacetylases in fungi is 50 to 60°C (11), the enzyme in this genome may have high activity at low temperatures as a consequence of host adaptation to low temperatures. Moreover, there are four cold shock proteins (D7004_00650, D7004_02470, D7004_11185, and D7004_17470) and six osmoprotectant-transporting proteins

Citation Cho A, Cho Y-J, Kim O-S. 2021. Draft genome sequence of the chitin-degrading psychrotolerant bacterium *Pedobacter jejuensis* TN23, isolated from Antarctic soil. Microbiol Resour Announc 10:e00523-21. <https://doi.org/10.1128/MRA.00523-21>.

Editor Frank J. Stewart, Montana State University

Copyright © 2021 Cho et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Ok-Sun Kim, oskim@kopri.re.kr.

Received 28 May 2021

Accepted 8 June 2021

Published 1 July 2021

(D7004_11055 to D7004_11060); glycine betaine accumulation against low temperatures functions as an osmoprotectant. This report is the first genome sequencing result for a *P. jejuensis* strain, and the information about the enzymes that degrade chitin should provide insights into a soil ecosystem under extreme environmental conditions.

Data availability. The raw Illumina sequence reads were deposited in the Sequence Read Archive (SRA) under the accession number [SRP260158](https://www.ncbi.nlm.nih.gov/sra/SRP260158). The genome sequences and annotations were deposited under the GenBank accession number [RBEE01000000.1](https://www.ncbi.nlm.nih.gov/genbank/RBEE01000000.1) and the BioProject accession number [PRJNA494037](https://www.ncbi.nlm.nih.gov/bioproject/PRJNA494037).

ACKNOWLEDGMENT

This work was supported by Korea Polar Research Institute (grant numbers PE19150 and PE21130).

REFERENCES

- Margesin R, Shivaji S. 2015. *Pedobacter*, p 1–17. In Whitman WB (ed), Bergey's manual of systematics of archaea and bacteria. John Wiley & Sons, Inc, Innsbruck, Austria. <https://doi.org/10.1002/9781118960608.gbm00360>.
- Kook M, Park Y, Yi T-H. 2014. *Pedobacter jejuensis* sp. nov., isolated from soil of a pine grove, and emended description of the genus *Pedobacter*. *Int J Syst Evol Microbiol* 64:1789–1794. <https://doi.org/10.1099/ijs.0.058024-0>.
- Kim O-S, Cho Y-J, Lee K, Yoon S-H, Kim M, Na H, Park S-C, Jeon YS, Lee J-H, Yi H, Won S, Chun J. 2012. Introducing EzTaxon-e: a prokaryotic 16S rRNA gene sequence database with phylotypes that represent uncultured species. *Int J Syst Evol Microbiol* 62:716–721. <https://doi.org/10.1099/ijs.0.038075-0>.
- Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* 30:2114–2120. <https://doi.org/10.1093/bioinformatics/btu170>.
- Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Prjibelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 19:455–477. <https://doi.org/10.1089/cmb.2012.0021>.
- Tatusova T, DiCuccio M, Badretdin A, Chetvernin V, Nawrocki EP, Zaslavsky L, Lomsadze A, Pruitt KD, Borodovsky M, Ostell J. 2016. NCBI Prokaryotic Genome Annotation Pipeline. *Nucleic Acids Res* 44:6614–6624. <https://doi.org/10.1093/nar/gkw569>.
- Richter M, Rosselló-Móra R, Oliver Glöckner F, Peplies J. 2016. JSpeciesWS: a Web server for prokaryotic species circumscription based on pairwise genome comparison. *Bioinformatics* 32:929–931. <https://doi.org/10.1093/bioinformatics/btv681>.
- Zhou L-Y, Chen X-Y, Du Z-J, Mu D-S. 2019. *Pedobacter chinensis* sp. nov., a cellulose-decomposing bacterium from Arctic tundra soil. *Int J Syst Evol Microbiol* 69:1926–1933. <https://doi.org/10.1099/ijsem.0.003403>.
- He R-H, Liu Z-W, Yu Y, Li H-R, Du Z-J. 2019. *Pedobacter changchengzhani* sp. nov., isolated from soil of Antarctica. *Antonie Van Leeuwenhoek* 112:1747–1754. <https://doi.org/10.1007/s10482-019-01305-4>.
- Zhang L-L, Gan L-Z, Xu Z-B, Yang F, Li Y, Fan X-L, Liu X-F, Tian Y-Q, Dai Y-M. 2018. *Pedobacter chitinilyticus* sp. nov., a chitin-degrading bacterium isolated from wheat leaf tissue. *Int J Syst Evol Microbiol* 68:3713–3719. <https://doi.org/10.1099/ijsem.0.003017>.
- Zhao Y, Park R-D, Muzzarelli RAA. 2010. Chitin deacetylases: properties and applications. *Mar Drugs* 8:24–46. <https://doi.org/10.3390/md8010024>.