

RESEARCH

Open Access

Concentration of acrylamide in a polyacrylamide gel affects VP4 gene coding assignment of group A equine rotavirus strains with P[12] specificity

LaShanda M Long-Croal^{1,2}, Xiaobo Wen¹, Eileen N Ostlund^{†3} and Yasutaka Hoshino^{*1}

Abstract

Background: It is universally acknowledged that genome segment 4 of group A rotavirus, the major etiologic agent of severe diarrhea in infants and neonatal farm animals, encodes outer capsid neutralization and protective antigen VP4.

Results: To determine which genome segment of three group A equine rotavirus strains (H-2, FI-14 and FI-23) with P[12] specificity encodes the VP4, we analyzed dsRNAs of strains H-2, FI-14 and FI-23 as well as their reassortants by polyacrylamide gel electrophoresis (PAGE) at varying concentrations of acrylamide. The relative position of the VP4 gene of the three equine P[12] strains varied (either genome segment 3 or 4) depending upon the concentration of acrylamide. The VP4 gene bearing P[3], P[4], P[6], P[7], P[8] or P[18] specificity did not exhibit this phenomenon when the PAGE running conditions were varied.

Conclusions: The concentration of acrylamide in a PAGE gel affected VP4 gene coding assignment of equine rotavirus strains bearing P[12] specificity.

Background

Diarrheal disease is one of the principal causes of morbidity and mortality among young children in the developing world. Infectious diarrhea of neonatal animals is also one of the most common and economically devastating conditions encountered in the animal agriculture industry. Among an array of infectious agents including bacteria, viruses and parasites, group A rotaviruses are the single most important etiologic agents of diarrhea in infants and young children worldwide and in addition, they are the most commonly identified viral cause of diarrhea in neonatal food animals [1-4]. In 1975, rotaviruses were first demonstrated being involved in foal diarrhea [5], and later established as the major cause of diarrhea in young foals [6-8].

The genome of group A rotavirus, a member of *Reoviridae* family, consists of eleven segments of double-stranded RNA numbered 1-11 according to their order of migration in polyacrylamide gels, segment 1 being the

slowest and segment 11 the fastest [9]. The rotavirus genome encodes six structural (VP1-VP4, VP6 and VP7) and six nonstructural (NSP1-NSP6) proteins [3]. Since two outer capsid proteins VP7 and VP4 are independent neutralization and protective antigens, a binary system of classification and nomenclature to designate the two neutralization specificities has been adopted: VP7 or G (because VP7 is a glycoprotein) serotype and VP4 or P (because VP4 is protease-sensitive) serotype [3]. Since (i) antibodies to the VP7 and VP4 have been demonstrated to confer resistance to virulent rotavirus in a type-specific manner in experimental animals; and (ii) observations made in various rotavirus vaccine trials have suggested that the induction of serotype-specific immunity may be important for optimal protection, serotypic-genotypic analyses of the VP7 and VP4 of a rotavirus derived from various animal species have been performed [3,10,11]. Such studies have established at least 14 G serotypes (21 G genotypes) and 14 P serotypes (29 P genotypes) [12]. Among equine rotaviruses, five G types (G3, G5, G10, G13 and G14) and three P types (P[7], P[12] and P[18]) have been identified.

In general, each rotavirus strain displays a dsRNA migration pattern (electropherotype) on polyacrylamide

* Correspondence: thoshino@niaid.nih.gov

¹ Rotavirus Vaccine Development Section, Laboratory of Infectious Diseases, NIAID, National Institutes of Health, Bethesda, MD 20892, USA

[†] Contributed equally

Full list of author information is available at the end of the article

gels distinct from that of other strains [9,13]. Hence analysis of such genomic polymorphism as determined by polyacrylamide gel electrophoresis (PAGE) as well as gene sequencing have been routinely used for gene coding assignments. Such studies have established that the VP7 protein is encoded by genome segment 7, 8 or 9 depending upon the rotavirus strain. For example, the VP7 is encoded by the 7th segment of rhesus rotavirus MMU18006 strain in a 12% gel [14], the 8th segment of human rotavirus DS-1 strain in a 7.5% gel [15], and the 9th segment of human rotavirus Wa strain in a 12% gel [16]. With regard to the VP4 protein, on the other hand, it is universally acknowledged that it is encoded by the genome segment 4 regardless of the rotavirus strain. During the course of generating various single gene substitution reassortants and hyperimmune antisera to them in an attempt to characterize and establish VP4 serotypes of selected equine rotaviruses [17], we found unexpectedly that the VP4 gene of equine rotavirus strains H-2, FI-14 and FI-23 was not the fourth segment but the third segment as determined by a standard 12% PAGE.

Results and discussion

Concentration of acrylamide affects the relative position of VP4 gene of equine rotavirus strains H-2, FI-14 and FI-23 in a PAGE gel

During the characterization by a standard 12% PAGE gel analysis of selected equine-human rotavirus reassortants that were generated between equine rotavirus (strain H-2 [18], FI-14 [19] or FI-23 [20]) and human rotavirus (strain DS-1 [21]), we noticed that the VP4-encoding gene of each of the three equine rotavirus strains was at the third position (Figure 1). This was unexpected since the fourth genome segment was the VP4-encoding gene of human rotavirus strains Wa (P[8]) [21], DS-1 (P[4]), ST3 (P[6]) [22] as well as rhesus rotavirus strain MMU18006 (P[3]) [23] under the same PAGE running condition. Since we reported previously that the acrylamide concentration in a PAGE gel affected the relative position of the VP7 gene of G2 rotavirus strains [24], we analyzed the effects of acrylamide concentration by using H-2 strain and its reassortant rotavirus strain. The VP4 gene of the H-2 strain was in the 4th position in a 5% (not shown) or 7.5% (Figure 2) gel, the 3rd or 4th position in a 10% (Figure 3) gel, however, it was in the 3rd position in a 12% (Figure 1) or 15% (Figure 4) gel. These findings demonstrated that the H-2 VP4 gene "flipped over" (i.e., the H-2 VP4 gene shifted to the 3rd position from its previous 4th position) in a PAGE gel containing acrylamide concentration between 7.5% and 12% (Table 1). Similarly, the FI-14 and FI-23 VP4 genes exhibited the "flip over" phenomenon between a 7.5% gel and a 12% gel (not shown, summarized in Table 1). Thus, we demonstrated that the concen-

tration of acrylamide played a critical role in determining the VP4 gene coding assignment of equine rotavirus strains H-2, FI-14 and FI-23. As we reported previously, the different PAGE running conditions affected not only the VP4 gene but also other genes as well. For example, although segments 2 and 3 of the DS-1 strain comigrated in a 7.5% gel (Figure 2), they were well separated in a 15% gel (Figure 4).

VP4 gene encoding P[12] specificity appeared to be affected most by the concentration of acrylamide in a PAGE gel

Next, we investigated whether the "flip-over" phenomenon was unique to P[12] equine rotavirus strains or common to any equine rotavirus strains. Previously [24], we showed that the VP4 gene of human rotavirus strains Wa (P[8]), DS-1 (P[4]), ST3 (P[6]) or rhesus rotavirus strain MMU18006 (P[3]) was at the 4th position regardless of acrylamide concentration in a PAGE gel (Table 1). We found in this study that the relative position of the VP4 gene of equine rotavirus strain H-1 [25] with P[7] specificity and strain L338 [26] with P[18] specificity was not affected by the varying concentration of acrylamide in a PAGE gel (data not shown, summarized in Table 1). Thus, the "flip-over" phenomenon of the VP4 gene observed in the present study appeared to be unique to equine rotavirus VP4 genes bearing P[12] specificity.

The mechanisms underlying this "flip-over" phenomenon displayed by the VP4 gene with P[12] specificity are unknown. Since the observed VP4 gene migration shift appears to be a function of acrylamide concentration (all other factors being equal), this would indicate the size of the pores in the gel is what is generating the shift. This argues for the shift being the result of a change in the tertiary structure of the molecule. Unfortunately, tools do not exist at present for predicting secondary or tertiary structures for double-stranded nucleic acid sequences. We analyzed predicted secondary structures of single-stranded RNA of VP4 gene of selected rotavirus strains including equine rotavirus strains with P[12] specificity, however, we did not find any predicted structures that were different between the equine VP4 sequences and the others (data not shown). In addition, we examined the VP4 sequences of selected rotavirus strains to look for potential changes in the equine VP4 sequence that might induce some sort of "pairing" of the ends of the molecule, however, we did not find any good candidate sequences.

Conclusions

The relative position of the VP4 gene of three equine P[12] strains (H-2, FI-14, FI-23) varied (either genome segment 3 or 4) depending upon the concentration of acrylamide. The VP4 gene bearing P[3], P[4], P[6], P[7],

Table 1: The concentration of acrylamide affects VP4-gene coding assignment of group A equine rotavirus strains H-2, FI-14, and FI-23 bearing P[12] specificity.

Rotavirus			Species of origin	VP4-gene coding assignment in a PAGE gel containing acrylamide at indicated concentration				
Strain [ref.]	P (VP4)type	G (VP7)type		5%	7.5%	10%	12%	15%
H-2 [18]	P[12]	G3	horse	4	4	3 or 4	3	3
FI-14 [19]	P[12]	G3	horse	ND ^a	4	ND	3	ND
FI-23 [20]	P[12]	G14	horse	ND	4	ND	3	ND
H-1 [25]	P[7]	G5	horse	4	4	4	4	4
L338 [26]	P[18]	G13	horse	4	4	4	4	4
Wa [21]	P[8]	G1	human	4	4	4	4	4
DS-1 [21]	P[4]	G2	human	4	4	4	4	4
ST3 [22]	P[6]	G4	human	4	4	4	4	4
MMU18006 [23]	P[3]	G3	rhesus	4	4	4	4	4

^aND = not done

P[8] or P[18] did not exhibit this phenomenon when the PAGE running conditions were varied. Caution needs to be exercised when PAGE analyses are used for VP4 gene coding assignment of rotaviruses.

Methods

Rotavirus strains, cell culture, and genetic reassortment

Table 1 summarizes group A human and animal rotavirus strains that were employed in this study. Each of the rotavirus strains used was plaque purified three times prior to use. Reassortant rotaviruses between equine rotavirus strain H-2, FI-14 or FI-23 and human rotavirus strain DS-1 were constructed by a procedure described previously [27]. Briefly, roller tube cultures of monkey kidney cell line MA104 were coinfecting at a multiplicity of infection of approximately one with the H-2 strain, FI-14 strain or FI-23 strain and the DS-1 strain. When approximately 75% of the infected cells displayed cytopathic effects, the cultures were frozen and thawed once and the lysate was plated on MA104 cells in a six-well plate (Costar, Corning Inc., Corning, NY) in the presence of G serotype cross-reactive neutralizing monoclonal antibody 57/8 [20] for selection of the desired H-2 × DS-1 and FI-14 × DS-1 and FI-23 × DS-1 reassortants. A plaque displaying a desired gene constellation (i.e., VP4 gene from the H-2, FI-14 or FI-23 strain and the VP7 gene from the DS-1 strain) was plaque purified three times prior to use. Reassortant rotaviruses between equine rotavirus strain H-1 or strain L338 and human rotavirus strain DS-1 were generated in a similar manner except that polyclonal antibodies raised against (i) porcine rotavirus OSU (P[7]G5) strain was used for selection of H-1 × DS-1 (P[7]G2) reassortant and (ii) L338 (P[18]G13) strain was used for selection of L338 × DS-1 (P[18]G2) reassortant. Eagle's minimum essential medium supplemented with 0.5 µg/ml trypsin (Sigma type IX trypsin, Sigma Chemical, St. Louis, MO) and antibiotics was used as maintenance medium and Leibovitz L-15 medium supplemented with antibiotics was




Figure 1 Electrophoretic migration patterns in a 12% PAGE gel of equine rotavirus H-2 strain, H-2 × DS-1 reassortant, and human rotavirus DS-1 strain; equine rotavirus FI-14 strain, FI-14 × DS-1 reassortant and DS-1 strain; and equine rotavirus FI-23 strain, FI-23 × DS-1 reassortant, and DS-1 strain. Arrows indicate the VP4 gene (3rd genome segment) of each of the 3 equine parental rotavirus strains.




Figure 2 Electrophoretic migration patterns in a 7.5% PAGE gel of equine rotavirus H-2 strain, H-2 x DS-1 reassortant and human rotavirus DS-1 strain. Arrow indicates the VP4 gene (4th genome segment) of the H-2 strain.




Figure 3 Electrophoretic migration patterns in a 10% PAGE gel of equine rotavirus H-2 strain, H-2 x DS-1 reassortant and human rotavirus DS-1 strain. Arrow indicates the VP4 gene of the H-2 strain. Note the 3rd and 4th genome segments of the H-2 strain comigrate.




Figure 4 Electrophoretic migration patterns in a 15% PAGE gel of equine rotavirus H-2 strain, H-2 x DS-1 reassortant and human rotavirus DS-1 strain. Arrow indicates the VP4 gene (3rd genome segment) of the H-2 strain.

employed when making virus dilutions. Agarose (SeaKem ME, BME, Rockland, ME) was used as a solidifying reagent in the overlay medium.

Rotavirus RNA extraction and PAGE analysis

The standard phenol-chloroform method or TRIzol method was employed to extract rotavirus genomic dsRNA as previously reported [28,29]. Analysis of rotavirus dsRNA was carried out at room temperature (approximately 26°C) in a discontinuous 5%, 7.5%, 10% 12% or 15%, acrylamide resolving slab gel (acrylamide:bisacrylamide 29:1, Bio-Rad Laboratories, Hercules, CA. 18 × 16 × 0.075 cm) with a 3.5% acrylamide stacking gel in the buffer system of Laemmli [30] without SDS using a SE600 gel apparatus (Amersham Biosciences, San Francisco, CA) and Tris-Glycine running buffer (pH 8.3) (Bio-Rad Laboratories). Since the polymerization temperature of acrylamide/bisacrylamide gels has been reported to affect the tertiary structure of the gel thereby influencing electrophoretic mobilities of selected RNA species [31], the polymerization of the PAGE gels used in this study was performed at a single temperature of 37°C in an incubator. In addition, since heat generated during electrophoresis has been reported to affect the mobilities of rotavirus genomic dsRNA [32], a water chiller (Lauda WKL230, Brinkmann Instruments, Westbury, NY) was used, if necessary, to maintain the desired temperature of running buffer especially when evaluating a gel with a high percentage of acrylamide/bisacrylamide. After electrophoresis, viral RNA bands were visualized by staining of the gel with silver nitrate [33].

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

All authors read and approved the final manuscript. LML, XW and ENO carried out the PAGE analyses. YH participated in the design of the study and drafted the manuscript.

Acknowledgements

We thank Dr. Albert Z. Kapikian for continuing support of the project and Ronald Jones for his excellent technical support. This work was supported by the Intramural Research Program of the National Institute of Allergy and Infectious Diseases, National Institutes of Health, USA.

Author Details

¹Rotavirus Vaccine Development Section, Laboratory of Infectious Diseases, NIAID, National Institutes of Health, Bethesda, MD 20892, USA, ²Center for Devices and Radiological Health, Food and Drug Administration, Silver Spring, MD 20994, USA and ³Diagnostic Virology Laboratory, National Veterinary Services Laboratories, Animal and Plant Health Inspection Service, USDA, Ames, IA 50010, USA

Received: 30 April 2010 Accepted: 23 June 2010

Published: 23 June 2010

References

1. Cheng AC, McDonald JR, Thielman NM: Infectious diarrhea in developed and developing countries. *Clin Gastroenterol* 2005, **39**:757-73.

2. Magdesian KG: **Neonatal foal diarrhea.** *Vet Clin North Am Equine Pract* 2005;295-312.
3. Estes MK, Kapikian AZ: **Rotaviruses.** In *Fields virology* 5th edition. Edited by: Knipe DM, Howley PM, Griffin DE, Lamb RA, Martin MA, Roizman B, Straus SE. Philadelphia, PA: Lippincott Williams and Wilkins; 2007:1917-74.
4. Saif LJ, Rosen BI, Parwani AVL: **Animal rotaviruses.** In *Viral Infections of the Gastrointestinal Tract* Edited by: Kapikian AZ. New York: Marcel Dekker; 1994:279-367.
5. Flewett TH, Bryden AS, Davies H: **Virus diarrhoea in foals and other animals.** *Vet Rec* 1975, **96**:477.
6. Conner ME, Darlington RW: **Rotavirus infection in foals.** *Am J Vet Res* 1980, **41**:1699-703.
7. Dwyer RM: **Rotaviral diarrhea.** *Vet Clin North Am Equine Pract* 1993, **9**:311-9.
8. Frederick J, Giguère S, Sanchez LC: **Infectious agents detected in the feces of diarrheic foals: a retrospective study of 233 cases (2003-2008).** *J Vet Intern Med* 2009, **23**:1254-60.
9. Rodger SM, Holmes IH: **Comparison of the genomes of simian, bovine, and human rotaviruses by gel electrophoresis and detection of genomic variation among bovine isolates.** *J Virol* 1979, **30**:839-46.
10. Hoshino Y, Kapikian AZ: **Rotavirus serotypes: classification and importance in epidemiology, immunity, and vaccine development.** *J Health Popul Nutr* 2000, **18**:5-14.
11. Kapikian AZ, Hoshino Y: **To serotype or not to serotype: that is still the question.** *J Infect Dis* 2007, **195**:611-4.
12. Abe M, Ito N, Morikawa S, Takasu M, Murase T, Kawashima T, Kawai Y, Kohara J, Sugiyama M: **Molecular epidemiology of rotaviruses among healthy calves in Japan: Isolation of a novel bovine rotavirus bearing new P and G genotypes.** *Virus Res* 2009, **144**:250-7.
13. Kalica AR, Sereno MM, Wyatt RG, Mebus CA, Chanock RM, Kapikian AZ: **Comparison of human and animal rotavirus strains by gel electrophoresis of viral RNA.** *Virology* 1978, **87**:247-55.
14. Broome RL, Vo PT, Ward RL, Clark HF, Greenberg HB: **Murine rotavirus genes encoding outer capsid proteins VP4 and VP7 are not major determinants of host range restriction and virulence.** *J Virol* 1993, **67**:448-55.
15. Greenberg HB, Flores J, Kalica AR, Wyatt RG, Jones R: **Gene coding assignments for growth restriction, neutralization and subgroup specificities of the W and DS-1 strains of human rotavirus.** *J Gen Virol* 1983, **64**:313-20.
16. Kalica AR, Greenberg HB, Wyatt RG, Flores J, Sereno MM, Kapikian AZ, Chanock RM: **Genes of human (strain Wa) and bovine (strain UK) rotaviruses that code for neutralization and subgroup antigens.** *Virology* 1981, **112**:385-90.
17. Ostlund EN, Hoshino Y: **Identification of distinct equine rotavirus VP4 serotypes and cold adaptation of equine rotavirus H2.** *Equine Infectious Diseases VIII* 1994:297.
18. Hoshino Y, Wyatt RG, Greenberg HB, Kalica AR, Flores J, Kapikian AZ: **Isolation, propagation, and characterization of a second equine rotavirus serotype.** *Infect Immun* 1983, **41**:1031-7.
19. Hoshino Y, Gorziglia M, Valdesuso J, Askaa J, Glass RI, Kapikian AZ: **An equine rotavirus (FI-14 strain) which bears both subgroup I and subgroup II specificities on its VP6.** *Virology* 1987, **157**:488-96.
20. Browning GF, Fitzgerald TA, Chalmers RM, Snodgrass DR: **A novel group A rotavirus G serotype: serological and genomic characterization of equine isolate FI23.** *J Clin Microbiol* 1991, **29**:2043-6.
21. Wyatt RG, Greenberg HB, James WD, Pittman AL, Kalica AR, Flores J, Chanock RM, Kapikian AZ: **Definition of human rotavirus serotypes by plaque reduction assay.** *Infect Immun* 1982, **37**:110-5.
22. Wyatt RG, James HB Jr, Pittman AL, Hoshino Y, Greenberg HB, Kalica AR, Flores J, Kapikian AZ: **Direct isolation in cell culture of human rotaviruses and their characterization into four serotypes.** *J Clin Microbiol* 1983, **18**:310-7.
23. Stuker G, Oshiro LS, Schmidt NJ: **Antigenic comparisons of two new rotaviruses from rhesus monkeys.** *J Clin Microbiol* 1980, **11**:202-3.
24. Ross J, Ostlund EN, Cao D, Tatsumi M, Hoshino Y: **Acrylamide concentration affects the relative position of VP7 gene of serotype G2 strains as determined by polyacrylamide gel electrophoresis.** *J Clin Virol* 2008, **42**:374-80.
25. Hoshino Y, Wyatt RG, Greenberg HB, Kalica AR, Kapikian AZ: **Isolation and characterization of an equine rotavirus.** *J Clin Microbiol* 1983, **18**:585-91.
26. Browning GF, Chalmers RM, Fitzgerald TA, Snodgrass DR: **Serological and genomic characterization of L338, a novel equine group A rotavirus G serotype.** *J Gen Virol* 1991, **72**:1059-64.
27. Hoshino Y, Jones RW, Ross J, Kapikian AZ: **Porcine rotavirus strain Gottfried-based human rotavirus candidate vaccines: Construction and characterization.** *Vaccine* 2005, **23**:3791-9.
28. Jones RW, Ross J, Hoshino Y: **Identification of parental origin of cognate dsRNA genome segment(s) of rotavirus reassortants by constant denaturant gel electrophoresis.** *J Clin Virol* 2003, **26**:347-54.
29. Santos N, Honma S, Timenetsky MCST, Linhares AC, Ushijima H, Armah GE, Gentsch JR, Hoshino Y: **Development of a microtiter plate hybridization-based PCR-enzyme-linked immunosorbent assay for identification of clinically relevant human group A rotavirus G and P genotypes.** *J Clin Microbiol* 2008, **46**:462-9.
30. Laemmli UK: **Cleavage of structural proteins during the assembly of the head of bacteriophage T4.** *Nature* 1970, **227**:680-5.
31. Gressel J, Rosner A, Cohen N: **Temperature of acrylamide polymerization and electrophoretic mobilities of nucleic acids.** *Anal Biochem* 1975, **69**:84-91.
32. Espejo RT, Puerto F: **Shifts in the electrophoretic pattern on the RNA genome of rotaviruses under different electrophoretic conditions.** *J Virol Methods* 1984, **8**:293-9.
33. Herring AJ, Inglis NF, Ojeh CK, Snodgrass DR, Menzies JD: **Rapid diagnosis of rotavirus infection by direct detection of viral nucleic acid in silver-stained polyacrylamide gels.** *J Clin Microbiol* 1982, **16**:473-7.

doi: 10.1186/1743-422X-7-136

Cite this article as: Long-Croal *et al.*, Concentration of acrylamide in a polyacrylamide gel affects VP4 gene coding assignment of group A equine rotavirus strains with P[12] specificity *Virology Journal* 2010, **7**:136

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

