



Bellini duct carcinoma revealed by cutaneous metastasis: A case report

Amine Oueslati^{a,*}, Ahmed Saadi^a, Marouene Chakroun^a, Selim Zaghbib^a, Ahlem Blel^b, Meriem Ksentini^b, Abderrazak Bouzouita^a, Amine Derouiche^a, Mohamed Riadh Ben Slama^a, Soumaya Rammeh^b, Haroun Ayed^a, Mohamed Chebil^a

^a Department of Urology, Charles Nicolle Hospital, Tunis, Tunisia

^b Department of Pathology, Charles Nicolle Hospital, Tunis, Tunisia



ARTICLE INFO

Article history:

Received 13 July 2020

Received in revised form

22 September 2020

Accepted 22 September 2020

Available online 28 September 2020

Keywords:

Bellini duct carcinoma

Renal tumor

Cutaneous metastasis

Case report

ABSTRACT

INTRODUCTION: Cutaneous metastasis of renal cell carcinoma is rare and the majority of these metastases are asynchronous. The scalp and face are the major sites of metastases, followed by the chest and abdomen. However, the entire body surface can be affected. When diagnosed, patients are multi-metastatic in 50–80% of cases during follow-up post nephrectomy.

CASE PRESENTATION: We report here a patient who consulted a dermatologist for multiple skin nodules that appeared 3 months prior. A skin biopsy of a nodule was performed and the pathological examination and immunohistochemistry profile confirmed a metastasis of Bellini Carcinoma, which is a renal cell carcinoma of the collecting duct. A thoraco-abdomino-pelvic scan showed a left renal tumor locally advanced with lung and liver metastases. Chemotherapy was indicated and the patient died four months after diagnosis.

DISCUSSION: Bellini carcinoma is a very rare type of carcinoma of renal cell origin with a very poor prognosis as it is diagnosed already at a metastatic state in the vast majority of cases. After analysis of the data from the literature, our case is the second reported case of a Bellini carcinoma revealed by cutaneous metastases. The peculiarity of our observation is metastases occurred on all four limbs and at the trunk level, and the asymptomatic characteristic of Bellini's carcinoma, which is a rare situation.

CONCLUSION: The originality of this observation is based on the mode of presentation of a rare renal tumor by an even rarer metastasis of the skin.

© 2020 Published by Elsevier Ltd on behalf of IJS Publishing Group Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Bellini Carcinoma, also known as collecting duct carcinoma is a rare tumor characterized by its aggressivity behavior and has a poor prognosis [1,2]. Most cases are metastatic at the time of diagnosis [1,2]. However, the cutaneous location of metastases is rare and the majority of these metastases are asynchronous. The entire body surface can be affected but the scalp and face are the major sites of metastases, followed by the chest and abdomen [1,2].

We report here the second documented case of Bellini carcinoma revealed by cutaneous metastases. This work has been reported in line with the SCARE criteria [3].

2. Observation

We report a case of a 52-year-old patient with no previous medical history who consulted a dermatologist for multiple skin lesions that have appeared 3 months prior. The patient had not reported any urinary symptom (lower back pain, hematuria...), but he noticed a weight loss with asthenia.

On examination, several cutaneous lesions of varied size and aspect were located on the limbs and trunk (12 lesions in total). At first, the lesions appeared to be hard, infiltrated, and erythematous (Fig. 1), and rapidly became necrotic and infected ulcers (Fig. 2). No abdominal mass was found and the rest of the examination was normal.

A skin biopsy was carried out. Histological examination showed an undifferentiated carcinoma. The diagnosis of multiple cutaneous metastases was suggested and was confirmed by a thoraco-abdomino-pelvic CT scan looking for primary cancer, finding a 52 × 64 mm left renal tumor enhancing at arterial phase contrast. The tumor was described as infiltrating the peri-renal fat with loss of the fatty border within the pancreas, spleen, and the left colic

* Corresponding author.

E-mail address: damine.oueslati.ao@gmail.com (A. Oueslati).



Fig. 1. a: Subcutaneous swelling of right arm coated with erythematous skin dotted with infracentimetric whitening nodules.
b: Erythematous lesion of the left infiltrated and indurated left shoulder, 3 × 4 cm, dotted with infracentimetric whitish nodules.



Fig. 2. Ulceration of the right leg, oval shaped with a 5 cm limited diameter with a necrotic border and erythematous purplish infiltrated edges, indurated and raised.

flexure (Fig. 3a). It also showed multiple mediastinal, lateroaortic, and left renal pedicle lymph nodes metastasis (Fig. 3b) associated with multiple liver and pulmonary metastases.

A second cutaneous biopsy was performed, the histological examination and the immunohistochemistry profile confirmed a metastasis of a renal Bellini carcinoma (Fig. 4).

The patient was in poor general condition, had no hematuria or low back pain. A left nephrectomy was deemed unnecessary to perform and he was instead considered for palliative chemotherapy with gemcitabine-cisplatin.

The patient died after 4 months of diagnosis.

3. Discussion

Renal cell carcinoma of the Bellini collecting ducts is included in the 2004 WHO classification of renal epithelial tumors and accounts for less than 1% of renal carcinomas worldwide [2,4].

The mean age of diagnosis is 66 years old with a standard deviation of range of ages from 20 to 80 years [2].

Unlike most renal cell carcinomas that are discovered fortuitously, Bellini's carcinoma is often symptomatic and most patients have lower back pain and general deterioration [2,4,5].

On the other hand, hematuria may be absent (intra-medullary development of the tumor has a hypovascular characteristic) [6,7]. Our patient had no low back pain nor hematuria, but there was a marked alteration of the general state of health.

This tumor is often aggressive, of high grade and advanced stage at the time of diagnosis and 80% of Bellini carcinoma cases are T3 N+ at the time of diagnosis with an average size of 80 mm [6].

At the time of diagnosis, most patients are metastatic at the regional lymph nodes. This is seen in 80% of the cases, but some cases also have a distant spread with 40% of pulmonary metastases, 30% of liver metastases, and 20% of bone metastases [1,2].

The peculiarity of our observation is twofold. On one hand, our patient had multiple cutaneous metastases on all four limbs and at the trunk level (12 lesions in total) associated with a diffuse ganglionary, hepatic and pulmonary metastases at the time of diagnosis. A single case of Bellini's carcinoma with cutaneous metastases as the initial clinical manifestation was reported by Grande et al. [1]. The patient was a 38-year-old man presenting a 6 cm left mediastinal tumor with unique cutaneous metastasis on his scalp. On the other hand, the second feature is the asymptomatic characteristic of Bellini's carcinoma, which is a rare situation.

Histologically, it is a minimally differentiated adenocarcinoma, predominantly of tubular architecture, often tubulo-papillary. However, its histological presentation may vary [4]. The carcinoma of the collecting tubules is a high-grade carcinoma with marked atypia. Therefore, it's not necessary to specify the Fuhrman grade according to the latest recommendations of the International Society of Urological Pathology [8].

The immunohistochemical study differentiate Bellini duct carcinoma from a high-grade papillary carcinoma, adenocarcinoma




Fig. 3. a: Cross-sectional scan showing a 52 × 64 mm left medial renal tissue mass significantly locally advanced.
b: Coronal section scan showing multiple latero-aortic and the left renal pedicle ganglionary metastasis.




Fig. 4. a: Tumor proliferation in the deeper layer of the dermis (HEx 25).
A: The Epidermis B: Tumor Proliferation.
b: Carcinomatous proliferation of irregular tubulopapillary structures within a fibro-inflammatory stroma (HEx200).
A: Cellular atypias: hyperchromatic volcanic nuclei or highly nucleated vesicular nuclei. B: Diffuse and intense expression of cytokeratin 34βE12.

metastasis, with a possible aid of the use of PAX8 and p63 to distinguish it from an intra-renal infiltrating urothelial carcinoma [4].

The morphology of the renal carcinoma of the collecting tubules and the immunohistochemistry phenotype can be very variable, making its diagnosis sometimes difficult, especially on biopsy specimens [3]. It explains the difficulty in diagnosing Bellini carcinoma on the first biopsy in our case.

Bellini carcinoma has what can be considered a very poor prognosis [1,2,6,9]. The majority of patients are metastatic at the time of diagnosis and half of the patients die within the first year [2,6,7]. The overall survival rate, generally reported, is approximately 60% for the first 6 months, 50% for 1 year and 20% for 2 years after diagnosis [6,7]. In metastatic patients the median survival rate is 6 months [2,6].

Nevertheless, there's an entity that should be differentiated from Bellini's carcinomas, referred as low-grade tubulocystic carcinoma, since it is has a good prognosis. It presents as a spongy and cystic medullary tumor, containing cysts lined with eosinophilic cells [9].

The treatment of Bellini's carcinomas must take into account the fact that it is a very aggressive tumor, with a clinical progression similar to urothelial tumors.

For localized Bellini carcinomas, minimal data is available in published literature for its management and there are only isolated

cases or a series of cases [11]. Surgery is indicated allowing durable remission for some cases [6,7].

However, most patients are metastatic at the time of diagnosis. Cytoreduction nephrectomy, which has shown its benefits for clear cell carcinomas, does not seem to improve the prognosis [1,11,12]. It should be reserved for patients in good general health condition and very symptomatic (back pain or hematuria) [1,4]. Some authors believe that it is completely useless and can even be dangerous with immediate death due to the alteration of the general state of health often observed in these patients [6,11]. They propose a tumor biopsy to avoid surgery [2,6].

In our case, nephrectomy was not performed because the patient was in poor general health condition upon entry. He had no hematuria nor low back pain. Grande et al. performed a nephrectomy with excision of the cutaneous metastasis but the patient died after 7 months [1].

Adjuvant treatments, immunotherapy and radiotherapy have not shown efficacy in the treatment of Bellini carcinoma [2].

Chemotherapy is relatively inactive, out of 23 patients treated with gemcitabine and cisplatin, the response rate was 26%, with progression-free survival and overall survival of 7.1 months and 10.5 months respectively [10]. There are no published studies on the efficacy of tyrosine kinase inhibitors [9].

Thus, in a metastatic situation, chemotherapy with gemcitabine-platinum salts can be proposed while waiting for evidence of targeted therapies [11].

4. Conclusion

Bellini carcinoma is a very rare type of carcinoma of renal cell origin with a very poor prognosis as it is diagnosed at a metastatic state in the vast majority of cases. After analysis of the data from the literature, our case is the second reported case of a Bellini carcinoma revealed by cutaneous metastases. Extended nephrectomy is found to not improve the prognosis of this metastatic carcinoma and chemotherapy with gemcitabine and cisplatin may be suggested.

Declaration of Competing Interest

The authors report no declarations of interest.

Funding

The authors received no specific funding for this work.

Ethical approval

Since this was just a case report and patient's confidentiality was assured, there was no need for ethical approval based on our institution regulations.

Consent

Written informed consent was obtained from the patient for publication of this case report and accompanying images. A copy of the written consent is available for review by the Editor-in-Chief of this journal on request.

Author contribution

A. Oueslati: Writing - original draft.
 A. Saadi: Writing - review & editing.
 M. Chakroun: Writing - review & editing.
 S. Zaghbib: Writing - original draft.
 A Blel Writing - original draft.
 M Ksentini Writing - original draft.
 A. Bouzouita: Supervision ; Reviewing and editing.

A. Derouiche: Supervision ; Reviewing and editing.
M.R. Ben Slama: Supervision ; Reviewing and editing.
S Rammeh Supervision ; Reviewing and editing.
H. Ayed: Supervision ; Reviewing and editing.
M. Chebil: Supervision ; Reviewing and editing.

Registration of research studies

N/A.

Guarantor

Amine Oueslati.

Provenance and peer review

Not commissioned, externally peer-reviewed.

References

- [1] M. Grande, F. Facchini, M. La Rosa, et al., Carcinoma of the collecting ducts of Bellini: a case report with cutaneous metastasis as the initial clinical manifestation, *Urologia* 77 (2010) 51–54.
- [2] M. Roupert, M. Peyromaure, V. Hupertan, et al., Carcinome à cellules rénales de Bellini: diagnostic et traitement, *Prog. Urol.* 14 (2004) 564–567.
- [3] R.A. Agha, M.R. Borrelli, R. Farwana, K. Koshy, A. Fowler, D.P. Orgill, For the SCARE Group, The SCARE 2018 statement: updating Consensus Surgical CAse REport (SCARE) guidelines, *Int. J. Surg.* 60 (2018) 132–136.
- [4] V. Lindner, Tumeursrare du rein. Cas no 1. Carcinoméral des tubes collecteurs de Bellini, *Ann. Pathol.* 34 (2014) 136–140.
- [5] D. Chao, A. Zisman, A.J. Pantuck, et al., Collecting duct renal cell carcinoma: clinical study of a rare tumor, *J. Urol.* 167 (2002) 71–74.
- [6] A. Méjean, M. Roupert, F. Larousserie, et al., Is there a place for radical nephrectomy in the presence of metastatic collecting duct (Bellini) carcinoma? *J. Urol.* 169 (2003) 1287–1290.
- [7] M. Peyromaure, N. Thiounn, F. Scotté, et al., Collecting duct carcinoma of the kidney: a clinicopathological study of 9 cases, *J. Urol.* 170 (2003) 1138–1140.
- [8] J.R. Srigley, B. Delahunt, J.N. Eble, et al., The International Society of Urological Pathology (ISUP) Vancouver classification of renal neoplasia, *Am. J. Surg. Pathol.* 37 (2013) 1469–1489.
- [9] Y. Neuzillet, N. Rioux-Leclercq, B. Escudier, Le point sur les différents types histologiques de cancer du rein et leur traitementspécifique, *Prog. Urol.* 21 (March (Suppl. 2)) (2011) S23–6.
- [10] M.B. Amin, G.T. MacLennan, R. Gupta, et al., Tubulocystic carcinoma of the kidney: clinicopathologic analysis of 31 cases of a distinctive rare subtype of renal cell carcinoma, *Am. J. Surg. Pathol.* 33 (2009) 384–392.
- [11] S. Dason, C. Allard, A. Sheridan-Jonah, et al., Management of renal collecting duct carcinoma: a systematic review and the McMaster experience, *Curr. Oncol.* 20 (2013) 223–232.
- [12] D. You, I.G. Jeong, J.H. Ahn, et al., The value of cytoreductive nephrectomy for metastatic renal cell carcinoma in the era of targeted therapy, *J. Urol.* 185 (2011) 54–59.

Open Access

This article is published Open Access at [sciencedirect.com](https://www.sciencedirect.com). It is distributed under the [IJSCR Supplemental terms and conditions](#), which permits unrestricted non commercial use, distribution, and reproduction in any medium, provided the original authors and source are credited.