

**(*N,N*-Dimethylformamide- $\kappa$ O)bis-(3-hydroxypicolinato- $\kappa^2$ *N,O*<sup>2</sup>)phenylbismuth(III)**

Vitalie Stavila and Kenton H. Whitmire\*

Department of Chemistry, Rice University, 6100 Main Street, Houston, Texas 77005, USA

Correspondence e-mail: whitmir@rice.edu


Received 20 October 2010; accepted 28 October 2010

Key indicators: single-crystal X-ray study;  $T = 294$  K; mean  $\sigma(\text{C-C}) = 0.006$  Å;  $R$  factor = 0.021;  $wR$  factor = 0.048; data-to-parameter ratio = 12.3.

The title organometallic complex,  $[\text{Bi}(\text{C}_6\text{H}_5)(\text{C}_6\text{H}_4\text{NO}_3)_2(\text{C}_3\text{H}_7\text{NO})]$ , features a Bi<sup>III</sup> atom in a distorted pentagonal-pyramidal coordination by two *N,O*-donating bidentate 3-hydroxypicolinate (3-hpic) ligands, one monodentate dimethylformamide (dmf) molecule and one phenyl ring. The C atom of the aryl ligand occupies the apical position of the  $\text{BiCN}_2\text{O}_3$  coordination polyhedron, while the equatorial plane is formed by one O atom of the dmf ligand and two sets of N and O atoms from the chelating 3-hpic ligands. Intermolecular secondary  $\text{Bi}\cdots\text{O}$  [3.485 (3) Å] and  $\text{O}\cdots\text{O}$  hydrogen-bonding interactions connect the complexes into a three-dimensional network. Intramolecular O—H···O hydrogen bonds are also observed.

## Related literature

For a review on the structural chemistry of organobismuth derivatives, see Silvestru *et al.* (1999). For the crystal structures of related arylbismuth(III) compounds, see: Stavila *et al.* (2007, 2009); Stavila & Dikarev (2009); Andrews *et al.* (2006); Yu *et al.* (2004). For bismuth(III) picolinate complexes, see: Callens *et al.* (2008). For a review on biomedical applications of bismuth(III) compounds, see: Briand & Burford (1999).



## Experimental

### Crystal data

$[\text{Bi}(\text{C}_6\text{H}_5)(\text{C}_6\text{H}_4\text{NO}_3)_2(\text{C}_3\text{H}_7\text{NO})]$	$V = 2173.6$ (7) Å <sup>3</sup>
$M_r = 635.38$	$Z = 4$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation
$a = 8.2377$ (16) Å	$\mu = 8.16$ mm <sup>-1</sup>
$b = 21.989$ (4) Å	$T = 294$ K
$c = 12.380$ (3) Å	$0.14 \times 0.11 \times 0.10$ mm
$\beta = 104.24$ (3)°	

### Data collection

Bruker SMART 1000 CCD diffractometer	15126 measured reflections
Absorption correction: multi-scan ( <i>SADABS</i> ; Bruker, 2004)	3669 independent reflections
$T_{\min} = 0.356$ , $T_{\max} = 0.450$	3314 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.031$

### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.021$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.048$	$\Delta\rho_{\text{max}} = 0.72$ e Å <sup>-3</sup>
$S = 1.11$	$\Delta\rho_{\text{min}} = -0.94$ e Å <sup>-3</sup>
3669 reflections	
299 parameters	
4 restraints	

**Table 1**  
Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
O13—H13A···O12	0.84 (3)	1.80 (3)	2.541 (5)	147 (4)
O23—H23A···O22	0.84 (3)	1.79 (3)	2.555 (5)	150 (3)
O23—H23A···O13 <sup>i</sup>	0.84 (3)	2.52 (2)	2.917 (6)	110 (2)

Symmetry code: (i)  $x + 1, y, z + 1$ .

Data collection: *SMART* (Bruker, 2004); cell refinement: *SAINT-Plus* (Bruker, 2004); data reduction: *SAINT-Plus* and *XPREP* (Bruker, 2004); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

The authors thank the CRDF for financial support (award No. MOE2-2850-CS-06).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: SJ5046).

## References

- Andrews, P. C., Deacon, G. B., Junk, P. C., Kumar, I. & Silberstein, M. (2006). *Dalton Trans.* pp. 4852–4858.
- Briand, G. G. & Burford, N. (1999). *Chem. Rev.* **99**, 2601–2658.
- Bruker (2004). *SMART, SAINT-Plus, XPREP* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Callens, E., Burton, A. J., White, A. J. P. & Barrett, A. G. M. (2008). *Tetrahedron Lett.* **49**, 3709–3712.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Silvestru, C., Breunig, H. J. & Althaus, H. (1999). *Chem. Rev.* **99**, 3277–3327.
- Stavila, V. & Dikarev, E. V. (2009). *J. Organomet. Chem.* **694**, 2956–2964.
- Stavila, V., Fettinger, J. C. & Whitmire, K. H. (2007). *Organometallics*, **26**, 3321–3328.
- Stavila, V., Thurston, J. H. & Whitmire, K. H. (2009). *Inorg. Chem.* **48**, 6945–6951.
- Yu, L., Ma, Y.-Q., Liu, R.-C., Wang, G.-C. & Li, J.-S. (2004). *Inorg. Chem. Commun.* **7**, 410–411.

## **supplementary materials**

Acta Cryst. (2010). E66, m1547-m1548 [doi:10.1107/S1600536810044235]

**(*N,N*-Dimethylformamide- $\kappa O$ )bis(3-hydroxypicolinato- $\kappa^2 N,O^2$ )phenylbismuth(III)**

**V. Stavila and K. H. Whitmire**

**Comment**

The title compound, (I), was obtained from the solvent-free reaction of BiPh<sub>3</sub> and 3-hydroxypicolinic acid (3-hpicH) with subsequent recrystallization from dimethylformamide (dmf) (see Experimental). Single-crystals suitable for X-Ray crystallography were obtained at room temperature from the concentrated dmF solution. The structure determination of (I) revealed a mononuclear compound, in which the Bi<sup>III</sup> atoms are hexa-coordinated in a distorted pentagonal pyramidal geometry with two N and two O atoms of N,O-chelating 3-hpic ligands and one dmF O donor in the equatorial plane (Figure 1). The axial position of the pentagonal pyramid is occupied by a carbon atom of the aryl group (Bi1—C41 = 2.245 (4) Å). Both picolinate ligands are monodeprotonated and display N,O-chelation through the pyridine N and carboxylate O atoms. There is an important asymmetry in the 3-hpic coordination to Bi<sup>III</sup> (Bi1—N1 = 2.660 (3) Å, Bi1—O11 = 2.348 (3) Å; Bi1—N2 = 2.488 (3) Å, Bi1—O21 = 2.382 (3) Å). The O atom of the coordinated dmF molecule (Bi1—O31 = 2.534 (3) Å) completes the equatorial plane of the pyramid.

There is a relatively large variation in the equatorial angles of the pyramid (64.97–77.37°) due to the difference in Bi—N and Bi—O bond lengths. Although the atoms Bi1, N1, O11, N2, O21 and O31 are not exactly coplanar, the sum of the corresponding angles is close to 360°, 359.80 (9)° (N1—Bi1—O11 = 64.98 (9)°, O11—Bi1—N2 = 74.25 (9)°, N2—Bi1—O21 = 67.72 (9)°, O21—Bi1—O31 = 77.37 (9)°, O31—Bi1—N1 = 75.49 (9)°). The C—Bi—O and C—Bi—N angles deviate from 90° (83.65–92.00°), contributing to the distortion of the pentagonal pyramidal coordination around the Bi<sup>III</sup> atom (Figure 2). Similar to other structurally characterized arylbismuth(III) compounds, the coordination sphere of Bi<sup>III</sup> is *hemidirected* (Stavila *et al.*, 2007, 2009), (Stavila & Dikarev, 2009), suggesting that a stereochemically active lone electron pair is present.

Generally, secondary bonding interactions are rather common for monoaryl-bismuth(III) complexes. Thus, intermolecular secondary bonding interactions have been found in a number of aryl-bismuth diketonates (Stavila & Dikarev, 2009) and carboxylates (Stavila *et al.*, 2007). In (I), the oxygen atom of one of the hydroxyl groups, O23, is involved in a weak secondary bond (Bi1···O23 = 3.485 (3) Å with an adjacent Bi<sup>III</sup> complex (Figure 3). The complex also displays intra- and intermolecular hydrogen bonds between the OH groups and oxygen atoms of the carboxylate groups (O13—H13A···O12, O13···O12 = 2.541 (5) Å; O23—H23A···O22, O23···O22 = 2.555 (5) Å; O23—H23A···O13<sup>i</sup>, O23···O13<sup>i</sup> = 2.917 (6) Å, (i)  $x + 1, y, z + 1$ ).

Structures containing aryl bismuth(III) complexes with O or O/N donors typically display pentagonal pyramidal geometries and are comparatively rare (Andrews *et al.*, 2006; Stavila & Dikarev, 2009; Stavila *et al.*, 2007; Yu *et al.*, 2004). In the same way, structures of bismuth(III) complexes with chelating picolinate ligands are uncommon (Callens *et al.*, 2008).

# supplementary materials

---

## Experimental

The initial reactants used were obtained commercially from Strem and Sigma-Aldrich. In a nitrogen filled glove-box, triphenylbismuth (440 mg, 1.0 mmol) and 3-hydroxypicolinic acid (420 mg, 3.0 mmol) were ground together for 30 min resulting in a light-grey powder. The mixture was placed in a Schlenk tube and heated upon stirring at 120 °C for 90 min. The resulting grey powder is treated with dry dmf, then filtered. The filtered solution is concentrated to ~1/4 of its initial volume and left for crystallization at room temperature. Crystals suitable for single-crystal X-ray crystallography were formed in 4 weeks.

## Refinement

The H atoms bound to O13 and O23 were located in a difference map and their coordinates were refined with  $U_{\text{iso}}(\text{H})$  values of  $1.2U_{\text{eq}}(\text{O})$ . C-bound H atoms were located in calculated positions and constrained to ride on their parent atoms at distances of  $d(\text{C}-\text{H}) = 0.93\text{\AA}$ ,  $U_{\text{iso}}=1.2U_{\text{eq}}(\text{C})$  for aromatic and  $0.96\text{\AA}$ ,  $U_{\text{iso}} = 1.5U_{\text{eq}}(\text{C})$  for  $\text{CH}_3$  atoms

## Figures




Fig. 1. An anisotropic displacement ellipsoid plot of the title compound (I), showing 40% probability displacement ellipsoids.




Fig. 2. The coordination polyhedron of the  $\text{Bi}^{\text{III}}$  ion in compound (I).




Fig. 3. A packing diagram of compound (I) viewed down the  $a$  axis.

## $(N,N\text{-Dimethylformamide-}\kappa O)\text{bis}(3\text{-hydroxypicolinato-}\kappa^2 N,O^2)\text{phenylbismuth(III)}$

### Crystal data

$[\text{Bi}(\text{C}_6\text{H}_5)(\text{C}_6\text{H}_4\text{NO}_3)_2(\text{C}_3\text{H}_7\text{NO})]$

$F(000) = 1224$

$M_r = 635.38$

$D_x = 1.942 \text{ Mg m}^{-3}$

Monoclinic,  $P2_1/c$

Mo  $K\alpha$  radiation,  $\lambda = 0.71073 \text{ \AA}$

Hall symbol: -P 2ybc

Cell parameters from 7774 reflections

$a = 8.2377 (16)$ Å	$\theta = 2.5\text{--}24.7^\circ$
$b = 21.989 (4)$ Å	$\mu = 8.16 \text{ mm}^{-1}$
$c = 12.380 (3)$ Å	$T = 294$ K
$\beta = 104.24 (3)^\circ$	Block, light-yellow
$V = 2173.6 (7)$ Å <sup>3</sup>	$0.14 \times 0.11 \times 0.10$ mm
$Z = 4$	

*Data collection*

Bruker SMART 1000 CCD diffractometer	3669 independent reflections
Radiation source: fine-focus sealed tube graphite	3314 reflections with $I > 2\sigma(I)$
$\omega$ scans	$R_{\text{int}} = 0.031$
Absorption correction: multi-scan ( <i>SADABS</i> ; Bruker, 2004)	$\theta_{\text{max}} = 24.8^\circ, \theta_{\text{min}} = 1.9^\circ$
$T_{\text{min}} = 0.356, T_{\text{max}} = 0.450$	$h = -9 \rightarrow 9$
15126 measured reflections	$k = -25 \rightarrow 25$
	$l = -14 \rightarrow 13$

*Refinement*

Refinement on $F^2$	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.021$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.048$	H atoms treated by a mixture of independent and constrained refinement
$S = 1.11$	$w = 1/[\sigma^2(F_o^2) + (0.0202P)^2 + 1.6444P]$
3669 reflections	where $P = (F_o^2 + 2F_c^2)/3$
299 parameters	$(\Delta/\sigma)_{\text{max}} < 0.001$
4 restraints	$\Delta\rho_{\text{max}} = 0.72 \text{ e \AA}^{-3}$
	$\Delta\rho_{\text{min}} = -0.94 \text{ e \AA}^{-3}$

*Special details*

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) etc. and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

## supplementary materials

---

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Bi1	0.163597 (16)	0.410112 (6)	0.270490 (11)	0.02944 (6)
N1	-0.0947 (4)	0.39564 (14)	0.0950 (3)	0.0350 (7)
N2	0.4598 (3)	0.44505 (13)	0.3132 (2)	0.0296 (7)
N3	-0.1613 (4)	0.30791 (16)	0.4752 (3)	0.0484 (9)
O11	0.2210 (3)	0.43021 (14)	0.0973 (2)	0.0460 (7)
O12	0.1580 (4)	0.43682 (17)	-0.0856 (2)	0.0616 (9)
O13	-0.1422 (4)	0.42463 (17)	-0.1993 (3)	0.0700 (10)
H13A	-0.047 (2)	0.4405 (17)	-0.179 (3)	0.084*
O21	0.3075 (3)	0.39088 (11)	0.4591 (2)	0.0345 (6)
O22	0.5381 (3)	0.40081 (12)	0.5967 (2)	0.0431 (7)
O23	0.7918 (3)	0.45888 (13)	0.5663 (2)	0.0424 (7)
H23A	0.721 (3)	0.4455 (17)	0.599 (2)	0.051*
O31	-0.0596 (3)	0.35957 (13)	0.3485 (2)	0.0463 (7)
C11	0.1176 (5)	0.42747 (18)	0.0033 (3)	0.0384 (9)
C12	-0.0582 (5)	0.41051 (16)	-0.0012 (3)	0.0342 (8)
C13	-0.1795 (6)	0.41097 (19)	-0.1030 (4)	0.0483 (11)
C14	-0.3443 (6)	0.3973 (2)	-0.1012 (4)	0.0602 (13)
H14A	-0.4287	0.3973	-0.1668	0.072*
C15	-0.3797 (5)	0.3840 (2)	-0.0021 (4)	0.0568 (12)
H15A	-0.4893	0.3757	0.0007	0.068*
C16	-0.2524 (5)	0.3828 (2)	0.0945 (4)	0.0446 (10)
H16A	-0.2781	0.3727	0.1614	0.053*
C21	0.4577 (5)	0.40784 (15)	0.4986 (3)	0.0311 (8)
C22	0.5437 (4)	0.43992 (15)	0.4213 (3)	0.0279 (8)
C23	0.7050 (4)	0.46376 (16)	0.4596 (3)	0.0310 (8)
C24	0.7768 (4)	0.49391 (17)	0.3847 (3)	0.0372 (9)
H24A	0.8836	0.5105	0.4082	0.045*
C25	0.6894 (5)	0.49916 (19)	0.2758 (3)	0.0405 (10)
H25A	0.7360	0.5194	0.2247	0.049*
C26	0.5304 (5)	0.47394 (18)	0.2424 (3)	0.0381 (9)
H26A	0.4716	0.4774	0.1682	0.046*
C31	-0.0559 (5)	0.34549 (19)	0.4457 (4)	0.0421 (10)
H31A	0.024 (5)	0.3600 (19)	0.512 (4)	0.056 (13)*
C32	-0.1501 (7)	0.2935 (3)	0.5913 (4)	0.0744 (16)
H32A	-0.0565	0.3146	0.6378	0.112*
H32B	-0.1352	0.2505	0.6026	0.112*
H32C	-0.2513	0.3060	0.6104	0.112*
C33	-0.2927 (7)	0.2800 (3)	0.3906 (5)	0.0903 (19)
H33A	-0.2842	0.2932	0.3183	0.135*
H33B	-0.3997	0.2916	0.4017	0.135*
H33C	-0.2814	0.2365	0.3955	0.135*
C41	0.2374 (4)	0.31401 (17)	0.2439 (3)	0.0341 (8)
C42	0.2978 (6)	0.2968 (2)	0.1537 (4)	0.0550 (12)
H42A	0.3107	0.3258	0.1019	0.066*
C43	0.3391 (6)	0.2369 (3)	0.1397 (4)	0.0666 (14)

H43A	0.3812	0.2263	0.0791	0.080*
C44	0.3194 (6)	0.1938 (3)	0.2124 (4)	0.0676 (16)
H44A	0.3451	0.1535	0.2011	0.081*
C45	0.2610 (6)	0.2096 (2)	0.3034 (4)	0.0602 (13)
H45A	0.2486	0.1800	0.3545	0.072*
C46	0.2204 (5)	0.26970 (18)	0.3191 (4)	0.0453 (10)
H46A	0.1814	0.2802	0.3810	0.054*

*Atomic displacement parameters ( $\text{\AA}^2$ )*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Bi1	0.03053 (9)	0.03335 (9)	0.02336 (9)	-0.00432 (6)	0.00457 (6)	0.00018 (6)
N1	0.0321 (17)	0.0393 (18)	0.0307 (19)	0.0017 (13)	0.0023 (14)	0.0009 (14)
N2	0.0325 (16)	0.0342 (16)	0.0216 (16)	-0.0069 (13)	0.0061 (13)	0.0022 (13)
N3	0.0418 (19)	0.053 (2)	0.052 (2)	-0.0026 (16)	0.0139 (17)	0.0137 (18)
O11	0.0412 (15)	0.0690 (19)	0.0247 (15)	-0.0184 (14)	0.0023 (12)	0.0043 (13)
O12	0.062 (2)	0.095 (2)	0.0273 (17)	-0.0225 (18)	0.0102 (15)	0.0074 (16)
O13	0.071 (2)	0.105 (3)	0.0252 (18)	-0.019 (2)	-0.0061 (16)	0.0155 (17)
O21	0.0361 (14)	0.0410 (14)	0.0254 (14)	-0.0129 (11)	0.0057 (11)	0.0031 (11)
O22	0.0423 (16)	0.0582 (18)	0.0249 (16)	-0.0082 (12)	0.0006 (12)	0.0097 (12)
O23	0.0358 (15)	0.0568 (18)	0.0301 (16)	-0.0098 (13)	-0.0005 (12)	0.0040 (13)
O31	0.0390 (15)	0.0590 (18)	0.0429 (18)	-0.0015 (13)	0.0141 (13)	0.0088 (14)
C11	0.048 (2)	0.040 (2)	0.026 (2)	-0.0058 (17)	0.0056 (18)	0.0034 (17)
C12	0.040 (2)	0.033 (2)	0.025 (2)	-0.0022 (16)	-0.0009 (16)	0.0041 (15)
C13	0.056 (3)	0.049 (3)	0.032 (3)	-0.005 (2)	-0.002 (2)	0.0060 (19)
C14	0.043 (3)	0.070 (3)	0.052 (3)	-0.005 (2)	-0.018 (2)	0.005 (2)
C15	0.036 (2)	0.072 (3)	0.056 (3)	-0.005 (2)	-0.001 (2)	0.004 (3)
C16	0.038 (2)	0.054 (3)	0.041 (3)	-0.0027 (19)	0.0094 (19)	0.000 (2)
C21	0.036 (2)	0.0296 (19)	0.027 (2)	-0.0038 (15)	0.0065 (17)	-0.0030 (15)
C22	0.0313 (18)	0.0269 (18)	0.025 (2)	0.0009 (14)	0.0069 (15)	-0.0014 (15)
C23	0.0310 (18)	0.0309 (19)	0.030 (2)	0.0008 (15)	0.0058 (16)	-0.0020 (16)
C24	0.0295 (19)	0.043 (2)	0.039 (2)	-0.0086 (16)	0.0087 (17)	-0.0041 (18)
C25	0.041 (2)	0.049 (2)	0.035 (2)	-0.0106 (18)	0.0179 (19)	0.0022 (18)
C26	0.041 (2)	0.048 (2)	0.025 (2)	-0.0090 (18)	0.0059 (17)	0.0032 (17)
C31	0.033 (2)	0.045 (2)	0.048 (3)	0.0026 (18)	0.008 (2)	0.003 (2)
C32	0.068 (3)	0.095 (4)	0.064 (4)	0.007 (3)	0.022 (3)	0.037 (3)
C33	0.082 (4)	0.092 (4)	0.093 (5)	-0.042 (3)	0.014 (3)	0.005 (4)
C41	0.0299 (19)	0.040 (2)	0.029 (2)	0.0004 (16)	0.0014 (16)	-0.0051 (17)
C42	0.059 (3)	0.066 (3)	0.042 (3)	0.012 (2)	0.015 (2)	-0.002 (2)
C43	0.074 (3)	0.078 (4)	0.048 (3)	0.027 (3)	0.014 (3)	-0.018 (3)
C44	0.071 (3)	0.059 (3)	0.060 (4)	0.027 (3)	-0.007 (3)	-0.021 (3)
C45	0.069 (3)	0.046 (3)	0.061 (3)	0.009 (2)	0.005 (3)	0.005 (2)
C46	0.054 (2)	0.040 (2)	0.041 (2)	0.0068 (19)	0.011 (2)	-0.002 (2)

*Geometric parameters ( $\text{\AA}$ ,  $^\circ$ )*

Bi1—C41	2.245 (4)	O31—C31	1.235 (5)
Bi1—O11	2.348 (3)	C11—C12	1.483 (5)
Bi1—O21	2.382 (3)	C12—C13	1.403 (6)

## supplementary materials

---

Bi1—N2	2.488 (3)	C13—C14	1.396 (7)
Bi1—O31	2.534 (3)	C14—C15	1.360 (7)
Bi1—N1	2.660 (3)	C15—C16	1.384 (6)
N1—C16	1.328 (5)	C21—C22	1.499 (5)
N1—C12	1.338 (5)	C22—C23	1.398 (5)
N2—C26	1.327 (4)	C23—C24	1.386 (5)
N2—C22	1.352 (4)	C24—C25	1.368 (5)
N3—C31	1.313 (5)	C25—C26	1.388 (5)
N3—C33	1.445 (6)	C41—C46	1.378 (6)
N3—C32	1.453 (6)	C41—C42	1.382 (5)
O11—C11	1.264 (5)	C42—C43	1.381 (7)
O12—C11	1.243 (5)	C43—C44	1.345 (7)
O13—C13	1.336 (5)	C44—C45	1.374 (7)
O21—C21	1.269 (4)	C45—C46	1.389 (6)
O22—C21	1.242 (5)	O13—H13A	0.84 (3)
O23—C23	1.342 (4)	O23—H23A	0.84 (3)
C41—Bi1—O11	85.44 (12)	N1—C12—C11	117.3 (3)
C41—Bi1—O21	83.65 (11)	C13—C12—C11	120.3 (4)
O11—Bi1—O21	139.90 (9)	O13—C13—C14	120.1 (4)
C41—Bi1—N2	92.01 (11)	O13—C13—C12	122.3 (4)
O11—Bi1—N2	74.25 (9)	C14—C13—C12	117.7 (4)
O21—Bi1—N2	67.72 (9)	C15—C14—C13	119.1 (4)
C41—Bi1—O31	83.76 (12)	C14—C15—C16	119.9 (4)
O11—Bi1—O31	139.33 (9)	N1—C16—C15	122.1 (4)
O21—Bi1—O31	77.37 (9)	O22—C21—O21	125.1 (3)
N2—Bi1—O31	145.09 (9)	O22—C21—C22	117.4 (3)
C41—Bi1—N1	87.62 (11)	O21—C21—C22	117.5 (3)
O11—Bi1—N1	64.98 (10)	N2—C22—C23	121.1 (3)
O21—Bi1—N1	152.21 (9)	N2—C22—C21	117.7 (3)
N2—Bi1—N1	139.13 (10)	C23—C22—C21	121.3 (3)
O31—Bi1—N1	75.49 (10)	O23—C23—C24	119.0 (3)
C16—N1—C12	118.8 (3)	O23—C23—C22	122.4 (3)
C16—N1—Bi1	127.9 (3)	C24—C23—C22	118.6 (3)
C12—N1—Bi1	112.8 (2)	C25—C24—C23	119.5 (3)
C26—N2—C22	119.6 (3)	C24—C25—C26	119.3 (3)
C26—N2—Bi1	125.0 (2)	N2—C26—C25	121.9 (3)
C22—N2—Bi1	114.9 (2)	O31—C31—N3	124.7 (4)
C31—N3—C33	119.6 (4)	C46—C41—C42	117.9 (4)
C31—N3—C32	121.8 (4)	C46—C41—Bi1	119.3 (3)
C33—N3—C32	118.6 (4)	C42—C41—Bi1	122.7 (3)
C11—O11—Bi1	126.2 (2)	C43—C42—C41	120.8 (5)
C21—O21—Bi1	121.9 (2)	C44—C43—C42	120.9 (5)
C31—O31—Bi1	129.4 (3)	C43—C44—C45	119.7 (5)
O12—C11—O11	122.8 (4)	C44—C45—C46	120.0 (5)
O12—C11—C12	118.7 (3)	C41—C46—C45	120.7 (4)
O11—C11—C12	118.5 (3)	O13—H13A—O12	147 (4)
N1—C12—C13	122.4 (4)		

*Hydrogen-bond geometry (Å, °)*

$D\text{---H}\cdots A$	$D\text{---H}$	$H\cdots A$	$D\cdots A$	$D\text{---H}\cdots A$
O13—H13A···O12	0.84 (3)	1.80 (3)	2.541 (5)	147 (4)
O23—H23A···O22	0.84 (3)	1.79 (3)	2.555 (5)	150 (3)
O23—H23A···O13 <sup>i</sup>	0.84 (3)	2.52 (2)	2.917 (6)	110 (2)

Symmetry codes: (i)  $x+1, y, z+1$ .

## supplementary materials

---

Fig. 1





Fig. 2



## supplementary materials

---

Fig. 3

