

Fukushima-derived radionuclides in sediments of the Japanese Pacific Ocean coast and various Japanese water samples (seawater, tap water, and coolant water of Fukushima Daiichi reactor unit 5)

Katsumi Shozugawa¹ · Beate Riebe² · Clemens Walther² · Alexander Brandl³ · Georg Steinhauser^{2,3}

Received: 3 June 2015 / Published online: 23 August 2015
© The Author(s) 2015. This article is published with open access at Springerlink.com

Abstract We investigated Ocean sediments and seawater from inside the Fukushima exclusion zone and found radiocesium (^{134}Cs and ^{137}Cs) up to 800 Bq kg^{-1} as well as ^{90}Sr up to 5.6 Bq kg^{-1} . This is one of the first reports on radiostrontium in sea sediments from the Fukushima exclusion zone. Seawater exhibited contamination levels up to 5.3 Bq kg^{-1} radiocesium. Tap water from Tokyo from weeks after the accident exhibited detectable but harmless activities of radiocesium (well below the regulatory limit). Analysis of the Unit 5 reactor coolant (finding only ^3H and even low ^{129}I) leads to the conclusion that the purification techniques for reactor coolant employed at Fukushima Daiichi are very effective.

Keywords Fukushima · Environmental radioactivity · Pacific Ocean · Cesium · ^{129}I · ^{90}Sr · ^3H

Introduction

Following the earthquake on March 11, 2011, a gigantic tsunami destroyed the cooling systems of Fukushima Daiichi nuclear power plant (NPP)(Japan) and caused partial melt-

down of 3 reactor cores. In the course of the nuclear accident a total of 520 PBq (excl. noble gases) were released to the atmosphere [1]. More than 99 % of the released substances were radionuclides of Kr, Te, I, Xe, and Cs. These nuclides have been monitored globally in air [2, 3]. Less volatile radionuclides such as radiostrontium [4, 5], plutonium [6–8] or radionuclides that are difficult to measure, such as ^3H [9, 10], ^{135}Cs [11] or ^{35}S [12] have been monitored much less frequently [13]. In any case, at least 80 % of the airborne radionuclides were transported offshore by the wind [14, 15], and a yet unknown amount has leaked directly into the Pacific Ocean. Although several studies have addressed the impact of the accident on the Pacific Ocean and its organisms [16–18], our knowledge on the impact of the Fukushima nuclear accident on the marine environment is yet far from complete.

In the current study, we investigated several interesting and unique sample materials, many of which were taken directly after the accident and/or inside the “exclusion zone” around the crippled reactors. Target nuclides were radiocesium [^{134}Cs ($T_{1/2} = 2.1$ years) and ^{137}Cs ($T_{1/2} = 30.2$ years)] as well as the understudied nuclides ^{90}Sr ($T_{1/2} = 28.6$ years) and ^3H ($T_{1/2} = 12.3$ years). For two samples, also ^{129}I analysis was performed. It is a long-lived ($T_{1/2} = 15.7$ million years) fission product that can be used as an (environmental) tracer nuclide [19] that is also useful for retrospective dosimetry after the decay of short-lived ^{131}I ($T_{1/2} = 8$ days) [20] which accumulates in the thyroid [21] and is mostly responsible for thyroid cancer cases after Chernobyl.

Materials and methods

Environmental materials analyzed in this study included sediments and seawater from the Japanese Pacific Ocean coast inside the exclusion zone, tap water from Tokyo,

✉ Georg Steinhauser
georg.steinhauser@ati.ac.at;
georg.steinhauser@colostate.edu; steinhauser@irs.uni-hannover.de

¹ Graduate School of Arts and Sciences, The University of Tokyo, Meguro-Ku, Tokyo, Japan

² Institute of Radioecology and Radiation Protection, Leibniz Universität Hannover, Herrenhäuser Straße 2, 30419 Hannover, Germany

³ Environmental and Radiological Health Sciences, Colorado State University, 1618 Campus Delivery, Fort Collins, CO 80523, USA

seawater from Hawaii as well as a sample of the reactor coolant water of the undamaged Unit 5 of Fukushima Daiichi NPP. The coolant was investigated also for ^{129}I to scrutinize the overall efficiency of the techniques currently employed for radionuclide removal from water at the Fukushima Daiichi NPP; most fission and activation products are cations that can be removed using natural or artificial cation exchanger. Iodine, however, is usually anionic (iodide or iodate) and thus more challenging to remove. A summary of all samples, the exact location and date of sampling, distance to the Fukushima Daiichi NPP as well as target nuclides is given in Table 1.

A map of the sample locations (excluding the seawater sample from Hawaii), is given in Fig. 1.

Gamma measurements for ^{134}Cs and ^{137}Cs were conducted with an ORTECTM 364 cm³ HPGe detector with a 0.76 mm Be window (2.32 keV resolution at the 1332 keV ^{60}Co peak; 87.4 % relative efficiency). The samples were weighed into 60 or 125 mL polypropylene containers (typically 100 mL for water samples; 70 g for sediment samples) that were placed on top of the detector. The samples were measured for at least 24 h. Detector efficiency calibration was done with an Eckert and Ziegler[®] multinuclide standard solution that was diluted in 2 M HCl (aqueous samples) or dispersed in quartz sand (sediment samples) (see [22]). Tritium (^3H) and ^{90}Sr was conducted with liquid scintillation counting (LSC). The LSC measurements were performed using a LabLogic 300SL Liquid Scintillation Counter with TDCR Technology (Hidex, Mustionkatu 2, FIN-20750 Turku, Finland). The user interface software is provided by MikroWinTM (Mikrotek




Fig. 1 Geographical setting of the sampling sites in Japan

Laborsysteme GmbH, Olper Strasse 35, D-51491 Overath, Germany). Instrument output was transferred to a vendor-provided PC-application spreadsheet with macros where the raw data could be analyzed and a graphic routine provided for convenient data visualization [23].

For radiocesium measurements, the sediment samples were dried at 100 °C, weighed and filled into calibrated geometries at the gamma detector station at Colorado State University. Water samples were weighed and measured for gamma emitters without further pre-treatment.

Table 1 Samples investigated in this study

Sample type	Sample no./code	Sample location	Sampling date	Distance to FDNPP (km)	Radionuclides analyzed
Ocean sediment	Sed 1	37.4040258; 141.033522	2011-12-20	1.95	^{90}Sr , $^{134,137}\text{Cs}$
Ocean sediment	Sed 2	37.478988; 141.039604	2011-12-20	6.54	^{90}Sr , $^{134,137}\text{Cs}$
Ocean sediment	Sed 3	37.567682; 141.026177	2011-12-20	16.1	^{90}Sr , $^{134,137}\text{Cs}$
Ocean sediment	Sed 4	37.21161; 141.005861	2011-12-20	23.5	^{90}Sr , $^{134,137}\text{Cs}$
Ocean sediment	Sed 5	37.13266; 140.998351	2011-12-20	32.2	^{90}Sr , $^{134,137}\text{Cs}$
Ocean water	OW 1	37.4040258; 141.033522	2011-12-20	1.95	^3H , ^{90}Sr , $^{134,137}\text{Cs}$
Ocean water	OW 2	37.478988; 141.039604	2011-12-20	6.54	^3H , ^{90}Sr , $^{134,137}\text{Cs}$
Ocean water	OW 3	37.567682; 141.026177	2011-12-20	16.1	^3H , ^{90}Sr , $^{134,137}\text{Cs}$
Ocean water	OW 4	37.13266; 140.998351	2011-12-20	23.5	^3H , ^{90}Sr , $^{134,137}\text{Cs}$
Ocean water	OW 5	37.21161; 141.005861	2011-12-20	32.2	^3H , ^{90}Sr , $^{134,137}\text{Cs}$
Tap water	TW 6	Tokyo, Japan	2011-03-21	225	^3H , ^{90}Sr , $^{134,137}\text{Cs}$
Tap water	TW 7	Tokyo, Japan	2011-03-24	225	^3H , ^{90}Sr , $^{134,137}\text{Cs}$
Tap water	TW 8	Tokyo, Japan	2011-03-27	225	^3H , ^{90}Sr , $^{134,137}\text{Cs}$
Tap water	TW 9	Tokyo, Japan	2011-03-31	225	^3H , ^{90}Sr , $^{134,137}\text{Cs}$
Hawaii Ocean water	HI W	Kona, Big Island, HI	2012-03-31	6350	^{129}I
Reactor unit 5 coolant water	CW #5	37.428526; 141.032841	2011-10-22	<0.75	^3H , ^{90}Sr , ^{129}I , $^{134,137}\text{Cs}$

Pure beta emitters such as ^{90}Sr or ^3H require chemical separation and purification prior to beta counting in the LSC. Water samples were carefully distilled to dryness; the distillate was mixed with Ultima GoldTM low level tritium (LLT) scintillation cocktail for measurement. Ocean sediments were treated with nitric acid for Sr leaching and purified using the EichromTM SR resin (2 mL cartridges), as described elsewhere [4]. In brief, the separation process involved weighing of the dried sand into a flask, addition of 8 mL HNO_3 (8 M), 2 mL H_2O_2 (30 %), 1 mL Sr carrier solution ($c_{\text{Sr}} = 1.2 \text{ mg mL}^{-1}$), and 3 mL of concentrated HNO_3 (70 %). This solution was boiled under reflux for 30 min and then filtered through a paper filter. The filtrate was loaded onto preconditioned (with 8 M HNO_3) SR resin cartridges. The flask was rinsed with $3 \times 2 \text{ mL HNO}_3$ (8 M). The resin was then rinsed with $10 \times 1 \text{ mL}$ of a mixture of HNO_3 (3 M) and oxalic acid (0.05 M). Then the Sr fraction was eluted with 0.01 M HNO_3 , which we found to be sufficiently acidic to not eluate any radiolabel from the column. Water samples (50 mL) were either brought to 8 M HNO_3 by addition of equal amounts of 16 M HNO_3 , or the solid residue after tritium distillation was taken up in 8 M HNO_3 . Since the natural Sr content in water is much lower than from leached sediments, the amount of Sr carrier was increased to 1.7 mg. The loading, rinsing and eluting processes were conducted as described above.

We found that the acidic eluate (0.01 M HNO_3) does not mix well with the Ultima GoldTM LSC cocktail. In order to remove the acid, the eluate was evaporated to almost dryness and then taken up in 1 mL H_2O again. This step was repeated 10 times. Then, the Sr fraction was transferred to LSC vials and the flask was rinsed with $4 \times 0.5 \text{ mL H}_2\text{O}$. Finally 18 mL of scintillation cocktail were added.

Addition of HNO_3 to sediments should be performed carefully as the carbonate fraction of the sand will vigorously decompose under formation of CO_2 gas which could cause the overflow of the flask with acidic foam. In our case, the sediments proved to have a very low carbonate content. Also organic substances may react violently upon exposure to the highly oxidizing $\text{HNO}_3/\text{H}_2\text{O}_2$ mixture, especially when heated. If in doubt, the use of protective measures (such as gloves, face shields) is recommended [24–26].

For the accelerator mass spectrometry (AMS) measurements of ^{129}I , iodine was separated from the matrix in two steps. First, Woodward iodine carrier was added and all the iodine was transferred into iodide. All iodine species were oxidized with $\text{Ca}(\text{OCl})_2$ to iodate and afterwards reduced with NH_3OHCl and NaHSO_3 to iodide. The second step consisted in an ion exchange separation using a DOWEX[®] 1 \times 8 analytical grade ion exchange resin, which was preconditioned with KNO_3 . After rinsing the ion

exchange columns with high purity water and a $0.5 \text{ mol L}^{-1} \text{ KNO}_3$ solution, the iodine was eluted with concentrated potassium nitrate solution (2.25 mol L^{-1}). The iodine was precipitated as AgI , which was dried, mixed with silver powder ($\text{AgI}:\text{Ag}$ 1:4 by weight), and pressed into titanium targets for AMS measurement. Measurements were performed with low-energy accelerator mass spectrometry (AMS) at the 0.5 MV TANDY facility at ETH Zurich, Switzerland.

For the ICP-MS measurement of ^{127}I no iodine matrix separation was necessary. The samples were diluted and tetramethylammoniumhydroxide (TMAH) was added, which causes hydrolysis of any organic compounds and reduces the redox potential. The stability of I^- increases in the sample relative to I_2 and IO_3^- . Moreover, iodide and iodate have not significantly different sensitivities in the ICP-MS measurements. A Thermo X7 (Thermo Fisher Scientific) was employed for the detection and quantification of stable ^{127}I [27].

Results and discussion

Results of the radionuclide analysis are summarized in Table 2. Tritium, ^{90}Sr , ^{134}Cs , and ^{137}Cs were obtained by radiometric means, ^{129}I by AMS. Please note that the seawater sample from Hawaii was only analyzed for ^{129}I , as radiometric methods were not deemed successful for the presumably very low contamination levels.

Ocean sediments

As expected, Pacific Ocean sediments from inside the exclusion zone exhibited the highest contamination levels. The correlation between radiostrontium and ^{134}Cs as well as ^{137}Cs is illustrated in Fig. 2.

Although a slight distance dependency can be observed (at least sample Sed 1 that was taken closest to the Fukushima Daiichi NPP exhibited the highest activity concentration), it was surprising to note that sample Sed 3 (taken 16.1 km north of the NPP) exhibited the lowest activities. It is also interesting that the radiocesium activity concentrations fluctuate by more than an order of magnitude within the samples, the ^{90}Sr activity concentrations, however, only by a factor of 2. The $^{90}\text{Sr}/^{137}\text{Cs}$ activity ratios (where calculable) were 0.012 ± 0.002 (Sed 1), 0.036 ± 0.005 (Sed 2), 0.15 ± 0.03 (Sed 3), 0.012 ± 0.002 (Sed 4), and 0.024 ± 0.004 (Sed 5). It is unclear why Sed 3 has a ten times higher $^{90}\text{Sr}/^{137}\text{Cs}$ ratio than the other sediments (which fluctuate within a factor of 3). This odd ratio is not so much due to a high ^{90}Sr concentration, which is in the range of the other samples, but rather due to an exceptionally low ^{137}Cs concentration. The

Table 2 Results of the radionuclide analyses on Pacific Ocean sediments and water samples in Bq·kg⁻¹ at the time of sampling


Sample no.	³ H	⁹⁰ Sr	¹²⁹ I	¹³⁴ Cs	¹³⁷ Cs
Sed 1	N/A	5.4 ± 0.8	N/A	357 ± 20	436 ± 11
Sed 2	N/A	5.6 ± 0.8	N/A	142 ± 12	156 ± 5
Sed 3	N/A	3.8 ± 0.6	N/A	19 ± 6	25.3 ± 2.4
Sed 4	N/A	2.8 ± 0.4	N/A	177 ± 2	229 ± 1
Sed 5	N/A	3.2 ± 0.5	N/A	102 ± 2	134 ± 2
OW 1	<0.1	<0.4	N/A	2.5 ± 0.7	2.8 ± 0.3
OW 2	<0.1	<0.4	N/A	<1.3	1.2 ± 0.4
OW 3	<0.1	<0.4	N/A	<2.9	<0.8
OW 4	<0.1	<1	N/A	1.8 ± 0.6	<0.4
OW 5	<0.1	<0.4	N/A	1.7 ± 0.6	1.1 ± 0.3
TW 6	<0.1	<0.4	N/A	< 0.32	1.6 ± 0.6
TW 7	<0.1	<0.4	N/A	1.52 ± 0.40	1.69 ± 0.14
TW 8	<0.1	<0.4	N/A	3.71 ± 0.35	4.27 ± 0.13
TW 9	<0.1	<0.4	N/A	1.57 ± 0.39	0.95 ± 0.25
HI W	N/A	N/A	2.37E-8 ± 1.9E-9*	N/A	N/A
CW #5	2260 ± 230	< 0.4	1.29E-6 ± 3.4E-8**	<4	<1.3

Activity concentrations for sediment samples are given per dry mass. Uncertainties of radiometric measurements are due to counting statistics

N/A not analyzed

* ¹²⁹I/¹²⁷I ratio 1.10E-10

** ¹²⁹I/¹²⁷I ratio 2.63E-8

**Fig. 2** Activity concentrations of ¹³⁴Cs, ¹³⁷Cs and ⁹⁰Sr in Fukushima Daiichi exclusion zone Pacific Ocean sediments (dry mass)

reason for this anomaly (which has been observed in similar manner in terrestrial environmental samples from north of the NPP [4]) is yet unclear. Although, from a radioecological point of view, sediment is not comparable to food, the high ⁹⁰Sr/¹³⁷Cs ratio of 0.15 challenges the governmental assumption for the regulatory limits in food,

namely a constant ⁹⁰Sr/¹³⁷Cs ratio in food of ≤0.1 [28]. Seafood organisms exposed to high-⁹⁰Sr effluents may exhibit higher ⁹⁰Sr concentrations than covered by this governmental assumption.

The ¹³⁴Cs/¹³⁷Cs activity ratios are reliable source identifiers [29]. The average ¹³⁴Cs/¹³⁷Cs ratio at the time of the accident (March 11, 2011) was found to be 0.98 [28]. Due to the shorter half-life of ¹³⁴Cs, this average ratio has been calculated to have gone down to 0.77 by the time of sampling (Dec 20, 2011). The ratios found were 0.81 ± 0.05 (Sed 1), 0.91 ± 0.08 (Sed 2), 0.75 ± 0.25 (Sed 3), 0.77 ± 0.01 (Sed 4), and 0.76 ± 0.02 (Sed 5). Sed 2 seems to exhibit a somewhat higher ¹³⁴Cs/¹³⁷Cs ratio than the other sediments, however, with the given uncertainty margins, this outlier seems not significant. It would be interesting to study the ¹³⁵Cs/¹³⁷Cs ratio [11] to possibly identify the source of these contaminations.

Other studies [30] for radiocesium in sea sediments found up to 250 Bq kg⁻¹-dry in Sendai (north of the Fukushima Daiichi NPP), which is naturally somewhat lower than the levels reported in this study, but in good agreement with our results. Our findings are also in good agreement with the activity concentrations reported by Yamamoto et al. [31]. A study on sediments from the Sea of Japan naturally found much lower radiocesium activity concentrations (0.25 Bq kg⁻¹ ¹³⁷Cs and 0.15 Bq kg⁻¹ ¹³⁴Cs in the Sado Basin) [32].

For comparison, it may be interesting to note that the activity levels found in these Ocean Sediments are roughly comparable to the radiocesium inventory in soil in Central Europe, e.g., in Austria [22], which is still significantly polluted from the Chernobyl accident and the fallout from 20th century's atmospheric explosions.

Seawater

With only small volumes available for analysis, it was not expected to detect any radionuclides. However, low activities of radiocesium were found (with high uncertainties), as summarized in Table 2. Given the high uncertainties $^{134}\text{Cs}/^{137}\text{Cs}$ ratios are less meaningful; they are 0.89 ± 0.3 (OW 1), <1.08 (OW2), >4.5 (OW4), and 1.5 ± 0.7 (OW5). In any case, our findings for seawater (up to 2.8 Bq kg^{-1} ^{137}Cs and 2.5 Bq kg^{-1} ^{134}Cs) are in good agreement with previously reported data [30] for seawater.

No tritium was detected in any of the seawater samples. The ^{129}I activity concentration in the seawater sample from Hawaii from 2012 ($2.37 \times 10^{-8} \pm 1.9 \times 10^{-9}$; $^{129}\text{I}/^{127}\text{I}$ ratio 1.10×10^{-10}) are in the range of the pre-Fukushima background reported by Stan-Sion et al. [33].

Tap water

Tap water from Tokyo exhibited contamination levels that were of no health concern (up to 4.3 Bq kg^{-1} ^{137}Cs). The activity concentrations found in this study are in good agreement with previously reported tap water results [28, 29]. The radiocesium activity concentrations were clearly below the regulatory limit for radiocesium in tap water of 200 Bq kg^{-1} . Previous studies [28] showed that ^{131}I was probably also present in Tokyo's tap water at the time of sampling; however, the measurements of this study were performed too late for the detection of this short-lived radionuclide. Strontium-90 and tritium could not be detected in tap water because of too low sample volumes and the limited detection limit of the LSC, respectively.

The $^{134}\text{Cs}/^{137}\text{Cs}$ activity ratios slightly deviated from the average ratio [28] of 0.98 at the time of the accident (0.97–0.96 at the time of sampling would have suggested): we found ratios of <0.2 (TW6), 0.90 ± 0.25 (TW7), 0.87 ± 0.09 (TW8), and 1.7 ± 0.6 (TW9). Like TW6, TW7, and TW8, also the ratios found in Tokyo tap water in a previous study [28] were consistently lower than the average; only TW9 exhibited a higher ratio, which raises the question of the source of these contaminations.

Reactor coolant

Without a doubt, the sample of reactor coolant was the most “unusual” sample of this study. The analysis of this

sample allows the investigation of the performance of the water purification techniques currently employed at the Fukushima Daiichi NPP. Purification is performed in multiple steps, including the application of charcoal and zeolite as adsorbent materials as well as reverse osmosis techniques. This purification method proves to be extremely efficient: No radiocesium and no radiostrontium could be detected. Also, the content of ^{129}I is surprisingly low; it is just two orders of magnitude higher than 2012 Pacific Ocean water from Hawaii (see Table 2). By far the dominant activity in the reactor coolant is, as expected, tritium (HTO) that cannot be removed from water by chemical means.

Conclusions

The sea sediments investigated in this study were found to be contaminated with radiocesium (up to almost 800 Bq kg^{-1}) and radiostrontium (up to 5.6 Bq kg^{-1}). This is one of the first reports on radiostrontium in sea sediments from the Fukushima exclusion zone. Seawater exhibited radiocesium contamination levels up to 5.3 Bq kg^{-1} , which is also in agreement with previous findings. Tap water from Tokyo weeks after the accident exhibited detectable but harmless activities of radiocesium (well below the regulatory limit). The investigation of the reactor coolant (finding only ^3H and even low ^{129}I) leads to the conclusion that the purification techniques for reactor coolant employed at Fukushima Daiichi are very effective.

Acknowledgments This work was supported by JSPS KAKENHI (Grant Number 25870158), from CDC NIOSH Mountain and Plains Education and Research Center (Grant Number T42OH009229-07), and the US Nuclear Regulatory Commission (NRC) (Grant Number NRC-HQ-12-G-38-0044). Its contents are solely the responsibility of the authors and do not necessarily represent the official views of the CDC NIOSH and MAP ERC.

Open Access This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

References

- Steinhauser G, Brandl A, Johnson TE (2014) Comparison of the Chernobyl and Fukushima nuclear accidents: a review of the environmental impacts. *Sci Total Environ* 470–471:800–817
- Masson O, Baeza A, Bieringer J, Brudecki K, Bucci S, Cappai M, Carvalho FP, Connan O, Cosma C, Dalheimer A, Didier D, Depuydt G, De Geer LE, De Vismes A, Gini L, Groppi F, Gudnason K, Gurriaran R, Hainz D, Halldorsson O, Hammond D,

- Hanley O, Holey K, Homoki Z, Ioannidou A, Isajenko K, Jan-kovic M, Katzlberger C, Kettunen M, Kierepko R, Kontro R, Kwakman PJM, Lecomte M, Leon Vintro L, Leppanen AP, Lind B, Lujaneni G, McGinnity P, McMahon C, Mala H, Manenti S, Manolopoulou M, Mattila A, Mairing A, Mietelski JW, Moller B, Nielsen SP, Nikolic J, Overwater RMW, Palsson SE, Papastefanou C, Penev I, Pham MK, Povinec PP, Rameback H, Reis MC, Ringer W, Rodriguez A, Rulik P, Saey PRJ, Samsonov V, Schlosser C, Sgorbati G, Silobritiene BV, Soderstrom C, Sogni R, Solier L, Sonck M, Steinhäuser G, Steinkopff T, Steinmann P, Stoulos S, Sykora I, Todorovic D, Tooloutalaie N, Tositti L, Tschiersch J, Ugron A, Vagenä E, Vargas A, Wershofen H, Zhukova O (2011) Tracking of airborne radionuclides from the damaged Fukushima Dai-ichi nuclear reactors by European networks. *Environ Sci Technol* 45:7670–7677
3. Thakur P, Ballard S, Nelson R (2013) An overview of Fukushima radionuclides measured in the northern hemisphere. *Sci Total Environ* 458–460:577–613
 4. Steinhäuser G, Schauer V, Shozugawa K (2013) Concentration of ^{90}S at selected hot spots in Japan. *PLoS ONE* 8:e57760
 5. Casacuberta N, Masque P, Garcia-Orellana J, Garcia-Tenorio R, Buesseler KO (2013) ^{90}Sr and ^{89}Sr in seawater off Japan as a consequence of the Fukushima Dai-ichi nuclear accident. *Biogeosciences* 10:3649–3659
 6. Zheng J, Tagami K, Watanabe Y, Uchida S, Aono T, Ishii N, Yoshida S, Kubota Y, Fuma S, Ihara S (2012) Isotopic evidence of plutonium release into the environment from the Fukushima DNPP accident. *Sci Rep* 2:304
 7. Zheng J, Tagami K, Uchida S (2013) Release of plutonium isotopes into the environment from the Fukushima Daiichi nuclear power plant accident: what is known and what needs to be known. *Environ Sci Technol* 47:9584–9595
 8. Schneider S, Walther C, Bister S, Schauer V, Christl M, Sinal H-A, Shozugawa K, Steinhäuser G (2013) Plutonium release from Fukushima Daiichi fosters the need for more detailed investigations. *Sci Rep* 3:2988
 9. Kakiuchi H, Akata N, Hasegawa H, Ueda S, Tokonami S, Yamada M, Hosoda M, Sorimachi A, Tazoe H, Noda K, Hisamatsu S (2012) Concentration of ^3H in plants around Fukushima Daiichi nuclear power station. *Sci Rep* 2:947
 10. Matsumoto T, Maruoka T, Shimoda G, Obata H, Kagi H, Suzuki K, Yamamoto K, Mitsuguchi T, Hagino K, Tomioka N, Sambandam C, Brummer D, Klaus PM, Aggarwal P (2013) Tritium in Japanese precipitation following the March 2011 Fukushima Daiichi nuclear plant accident. *Sci Total Environ* 445–446:365–370
 11. Zheng J, Tagami K, Bu W, Uchida S, Watanabe Y, Kubota Y, Fuma S, Ihara S (2014) Isotopic ratio of $^{135}\text{Cs}/^{137}\text{Cs}$ as a new tracer of radiocesium released from the Fukushima nuclear accident. *Environ Sci Technol* 48:5433–5438
 12. Priyadarshi A, Dominguez G, Thiemens MH (2011) Evidence of neutron leakage at the Fukushima nuclear plant from measurements of radioactive ^{35}S in California. *Proc Natl Acad Sci USA* 108:14422–14425
 13. Steinhäuser G (2014) Fukushima's forgotten radionuclides: a review of the understudied radioactive emissions. *Environ Sci Technol* 48:4649–4663
 14. Morino Y, Ohara T, Nishisawa M (2011) Atmospheric behavior, deposition, and budget of radioactive materials from the Fukushima Daiichi nuclear power plant in March 2011. *Geophys Res Lett* 38:[L00G11/1-7]
 15. Kawamura H, Kobayashi T, Furuno A, In T, Ishikawa Y, Nakayama T, Shima S, Awaji T (2011) Preliminary numerical experiments on oceanic dispersion of ^{131}I and ^{137}Cs discharged into the ocean because of the Fukushima Daiichi nuclear power plant disaster. *J Nucl Sci Technol* 48:1349–1356
 16. Buesseler K, Aoyama M, Fukasawa M (2011) Impacts of the Fukushima nuclear power plants on marine radioactivity. *Environ Sci Technol* 45:9931–9935
 17. Buesseler KO, Jayne SR, Fisher NS, Rypina II, Baumann H, Baumann Z, Breier CF, Douglass EM, George J, MacDonald AM, Miyamoto H, Nishikawa J, Pike SM, Yoshida S (2012) Fukushima-derived radionuclides in the ocean and biota off Japan. *Proc Natl Acad Sci USA* 109:5984–5988
 18. Tumey SJ, Guilderson TP, Brown TA, Broek T, Buesseler KO (2013) Input of ^{129}I into the western Pacific Ocean resulting from the Fukushima nuclear event. *J Radioanal Nucl Chem* 296:957–962
 19. Povinec PP, Aoyama M, Biddulph D, Breier R, Buesseler K, Chang CC, Golser R, Hou XL, Jeřkovský M, Jull AJT, Kaizer J, Nakano M, Nies H, Palcsu L, Papp L, Pham MK, Steier P, Zhang LY (2013) Cesium, iodine and tritium in NW Pacific waters—a comparison of the Fukushima impact with global fallout. *Biogeosciences* 10:5481–5496
 20. Michel R, Handl J, Ernst T, Botsch W, Szidat S, Schmidt A, Jakob D, Beltz D, Romantschuk LD, Sinal HA, Schnabel C, Lopez-Gutierrez JM (2005) ^{129}I in soils from Northern Ukraine and the retrospective dosimetry of the ^{131}I exposure after the Chernobyl accident. *Sci Total Environ* 340:35–55
 21. Steinhäuser G, Merz S, Küber-Heiss A, Katzlberger C (2012) Using animal thyroids as ultra-sensitive biomonitors for environmental radioiodine. *Environ Sci Technol* 46:12890–12894
 22. Steinhäuser G, Merz S, Hainz D, Sterba JH (2013) Artificial radioactivity in environmental media (air, rainwater, soil, vegetation) in Austria after the Fukushima nuclear accident. *Environ Sci Pollut Res* 20:2527–2537
 23. Atkinson R, Eddy T, Kuhne W, Jannik T, Brandl A (2014) Measurement of the tritium concentration in the fractionated distillate from environmental water samples. *J Environ Radioact* 135:113–119
 24. Steinhäuser G, Evers J, Jakob S, Klapötke TM, Oehlinger G (2008) A review on fulminating gold (Knallgold). *Gold Bull* 41:305–317
 25. Klapötke TM, Krumm B, Steemann FX, Steinhäuser G (2010) Hands on explosives: safety testing of protective measures. *Saf Sci* 48:28–34
 26. Steinhäuser G, Klapötke TM (2010) Using the chemistry of fireworks to engage students in learning basic chemical principles: a lesson in eco-friendly pyrotechnics. *J Chem Educ* 87:150–156
 27. Michel R, Daraoui A, Gorny M, Jakob D, Sachse R, Tosch L, Nies H, Goroncy I, Herrmann J, Sinal H-A, Stocker M, Alfimov V (2012) ^{129}I and ^{127}I in European seawaters and in precipitation from Northern Germany. *Sci Total Environ* 419:151–169
 28. Merz S, Shozugawa K, Steinhäuser G (2015) Analysis of Japanese radionuclide monitoring data of food before and after the Fukushima nuclear accident. *Environ Sci Technol* 49:2875–2885
 29. Merz S, Steinhäuser G, Hamada N (2013) Anthropogenic radionuclides in Japanese food: environmental and legal implications. *Environ Sci Technol* 47:1248–1256
 30. Kofuji H (2015) In situ measurement of ^{134}Cs and ^{137}Cs in seabed using underwater gamma-spectrometry systems: application in surveys following the Fukushima Dai-ichi Nuclear Power Plant accident. *J Radioanal Nucl Chem* 303:1575–1579
 31. Yamamoto K, Tagami K, Uchida S, Ishii N (2015) Model estimation of ^{137}Cs concentration change with time in seawater and sediment around the Fukushima Daiichi Nuclear Power Plant site considering fast and slow reactions in the seawater-sediment systems. *J Radioanal Nucl Chem* 304:867–881
 32. Inoue M, Yoneoka S, Ochiai S, Oikawa S, Fujimoto K, Yagi Y, Honda N, Nagao S, Yamamoto M, Hamajima Y, Murakami T, Kofuji H, Misonoo J (2015) Lateral and temporal variations in

- Fukushima Dai-ichi NPP-derived ^{134}Cs and ^{137}Cs in marine sediments in/around the Sado Basin, Sea of Japan. *J Radioanal Nucl Chem* 303:1313–1316
33. Stan-Sion C, Enachescu M, Petre AR (2015) AMS analyses of ^{129}I from the Fukushima Daiichi nuclear accident in the pacific ocean water of the coast of La Jolla—San Diego, USA. *Environ Sci Proc Impacts* 17:932–938