



WORKSHOP PRESENTATION

Open Access

Quantitative assessment of myocardial motion from velocity encoded MRI

Anja Lutz¹, Jan Paul¹, Patrick Etyngier³, Axel Bornstedt¹, Peter Bernhardt¹, Gerd Ulrich Nienhaus², Wolfgang Rottbauer¹, Volker Rasche^{1*}

From 15th Annual SCMR Scientific Sessions
Orlando, FL, USA. 2-5 February 2012

Summary

The objective of this study is to investigate the potential role of different automatically derived quantitative parameters derived from velocity encoded MRI for the identification of asynchronous patients.

Background

About 30% of patients treated with cardiac resynchronization therapy (CRT) do not benefit from the procedure. Quantitative analysis of motion encoded MRI data may provide helpful parameters for the identification of CRT patients and prediction of the therapy outcome.

Methods

11 Volunteers (30 ± 8 years) and 3 patients (41 ± 11 years) were investigated at a 3T whole body MR scanner (Achieva, Philips) with a 32 channel cardiac coil. The patients suffered from DCM, asynchrony and/or LBBB. A velocity encoded (TPM) navigated segmented gradient echo sequence was applied in the apical, equatorial and basal slice. The acquisition parameters were: FOV= 340^2 mm 2 , in-plane resolution = 2.5^2 mm 2 , slice thickness=8mm, acquisition matrix MxP=172x168, TR/TE=6.3ms/4.6ms, $\alpha=15^\circ$, 3 k-lines per segment, VENC=30cm/s, nominal scan duration =5:51 minutes, black blood imaging with alternating presaturation pulses [1] and a SENSE acceleration factor of 2. For 60 bpm 32 cardiac phases were measured with a phase interval of 29.1ms.

From the TPM data, the longitudinal and radial standard deviation of time to peak systolic and diastolic velocities SD(TTP_{l,sys}), SD(TTP_{l,dias}), SD(TTP_{r,sys}), SD(TTP_{r,dias}) over 6 segments [2], the radial, circumferential

and longitudinal asynchrony correlation coefficient (ACC)[3], the longitudinal and radial velocity range $\Delta v_l = v_{l,max} - v_{l,min}$, $\Delta v_r = v_{r,max} - v_{r,min}$ and the temporal uniformity of velocity (TUV) in radial, longitudinal and circumferential direction were derived. The latter one was defined in analogy to the temporal uniformity of strain [4,5].

Results

In all patients, a substantially deterioration of the motion curve from the healthy volunteers was observed (Figure 1). Most obvious, a clear reduction of the peak velocities can be appreciated.

The derived quantitative motion parameters are listed in table 1. All mean values of the derived parameters show differences between the patient and the volunteer group. Δv_l and Δv_r show large differences for all investigated motion directions, SD(TTP) appears increased, the mean ACC and the radial and longitudinal TUV reduced. Significances of the differences could not be calculated due to the small number of patients.

Conclusions

Several quantitative motion parameters show substantial differences between patients and volunteers and may be applied for automatic identification left ventricular asynchrony. Whether the investigated parameters can be applied for CRT patient selection and outcome prediction must be proven in a larger clinical study.

Funding

AL and VR have a research agreement with Philips Medical. PE is employed by Philips Healthcare.

¹Department of Internal Medicine II, University Hospital of Ulm, Ulm, Germany

Full list of author information is available at the end of the article




Figure 1 Apical, equatorial and basal longitudinal motion exemplarily for one volunteer.

Table 1 Velocity based motion parameters evaluated for all volunteers (mean \pm standard deviation) as well for the 3 investigated patients.

parameter	volunteers		patients	
	mean	σ	mean	σ
SD(TTP _{I,sys} [ms])	36.84	22.53	48.94	27.77
SD(TTP _{I,dias} [ms])	15.72	4.64	45.68	22.23
SD(TTP _{r,sys} [ms])	46.40	12.48	61.78	17.07
SD(TTP _{r,dias} [ms])	33.85	8.75	49.75	21.39
mean ACC _I	0.93	0.03	0.70	0.09
mean ACC _r	0.79	0.04	0.52	0.07
mean ACC _c	0.72	0.07	0.53	0.16
$\Delta v_{l,apical}$ [cm/s]	10.77	2.19	5.12	1.37
$\Delta v_{l,equatorial}$ [cm/s]	16.13	2.54	7.14	3.27
$\Delta v_{l,basal}$ [cm/s]	20.71	2.50	7.26	3.11
$\Delta v_{r,apical}$ [cm/s]	8.31	1.05	4.51	1.21
$\Delta v_{r,equatorial}$ [cm/s]	7.97	0.97	4.26	0.40
$\Delta v_{r,basal}$ [cm/s]	7.51	1.23	5.15	1.04
TUV _I	0.87	0.02	0.78	0.04
TUV _r	0.80	0.02	0.70	0.02
TUV _c	0.77	0.03	0.69	0.06

Author details

¹Department of Internal Medicine II, University Hospital of Ulm, Ulm, Germany. ²Institute of Applied Physics and Center for Functional Nanostructures, Karlsruhe Institute of Technology, Karlsruhe, Germany.

³Medisys Research Lab, Philips Healthcare, Suresnes, France.

Published: 1 February 2012

References

1. Lutz , et al: MAGMA. 2011, 24(3):127-35.
2. Föll , et al: J Magn Reson Imaging. 2011.
3. Schneider , et al: J Comput Assist Tomogr. 2001, 25:550-557.
4. Bilchick , et al: JACC Cardiovasc Imaging. 2008, 1:561-568.
5. Leclercq , et al: Circulation. 2002, 106:1760-1763.

doi:10.1186/1532-429X-14-S1-W39

Cite this article as: Lutz et al: Quantitative assessment of myocardial motion from velocity encoded MRI. *Journal of Cardiovascular Magnetic Resonance* 2012 14(Suppl 1):W39.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

