

Research

Open Access

The proteome of neural stem cells from adult rat hippocampus

Martin H Maurer*^{†1}, Robert E Feldmann Jr^{†1}, Carsten D Fütterer² and Wolfgang Kuschinsky¹

Address: ¹Dept. of Physiology and Pathophysiology, University of Heidelberg, Im Neuenheimer Feld 326, 69120 Heidelberg, Germany and ²Dept. of Anesthesiology and Critical Care Medicine, University of Heidelberg, Faculty of Clinical Medicine Mannheim, Theodor-Kutzer-Ufer, 68167 Mannheim, Germany

Email: Martin H Maurer* - maurer@uni-hd.de; Robert E Feldmann - robert_feldmann@gmx.li; Carsten D Fütterer - c.fuetterer@pio1.uni-heidelberg.de; Wolfgang Kuschinsky - wolfgang.kuschinsky@pio1.uni-heidelberg.de

* Corresponding author †Equal contributors

Published: 12 June 2003

Received: 7 May 2003

Proteome Science 2003, 1:4

Accepted: 12 June 2003

This article is available from: <http://www.Proteomesci.com/content/1/1/4>

© 2003 Maurer et al; licensee BioMed Central Ltd. This is an Open Access article: verbatim copying and redistribution of this article are permitted in all media for any purpose, provided this notice is preserved along with the article's original URL.

Abstract

Background: Hippocampal neural stem cells (HNSC) play an important role in cerebral plasticity in the adult brain and may contribute to tissue repair in neurological disease. To describe their biological potential with regard to plasticity, proliferation, or differentiation, it is important to know the cellular composition of their proteins, subsumed by the term proteome.

Results: Here, we present for the first time a proteomic database for HNSC isolated from the brains of adult rats and cultured for 10 weeks. Cytosolic proteins were extracted and subjected to two-dimensional gel electrophoresis followed by protein identification through mass spectrometry, database search, and gel matching. We could map about 1141 ± 209 ($N = 5$) protein spots for each gel, of which 266 could be identified. We could group the identified proteins into several functional categories including metabolism, protein folding, energy metabolism and cellular respiration, as well as cytoskeleton, Ca^{2+} signaling pathways, cell cycle regulation, proteasome and protein degradation. We also found proteins belonging to detoxification, neurotransmitter metabolism, intracellular signaling pathways, and regulation of DNA transcription and RNA processing.

Conclusions: The HNSC proteome database is a useful inventory which will allow to specify changes in the cellular protein expression pattern due to specific activated or suppressed pathways during differentiation or proliferation of neural stem cells. Several proteins could be identified in the HNSC proteome which are related to differentiation and plasticity, indicating activated functional pathways. Moreover, we found a protein for which no expression has been described in brain cells before.

Background

Stem cells are cells found in nearly all tissues [1], although generally in small numbers. They are defined by several unique properties [2,3]: Stem cells are unspecialized cells, they are capable of dividing and renewing themselves for long periods of time, and they can give rise to many types of specialized cells, such as blood, nerve, and muscle cells.

Whereas embryonic stem cells, which are derived from very early embryos, are totipotent – that is, they are capable of generating all types of cells in the body during normal development – adult stem cells have lost this potential. When adult stem cells differentiate, they seem to be restricted to produce cells from the tissue they

originate, though there are recent publications suggesting a transdifferentiation potential [4–7].

Adult neural stem cells have been isolated from various regions of the adult mammalian brain, where the highest densities of neural stem cells have been found in the hippocampus, the subventricular zone, and the olfactory bulb [8,9]. It seems that adult neural stem cells have the ability to develop into functional mature neurons [10]. These regions are of special interest as they reveal spontaneous neurogenesis throughout the entire lifetime, suggesting to play a functional role in physiological cell replacement in aging, learning and cognition, as well as proposing a therapeutic potential in neurological disease [5,11,12], including neurodegenerative disorders like Alzheimer's and Parkinson's disease, cerebrovascular insults such as stroke, or developmental impairments.

Although proteomic technology has been widely used with regard to different aspects of multiple neurological diseases [13,14], neural stem cells isolated from the brains of adult mammals have not been subjected to profound proteome analysis. On the other hand, genome-wide approaches have been published recently which were analysing neural stem cells using DNA microarrays or differential display methods [15–18].

To elucidate the functional role of protein interaction in HNSC with regard to plasticity, proliferation, or differentiation, proteomics, based on high-resolution two-dimensional gel electrophoresis (2-DE) combined with mass spectrometry, is a useful tool. Prerequisite of any functional HNSC experiment is the knowledge of the cellular proteome which then allows assessment of differential protein expression. In the present study, we propose a reference database and reference map for neural stem cells from adult rat hippocampus.

Results

Protein expression standard pattern

The two-dimensional (2-D) standard pattern of HNSC isolated from adult rat brain is shown in Fig. 1 [see also additional file 2] as revealed by scanning, digitizing, densitometry, and image analysis. Each spot detected by the 2-D software was assigned a unique number to identify spots in a gel matching process. Spots of 5 different experiments were compared to a reference map. Technical replicates of gels from the same experiments showed a high reproducibility when run under the same electrophoretic conditions. In the 2-DE gels analysed, we could map an average of 1141 ± 209 ($N = 5$) protein spots representing the cellular proteome of HNSC, ranging from 967 to 2066 assigned protein spots. Thus we could create a proteome database of 2472 unique spots being present in at least 1 gel.

Protein identification

Of the 2472 unique spots in the HNSC database, we could identify 266 protein spots using MS, database search and subsequent gel matching, representing 109 individual proteins.

Table 1 [see additional file 1] lists the identified proteins with their respective SwissProt database accession number, which can be assessed via the internet URL <http://www.expasy.ch/sprot>, the NCBI GenBank accession number, which can be assessed via the internet URL <http://www.ncbi.nlm.nih.gov/Entrez>, and the theoretical and experimental pI and molecular weight. Additionally, we added the Mascot score results quantifying the identification probability of the fragment match [19]. A definition and interpretation of the meaning of the Mascot score can be found at the internet URL <http://www.matrix-science.com>. We have only included statistically significant Mascot score results ($P < 0.05$) in the table, protein spots with statistically not significant Mascot scores remained unidentified. For inclusion into the HNSC database, we then added only the protein search result with the highest Mascot score.

Classification of spot families

We grouped the identified proteins into several functional categories (Fig. 2). Most proteins belonged to metabolism pathways, which include glycolysis, tricarboxylic acid cycle, amino acid metabolism, and protein synthesis. The second largest group comprised proteins involved in protein folding. Most proteins in this group were represented chaperones, chaperonins, heat shock proteins, and foldases. A third major group comprehended energy metabolism and cellular respiration, necessary for the cell's ATP turnover. The cytoskeletal group comprehended not only proteins constitutive for the cellular cytoskeleton like actin or tubulin, but also proteins modifying and rearranging the cytoskeleton.

Several other proteins belonged to Ca^{2+} signaling pathways, cell cycle regulation, proteasome and protein degradation, detoxification, neurotransmitter metabolism with emphasis of glutamate and glycine metabolism, intracellular signaling mechanisms like protein kinase pathways, and regulation of DNA transcription and RNA processing.

Discussion

The aim of the present study was to establish a proteomic database for neural stem cells. We isolated neural stem cells from hippocampi of adult rat brains, as the hippocampal formation is a neurogenic region in the mammalian brain throughout the whole life [20–23], contributing to learning and memory processes.




Figure 1


Two-dimensional gel of adult rat hippocampal stem cell lysate. (A) Representative original gel image. (B) Scout to find annotated spots in magnified gel sectors (see supplemental material).

To our knowledge, nothing is presently known about the cellular proteome of HNSC. With the exception of a preliminary study on human fetal cortical stem cells [24], where only two proteins, i. e. nestin and microtubule-associated protein MAP-2c, have been identified and described, there is no proteomic reference material for adult neural stem cells. Recently, data became available for a 2-DE study in mouse embryonic stem cells differentiated *in vitro* to neural phenotypes, where 24 protein spots could be identified [25]. In the present study, we describe a comprehensive picture of the HNSC proteome, and, looking into detail, we could identify nearly all of the most abundantly expressed proteins in the neural precursor cells and include these in the database.

Since no study has been performed investigating the protein expression pattern of neural stem cells, we matched our results with murine brain proteome databases. Several studies investigated murine brain proteomes [26,27], and

there is a certain overlap of proteins found both in neural stem cells and mature brain cells. Gauss et al. [26] set up a standard spot pattern for mouse brain consisting of 8767 spots, where they could identify 166 (2%) spots representing 90 different proteins. Tsugita et al. [27] mapped about 1188 spots in the mouse brain, of which 122 could be identified (10%), representing 61 different proteins. Compared to these studies, we could identify 266 out of 2472 protein spots (11%).

Moreover, we could identify additional proteins, which are characteristic to the precursor cells. These include the Neuronal Differentiation-Related Protein (NDRP) [28]. This protein is predominantly expressed in developing and regenerating sensory neurons and has been found in the embryonic retina and olfactory epithelium, as well as in perinatal dorsal root ganglia. Of note, NDRP is inducible in motor neurons after axotomy [28]. It contains 6 WD (tryptophan-aspartate) repeats, which can be

**Figure 2**

Distribution of the identified protein spots into functional groups. This figure shows the percentual distribution of the identified proteins in functional groups. The largest group, which comprehends about a third of the proteins, is constituted by cellular metabolism. But also cytoskeletal rearrangement and protein folding were major contributors.

found in proteins involved in signal transduction, RNA processing, gene regulation, vesicular traffic and regulation of cytoskeletal assembly and/or cell cycle [29,30]. With regard to our data, this protein might be involved also in the generation of hippocampal neurons in neural progenitors during the process of differentiation.

We have identified another differentiation-related protein, the Phosphatidyl Ethanolamine-Binding Protein (PEBP) which is highly expressed in the brain but also a major component of epididymal secretions and sperm plasma membranes [31]. PEBP binds ATP, opioids and phosphatidylethanolamine and -with lower affinity- phosphatidylinositol and phosphatidylcholine. Similarity search has revealed that PEBP might be a serine protease inhibitor which inhibits thrombin, neuropsin and chymotrypsin but not trypsin, tissue type plasminogen activator and elastase. This protein was described as hippocampal precursor peptide stimulating cholinergic phenotypes in the developing brain in cooperation with nerve growth factor NGF, where it increases the produc-

tion of choline acetyltransferase but not acetylcholinesterase [32,33]. This effect seems to be mediated by a specific receptor [34]. Together with the finding of the present study, that PEBP is also expressed in HNSC, is concordant with the finding that hippocampal neural stem and progenitor cells have the ability to produce cholinergic neurons [35–37]. PEBP was associated with memory, learning disorders, dementia [38].

Additionally, we have found the expression of CAP1, or synonymously Sperm Protein SP22, in HNSC. This protein has not been described to be expressed in the brain, or nervous tissue before. This putative intracellular protease/amidase has been found originally in sperm, where it is thought to play a role in fertilization and development [39]. Speculating in neural progenitors, this protein could also contribute to cell differentiation and development.

In summary, this database includes a wide variety of the most abundantly expressed proteins of rat hippocampal

neural stem cells. Therefore this database will serve as reference database for future stem cell proteomic experiments.

Conclusions

In the present study, we describe a proteome database of neural stem cells isolated from adult rat hippocampus. This database provides a protein inventory which will allow to specify changes in the protein expression pattern due to specific activated or suppressed pathways during differentiation or proliferation of the stem cells. Therefore, the present database is a useful tool for further stem cell research.

Moreover, we have identified several proteins which are associated with pathways involved in differentiation and plasticity, indicating a functional role for these proteins in HNSC. Additionally, the expression of the fertility protein CAP1 has not been described in brain cells before, indicating a role in cellular maturation.

Methods

Cell culture

Protocols are concordant with the policy on the use of laboratory animals, as endorsed by the National Research Council of the U.S.A., and fulfill the requirements of German law. Six male Wistar rats (4–6 weeks old) were anesthetized by short isoflurane inhalation and sacrificed by decapitation. Primary culture of rat neurospheres were obtained as described [20,40,41]. Brains were removed quickly and washed in 50 mL ice-cold Dulbecco's Phosphate Buffered Saline (DPBS) supplemented with 4.5 g/L glucose (DPBS/Glc). The dissected hippocampi were washed in 10 mL DPBS/Glc and centrifuged for 5 min at $1600 \times g$ at 4°C . After removal of the supernatant, the tissue was homogenized with scissors and scalpels. The pieces were washed with DPBS/Glc for 5 min at $800 \times g$, and the pellet was resuspended in 0.01% (w/v) papain, 0.1% (w/v) dispase II (neutral protease), 0.01% (w/v) DNase I, and 12.4 mM manganese sulfate in Hank's Balanced Salt Solution. The tissue was triturated with plastic pipet tips and incubated for 40 min at room temperature, but every 10 min the solution was mixed well. Following centrifugation at $800 \times g$ for 5 min at 4°C , pellets were washed three times in 10 mL Dulbecco's Modified Eagle's Medium-Ham's F-12 medium supplemented with 2 mM L-glutamine, 100 units/mL penicillin and 100 units/mL streptomycin. They were resuspended in 1 mL Neurobasal medium supplemented with B27 (Invitrogen, Carlsbad, CA, USA), 2 mM L-glutamine, 100 units/mL penicillin and 100 units/mL streptomycin, 20 ng/mL Epidermal Growth Factor, 20 ng/mL Fibroblast Growth Factor-2, and 2 $\mu\text{g}/\text{mL}$ heparin. Cells were plated under sterile conditions in 6-well dishes in a concentration of 25,000–100,000 cells/mL. Dishes were incubated at 37°C in 5%

CO_2 . Cell culture medium was changed once a week, where about two thirds of the volume were replaced.

Sample preparation

For obtaining high-yield protein extracts, we developed a specific protocol: After 5 passages of 14 days each, cells were removed from the medium, washed 3 times in 300 mosmol/L Tris-HCl sucrose, pH 7.4, and dissolved in a detergent lysis buffer containing 7 M urea, 2 M thiourea, 4% (w/v) CHAPS, 0.5% (v/v) Triton X-100, 0.5% (v/v) IPG buffer pH 3–10 (Amersham Biosciences, Uppsala, Sweden), 100 mM DTT and 1.5 mg/mL Complete protease inhibitor (Roche, Mannheim, Germany) for 1 hour at 18°C in an orbital shaker. The lysate was then centrifuged at $21,000 \times g$ for 30 min. Protein content of the supernatant was measured by the Bradford assay [42,43].

Two-dimensional gel electrophoresis

Protein extracts were separated by two-dimensional SDS-PAGE essentially as described (.). [44–46]. Briefly, for first dimensional isoelectric focussing, 500 μg of the cell extract were run in 6 M urea, 2 M thiourea, 1 M DTT, 2% (w/v) CHAPS, 0.5% (v/v) IPG buffer, on 18 cm immobilized non-linear pH 3–10 gradient IPG strips (Immobiline DryStrip pH 3–10 NL, Amersham Biosciences, Uppsala, Sweden), using the IPGphor apparatus (Amersham Biosciences, Uppsala, Sweden). After 12 hours of reswelling time at 30 V, voltages of 200 V, 500 V, and 1,000 V were applied for 1 hour each. Then voltage was increased to 8,000 V within 30 min and kept constant at 8,000 V for 12 hours, resulting in a total of 100,300 Vh. For the second dimension, electrophoretic mass separation was achieved using $20 \times 18 \times 0.2 \text{ cm}^3$, 12.5% polyacrylamide gels. Protein spots in five replicates of the protein extracts were visualized by ultra-sensitive silver staining [47] and detected by the Phoretix 2D Advanced v6.01c software (Nonlinear Dynamics, Newcastle-upon-Tyne, UK).

Mass spectrometry and protein identification

Protein spots were identified by mass spectrometry. Mass spectrometry, spectral data acquisition, and database queries were performed by Proteosys AG (Mainz, Germany). Briefly, protein spots were excised with an automated spot-picker and destained as described [48]. In-gel digestion with trypsin (Promega, Madison, WI, USA) was employed using a modified protocol as described [49]. For MALDI-TOF, 1 μL of the peptide mixture was spotted onto a metal MALDI-TOF target (SCOUT-MTP 384, Bruker-Daltonics, Bremen, Germany) and allowed to air-dry. Thereafter, 1 μL of the α -cyano-4-hydroxy-cinnamic acid matrix solution consisting of 10 mg/mL in 50% acetonitrile and 0.1% trifluoroacetic acid was applied to the dried sample and again allowed to air-dry. Peptide mass spectra were obtained using a Bruker Autoflex MALDI-TOF (Bruker-Daltonics, Bremen, Germany) in the reflect-

tron operation mode. Resulting fragment masses were the basis of mining the NCBI, Swiss-Prot and TrEMBL databases for protein identification via Mascot query <http://www.matrixscience.com>. Additional spots were identified by gel matching with published references for neural tissue [13,26,27,50,51].

List of abbreviations

2-D, two-dimensional;

2-DE, two-dimensional electrophoresis;

CHAPS, 3-[(3-cholamidopropyl)dimethylammonio]-1-propanesulfonate;

DTT, dithiothreitol;

HCNP, hippocampal cholinergic neurostimulating peptide;

HNSC, hippocampal neural stem cell;

IEF, isoelectric focussing;

MALDI-TOF, matrix-assisted laser desorption/ionization-time of flight mass spectrometry;

MS, mass spectrometry;

NDRP, Neuronal differentiation-related protein;

PEBP, phosphatidylethanolamine-binding protein;

SDS-PAGE, sodium dodecylsulfate polyacrylamide gel electrophoresis;

Competing interests

The authors declare no competing financial interests in this study.

Authors' contributions

MHM established 2-DE and neural stem cell culture techniques and isolated neurospheres for cell culture. REF performed cell culture, developed protein extraction and supervised 2-DE and mass spectrometry. MHM and REF improved and performed image analysis and created the database. CDF supported the design of the study and helped with 2-DE and gel matching. WK conceived the study and participated in the design and coordination of the experiments. All authors have read and approved the final manuscript.

Additional material

Additional File 2

Fig. 1-01 to Fig. 1-12 – HNSC reference map. PDF file containing the HNSC proteome map according to the scout's sectors (Fig. 1B). The spot numbers correlate with table 1 to allow easy allocation of the spots.

Click here for file

[<http://www.biomedcentral.com/content/supplementary/1477-5956-1-4-S2.pdf>]

Additional File 1

Table 1 – HNSC database. Microsoft Excel spreadsheet (XLS format) containing the proteome database for neural stem cells isolated from adult rat hippocampus.

Click here for file

[<http://www.biomedcentral.com/content/supplementary/1477-5956-1-4-S1.xls>]

Acknowledgements

We are indebted to Mrs I Keller, Mrs M Harlacher and Mrs T Lorenz for technical assistance. The authors thank C Hunzinger for mass spectrometry. This study was supported by the German Ministry for Education and Research (BMBF, Kompetenznetz Schlaganfall, B2).

References

- National Institutes of Health: **Stem cells: a primer**. 2000 [<http://www.nih.gov/news/stemcell/primer.htm>].
- Hall PA and Watt FM: **Stem cells: the generation and maintenance of cellular diversity**. *Development* 1989, **106**:619-633.
- Weissman IL: **Stem cells: units of development, units of regeneration, and units in evolution**. *Cell* 2000, **100**:157-168.
- Price J and Williams BP: **Neural stem cells**. *Curr Opin Neurobiol* 2001, **11**:564-567.
- Tsai RY, Kittappa R and McKay RD: **Plasticity, niches, and the use of stem cells**. *Dev Cell* 2002, **2**:707-712.
- Vescovi A, Gritti A, Cossu G and Galli R: **Neural stem cells: plasticity and their transdifferentiation potential**. *Cells Tissues Organs* 2002, **171**:64-76.
- Wulf GG, Jackson KA and Goodell MA: **Somatic stem cell plasticity: current evidence and emerging concepts**. *Exp Hematol* 2001, **29**:1361-1370.
- Gage FH: **Mammalian neural stem cells**. *Science* 2000, **287**:1433-1438.
- Okano H: **Stem cell biology of the central nervous system**. *J Neurosci Res* 2002, **69**:698-707.
- van Praag H, Schinder AF, Christie BR, Toni N, Palmer TD and Gage FH: **Functional neurogenesis in the adult hippocampus**. *Nature* 2002, **415**:1030-1034.
- Brüstle O and McKay RD: **Neuronal progenitors as tools for cell replacement in the nervous system**. *Curr Opin Neurobiol* 1996, **6**:688-695.
- Abe K: **Therapeutic potential of neurotrophic factors and neural stem cells against ischemic brain injury**. *J Cereb Blood Flow Metab* 2000, **20**:1393-1408.
- Rohlf C: **Proteomics in molecular medicine: applications in central nervous systems disorders**. *Electrophoresis* 2000, **21**:1227-1234.
- Morrison RS, Kinoshita Y, Johnson MD, Uo T, Ho JT, McBee JK, Conrads TP and Veenstra TD: **Proteomic analysis in the neurosciences**. *Mol Cell Proteomics* 2002, **1**:553-560.
- Ivanova NB, Dimos JT, Schaniel C, Hackney JA, Moore KA and Lemischka IR: **A stem cell molecular signature**. *Science* 2002, **298**:601-604.
- Pazman C, Castelli JC, Wen X and Somogyi R: **Large-scale identification of differentially expressed genes during neurogenesis**. *Neuroreport* 2000, **11**:719-724.

17. Ramalho-Santos M, Yoon S, Matsuzaki Y, Mulligan RC and Melton DA: **"Stemness": transcriptional profiling of embryonic and adult stem cells.** *Science* 2002, **298**:597-600.
18. Burns CE and Zon LI: **Portrait of a stem cell.** *Dev Cell* 2002, **3**:612-613.
19. Perkins DN, Pappin DJ, Creasy DM and Cottrell JS: **Probability-based protein identification by searching sequence databases using mass spectrometry data.** *Electrophoresis* 1999, **20**:3551-3567.
20. Gage FH, Coates PW, Palmer TD, Kuhn HG, Fisher LJ, Suhonen JO, Peterson DA, Suhr ST and Ray J: **Survival and differentiation of adult neuronal progenitor cells transplanted to the adult brain.** *Proc Natl Acad Sci USA* 1995, **92**:11879-11883.
21. Johansson CB, Svensson M, Wallstedt L, Janson AM and Frisen J: **Neural stem cells in the adult human brain.** *Exp Cell Res* 1999, **253**:733-736.
22. Eriksson PS, Perfilieva E, Bjork-Eriksson T, Alborn AM, Nordborg C, Peterson DA and Gage FH: **Neurogenesis in the adult human hippocampus.** *Nat Med* 1998, **4**:1313-1317.
23. Antel JP, Nalbantoglu J and Olivier A: **Neuronal progenitors-learning from the hippocampus.** *Nat Med* 2000, **6**:249-250.
24. Pearce A and Svendsen CN: **Characterisation of stem cell expression using two-dimensional electrophoresis.** *Electrophoresis* 1999, **20**:969-970.
25. Guo X, Ying W, Wan J, Hu Z, Qian X, Zhang H and He F: **Proteomic characterization of early-stage differentiation of mouse embryonic stem cells into neural cells induced by all-trans retinoic acid in vitro.** *Electrophoresis* 2001, **22**:3067-3075.
26. Gauss C, Kalkum M, Löwe M, Lehrach H and Klose J: **Analysis of the mouse proteome. (I) Brain proteins: separation by two-dimensional electrophoresis and identification by mass spectrometry and genetic variation.** *Electrophoresis* 1999, **20**:575-600.
27. Tsugita A, Kawakami T, Uchida T, Sakai T, Kamo M, Matsui T, Watanabe Y, Morimasa T, Hosokawa K and Toda T: **Proteome analysis of mouse brain: two-dimensional electrophoresis profiles of tissue proteins during the course of aging.** *Electrophoresis* 2000, **21**:1853-1871.
28. Kato H, Chen S, Kiyama H, Ikeda K, Kimura N, Nakashima K and Taga T: **Identification of a novel WD repeat-containing gene predominantly expressed in developing and regenerating neurons.** *J Biochem (Tokyo)* 2000, **128**:923-932.
29. Smith TF, Gaitatzes C, Saxena K and Neer EJ: **The WD repeat: a common architecture for diverse functions.** *Trends Biochem Sci* 1999, **24**:181-185.
30. Li D and Roberts R: **WD-repeat proteins: structure characteristics, biological function, and their involvement in human diseases.** *Cell Mol Life Sci* 2001, **58**:2085-2097.
31. Katada E, Mitake S, Matsukawa N, Otsuka Y, Tsugu Y, Fujimori O and Ojika K: **Distribution of hippocampal cholinergic neurostimulating peptide (HCNP)-like immunoreactivity in organs and tissues of young Wistar rats.** *Histochem Cell Biol* 1996, **105**:43-51.
32. Ojika K, Mitake S, Tohdoh N, Appel SH, Otsuka Y, Katada E and Matsukawa N: **Hippocampal cholinergic neurostimulating peptides (HCNP).** *Prog Neurobiol* 2000, **60**:37-83.
33. Ojika K, Mitake S, Kamiya T, Kosuge N and Taiji M: **Two different molecules, NGF and free-HCNP, stimulate cholinergic activity in septal nuclei in vitro in a different manner.** *Brain Res Dev Brain Res* 1994, **79**:1-9.
34. Morishita M, Otsuka Y, Matsukawa N, Suzuki H, Nakazawa H, Maki M, Katou H, Ueda R and Ojika K: **Specific binding of 125I-hippocampal cholinergic neurostimulating peptide (HCNP) to rat brain membranes: characterization and regional distribution.** *Brain Res* 2003, **965**:194-202.
35. Gray JA, Grigoryan G, Virley D, Patel S, Sinden JD and Hodges H: **Conditionally immortalized, multipotential and multifunctional neural stem cell lines as an approach to clinical transplantation.** *Cell Transplant* 2000, **9**:153-168.
36. Doering LC and Snyder EY: **Cholinergic expression by a neural stem cell line grafted to the adult medial septum/diagonal band complex.** *J Neurosci Res* 2000, **61**:597-604.
37. Wu P, Tarasenko YI, Gu Y, Huang LY, Coggeshall RE and Yu Y: **Region-specific generation of cholinergic neurons from fetal human neural stem cells grafted in adult rat.** *Nat Neurosci* 2002, **5**:1271-1278.
38. Maki M, Matsukawa N, Yuasa H, Otsuka Y, Yamamoto T, Akatsu H, Okamoto T, Ueda R and Ojika K: **Decreased expression of hippocampal cholinergic neurostimulating peptide precursor protein mRNA in the hippocampus in Alzheimer disease.** *J Neuropathol Exp Neurol* 2002, **61**:176-185.
39. Welch JE, Barbee RR, Roberts NL, Suarez JD and Klinefelter GR: **SP22: a novel fertility protein from a highly conserved gene family.** *J Androl* 1998, **19**:385-393.
40. Ray J, Peterson DA, Schinstine M and Gage FH: **Proliferation, differentiation, and long-term culture of primary hippocampal neurons.** *Proc Natl Acad Sci USA* 1993, **90**:3602-3606.
41. Reynolds BA and Weiss S: **Generation of neurons and astrocytes from isolated cells of the adult mammalian central nervous system.** *Science* 1992, **255**:1707-1710.
42. Bradford MM: **A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding.** *Anal Biochem* 1976, **72**:248-254.
43. Ramagli LS: **Quantifying protein in 2-D PAGE solubilization buffers.** *Methods Mol Biol* 1999, **112**:99-103.
44. Görg A, Obermaier C, Boguth G, Harder A, Scheibe B, Wildgruber R and Weiss W: **The current state of two-dimensional electrophoresis with immobilized pH gradients.** *Electrophoresis* 2000, **21**:1037-1053.
45. Berkelman T and Stenstedt T: **2-D Electrophoresis using Immobilized pH Gradients: Principles & Methods.** 2nd edition. Uppsala, Sweden, Amersham Biosciences; 2002.
46. Link AJ: **2-D proteome analysis protocols.** Totowa, NJ, Humana Press; 1999.
47. Blum H, Beier H and Gross HJ: **Improved silver staining of plant proteins, RNA and DNA in polyacrylamide gels.** *Electrophoresis* 1987, **8**:93-99.
48. Gharahdaghi F, Weinberg CR, Meagher DA, Imai BS and Mische SM: **Mass spectrometric identification of proteins from silver-stained polyacrylamide gel: a method for the removal of silver ions to enhance sensitivity.** *Electrophoresis* 1999, **20**:601-605.
49. Shevchenko A, Wilm M, Vorm O and Mann M: **Mass spectrometric sequencing of proteins silver-stained polyacrylamide gels.** *Anal Chem* 1996, **68**:850-858.
50. Castegna A, Aksenov M, Aksenova M, Thongboonkerd V, Klein JB, Pierce WM, Booze R, Markesbery WR and Butterfield DA: **Proteomic identification of oxidatively modified proteins in Alzheimer's disease brain. Part I: creatine kinase BB, glutamine synthase, and ubiquitin carboxy-terminal hydrolase L-1.** *Free Radic Biol Med* 2002, **33**:562-571.
51. Tillemann K, Stevens I, Spittaels K, Haute CV, Clerens S, Van Den Bergh G, Geerts H, Van Leuven F, Vandesande F and Moens L: **Differential expression of brain proteins in glycogen synthase kinase-3 transgenic mice: a proteomics point of view.** *Proteomics* 2002, **2**:94-104.
52. Molloy MP, Herbert BR, Walsh BJ, Tyler MI, Traini M, Sanchez JC, Hochstrasser DF, Williams KL and Gooley AA: **Extraction of membrane proteins by differential solubilization for separation using two-dimensional gel electrophoresis.** *Electrophoresis* 1998, **19**:837-844.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

