

Pan-drug-resistant and biofilm-producing strain of *Burkholderia pseudomallei*: first report of melioidosis from a diabetic patient in Yogyakarta, Indonesia [Response to Letter]

This article was published in the following Dove Press journal:
International Medical Case Reports Journal

Titik Nuryastuti¹
Nusaibah Umaroh²
Rizka Humardewyanti Asdie³
Ika Puspita Sari⁴
Ahmad Musthafa¹

¹Department of Microbiology, Faculty of Medicine, Public Health and Nursing, Universitas Gadjah Mada, Yogyakarta, Indonesia; ²Pharmacy Installation of Dr. Kariadi General Hospital, Semarang, Center of Java, Indonesia; ³Department of Internal Medicine, Faculty of Medicine, Public Health and Nursing, Universitas Gadjah Mada/Sardjito Hospital, Yogyakarta, Indonesia; ⁴Department of Pharmacology and Clinical Pharmacy, Faculty of Pharmacy, Universitas Gadjah Mada, Yogyakarta, Indonesia

Dear editor

We thank Dr David Dance et al from the International Melioidosis Society Committee for their interest in our case report. This is a response to their letter, which argued that our paper¹ does not support convincing identification of *Burkholderia pseudomallei*. In our study, we performed culture on Ashdown agar and identification by Gram staining (Gram-negative bacilli), positive oxidase test, and biochemical testing by API 20NE. The identity of the bacterium was subsequently confirmed with nested PCR, using two sets of PCR amplification primers from the variable region of the 16S rRNA gene of *B. pseudomallei*. As reported by Inglis et al, the API 20NE method identified only 37% of the *B. pseudomallei* isolates.² *B. pseudomallei* has a high degree of phenotypic plasticity and the colony morphology varies greatly within and between samples.^{3,4} Therefore, the identification was continued by nested PCR of 16S rRNA, which has been shown to be highly sensitive,^{5,6} being able to detect as few as two organisms present in the reaction.⁶ However, we fully welcome the suggestion of Dance et al that further additional testing, eg, multilocus sequence typing, nucleotide sequence analysis, or PCR for the *TTS1* gene, is needed to verify the identification of *B. pseudomallei*. We have previously asked a member of the International Melioidosis Society Committee to review our identification method; however, that request was declined at the time.

Dance et al stated that the antibiotic susceptibility pattern of our clinical isolate of *B. pseudomallei* was unusual since it showed resistance against all antibiotics tested, particularly ceftazidime and meropenem. In our case, the patient had been suffering from chronic diabetic ulcer for more than 7 years, and had already been managed with various antibiotics, including cotrimoxazole, clindamycin, metronidazole, levofloxacin, ciprofloxacin, cefadroxil, meropenem, and cefixime. It has been shown that recurrent melioidosis correlates with increasing resistance to antibiotics owing to their frequent use.⁷ In addition, ceftazidime- and carbapenem-resistant isolates have been reported in Malaysia, Australia, and north-east Thailand.⁸⁻¹⁰ As mentioned in our report, reference guidelines for the threshold of the susceptibility assay for *B. pseudomallei* have not yet been published, and therefore the interpretation results were adjusted to the reference guidelines of

Correspondence: Titik Nuryastuti
Department of Microbiology, Faculty of Medicine, Public Health and Nursing, Universitas Gadjah Mada, Farmaco Street, Sekip, Yogyakarta 55281, Indonesia
Tel +6 227 458 0297
Email t.nuryastuti@ugm.ac.id

Burkholderia cepacia, *Pseudomonas aeruginosa*, or Enterobacteriaceae. Indeed, there are fundamental issues, apart from the antibiotic susceptibility pattern, that complicate the management of *Burkholderia* infection. We should be very careful in our interpretation of the results, and therefore we agree that confirmation by measuring minimum inhibitory concentrations by the broth dilution or gradient diffusion method, as well as detection of resistance genes, is needed. Moreover, since ceftazidime and carbapenems remain the drugs of choice for management therapy of melioidosis, the development of resistance to these antibiotics should be monitored periodically.

The authors would like to thank Dr Dance et al for offering to perform confirmatory identification of this pathogen. We intend to explore further the burden of infection of *B. pseudomallei* in our community, including both clinical and environmental strains.

Disclosure

The authors report no conflicts of interest in this communication.

References

- Nuryastuti T, Umaroh N, Asdi RH, Sari IP, Musthafa A. Pan-drug-resistant and biofilm-producing strain of *Burkholderia pseudomallei*: first report of melioidosis from a diabetic patient in Yogyakarta, Indonesia. *Int Med Case Rep J*. 2018;2018(11):319–323. doi:10.2147/IMCRJ.S173461
- Inglis TJ, Merritt A, Childlow G, et al. Comparison of diagnostic laboratory methods for identification of *Burkholderia pseudomallei*. *J Clin Microbiol*. 2007;43(5):2201–2206. doi:10.1128/JCM.43.5.2201-2206.2005
- Chantratita N, Wuthiekanun V, Boonbumrung K, et al. Biological relevance of colony morphology and phenotypic switching by *Burkholderia pseudomallei*. *J Bacteriol*. 2007;189(3):807–817. doi:10.1128/JB.01258-06
- Wikraiphat C, Saiprom N, Tandhavanant S, et al. Colony morphology variation of *Burkholderia pseudomallei* is associated with antigenic variation and O-polysaccharide modification. *Infect Immun*. 2015;83(5):2127–2138. doi:10.1128/IAI.02785-14
- Dutta S, Haq S, Vijayalakshmi N, Haq JA. Antimicrobial susceptibility pattern of clinical isolates of *Burkholderia pseudomallei* in Bangladesh. *BMC Res Notes*. 2017;10:299. doi:10.1186/s13104-017-2626-5
- Dharakul T, Songsivilai S, Viriyachitra S, Luangwedchakarn V, Tassaneetritap B, Chaowagul W. Detection of *Burkholderia pseudomallei* DNA in patients with septicemic melioidosis. *J Clin Microbiol*. 1996;34(3):609–614.
- Hayden HS, Lim R, Brittnacher MJ, et al. Evolution of *Burkholderia pseudomallei* in recurrent melioidosis. *PLoS One*. 2012;7:e36507. doi:10.1371/journal.pone.0036507
- Khosravi Y, Vellamy M, Mariappan V, Ng S-L, Vadivelu J. Antimicrobial susceptibility and genetic characterisation of *Burkholderia pseudomallei* isolated from Malaysian patients. *Sci World J*. 2014;2014:1–9. Article ID 132971. doi:10.1155/2014/132971
- Jenney AW, Lum G, Fisher DA, Currie BJ. Antibiotic susceptibility of *Burkholderia pseudomallei* from tropical northern Australia and implications for therapy of melioidosis. *Int J Antimicrob Agents*. 2001;17:109–113. doi:10.1016/S0924-8579(00)00334-4
- Wuthiekanun V, Amornchai P, Saiprom N, et al. Survey of antimicrobial resistance in clinical *Burkholderia pseudomallei* isolates over two decades in Northeast Thailand. *Antimicrob Agents Chemother*. 2011;55(11):5388–5391. doi:10.1128/AAC.0517-11

Dove Medical Press encourages responsible, free and frank academic debate. The content of the International Medical Case Reports Journal 'letters to the editor' section does not necessarily represent the views of Dove Medical Press, its officers, agents, employees, related entities or the International Medical Case Reports Journal editors. While all reasonable steps have been taken to confirm the content of each letter, Dove Medical Press accepts no liability in respect of the content of any letter, nor is it responsible for the content and accuracy of any letter to the editor.

International Medical Case Reports Journal

Dovepress

Publish your work in this journal

The International Medical Case Reports Journal is an international, peer-reviewed open-access journal publishing original case reports from all medical specialties. Previously unpublished medical posters are also accepted relating to any area of clinical or preclinical science. Submissions should not normally exceed 2,000 words or 4

published pages including figures, diagrams and references. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/international-medical-case-reports-journal-journal>