

SCIENTIFIC REPORTS



OPEN

The Sema3A receptor Plexin-A1 suppresses supernumerary axons through Rap1 GTPases

Nannan Wang¹, Pratibha Dhumale^{1,2,3}, Joanna Chiang¹ & Andreas W. Püschel^{1,2}

The highly conserved Rap1 GTPases perform essential functions during neuronal development. They are required for the polarity of neuronal progenitors and neurons as well as for neuronal migration in the embryonic brain. Neuronal polarization and axon formation depend on the precise temporal and spatial regulation of Rap1 activity by guanine nucleotide exchange factors (GEFs) and GTPases-activating proteins (GAPs). Several Rap1 GEFs have been identified that direct the formation of axons during cortical and hippocampal development *in vivo* and in cultured neurons. However little is known about the GAPs that limit the activity of Rap1 GTPases during neuronal development. Here we investigate the function of Sema3A and Plexin-A1 as a regulator of Rap1 GTPases during the polarization of hippocampal neurons. Sema3A was shown to suppress axon formation when neurons are cultured on a patterned substrate. Plexin-A1 functions as the signal-transducing subunit of receptors for Sema3A and displays GAP activity for Rap1 GTPases. We show that Sema3A and Plexin-A1 suppress the formation of supernumerary axons in cultured neurons, which depends on Rap1 GTPases.

Small GTPases of the Ras superfamily perform essential functions throughout neuronal development and in mature neurons¹. The highly conserved Rap1 GTPases encoded by the *Rap1a* and *Rap1b* genes in mammals are required for the polarity of neuronal progenitors and neurons as well as for neuronal migration in the embryonic brain^{1–5}. In the developing brain, newborn neurons that initially have a multipolar morphology become polarized by forming an axon and a leading process^{1,6–9}. In culture, dissociated neurons from the embryonic hippocampus or cortex undergo a similar differentiation but polarize without the need for a patterned exogenous signal^{7,10}. After attaching to the culture substrate neurons first extend several undifferentiated neurites (stage 2 of neuronal polarization) before one of them becomes an axon and extends rapidly. The inactivation of Rap1 GTPases impairs the formation of axons during cortical and hippocampal development *in vivo* and in cultured neurons^{1,11}. Neuronal polarization and axon formation depend on the precise temporal and spatial regulation of Rap1 activity by GEFs and GAPs. Rapgef1 (also called C3G), Rapgef2 and Rapgef6 have been identified as the Rap1 GEFs that are required for the development of the neocortex and hippocampus^{12–16}. However little is known about the GAPs that limit the activity of Rap1 GTPases during neuronal development¹⁴.

The plexins are integral membrane proteins with an intracellular domain that shows sequence similarity to GAPs with dual specificity for Ras and Rap GTPases^{17–21}. The mutation of conserved arginine residues in this GAP domain is sufficient to abolish their activity *in vitro* and *in vivo*^{20–23}. Plexins are receptors for the semaphorins, a large family of secreted and membrane-bound proteins that act as axon guidance signals but also perform important functions in other tissues^{24–26}. The nine plexins in mammals can be subdivided into four subfamilies (PlexinA to -D). *In vitro* assays first showed that Plexin-B1 acts as a GAP for R- and M-Ras but not H-Ras^{27–31}. Subsequently, a structural analysis of the GAP domain combined with biochemical assays demonstrated that Plexin-A1 and -C1 specifically regulate Rap1 and Rap2 GTPases^{20,21,32–34}. The phenotype of plexin mutants confirmed that GAP activity is essential for their function *in vivo* and provided evidence for a regulation of Ras and Rap1 GTPases^{22,23,35}.

The A-type plexins act as the signal-transducing subunit of receptors for the secreted Sema3A in a complex with Neuropilin-1 as the ligand binding subunit^{24,36–40}. Semaphorins perform important functions not only

¹Institut für Molekulare Zellbiologie, Westfälische Wilhelms-Universität, Schloßplatz 5, D-48149, Münster, Germany.

²Cells-in-Motion Cluster of Excellence, University of Münster, D-48149, Münster, Germany. ³Present address: Department of Cardiovascular and Renal Research, University of Southern Denmark, JB Winsløvs Vej 21, 5000, Odense, Denmark. Pratibha Dhumale and Joanna Chiang contributed equally. Correspondence and requests for materials should be addressed to A.W.P. (email: apuschel@uni-muenster.de)




Figure 1. Hippocampal neurons from *Sema3a^{-/-}* embryos extend supernumerary axons in culture. (a–c) Cultures of hippocampal or cortical neurons from E17 *Sema3a^{+/-}* or *Sema3a^{-/-}* embryos were analyzed at 3, 4, 5 and 6 d.i.v. by staining with an anti-MAP2 (green, dendrites) and the Tau-1 (a, red, axons) or an anti-AnkG antibody (b, red, AIS marked by arrow heads). Representative images of hippocampal neurons at the indicated times in culture are shown. The scale bar is 25 μm. (c) The percentage of unpolarized hippocampal neurons without an axon or AIS (0, red), polarized neurons with a single axon or AIS (1, blue), neurons with multiple axons or AIS (>1, green) and neurons with neurites that are positive for both axonal and dendritic markers (in, yellow) is shown (Student's t-test and two-way ANOVA; n = 3 independent experiments with >150 neurons per genotype; values are means ± s.e.m., **p < 0.01, ***p < 0.001 compared to control as indicated).

during axon guidance but also in the regulation of neuronal migration and the formation of axons, dendrites and synapses^{25,36,41–43}. *Sema3A* directs the orientation of axons and apical dendrites in the developing cortex^{44–47}. It can also regulate the establishment of neuronal polarity by promoting the formation of dendrites and suppressing the extension of axons when neurons are cultured on a patterned substrate with stripes of immobilized *Sema3A*^{48,49}. In cultures of sensory neurons from embryonic dorsal root ganglia, *Sema3A* accelerates the establishment of neuron polarity⁵⁰. It remains to be investigated if plexins regulate the activity of Rap1 GTPases during neuronal polarization.

Here we show that *Sema3A* and *Plexin-A1* suppress the formation of supernumerary axons in cultured neurons. Hippocampal but not cortical neurons from *Sema3A* knockout embryos form multiple axons in culture. While inactivation of *Plexin-A1* results in the formation of supernumerary axons constitutively active *Plexin-A1* inhibits axon formation. These effects can be rescued by the knockdown of *Rap1B* and the expression of active *Rap1B*, respectively, indicating that *Plexin-A1* suppresses axon formation by regulating *Rap1* GTPases.

Results

Hippocampal neurons from *Sema3a* knockout mice form multiple axons. To determine if the establishment of neuronal polarity is affected in the absence of *Sema3A* neurons were isolated from the hippocampus and cortex of E17 *Sema3a^{-/-}* and *Sema3a^{+/-}* embryos and analyzed by staining with an anti-MAP2 and the Tau-1 antibody as markers for dendrites and axons, respectively (Fig. 1a,c). The majority (60 ± 5%) of the *Sema3^{-/-}* neurons extended several long neurites that were positive for both axonal and dendritic markers

at day 3 days in culture (3 days *in vitro*, d.i.v.) compared to only $9 \pm 4\%$ in cultures from *Sema3a*^{+/-} embryos as control (Fig. 1c). To evaluate axon formation with a second axonal marker, cultures were analyzed at 4, 5 and 6 d.i.v. by staining with the Tau-1 or an anti-Ankyrin G (AnkG) antibody as a marker for the axonal initial segment (AIS) (Fig. 1b,c). A large proportion of the *Sema3a*^{-/-} neurons formed multiple Tau-1 positive axons ($44 \pm 1\%$) and multiple AIS ($43 \pm 1\%$) at 4 d.i.v. while only few were observed in *Sema3a*^{+/-} controls (axons: $11 \pm 2\%$; AIS: $8 \pm 1\%$). The number of *Sema3a*^{-/-} neurons with multiple Tau-1-positive axons increased to $63 \pm 3\%$ (AnkG: $58 \pm 1\%$) at 5 d.i.v. and $71 \pm 1\%$ (AnkG: $73 \pm 1\%$) at 6 d.i.v. (Fig. 1c). Thus, *Sema3A*-deficient neurons from the embryonic hippocampus showed defects in neuronal polarization. Neurons extended multiple neurites that initially display mixed axonal and dendritic characteristics at 3 d.i.v. but become supernumerary axons at 4 d.i.v. When cultures of cortical neurons from *Sema3a*^{+/-} and *Sema3a*^{-/-} embryos were analyzed no significant defect in neuronal polarization was observed at any of the analyzed time points (Fig. 1a,c). These results show that the knockout of *Sema3a* induces the formation of supernumerary axons by hippocampal but not cortical neurons.

The knockdown of *Sema3A* induces cell-autonomous defects. Secreted *Sema3A* tightly binds to the cell surface of the producing cells and can act as an autocrine signal⁵¹. To investigate this possibility we used an miRNA expression vector to knock down *Sema3A* in cultured neurons. Since only a small number of neurons were transfected, a knockdown reduces the concentration of *Sema3A* in the medium only minimally compared to cultures from knockout embryos. The efficiency of the *Sema3A* knockdown construct was confirmed by Western blot after co-expression with tagged *Sema3A* in HEK 293 T cells (Fig. 2a) and by immunofluorescence staining of cultured hippocampal neurons (Suppl. Fig. S1a). Hippocampal neurons were transfected with the *Sema3A* knockdown vector and analyzed at 3 d.i.v. (Fig. 2b,c). Suppression of *Sema3a* induced the formation of multiple Tau-1 positive axons in $45 \pm 1\%$ of the neurons compared to $8 \pm 2\%$ in controls. By contrast, non-transfected neurons in the same culture were not affected by the knockdown indicating a cell-autonomous function of *Sema3A* in cultured neurons (Suppl. Fig. S1b). The phenotype of the *Sema3A* knockdown could be rescued by the expression of an RNAi-resistant *Sema3A* construct (*Sema3A*-res; Fig. 2) confirming the specificity of the knockdown. Only $15 \pm 2\%$ of the neurons that were co-transfected with the RNAi vector and the expression vector for *Sema3A*-res extended multiple axons and $78 \pm 3\%$ formed a single axon. These results suggest that *Sema3A* acts as an autocrine signal to suppress the formation of supernumerary axons.

Plexin-A but not Plexin-B activity is required for neuronal polarity. *Sema3A* acts through receptors that contain an A-type plexin as the signal transducing subunit^{17,26,52,53}. To investigate whether plexins are involved in the regulation of neuronal polarization, we used a Plexin-A1 construct (Plexin-A1 Δ cyt) with a deletion of the intracellular domain that has a dominant-negative effect by forming non-functional receptor complexes^{19,39,52}. Neurons from the hippocampus of E18 rat embryos were transfected with vectors for Plexin-A1 Δ cyt and analyzed at 3 d.i.v. by staining with an anti-MAP2 and the Tau-1 antibody (Fig. 3a). Expression of Plexin-A1 Δ cyt increased the number of neurons with multiple axons from $7 \pm 1\%$ in controls to $61 \pm 2\%$ (Fig. 3b). Deletion of the semaphorin domain releases Plexin-A1 from its auto-inhibited state and results in a constitutively active receptor (Plexin-A1 Δ sema)⁵⁴. Expression of Plexin-A1 Δ sema had the opposite effect of dominant-negative Plexin-A1. The majority of neurons were negative for Tau-1 staining and the percentage of unpolarized neurons was increased to $69 \pm 3\%$ (control: $9 \pm 1\%$; Fig. 3b). These results indicate that interfering with the function of A-type plexins disrupts the establishment of neuronal polarity.

To test a possible involvement of B-type plexins in axon formation, we used dominant-negative Plexin-B1 Δ cyt. After expression of Plexin-B1 Δ cyt, hippocampal neurons polarize normally and extend a single Tau-1 positive axon (Fig. 3a). No significant difference was detectable between controls (neurons with a single axon: $85 \pm 4\%$) and the expression of Plexin-B1 Δ cyt ($82 \pm 25\%$; Fig. 3b). Thus, inhibition of Plexin-B1 has no effect on axon formation.

Suppression of Plexin-A1 induces multiple axons. The dominant-negative Plexin-A1 Δ cyt construct may affect all receptors that include a member of the Plexin-A subfamily. To identify which A-type plexin regulates neuronal polarity, we used established RNAi vectors for PlexinA1 - A4⁴¹. Rat hippocampal neurons were transfected with these RNAi vectors and axon formation was analyzed at 3 d.i.v. (Fig. 4a,b). After knockdown of Plexin-A1, the number of neurons with multiple axons increased from $7 \pm 2\%$ in controls to $51 \pm 4\%$ (Fig. 4b). In addition, $20 \pm 4\%$ of the neurons formed neurites that were positive for both axonal and dendritic markers. By contrast, after knockdown of Plexin-A2, -A3 or -A4 no significant increase in the extension of supernumerary axons was observed (neurons with a single axon: Plexin-A2: $77 \pm 4\%$, Plexin-A3: $74 \pm 3\%$; Plexin-A4: $72 \pm 2\%$; control: $77 \pm 5\%$). Staining with an anti-Plexin-A1 antibody confirmed that the knockdown efficiently suppressed the expression of Plexin-A1 that is present in all neurites of polarized hippocampal neurons (Fig. 4c). The formation of supernumerary axons after knockdown of Plexin-A1 could be reversed by co-expressing murine Plexin-A1 that contains mismatches in the RNAi target site compared to the rat sequence confirming the specificity of the Plexin-A1 knockdown in rat neurons (Suppl. Fig. S2). These results show that Plexin-A1 is required during neuronal polarization to prevent the formation of supernumerary axons.

Plexin-A1 acts upstream of Rap1 GTPases during neuronal polarization. Structural and biochemical analyses showed that Plexin-A1 regulates Rap1 GTPases that are required for the formation of axons^{11,20,21}. The inactivation of the Rap1 GAP Plexin-A1 may, therefore, result in the extension of supernumerary axons due to increased Rap1 activity. To investigate this possibility, we tested if the formation of multiple axons after expression of dominant-negative Plexin-A1 Δ cyt can be blocked by a knockdown of Rap1B. Hippocampal neurons were co-transfected with vectors for Plexin-A1 Δ cyt and an shRNA directed against Rap1B¹¹ and analyzed at 3 d.i.v. (Fig. 5a). After knockdown of Rap1B, only $19 \pm 4\%$ of the transfected neurons extended an axon compared to




Figure 2. Knockdown of Sema3A induces the formation of supernumerary axons. **(a)** HEK 293 T cells were transfected with vectors for FLAG-Sema3A (S3A) or RNAi-resistant FLAG-Sema3A-res (S3Ares) and an shRNA directed against Sema3A (S3A RNAi) or pcDNA6.2-GW/EmGFP-miR (Ctrl). The expression of Sema3A and GFP was analyzed by Western blot (WB) using an anti-FLAG antibody. The molecular weight is indicated in kDa. **(b)** Hippocampal neurons from E18 rat embryos were transfected with vectors for GFP (green), an shRNA against Sema3A (Sema3A RNAi) or pcDNA6.2-GW/EmGFP-miR (control), and a vector for RNAi-resistant FLAG-Sema3A-res (Sema3A-res) or pBK-CMV (control) as indicated. Neurons were analyzed at 3 d.i.v. by staining with an anti-MAP2 (red, dendrites) and the Tau-1 antibody (blue, axon). Representative images of transfected neurons are shown. The scale bar is 25 μ m. **(c)** The percentage of unpolarized neurons without an axon (0, red), polarized neurons with a single axon (1, blue) and neurons with multiple axons (>1, green) is shown (Student's t-test and two-way ANOVA; $n = 3$, independent experiments with >150 neurons per experiment; values are means \pm s.e.m., *** $p < 0.001$ compared to control as indicated; n.s., not significant).

84 \pm 3% in controls (Fig. 5b) as shown before¹¹. The induction of supernumerary axons by Plexin-A1 Δ cyt was prevented by the knockdown of Rap1B (unpolarized neurons: 26 \pm 5%, neurons with a single axon: 66 \pm 3%, neurons with multiple axons: 8 \pm 4%) indicating that it depends on Rap1B. The specificity of the Rap1B knockdown was verified by co-expression of an Rap1B construct with mismatches in the RNAi target site that rescued the loss of axons after Rap1B knockdown (Suppl. Fig. S3).

The expression of constitutively active Plexin-A1 Δ sema has an effect that is similar to that of a Rap1B knockdown and leads to the loss of axons. We tested if this phenotype can be rescued by the expression of active Rap1BV12. As reported before¹¹, expression of Rap1BV12 increased the number of hippocampal neurons with multiple axons (53 \pm 1%) compared to controls (17 \pm 1%) while expression of Plexin-A1 Δ sema increased the number of unpolarized neurons (51 \pm 2%, control: 11 \pm 1%; Fig. 6a,b). The suppression of axon formation by PlexinA1 Δ sema could be rescued by co-expression of Rap1BV12, which increased the percentage of polarized




Figure 3. A-type plexins suppress the formation of axons. (a,b) Hippocampal neurons from E18 rat embryos were transfected with vectors for GFP (green, control), dominant-negative Plexin-B1 Δ cyt (PlxnB1-DN), Plexin-A1 Δ cyt (PlxnA1-DN) or constitutively active PlexinA1 Δ sema (PlxnA1-CA) and analyzed at 3 d.i.v. by staining with an anti-MAP2 (red, dendrites) and the Tau-1 antibody (blue, axon). Representative images of transfected neurons are shown. The scale bar is 25 μ m. (b) The percentage of unpolarized neurons without an axon (0, red), polarized neurons with a single axon (1, blue) and neurons with multiple axons (>1, green) is shown (Student's t-test and two-way ANOVA; n = 3, independent experiments with >150 neurons per experiment for; values are means \pm s.e.m, ***p < 0.001 compared to control; n.s. not significant).

neurons with a single axon from $43 \pm 1\%$ (only PlexinA1 Δ sema) to $63 \pm 4\%$ (PlexinA1 Δ sema and Rap1BV12, Fig. 6b). These results indicate that Plexin-A1 acts upstream of Rap1 GTPases and restricts their activity.

To confirm that Rap1 GTPases act downstream of Plexin-A1 we tested if the formation of supernumerary axons caused by the knockdown of Plexin-A1 can be prevented by the knockdown of Rap1B. After transfection of hippocampal neurons with the shRNA vector directed against Plexin-A1, $48 \pm 1\%$ of the transfected neurons extended multiple axons (Fig. 7a,b). This number was reduced to $14 \pm 1\%$ when the shRNA against Rap1B was cotransfected with the Plexin-A1 shRNA and the majority of the neurons ($70 \pm 3\%$) extended a single Tau-1 positive axon. These results show that Rap1B is required for the induction of supernumerary axons by the knockdown of Plexin-A1. Taken together, our results show that Plexin-A1 acts upstream of Rap1B and is required to restrict its activity in hippocampal neurons.

Rap1B acts downstream of Sema3A in hippocampal neurons. Sema3A-deficient neurons extend multiple axons similar to the phenotype of neurons transfected with a vector for constitutively active Rap1BV12. To investigate if Rap1B acts downstream of Sema3A we tested if the knockdown of Rap1B prevents the formation of supernumerary axons after knockdown of Sema3A. $32 \pm 5\%$ of the transfected neurons extended multiple axons (Fig. 8) that were also positive for Rap1 after the knockdown of Sema3A (Suppl. Fig. S4). This number was reduced to $18 \pm 2\%$ with the majority of the neurons ($67 \pm 6\%$) extending a single Tau-1 positive axon when Rap1B was knocked down together with Sema3A. These results show that the formation of supernumerary axons in Sema3A-deficient neurons depends on Rap1B.

Discussion

Our results suggest that Sema3A acts as an autocrine signal through Plexin-A1, which restricts the activity of Rap1 and thereby prevents the formation of supernumerary axons. Hippocampal neurons from *Sema3a*^{-/-} knockout mice form multiple axons in culture. Suppressing Plexin-A1 function by a dominant-negative construct or a knockdown had a similar effect while expression of constitutively active Plexin-A1 blocked axon formation. The defects in neuronal polarity were rescued by the knockdown of Rap1B and the expression of active Rap1BV12, respectively, which is consistent with the activity of Plexin-A1 as a GAP for Rap1^{20,21}. These results




Figure 4. Knockdown of Plexin-A1 induces the formation of supernumerary axons. **(a,b)** Hippocampal neurons from E18 rat embryos were transfected with vectors for GFP (green) and shRNAs against Plexin-A1, -A2, -A3, or -A4 or pSUPER (control) and analyzed at 3 d.i.v. by staining with an anti-MAP2 (red, dendrites) and the Tau-1 antibody (blue, axon). Representative images of transfected neurons are shown. The scale bar is 25 μ m. **(b)** The percentage of unpolarized neurons without an axon (0, red), polarized neurons with a single axon (1, blue), neurons with multiple axons (>1, green) and neurons with neurites that are positive for both axonal and dendritic markers (in, yellow) is shown (Student's t-test and two-way ANOVA; $n = 3$, independent experiments with >150 neurons per experiment; values are means \pm s.e.m., ** $p < 0.01$ compared to control). **(c)** Hippocampal neurons from E18 rat embryos were transfected with vectors for GFP (green) and an shRNA against Plexin-A1 (RNAi) or pSUPER (control) and analyzed at 3 d.i.v. by staining with an anti-Plexin-A1 (red) and the Tau-1 antibody (blue, axon). Representative images of transfected neurons are shown. The scale bar is 25 μ m.

indicate that Rap1 GTPases act downstream of Plexin-A1 and are required for its function to suppress the formation of supernumerary axons.

Our results are consistent with previous reports that Sema3A suppresses axon formation^{48,49}. Hippocampal neurons express Sema3A^{44,55} that remains tightly bound to the surface of producing cells after secretion by binding to proteoglycans⁵¹. A knockdown of Sema3A induces the formation of supernumerary axons only in transfected neurons while non-transfected neurons in the same culture are not affected. The co-expression of an RNAi-resistant Sema3A rescues the Sema3A knockdown and prevents the formation of supernumerary axons. These results suggest that Sema3A acts primarily in a cell-autonomous manner as an autocrine factor as described for other cell types^{56–60}. *Sema3a* knockout neurons extend multiple long neurites that are initially positive for both axonal and dendritic markers at 3 d.i.v. before they become axons by 4 d.i.v. while most neurons extend




Figure 5. Knockdown of Rap1B counteracts the induction of supernumerary axons by dominant-negative Plexin-A1. **(a)** Hippocampal neurons from E18 rat embryos were transfected with vectors for GFP (green, control), an shRNA against Rap1B or pSHAG-1 (control) and Plexin-A1 Δ cyt (PlxnA1-DN) or pBK-CMV (control). The establishment of neuronal polarity was analyzed at 3 d.i.v. by staining with an anti-MAP2 (red, dendrites) and the Tau-1 antibody (blue, axons). Representative images of transfected neurons are shown. The scale bar is 25 μ m. **(b)** The percentage of unpolarized neurons without an axon (0, red), polarized neurons with a single axon (1, blue) and neurons with multiple axons (>1, green) is shown (Student's t-test and two-way ANOVA; $n = 3$, independent experiments with >150 neurons per experiment for; values are means \pm s.e.m, ** $p < 0.01$; *** $p < 0.001$ compared to control and as indicated).

multiple axons already at 3 d.i.v. after expression of dominant-negative Plexin-A1 and knockdown of Plexin-A1 or Sema3A. The major reason for this difference in the time course probably is the time point at which Sema3A/Plexin-A1 signaling is inhibited. Sema3A acts like a global inhibitory signal that suppresses axon formation^{48,61}. It is absent from the beginning in knockout neurons while it is transiently produced after a knockdown, which may explain why knockout neurons initially extend neurites with both dendritic and axonal properties before these become axons.

In contrast to Plexin-A1, we did not observe defects in neuronal polarity after expression of dominant-negative Plexin-B1. The analysis of Plexin-B1 suggested that it acts as a GAP for R-Ras and M-Ras²⁷⁻³⁰. Different Ras GTPases have been implicated in axon formation^{29,62,63}. The activation of Plexin-B1 by Sema4D down-regulates the activity of R- and M-Ras in the growth cones of axons and dendrites of cultured hippocampal neurons, respectively^{27,30-32}. However, neither single nor double knockouts of Ras GTPases show defects in brain development⁶⁴, indicating that they are not required for neuronal polarity *in vivo* or that their loss is compensated by other Ras GTPases. *In vivo* studies indicate that B-type plexins perform different functions during neuronal development^{22,65-69}. Knockout of *Plxnb1* and *Plxnb2* results in a reduced proliferation of neural progenitors and cortical thinning, which could indicate a function in regulating the orientation of the mitotic spindle as shown for epithelial cells^{22,67,69}. Further studies are required to elucidate which GTPases are regulated by the different plexins *in vivo* and what determines their specificity. It also remains to be investigated whether other GAPs are involved in mediating the function of Sema3A or the regulation of Rap1 GTPases during neuronal polarization.

Unlike in hippocampal neurons, axon formation was not affected in cortical neurons. The differential effects of the *Sema3a* knockout may result from differences in the expression of semaphorin receptors and other semaphorins like Sema3C may act in the developing neocortex. Sema3C is expressed in migrating neurons in the cortex and its overexpression interferes with their polarization and radial migration⁴³. Sema3A could be required




Figure 6. Active Rap1B rescues the suppression of axon formation by constitutively active Plexin-A1. **(a)** Hippocampal neurons from E18 rat embryos were transfected with vectors for GFP (green) and Plexin-A1 Δ sema (PlxnA1-CA), Rap1BV12 or a combination of both and analyzed at 3 d.i.v. by staining with an anti-MAP2 (red, dendrites) and the Tau-1 antibody (blue, axon). pBK-CMV and pSHAG-1 were transfected as control. Representative images of transfected neurons are shown. The scale bar is 25 μ m. **(b)** The percentage of unipolarized neurons without an axon (0, red), polarized neurons with a single axon (1, blue) and neurons with multiple axons (>1, green) is shown (Student's t-test and two-way ANOVA; $n = 3$, independent experiments with >150 neurons per experiment; values are means \pm s.e.m., ** $p < 0.01$; *** $p < 0.001$ compared to control and as indicated).

specifically in hippocampal neurons to suppress axon formation. A difference between hippocampus and cortex was observed also for the *Rap1a;Rap1b* knockout¹. The conditional knockout of Rap1 GTPases at different time points showed that an inactivation late during neuronal polarization interferes with axon formation in the hippocampus but not in the cortex. The hippocampus-specific functions of *Sema3A* and Rap1 GTPases could be linked to the extended time that hippocampal neurons remain in the multipolar phase of migration, which may require additional mechanisms to delay axon formation⁷⁰.

Materials and Methods

Sema3a knockout mice. *Sema3a*^{+/-} mice⁷¹ were maintained in a C57Bl/6 background. Genotyping was performed using the primers 5'-ATGGTTCTGA TAGGTGAGGC ATGG-3', 5'-GTTCTGCTCC CGGCTCTAAA TCTC-3' and 5'-AGGCAAAC TA TGCAAACGG AAAG-3'⁷². Mice were housed at four to five per cage with a 12 h light/dark cycle (lights on from 07:00 to 19:00 h) at constant temperature (23 °C) with *ad libitum* access to food and water. All animal protocols were carried out in accordance with the relevant guidelines and regulations and approved by the Landesamt für Natur, Umwelt und Verbraucherschutz Nordrhein-Westfalen.

Cell culture and transfection. Cultures of hippocampal and cortical neurons were prepared from E17 *Sema3a* mouse embryos as described before¹. Hippocampal neurons from E18 rat embryos were prepared and transfected by calcium phosphate co-precipitation as described previously¹. Dissociated neurons were plated at a density as 65,000 cells per well of a 24-well plate containing cover slips coated with poly-L-ornithine (15 μ g/ml, SigmaAldrich). Neurons were cultured at 37 °C and 5% CO₂ for 3–6 days in Neurobasal medium (Invitrogen) with supplements. An excess of the expression and knockdown vectors were combined with a small amount of pEGFP-N3 (Clontech) to label transfected neurons.




Figure 7. Rap1B is required for the induction of supernumerary axons by the knockdown of Plexin-A1 in hippocampal neurons. **(a)** Hippocampal neurons from E18 rat embryos were transfected with vectors for GFP (green) and Plexin-A1 Δ sema (PlxnA1-CA), Rap1BV12, pSHAG-1 (control) or a combination of both and analyzed at 3 d.i.v. by staining with an anti-MAP2 (red, dendrites) and the Tau-1 antibody (blue, axon). Representative images of transfected neurons are shown. The scale bar is 25 μ m. **(b)** The percentage of unpolarized neurons without an axon (0, red), polarized neurons with a single axon (1, blue) and neurons with multiple axons (>1, green) is shown (Student's t-test and two-way ANOVA; $n = 3$, independent experiments with >150 neurons per experiment for; values are means \pm s.e.m., ** $p < 0.01$; *** $p < 0.001$ compared to control and as indicated).

Immunofluorescence staining. Neurons were fixed with 4% paraformaldehyde/15% sucrose in phosphate buffered saline (PBS) for 15 min at RT and permeabilized with 0.1% Triton X-100/0.1% Na-Citrate/PBS for 10 min on ice. Cells were incubated in 10% normal goat serum (PAN Biotech) in PBS for 1 h at RT, stained with primary antibody overnight at 4 $^{\circ}$ C and secondary antibody for 90 min, and mounted using Mowiol (SigmaAldrich). Neuronal morphology and axon formation were analyzed as described before using a Zeiss Axiophot microscope equipped with a Visitron CCD camera and the SPOT Advanced Imaging software or a Zeiss LSM 700 using the ZEN (black edition) software¹. Image analysis was done using ImageJ 1.45 s (NIH), ZEN (black edition) and Adobe Photoshop CS5. All statistical data are means \pm s.e.m. from at least three times independent experiments. Statistical significance was determined using the Student's t-test.

Antibodies. The following antibodies were used: mouse Tau-1 (Chemicon, MAB3420, 1:200), mouse anti-MAP2 (SigmaAldrich, M4403, 1:1500), rabbit anti-MAP2 (Abcam, ab32454, 1:1000), mouse anti-Ankyrin-G (Antibodies Inc., 75–146, 1:100), rabbit anti-Plexin-A1 (Abcam, ab23391, 1:1000), mouse anti-GFP (Covance, MMS-118P, 1:1000), mouse anti-FLAG M2 (SigmaAldrich, F3165, 1:1000), anti-Rap1 (Upstate, #07–916, 1:200), anti-Sema3A (Abcam, ab23393, 1:200), SMI-312 (BioLegend, 837904, 1:200) and goat secondary antibodies labeled with Alexa-350 (Molecular Probes, 1:200), –488 (1:800) or –594 (1:800). Nuclei were stained with Hoechst 33342 (Molecular Probes, 1:6000).

Plasmids. Published shRNA vectors were used for a knockdown of different A-type plexins (target sequences: Plexin-A1: 5'-CCGTATTTAC AAGCTGTCG-3'; Plexin-A2: 5'-GCGCAAGTCT AGGGAAAAT-3'; Plexin-A3: 5'-GTGCGGGTTC GGCCTAATA-3'; Plexin-A4: 5'-AGATGCTGCT TATAGAC TA-3')⁴¹. The Plexin-A1 shRNA is specific for the rat sequence⁴¹. For rescue experiments, an expression vector for murine Plexin-A1¹⁹ that contains mismatches in the shRNA target site compared to the rat sequence was used. The shRNA vector targeting Rap1B and the expression vectors for Rap1BV12 and Flag-Sema3A have been described before^{11,73}. A vector for RNAi-resistant Rap1B (Rap1B-res) was constructed by site-directed mutagenesis of the shRNA target site¹¹ to introduce




Figure 8. Rap1B acts downstream of Sema3A in hippocampal neurons. **(a)** Hippocampal neurons from E18 rat embryos were transfected with vectors for GFP (green), an shRNA against Sema3A (S3A RNAi) or pcDNA6.2-GW/EmGFP-miR (control) and an shRNA against Rap1B (R1B RNAi) or pSHAG-1 (control) as indicated. Neurons were analyzed at 3 d.i.v. by staining with an anti-MAP2 (red, dendrites) and the Tau-1 antibody (blue, axon). Representative images of transfected neurons are shown. The scale bar is 25 μ m. **(b)** The percentage of unpolarized neurons without an axon (0, red), polarized neurons with a single axon (1, blue) and neurons with multiple axons (>1, green) is shown (Student's t-test and two-way ANOVA; $n = 3$, independent experiments with >150 neurons per experiment for; values are means \pm s.e.m, *** $p < 0.001$ compared to control as indicated).

mismatches using the QuikChange Site-Directed Mutagenesis kit (Stratagene) with the oligonucleotides 5'-GT TGTAGGAA AAGAACAGGG TCAAAACCTA GCAAGACAG-3' and 5'-CTGTCT TGCT AGGTTTTGAC CC TGTTCTTT TCCTACAAC-3'. The miRNA vector targeting Sema3A with the target sequence 5'-TTCCGGGAAC CAACAACACTATT-3' was generated using the BLOCK-iT Pol II miRNA Expression Vector Kit (Invitrogen), inserted into pcDNA6.2-GW/EmGFP-miR and confirmed by sequencing. A vector for RNAi-resistant FLAG-Sema3A (Sema3A-res) was constructed by site-directed mutagenesis using the QuikChange Site-Directed Mutagenesis kit (Stratagene) with the oligonucleotides 5'-GAAATGACCG TCTTCCGTGA ACCGACAACC ATTTTCAGCAATG-3' and 5'-CATTGCTGAA ATGGTTGTCG GTTCACGGAA GACGGTCATT TC-3'. pE GFP-N3 (Clontech) was used as transfection control. The dominant-negative Plexin-A1 construct pBK-CMV-VSV-Plexin-A1 Δ cyt with a deletion of the intracellular domain and the constitutively active pBK-CMV-VSV-Plexin-A1 Δ sema with a deletion of the semaphorin domain were constructed as described before^{19,39,52,54}.

Transfection of HEK 293T cells and Western blot. HEK 293T cells were transfected using the calcium phosphate co-precipitation method as described previously⁷⁴. Transfected HEK 293T cells were lysed in Tris/HCl 50 mM, pH 7.4, NaCl 150 mM, DTT 1 mM, MgCl₂ 1.5 mM, EDTA 4 mM, glycerol 10% (v/v), Triton X-100 1% (v/v), cComplete protease inhibitor (Sigma-Aldrich) and expression of Sema3A analyzed by Western blot using horseradish peroxidase conjugated secondary antibodies (Dianova, 1:3000). Peroxidase activity was visualized by the enhanced chemiluminescence detection system (Uptima, Interchim UP99619A) using the ChemiDocTM MP imaging system (Bio-Rad).

Quantification. The establishment of neuronal polarity was quantified by counting the number of transfected (GFP-positive) neurons that did not extend an axon (unpolarized neuron), formed a single axon positive for Tau-1 (polarized neuron), multiple axons positive for Tau-1 (multiple axons) or multiple axons positive for Tau-1 and MAP2 (indeterminate phenotype). Statistical analyses were done using the GraphPad Prism 6.0 software. Statistical significance was calculated for at least three independent experiments using two-way ANOVA with Tukey's multiple comparison test and Student's t-Test for parametric data sets. Significance was defined as: $p > 0.05$, n.s.; * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$.

References

- Shah, B. *et al.* Rap1 GTPases Are Master Regulators of Neural Cell Polarity in the Developing Neocortex. *Cereb Cortex* **27**, 1253–1269 (2017).
- Franco, S. J., Martinez-Garay, I., Gil-Sanz, C., Harkins-Perry, S. R. & Muller, U. Reelin regulates cadherin function via Dab1/Rap1 to control neuronal migration and lamination in the neocortex. *Neuron* **69**, 482–497 (2011).
- Jossin, Y. & Cooper, J. A. Reelin, Rap1 and N-cadherin orient the migration of multipolar neurons in the developing neocortex. *Nat Neurosci* **14**, 697–703 (2011).
- Ye, T., Ip, J. P., Fu, A. K. & Ip, N. Y. Cdk5-mediated phosphorylation of RapGEF2 controls neuronal migration in the developing cerebral cortex. *Nat Commun* **5**, 4826 (2014).
- Shah, B. & Püschel, A. W. *In vivo* functions of small GTPases in neocortical development. *Biol Chem* **395**, 465–476 (2014).
- Funahashi, Y., Namba, T., Nakamura, S. & Kaibuchi, K. Neuronal polarization *in vivo*: Growing in a complex environment. *Curr Opin Neurobiol* **27**, 215–223 (2014).
- Namba, T. *et al.* Extracellular and Intracellular Signaling for Neuronal Polarity. *Physiol Rev* **95**, 995–1024 (2015).
- Namba, T. *et al.* Pioneering axons regulate neuronal polarization in the developing cerebral cortex. *Neuron* **81**, 814–829 (2014).
- Sakakibara, A. & Hatanaka, Y. Neuronal polarization in the developing cerebral cortex. *Front Neurosci* **9**, 116 (2015).
- Schelski, M. & Bradke, F. Neuronal polarization: From spatiotemporal signaling to cytoskeletal dynamics. *Mol Cell Neurosci* (2017).
- Schwamborn, J. C. & Püschel, A. W. The sequential activity of the GTPases Rap1B and Cdc42 determines neuronal polarity. *Nat Neurosci* **7**, 923–929 (2004).
- Bilasy, S. E. *et al.* Dorsal telencephalon-specific RA-GEF-1 knockout mice develop heterotopic cortical mass and commissural fiber defect. *Eur J Neurosci* **29**, 1994–2008 (2009).
- Maeta, K. *et al.* Crucial Role of Rapgef2 and Rapgef6, a Family of Guanine Nucleotide Exchange Factors for Rap1 Small GTPase, in Formation of Apical Surface Adherens Junctions and Neural Progenitor Development in the Mouse Cerebral Cortex. *eNeuro* **3**, ENEURO.0142-16.2016 (2016).
- Shah, B. & Püschel, A. W. Regulation of Rap GTPases in mammalian neurons. *Biol Chem* **397**, 1055–1069 (2016).
- Voss, A. K., Krebs, D. L. & Thomas, T. C3G regulates the size of the cerebral cortex neural precursor population. *EMBO J* **25**, 3652–3663 (2006).
- Voss, A. K. *et al.* C3G regulates cortical neuron migration, preplate splitting and radial glial cell attachment. *Development* **135**, 2139–2149 (2008).
- Hota, P. K. & Buck, M. Plexin structures are coming: opportunities for multilevel investigations of semaphorin guidance receptors, their cell signaling mechanisms, and functions. *Cell Mol Life Sci* **69**, 3765–3805 (2012).
- Pascoe, H. G., Wang, Y. & Zhang, X. Structural mechanisms of plexin signaling. *Prog Biophys Mol Biol* **118**, 161–168 (2015).
- Rohm, B., Rahim, B., Kleiber, B., Hovatta, I. & Püschel, A. W. The semaphorin 3A receptor may directly regulate the activity of small GTPases. *FEBS Lett* **486**, 68–72 (2000).
- Wang, Y., Pascoe, H. G., Brautigam, C. A., He, H. & Zhang, X. Structural basis for activation and non-canonical catalysis of the Rap GTPase activating protein domain of plexin. *Elife* **2**, e01279 (2013).
- Wang, Y. *et al.* Plexins are GTPase-activating proteins for Rap and are activated by induced dimerization. *Sci Signal* **5**, ra6 (2012).
- Worzfeld, T. *et al.* Genetic dissection of plexin signaling *in vivo*. *Proc Natl Acad Sci USA* **111**, 2194–2199 (2014).
- Yang, T. & Terman, J. R. 14-3-3epsilon couples protein kinase A to semaphorin signaling and silences plexin RasGAP-mediated axonal repulsion. *Neuron* **74**, 108–121 (2012).
- Jongbloets, B. C. & Pasterkamp, R. J. Semaphorin signalling during development. *Development* **141**, 3292–3297 (2014).
- Koropouli, E. & Kolodkin, A. L. Semaphorins and the dynamic regulation of synapse assembly, refinement, and function. *Curr Opin Neurobiol* **27**, 1–7 (2014).
- Worzfeld, T. & Offermanns, S. Semaphorins and plexins as therapeutic targets. *Nat Rev Drug Discov* **13**, 603–621 (2014).
- Oinuma, I., Ito, Y., Katoh, H. & Negishi, M. Semaphorin 4D/Plexin-B1 stimulates PTEN activity through R-Ras GTPase-activating protein activity, inducing growth cone collapse in hippocampal neurons. *J Biol Chem* **285**, 28200–28209 (2010).
- Oinuma, I., Katoh, H. & Negishi, M. Molecular dissection of the semaphorin 4D receptor plexin-B1-stimulated R-Ras GTPase-activating protein activity and neurite remodeling in hippocampal neurons. *J Neurosci* **24**, 11473–11480 (2004).
- Oinuma, I., Katoh, H. & Negishi, M. R-Ras controls axon specification upstream of glycogen synthase kinase-3beta through integrin-linked kinase. *J Biol Chem* **282**, 303–318 (2007).
- Saito, Y., Oinuma, I., Fujimoto, S. & Negishi, M. Plexin-B1 is a GTPase activating protein for M-Ras, remodelling dendrite morphology. *EMBO Rep* **10**, 614–621 (2009).
- Tasaka, G., Negishi, M. & Oinuma, I. Semaphorin 4D/Plexin-B1-mediated M-Ras GAP activity regulates actin-based dendrite remodeling through Lamellipodin. *J Neurosci* **32**, 8293–8305 (2012).
- Ito, Y., Oinuma, I., Katoh, H., Kaibuchi, K. & Negishi, M. Sema4D/plexin-B1 activates GSK-3beta through R-Ras GAP activity, inducing growth cone collapse. *EMBO Rep* **7**, 704–709 (2006).
- Yukawa, K. *et al.* Sema4A induces cell morphological changes through B-type plexin-mediated signaling. *Int J Mol Med* **25**, 225–230 (2010).
- Yoo, S. K. *et al.* Plexins function in epithelial repair in both Drosophila and zebrafish. *Nat Commun* **7**, 12282 (2016).
- Ueda, Y. *et al.* Sema3e/Plexin D1 Modulates Immunological Synapse and Migration of Thymocytes by Rap1 Inhibition. *J Immunol* **196**, 3019–3031 (2016).
- Goshima, Y., Yamashita, N., Nakamura, F. & Sasaki, Y. Regulation of dendritic development by semaphorin 3A through novel intracellular remote signaling. *Cell Adh Migr* **10**, 627–640 (2016).
- He, Z. & Tessier-Lavigne, M. Neuropilin is a receptor for the axonal chemorepellent Semaphorin III. *Cell* **90**, 739–751 (1997).
- Kolodkin, A. L. *et al.* Neuropilin is a semaphorin III receptor. *Cell* **90**, 753–762 (1997).
- Tamagnone, L. *et al.* Plexins are a large family of receptors for transmembrane, secreted, and GPI-anchored semaphorins in vertebrates. *Cell* **99**, 71–80 (1999).
- Winberg, M. L. *et al.* Plexin A is a neuronal semaphorin receptor that controls axon guidance. *Cell* **95**, 903–916 (1998).
- Chen, G. *et al.* Semaphorin-3A guides radial migration of cortical neurons during development. *Nat Neurosci* **11**, 36–44 (2008).
- Sasaki, Y. *et al.* Fyn and Cdk5 mediate semaphorin-3A signaling, which is involved in regulation of dendrite orientation in cerebral cortex. *Neuron* **35**, 907–920 (2002).

43. Wiegrefe, C. *et al.* Bcl11a (Ctip1) Controls Migration of Cortical Projection Neurons through Regulation of Sema3c. *Neuron* **87**, 311–325 (2015).
44. Chedotal, A. *et al.* Semaphorins III and IV repel hippocampal axons via two distinct receptors. *Development* **125**, 4313–4323 (1998).
45. Polleux, F., Giger, R. J., Ginty, D. D., Kolodkin, A. L. & Ghosh, A. Patterning of cortical efferent projections by semaphorin-neuropilin interactions. *Science* **282**, 1904–1906 (1998).
46. Polleux, F., Morrow, T. & Ghosh, A. Semaphorin 3A is a chemoattractant for cortical apical dendrites. *Nature* **404**, 567–573 (2000).
47. Ruediger, T. *et al.* Integration of opposing semaphorin guidance cues in cortical axons. *Cereb Cortex* **23**, 604–614 (2013).
48. Shelly, M. *et al.* Semaphorin3A regulates neuronal polarization by suppressing axon formation and promoting dendrite growth. *Neuron* **71**, 433–446 (2011).
49. Nishiyama, M. *et al.* Semaphorin 3A induces Ca_v2.3 channel-dependent conversion of axons to dendrites. *Nat Cell Biol* **13**, 676–685 (2011).
50. Lerman, O., Ben-Zvi, A., Yagil, Z. & Behar, O. Semaphorin3A accelerates neuronal polarity *in vitro* and in its absence the orientation of DRG neuronal polarity *in vivo* is distorted. *Mol Cell Neurosci* **36**, 222–234 (2007).
51. De Wit, J., De Winter, F., Klooster, J. & Verhaagen, J. Semaphorin 3A displays a punctate distribution on the surface of neuronal cells and interacts with proteoglycans in the extracellular matrix. *Mol Cell Neurosci* **29**, 40–55 (2005).
52. Takahashi, T. *et al.* Plexin-neuropilin-1 complexes form functional semaphorin-3A receptors. *Cell* **99**, 59–69 (1999).
53. Wu, K. Y. *et al.* Semaphorin 3A activates the guanosine triphosphatase Rab5 to promote growth cone collapse and organize callosal axon projections. *Sci Signal* **7**, ra81 (2014).
54. Takahashi, T. & Strittmatter, S. M. PlexinA1 autoinhibition by the plexin sema domain. *Neuron* **29**, 429–439 (2001).
55. Nakamura, F. *et al.* Increased proximal bifurcation of CA1 pyramidal apical dendrites in sema3A mutant mice. *J Comp Neurol* **516**, 360–375 (2009).
56. Foley, K. *et al.* Semaphorin 3D autocrine signaling mediates the metastatic role of annexin A2 in pancreatic cancer. *Sci Signal* **8**, ra77 (2015).
57. Bagci, T., Wu, J. K., Pfannl, R., Ilag, L. L. & Jay, D. G. Autocrine semaphorin 3A signaling promotes glioblastoma dispersal. *Oncogene* **28**, 3537–3550 (2009).
58. Fukuda, T. *et al.* Sema3A regulates bone-mass accrual through sensory innervations. *Nature* **497**, 490–493 (2013).
59. Hamm, M. J., Kirchmaier, B. C. & Herzog, W. Sema3d controls collective endothelial cell migration by distinct mechanisms via Nr1 and PlxnD1. *J Cell Biol* **215**, 415–430 (2016).
60. Man, J. *et al.* Sema3C promotes the survival and tumorigenicity of glioma stem cells through Rac1 activation. *Cell Rep* **9**, 1812–1826 (2014).
61. Arimura, N. & Kaibuchi, K. Neuronal polarity: from extracellular signals to intracellular mechanisms. *Nat Rev Neurosci* **8**, 194–205 (2007).
62. Yoshimura, T. *et al.* Ras regulates neuronal polarity via the PI3-kinase/Akt/GSK-3beta/CRMP-2 pathway. *Biochem Biophys Res Commun* **340**, 62–68 (2006).
63. Fivaz, M., Bandara, S., Inoue, T. & Meyer, T. Robust neuronal symmetry breaking by Ras-triggered local positive feedback. *Curr Biol* **18**, 44–50 (2008).
64. Esteban, L. M. *et al.* Targeted genomic disruption of H-ras and N-ras, individually or in combination, reveals the dispensability of both loci for mouse growth and development. *Mol Cell Biol* **21**, 1444–1452 (2001).
65. Azzarelli, R. *et al.* An antagonistic interaction between PlexinB2 and Rnd3 controls RhoA activity and cortical neuron migration. *Nat Commun* **5**, 3405 (2014).
66. Deng, S. *et al.* Plexin-B2, but not Plexin-B1, critically modulates neuronal migration and patterning of the developing nervous system *in vivo*. *J Neurosci* **27**, 6333–6347 (2007).
67. Daviaud, N., Chen, K., Huang, Y., Friedel, R. H. & Zou, H. Impaired cortical neurogenesis in plexin-B1 and -B2 double deletion mutant. *Dev Neurobiol* **76**, 882–899 (2016).
68. Hirschberg, A. *et al.* Gene deletion mutants reveal a role for semaphorin receptors of the plexin-B family in mechanisms underlying corticogenesis. *Mol Cell Biol* **30**, 764–780 (2010).
69. Xia, J. *et al.* Semaphorin-Plexin Signaling Controls Mitotic Spindle Orientation during Epithelial Morphogenesis and Repair. *Dev Cell* **33**, 299–313 (2015).
70. Kitazawa, A. *et al.* Hippocampal pyramidal neurons switch from a multipolar migration mode to a novel “climbing” migration mode during development. *J Neurosci* **34**, 1115–1126 (2014).
71. Taniguchi, M. *et al.* Disruption of semaphorin III/D gene causes severe abnormality in peripheral nerve projection. *Neuron* **19**, 519–530 (1997).
72. Schlomann, U., Schwamborn, J. C., Muller, M., Fassler, R. & Puschel, A. W. The stimulation of dendrite growth by Sema3A requires integrin engagement and focal adhesion kinase. *J Cell Sci* **122**, 2034–2042 (2009).
73. Adams, R. H., Lohrum, M., Klostermann, A., Betz, H. & Puschel, A. W. The chemorepulsive activity of secreted semaphorins is regulated by furin-dependent proteolytic processing. *EMBO J* **16**, 6077–6086 (1997).
74. Yang, R., Kong, E., Jin, J., Hergovich, A. & Puschel, A. W. Rassf5 and Ndr kinases regulate neuronal polarity through Par3 phosphorylation in a novel pathway. *J Cell Sci* **127**, 3463–3476 (2014).

Acknowledgements

We thank Maria Wenning and Ina Kowsky for technical assistance and Drs. T. Yagi, M. Taniguchi, T. Holtmaat and J. Verhaagen for providing us with the *Sema3a*^{+/-} mice. This work was supported by the Deutsche Forschungsgemeinschaft (DFG) through the Cells-in-Motion Cluster of Excellence (EXC 1003 - CiM) and with a fellowship to N.W. by the China Scholarship Council.

Author Contributions

A.W.P. conceived the study, N.W., P.D. and J.C. performed the experiments, N.W., P.D., J.C. and A.W.P. contributed to the experimental design and interpretation of the results, and N.W. and A.W.P. wrote the manuscript.

Additional Information

Supplementary information accompanies this paper at <https://doi.org/10.1038/s41598-018-34092-5>.

Competing Interests: The authors declare no competing interests.

Publisher's note: Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2018