

BMJ Open Effect of eHealth-delivered exercise programmes on balance in people aged 65 years and over living in the community: a systematic review and meta-analysis of randomised controlled trials

Meghan Ambrens ,^{1,2} Stephanie Alley ,³ Juliana S Oliveira,^{4,5} Quyen To,³ Kim Delbaere ,^{2,6} Corneel Vandelanotte,³ Anne Tiedemann^{4,5}

To cite: Ambrens M, Alley S, Oliveira JS, *et al.* Effect of eHealth-delivered exercise programmes on balance in people aged 65 years and over living in the community: a systematic review and meta-analysis of randomised controlled trials. *BMJ Open* 2022;**12**:e051377. doi:10.1136/bmjopen-2021-051377

► Prepublication history and additional supplemental material for this paper are available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2021-051377>).

Received 19 March 2021
Accepted 08 May 2022



© Author(s) (or their employer(s)) 2022. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

Correspondence to

Dr Meghan Ambrens;
m.ambrens@neura.edu.au

ABSTRACT

Introduction Exercise that challenges balance is proven to prevent falls in community-dwelling older people, yet widespread implementation and uptake of effective programmes is low. This systematic review and meta-analysis synthesised the evidence and evaluated the effect of eHealth-delivered exercise programmes compared with control on balance in community-dwelling people aged ≥65 years.

Methods Nine databases including MEDLINE, CINAHL and Embase, were searched from inception to January 2022 to identify randomised controlled trials evaluating eHealth-delivered exercise programmes for community-dwelling people aged ≥65 years, published in English that included a balance outcome. Primary outcomes were static and dynamic balance. Secondary outcomes included fall risk and fear of falling. We calculated standardised mean differences (SMDs, Hedges' g) with 95% CIs from random effects meta-analyses.

Results We identified 14 eligible studies that included 1180 participants. Methodological quality ranged from 3 to 8 (mean, 5). The pooled effect indicated that eHealth-delivered exercise programmes have a medium significant effect on static balance (11 studies; SMD=0.62, 95% CI 0.27 to 0.72) with very low-quality evidence. There was small statistically significant effect on dynamic balance (14 studies; SMD=0.42, 95% CI 0.11 to 0.73) with very low-quality evidence, and fall risk (5 studies; SMD=0.32, 95% CI 0.00 to 0.64) with moderate-quality evidence. No significant effect of eHealth programmes on fear of falling was found (four studies; SMD=0.10, 95% CI -0.05 to 0.24; high-quality evidence).

Conclusion This review provides preliminary evidence that eHealth-delivered exercise programmes improved balance and reduced fall risk in people aged ≥65 years. There is still uncertainty regarding the effect of eHealth delivered exercise programmes on fear of falling.

PROSPERO registration number CRD42018115098.

STRENGTHS AND LIMITATIONS OF THIS STUDY

- ⇒ We conducted this systematic review in accordance with the Preferred Reporting Items for Systematic Reviews and Meta-Analyses guidelines and followed a prespecified protocol registered on PROSPERO.
- ⇒ We had specific criteria that allowed for the inclusion of studies with both generally healthy participants and those with selected clinical conditions.
- ⇒ We also ensured the inclusion of as many relevant studies as possible by searching across nine databases, conducting a thorough hand search of relevant published literature and consulting with experts in the field.
- ⇒ Included studies had to comprise a measure of balance and we may have missed relevant studies that included measures of falls.
- ⇒ We only included outcome data from the immediate postintervention time point, which limits the interpretation of results to the short-term.

INTRODUCTION

Ageing is associated with a decline in the physiological systems responsible for postural stability and hence an increase in the risk of falls.¹ A fall is defined as 'an unexpected event in which the participants come to rest on the ground, floor, or other lower level'.² Each year approximately one-third of community-dwelling people aged 65 years and over experience a fall.^{3,4} Forty per cent of injuries requiring hospital admission are due to falls, and in 2016–2017 more than 125 000 Australians aged 65 years or over were hospitalised due to a fall.^{5,6} Falls place a significant burden on health systems and can result in serious long-term costs to the individual.⁷ Falls can also result in the loss of independence, depression, social isolation and admission of

the older person to a care facility.⁷ As such, fall prevention is a public health priority.

There is clear evidence that exercise is crucial for preventing falls in community-dwelling older people.⁸ However, the effect on falls varies by exercise type. A Cochrane systematic review that included 108 studies of exercise, established that exercise that challenges balance has the greatest effect on both the rate of falls (24% reduction) and risk of falls (13% reduction) in community-dwelling older people.⁸ Despite the benefits of exercise for preventing falls, widespread implementation and adherence to effective programmes is poor, significantly reducing the population-wide impact.⁹ Furthermore, the 2020 WHO guidelines on physical activity and sedentary behaviour recommend that older adults should undertake multicomponent physical activity that emphasises functional balance and strength training at least three-times per week to enhance functional capacity and prevent falls.¹⁰ Therefore, exploring the effectiveness of novel exercise programmes with potential for wide reach that can improve balance is important.

Advances in technology have led to new ways to deliver exercise-based programmes. Such technology-based programmes, commonly referred to as electronic-Health (eHealth), that use the internet, websites, mobile applications (apps) or exergames, may provide effective alternatives to more traditional modes of delivering exercise-based programmes to improve balance and prevent falls, and increase access to such programmes. Previous studies show that eHealth interventions can successfully improve the health and physical activity of older people,^{11–16} and adherence by some older people is higher for technology-delivered interventions compared with traditional interventions, independent of study site, level of supervision and mode of delivery.¹⁷

eHealth-delivered exercise programmes are a safe way to exercise, and many older people perceive them as fun and enjoyable.^{18–22} This mode of delivery has also been successful for improving balance in younger adults, aged 18+ years, with a number of systematic reviews evaluating the effectiveness of specific, technology-based approaches to improving balance or reducing fall risk in adults.^{23–26}

Given the lack of systematic reviews and meta-analyses on the effectiveness of eHealth-delivered exercise programmes for improving balance in older people, further evaluation of the role of technology-driven platforms is needed. This systematic review and meta-analysis aimed to synthesise the evidence and evaluated the effect of eHealth-delivered exercise programmes on balance in people aged 65 years and older living in the community compared with a control.

METHODS

Protocol

This review was registered on the PROSPERO database and followed the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines.²⁷

The systematic review protocol has been published and provides a full outline of the methods.²⁸ A summary of the methods is reported in this paper.

Data sources

A database search was conducted from inception to January 2022 of MEDLINE, CINAHL Complete, Embase, PsychINFO, Scopus, Web of Science, PubMed Central, Cochrane Database Central and PEDro. The protocol²⁸ details the complete search strategy used. Online supplemental figure 1 provides the MEDLINE search strategy.

Eligibility criteria

Studies included in this systematic review met the following criteria: (1) published in English, (2) randomised controlled trials (RCTs), (3) participants were community-dwelling people aged ≥ 65 years, (4) reported data for a validated measure of balance, (5) included eHealth delivery of an exercise programme compared with no intervention, usual care or wait-list control. Studies that did not meet these criteria were excluded.

We included all RCT designs such as crossover, cluster, patient-randomised clinical trials that examined the effect of eHealth-delivered exercise programmes vs a control group. Single and multi-factorial interventions were also included. Studies published only as abstracts or yet to be published were excluded due to possible data inaccuracy and incompleteness.

Outcome measures

The primary outcome measure was balance, defined as staying upright and steady when stationary, such as when standing, or sitting, or during movement.²⁹ Technically, balance is defined as ‘the ability to maintain the projection of the body’s centre of mass (CoM) within manageable limits of the base of support, as in sitting or standing, or in transit to a new base of support, as in walking’.³⁰ The balance outcomes were further categorised as static or dynamic measures of balance in the analyses. Static balance refers to maintaining balance when the body has a constant or static base of support.²⁹ Whereas dynamic balance refers to maintaining balance during movement from one base of support to another, such as when walking.²⁹ In the absence of functional measures of balance, we included studies which reported direct measures of balance, such as those measured with a force platform. Fall risk, fear of falling and fall rate were included as secondary outcomes. We included studies that used either validated self-report questionnaires or performance-based measures for these outcomes.

Study selection

After pilot-testing criteria for full-text articles, screening for eligible studies was conducted independently by two reviewers (MA, KLA/RS). An electronic screening form was used, and screening occurred in stages: first, titles were screened, followed by abstracts and finally

full-text articles were screened. Conflicts were resolved by consensus from AT, KD and CV.

Data extraction

Data extraction was completed by two researchers independently from one another (MA, QT), and conflicts were resolved by a third reviewer (SA). Data were extracted using a piloted electronic data extraction form, and according to the PRISMA statement.²⁷ Where data were missing, study authors were contacted by email to provide further information. Where the authors did not reply within 2 weeks, a second email was sent as a reminder.

The following data were extracted from each study: author, year of publication, country, sample characteristics (sample size, age, sex of participants and health status), study design: including number of study arms, recruitment sources, eligibility criteria, setting, delivery method and technology used, intervention description, comparator, intervention duration and frequency, assessment time-points. Also extracted were data on drop out, attrition, adverse events and intervention features such as implementation fidelity, evidenced-based theory, tailoring, supervision, intervention acceptability. Primary and secondary outcome data were extracted for pre-intervention and postintervention timepoints. Where data were available for more than one postintervention timepoint, we included the data from the timepoint closest to intervention completion.

Methodological quality and risk of bias assessment

The PEDro scale (1–10) was used to assess the methodological quality of the included studies. PEDro scores were extracted from the PEDro database.³¹ The PEDro scale assesses the internal validity of an RCT by evaluating 11 items: participant eligibility criteria, random allocation, concealed allocation, homogeneity of groups at baseline, blinding of subjects, blinding of therapist, blinding of assessor, completeness of follow-up, intention-to-treat analysis, between-group statistical analysis, and variability and point measures.³² A score of 10 is considered to be methodologically excellent, whereas 0 demonstrates poor methodological quality.³² Methodological quality was not an inclusion criterion for this review.

In addition to the PEDro scale, we also used the Cochrane Risk of Bias to assess the risk of bias in each included study. The Cochrane Risk of Bias was undertaken by two independent reviewers (MA, SA) with conflicts resolved by a third reviewer (CV). Risk of bias is assessed across a number of domains as a judgement of low risk, high risk, or unclear.³³

Assessment of quality of evidence

To evaluate overall quality of evidence we used the Grading of Recommendations, Assessment, Development and Evaluations (GRADE) system. The GRADE appraisal was conducted by pairs of independent reviewers (MA, JSO) and guided by the Cochrane Handbook for Systematic Reviews.³³ This is a subjective evaluation of the quality

of the evidence as high, moderate, low or very low based on the presence or extent of the following factors: risk of bias, imprecision and inconsistency of the effect. The GRADE classification was downgraded from high quality by one level for each factor encountered: (1) design limitations (>25% of studies with low methodological quality based on the Cochrane Risk of bias), (2) inconsistency of results (large heterogeneity between the studies $I^2 > 50\%$), (3) imprecision (<400 participants for each outcome).³³ We did not consider indirectness as it encompasses a specific population (older people) with relevant outcome measures (balance) and direct comparisons. We were unable to consider publication bias for secondary outcome measures due to the limited number of studies that collected measures for the fall-related outcome measures.³³

Statistical analysis

We performed meta-analyses with Comprehensive Meta-Analysis software (V.3, Biostat, Englewood, New Jersey, USA) using the random effects models for primary and secondary outcome measures. The random effects model was chosen given the heterogeneity of the population and the interventions being evaluated. We calculated treatment effects for the continuous variables using standardised mean differences (SMDs, Hedges' g) standardised by postscore standard deviation (or its estimate) with 95% CIs, for either between-group differences in point estimates at the follow-up time points or between-group differences in change scores based on available data. SMDs were calculated using the premean and postmean and SD. Where this was not available, we used the mean change score. Effect sizes were categorised as small (0.2), medium (0.5) or large (0.8 or greater).³⁴

We visually inspected the forest plot for evidence of heterogeneity among studies with consideration of the I^2 and χ^2 tests. We determined clinical heterogeneity by consensus among the investigators on the basis of collective experience in the field.

RESULTS

Flow of studies included in this review

The initial search of the databases resulted in 1080 publications. An additional hand search, including the reference lists of relevant review articles found a further 40 publications. After removing duplicate papers, 783 publications were screened by title and abstract. Forty-five publications reported on potentially eligible studies before full-text screening. After the full-text screen, 14 studies, reported by 15 manuscripts, were identified as eligible and included in this study. Schoene *et al*³⁵ conducted a study involving interactive cognitive-motor step training, however this paper did not report the balance outcomes measured during this study. These balance outcomes were reported in Gschwind *et al*³⁶ as the step-mat-training (SMT) intervention group. This review extracted outcome data for the SMT group only from




Figure 1 Preferred Reporting Items for Systematic Reviews and Meta-Analyses flow of studies through the review. **RCT, randomised controlled trial.**

Gschwind *et al.*³⁶ paper, all other data pertaining to this study (including for the control group) were extracted from Schoene *et al.*³⁵ Figure 1 outlines the PRISMA study flow of studies included in this review.

We pooled all included studies in the primary meta-analysis evaluating the effect of interventions that use an eHealth technology to deliver an exercise programme to older people.

Risk of bias and quality

Table 1 reports the methodological quality of eligible studies. The total PEDro scores ranged from 3 to 8 (mean of 5). For the static balance outcome five studies out of 11 (45%) were of high methodological quality (a score ≥ 6).^{18 36–39} For the dynamic balance outcome, 5 studies out of 14 (36%) were of high methodological quality.^{18 36–39} All participants were randomly allocated and provided the calculation of point estimates and variability (PEDro items 2 and 11). Four studies out of 14 (29%) did not undertake an intention-to-treat analysis^{18 40–42} and 3 studies provided insufficient information to determine this.^{43–45} None of the studies included blinded participants or blinded therapists, however, blinding of participants or therapists is not possible for exercise interventions. Table 2 reports the quality of the evidence. Figure 2 presents the risk of bias.

Cohort characteristics

Studies included samples ranging from 9 to 503 participants (n=1180). The mean age of participants ranged from 65 to 89 years. Both males and females were included in 11 studies, all had a higher percentage of female participants.^{18 36–38 40 42 46 47} Two studies included only female participants,^{41 45} one study included only male participants.⁴⁸ Every study recruited participants from the general community.^{18 36–48} Two studies recruited

participants with a history of falls.^{41 45} Table 1 presents a summary of cohort characteristics.

Characteristics of included studies

Publication dates ranged from 2013 to 2021, with 11 (76%) published during or after 2015. Studies were conducted in eight different countries: single studies were conducted in Hong Kong,⁴⁶ Taiwan,⁴⁷ the USA,⁴² the UK,⁴⁰ Japan,⁴⁴ Malaysia⁴⁸ and Thailand⁴³; three studies were conducted in Australia^{18 36 39} and South Korea^{37 41 45}; and one study was conducted across three countries: Germany, Spain and Australia.³⁸ Details of study characteristics are summarised in table 1.

eHealth-delivered exercise programmes

The duration of the included interventions ranged from 2 to 52 weeks, with a mean duration of 12 weeks. Eleven studies (79%) used a commercially available exergame system to deliver the exercises: five (36%) used the Microsoft Xbox Kinect,^{38 43 44 46 48} four (29%) studies used the Nintendo Wii console,^{37 40 42 45} two (14%) used the Dance Dance Revolution StepMania.^{18 36} Three studies (21%) used customised technologies: an app-based intervention called *StandingTall*,³⁹ a web-based intervention called telepresence,⁴¹ and the Xavix Measured Step System.⁴⁷

Five studies (36%) used technology to provide a home-based intervention.^{18 36 38 39 41} Nine studies (64%) used technology to deliver a supervised intervention: participants attended a supervised group class,^{37 44 46} or a supervised one-on-one session.^{40 42 43 45 47 48}

In seven studies (50%), the control group received no information and were encouraged to continue with normal daily activities.^{40 42 47} Control participants in five studies (36%) received educational advice related to fall prevention, general health, nutrition and physical activity, in the form of a booklet or classes.^{36–39 41} Participants in one study continued with the regular, seated social games available at the senior's activities centre.⁴⁶

Effect of eHealth-delivered exercise programmes on balance

Eleven studies (79%) measured static balance using static balance,³⁹ single leg stance,^{37 46–48} tandem stance,³⁸ postural sway,^{18 36 40} functional reach⁴⁵ and the Fullerton Advanced Balance Scale.⁴² The pooled effect of eHealth-delivered exercise programmes on static balance indicates a medium, statistically significant effect compared with control (SMD=0.62, 95% CI 0.27 to 0.72; $I^2=82%$, $p=0.001$) (figure 3). The pooled results provide very low-quality evidence (GRADE).

All 14 studies measured dynamic balance with methods ranging from the Timed Up and Go,^{36–39 42 43 45–48} the Berg Balance Scale^{40 41 44} and the Alternative Step test.¹⁸ The pooled effect of eHealth-delivered exercise programmes on dynamic balance indicates there was a small, statistically significant effect compared with control (SMD=0.42, 95% CI 0.11 to 0.73; $I^2=79%$, $p=0.009$) (figure 4). The pooled results provide very low-quality evidence

Table 1 Summary of the characteristics of the included studies

Sample		eHealth exercise intervention							
Study author; country	Study design; sample size; PEDro score	Intervention content	Outcome measures: primary and secondary	Tailored progression of exercises	Dosage: duration; session length; frequency	Setting	Comparator	Adverse events	Intervention acceptability and how it was assessed
Bienyla and Dold; USA ⁴²	RCT; N=12; 3 81.5±5.5; 83%; healthy	Exercises: yoga (half-moon, chair, warrior), aerobic (torso twist), balance games (soccer heading, ski jump)	Primary: FAB, BBS	↑ intensity; ↑ challenge ↑ duration	3 wk; 30 min; 3x wk	Centre; supervised			Not assessed
Chow and Mann; Hong Kong ⁴⁶	RCT; N=20; 3 70.4; 65%; healthy	Exercises: Tiger Woods PGA Tour 13 golf game	Primary: SLS, TUG		2 wk; 30–45 min; daily	Centre; supervised	Table games		Not assessed
Delbaere et al; Australia ³⁹	RCT; N=503; 8 77.1±5.5; 67%; healthy	Exercises: balance and lower limb muscle strength; health education	Primary: standing balance, TUG Secondary: PPA, icon-FES	↑ intensity; ↑ challenge ↑ duration	52 wk (balance measures), 104 wk (fall measures); 120 min per wk	Home; unsupervised	Health education programme delivered via iPad	5 falls	SUS to assess usability
Gschwind et al; Australia, Germany, Spain ³⁶	RCT; N=153; 6 74.7±6.3; 61%; healthy	Exercise: balance and muscle strength; educational booklet on general health and fall prevention	Primary: tandem stance, TUG Secondary: icon-FES, PPA		16 wk; balance: 40 min; 3x wk. Strength: 15–20 min; 3x wk	Home; unsupervised	Educational booklet: general health and fall prevention	24 falls (IG: 8, CG: 15)	SUS to assess usability and enjoyment; DART to assess user acceptance.
Gschwind et al; Schoene et al; Australia ³⁵ ³⁶	RCT; n=90; 7 82±7; 66%; healthy	Exercises: trail-stepping – visual attention, set-shifting; stepping to connect numbers and letters, Tetris-stepping – visuo-spatial skills, problem-solving; stepping	Primary: standing balance, TUG Secondary: PPA	↑ intensity; ↑ challenge	16 wk; 20 min; 3x wk	Home; unsupervised	Educational booklet: general health and fall prevention		Not assessed
Hong et al; South Korea ⁴¹	RCT; N=30; 5 78.10; 100%; fall risk	Exercises: resistance, two-legged standing, tandem standing, one-legged standing, semi-tandem standing, tandem walking, turning in a circle, toe stands+nutrition and exercise education	Primary: BBS Secondary: FES-K	↑ intensity	12 wk; 20–40 min; 3x wk	Home; supervised	Nutrition and exercise education		Not assessed
Jung et al; Republic of Korea ⁴⁵	RCT; N=24; 4 74.3±2.1; 100%; fall risk	Exercises (NWS group): including the wakeboard, Frisbee dog, jet ski and canoe games	Primary: functional reach, TUG, BBS		8 wk; 30 min; 2x wk	Centre; supervised			Not assessed
Lai et al; Taiwan ⁴⁷	Cross-over; N=30; 4 70.6±3.5; 57%; healthy	Exercises: IVGB stepping exercise	Primary: SLS, BBS		6 wk; 30 min; 3x wk	Centre; supervised	Cross-over trial		Not assessed

Continued

Table 1 Continued

Sample		eHealth exercise intervention							
Study author; country	Study design; sample size; PEDro score	Intervention content	Outcome measures: primary and secondary	Tailored progression of exercises	Dosage: duration; session length; frequency	Setting	Comparator	Adverse events	Intervention acceptability and how it was assessed
Lee <i>et al</i> ; South Korea ³⁷	RCT; N=44; 6 76.15±4.55; 77%; healthy	Exercises: jogging for gait, swordplay for agility and balance, ski jump for balance, hula-hoop for balance and lower extremity strength, tennis for balance and agility, and step dance for gait for lower extremity strength	Primary: SLS, BBS		6 wk; 60 min; 2x wk	Centre; supervised	3x fall education sessions		Not assessed
Phirom <i>et al</i> ; Thailand ⁴³	RCT; N=40 70.21±4.18; healthy	Exercises: interactive game-based training which involved stepping on different targets and in different directions and balance training	Primary: TUG Secondary: PPA		12 wk; 60 min; 3x wk	Centre; supervised			Not assessed
Sadeghi <i>et al</i> ; Malaysia ⁴⁸	RCT; N=64; 5 71.8±6.09; 0%; healthy	Exercises (BT group): single-leg stance with eyes open and closed, standing on heels or toes, tandem and semi-tandem foot stance, tandem walking, walking backward and forward, and weight shifting	Primary: SLS, TUG	↑ intensity; ↑ challenge	8 wk; 40 min; 3x wk	Centre; supervised	Educational material covering cognitive enhancement and fall prevention strategies		Not assessed
Sato <i>et al</i> ; Japan ⁴⁴	RCT; N=54; 4 69.25±5.41; 80%; healthy	Complete four games – apple game, balloon popping game, tightrope game, one-leg standing game	Primary: BBS		9 wk (24 sessions); 40–60 min; 2–3x wk	Centre; supervised			Not assessed
Schoene <i>et al</i> ; Australia ¹⁸	RCT; N=37; 7 77.5±4.5; healthy	Exercises: a series of stepping challenges+cognitive load	Primary: SB, AST Secondary: icon-FES, PPA	↑ intensity	8 wk; 15–20 min; 2–3x wk	Home; unsupervised			Participants were asked whether they enjoyed playing the exergame (YES/NO response)
Whyatt <i>et al</i> ; UK ⁴⁰	RCT; N=82; 5 77.18±6.59; 70%; healthy	Exercises: games targeting components of balance (Apple Catch, Bubble Pop, Avoid the Shark, Smart Shrimp)	Primary: SLS, TUG Secondary: ABC Scale	↑ challenge	5 wk; 30 min; 2x wk	Centre; supervised	Activity diaries		Not assessed

Wk: week(s); PEDro: Physiotherapy Evidence Database (used to rate the methodological quality of studies – ‘poor’ PEDro score of 1–3, ‘moderate’ PEDro score of 4–5 and ‘high’ PEDro score of 6–10); IVGB: interactive video game based; +, plus; ↑, increase; Centre: exercise delivered within a gymnasium, clinic or other facility that is not the participants place of residence; Primary: primary outcome measure, Secondary: secondary outcome measure; Static balance: SLS, Single Leg Stand; SB, Standing Balance; FAB, Fullerton Advanced Balance Scale; Dynamic balance: BBS, Berg Balance Scale; TUG, Timed Up and Go; FR, functional reach; AST, Alternative Step Test; Fear of Falling: Icon-FES, Iconographic Falls Efficacy Scale; FES-K, Falls Efficacy Scale-International; Fall risk: PPA, Physiological Profile Assessment; IG, intervention group; CG: control group; NWS group: Nintendo Wii Sports group; NSW BT group: balance training group; SUS: System Usability Scale; DART: Dynamic Acceptable Model for the Re-evaluation of Technologies.

Table 2 Summary of the quality of evidence and strength of recommendation

Quality assessment					
Meta-analysis	Study limitations*	Inconsistency†	Imprecision‡	Publication bias	Overall GRADE
Static balance	↓	↓		↓	Very low
Dynamic balance	↓	↓		↓	Very low
Fall risk		↓			Moderate
Fear of falling					High

*Risk of bias: we downgraded the evidence if >25% of included studies had a high risk of bias.
 †Heterogeneity >50%.
 ‡We downgraded if there were <400 participants.
 ↓, downgraded; GRADE, Grading of Recommendations, Assessment, Development and Evaluations.

(GRADE). Online supplemental figure 2 provides the Funnel plot for static and dynamic balance.

Effect of eHealth-delivered exercise programmes on secondary outcomes

Five studies (36%) measured fall risk using the Physiological Profile Assessment.^{18 36 38 39 43} The pooled effect of eHealth-delivered exercise programmes on fall risk indicates a small effect compared with control (SMD=0.32, 95% CI 0.00 to 0.64; I²=69.6%, p=0.048). The pooled results provide moderate-quality evidence (GRADE).

Four studies (29%) reported measures of fear of falling using the shortened Iconographical Falls Efficacy Scale,^{18 38 39} or the Fear of Falling Questionnaire.⁴¹ The pooled effect indicates no significant effect of eHealth-delivered exercise programmes on fear of falling compared with control (SMD=0.10, 95% CI -0.05 to 0.24; I²=0.0%, p=0.201). The pooled results provide high-quality evidence (GRADE).

Two studies collected data for fall rate.^{37 39} However, Lee *et al*³⁷ only reported a baseline measure for fall rate without providing further follow-up data. We were therefore unable to report on fall rate.

Adverse events

Eight included studies (57%) measured adverse events.^{35-41 46} Of those reported, no major adverse events were related to the intervention.

DISCUSSION

This systematic review and meta-analysis provides preliminary evidence that eHealth-delivered exercise programmes improve balance (static and dynamic) and provide an alternative method of delivering an exercise

programme to people aged 65 years or over living in the community. This reviews also demonstrates that an eHealth-delivered exercise programme may improve fall risk in people aged 65 years or over living in the community. However, we are uncertain whether an eHealth-delivered exercise programme improves fear of falling or fall rate in people aged 65 years or over living in the community.

There are a number of factors that may have influenced these results. First, the dose and intensity of the prescribed exercise in many of the studies may have been insufficient to substantially improve dynamic balance. A Cochrane review examining the effects of exercise interventions on balance found effective programmes were those attended three times per week for 3 months and involved dynamic exercises in a standing position.²⁹ While in 79% of studies (11/14) participants completed intervention exercises three times per week, only five studies had an exercise duration of 12 or more weeks. Furthermore, a systematic review on falls prevention




Figure 2 Risk of bias .




Figure 3 .

Forest plot: Effect size (95% CI) of e-Health interventions on static balance outcome by pooling data from 11 studies comparing e-Health versus control using random-effects meta-analysis (n = 1,056)




Figure 4

Forest plot: Effect size (95% CI) of e-Health interventions on dynamic balance outcome by pooling data from 14 studies comparing e-Health versus control using random-effects meta-analysis ($n = 1,180$)

found greater effects from interventions that challenged balance and included >3 hours per week.⁴⁹ Six studies (43%) included tailoring (increases in intensity and challenge). In addition, only four studies engaged participants in at least 180 min of exercise per week for the duration of the intervention. Most studies (71%) only engaged participants for between 30 and 120 min of exercise per week. This suggests that the challenge to balance may not have been of a sufficiently high dose. Therefore, our results, which indicate a small but statistically significant effect on dynamic balance, are promising. Further research is needed to explore the effect higher dosage (i.e., tailoring exercises to achieve increased intensity and challenge over the duration of the intervention) has on dynamic balance.

Finally, the tools used to measure dynamic balance may not be the most appropriate for the healthy older people. The most frequently used measure of dynamic balance was the Timed-up-and-go (TUG) (10/14). While the TUG is a validated tool and is recommended by the National Institute of Clinical Evidence for the assessment of gait and balance in the prevention of falls in older people,³ research has found the TUG may be more appropriate for frail older people who use walking aids rather than healthy older people.⁵⁰

The results for the secondary measures related to falls are mixed. While there was a small but significant effect on fall risk compared with the control, there is still uncertainty around the effect eHealth-delivered exercise programmes have on fear of falling. This is despite strong evidence that exercise interventions reduce falls in older community-dwellers.⁸ There are two possible explanations for these findings. First, to be eligible the included studies had to report on balance, resulting in only a small

number ($n=5$ fall risk; $n=4$ fear of falling) of studies also reporting a fall-related measure. This suggests we may have missed studies that measured the other fall-related outcomes of interest. Second, it is likely that the studies that reported a fall-related outcome, were not powered to detect an impact on falls (sample sizes ranged from 30 to 503 participants).²

Implications for clinicians and policymakers

Given the often low levels of adherence to exercise-based fall prevention programmes among older people, new delivery methods that improve access and encourage uptake of programmes designed to improve balance and reduce falls are needed. This review demonstrates that eHealth platforms are an effective mode of delivering exercises to improve balance. Although we identified an intervention effect on balance (static and dynamic), not only does this need to be considered in the context of the quality of the evidence but also in the ability to scale up and implement such interventions to large populations where resources are available. Clinicians should consider the use of eHealth platforms for delivering exercise programmes to older people living in the community.

Unanswered questions

Our results identified evidence of the benefit of eHealth-delivered exercise programmes on balance (static and dynamic) in older people living in the community. However, what remains unclear is the effect of eHealth-delivered exercise programmes on fall risk, fear of falling and fall rate in this population. Future research should focus on high-quality studies that deliver the recommended intensity and duration of exercise to provide a sufficiently high challenge to balance and impact on fall-related outcomes in a safe manner. Furthermore, future research needs to explore the long-term impact, cost-effectiveness and sustainability of eHealth-delivered programmes on balance and fall-related outcomes in older people living in the community.

Author affiliations

¹School of Health, Medical and Applied Sciences, CQUniversity, Rockhampton, Queensland, Australia

²Neuroscience Research Australia, Randwick, New South Wales, Australia

³Physical Activity Research Group, Appleton Institute, CQUniversity, Rockhampton, Queensland, Australia

⁴Institute for Musculoskeletal Health, The University of Sydney and Sydney Local Health District, Camperdown, New South Wales, Australia

⁵School of Public Health, Faculty of Medicine & Health, The University of Sydney, Sydney, New South Wales, Australia

⁶School of Public Health & Community Medicine, University of New South Wales, Sydney, New South Wales, Australia

Twitter Meghan Ambrens @meghan_ambrens and Corneel Vandelanotte @corneelvd

Acknowledgements We acknowledge the assistance of Kristie-Lee Alfrey (KLA) and Rachael Smallwood (RS) for helping to screen the articles.

Contributors MA, CV, AT, SA and KD were involved in the conception of the design of this systematic review, including the scope of the review; development of selection criteria; and development of the data analysis strategy. MA and QT were involved in the screening and data extraction. MA and JSO were involved with

the GRADE. MA, JSO and AT were involved in the data analysis. MA drafted the manuscript. MA, CV, AT, SA and KD were involved in all revisions and edits made to the manuscript. MA, CV, AT, SA, KD, QT and JSO were involved in the final approval of the version to be published. We acknowledge the assistance of KLA and RS for the screening of articles. MA and AT acted as guarantor.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient and public involvement Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

Patient consent for publication Not applicable.

Ethics approval Not applicable.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement All data relevant to the study are included in the article or uploaded as supplementary information. We agree to share any additionally relevant data that has not been included in the article or uploaded as supplementary information.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iDs

Meghan Ambrens <http://orcid.org/0000-0001-9117-7722>

Stephanie Alley <http://orcid.org/0000-0001-9666-5071>

Kim Delbaere <http://orcid.org/0000-0002-5655-0234>

REFERENCES

- Tinetti ME, Speechley M, Ginter SF. Risk factors for falls among elderly persons living in the community. *N Engl J Med* 1988;319:1701–7.
- Lamb SE, Jørstad-Stein EC, Hauer K, et al. Development of a common outcome data set for fall injury prevention trials: the prevention of falls network Europe consensus. *J Am Geriatr Soc* 2005;53:1618–22.
- National Institute for Health and Clinical Excellence [NICE]. *Falls in older people: assessing risk and prevention [CG161]*. United Kingdom: National Institute for Health and Clinical Excellence, 2013.
- Campbell AJ, Borrie MJ, Spears GF, et al. Circumstances and consequences of falls experienced by a community population 70 years and over during a prospective study. *Age Ageing* 1990;19:136–41.
- Australian Institute of Health & Welfare [AIHW]. Trends in hospitalised injury due to falls in older people 2007–08 to 2016–17. In: *Injury research and statistics series*. Australian Government: Australian Institute of Health & Welfare, 2019.
- Australian Institute of Health & Welfare [AIHW]. Injury, 2020. Available: <https://www.aihw.gov.au/reports-data/health-conditions-disability-deaths/injury/overview> [Accessed 31 Jul 2020].
- Owens PL, Russo CA, Spector W. *Healthcare cost and utilization (HCUP) statistical brief #80: Emergency department visits for injurious falls among the elderly*. Rockville, MD: Agency for Health Care Policy and Research, 2009.
- Sherrington C, Fairhall NJ, Wallbank GK, et al. Exercise for preventing falls in older people living in the community. *Cochrane Database Syst Rev* 2019;1:CD012424.
- Merom D, Pye V, Macniven R, et al. Prevalence and correlates of participation in fall prevention exercise/physical activity by older adults. *Prev Med* 2012;55:613–7.
- Bull FC, Al-Ansari SS, Biddle S, et al. World Health organization 2020 guidelines on physical activity and sedentary behaviour. *Br J Sports Med* 2020;54:1451–62.
- Alley SJ, Kolt GS, Duncan MJ, et al. The effectiveness of a web 2.0 physical activity intervention in older adults – a randomised controlled trial. *Int J Behav Nutr Phys Act* 2018;15:4.
- Mouton A, Cloes M. Efficacy of a web-based, Center-Based or combined physical activity intervention among older adults. *Health Educ Res* 2015;30:422–35.
- Bickmore TW, Silliman RA, Nelson K, et al. A randomized controlled trial of an automated exercise coach for older adults. *J Am Geriatr Soc* 2013;61:1676–83.
- Wijman CA, Westendorp RG, Verhagen EA, et al. Effects of a web-based intervention on physical activity and metabolism in older adults: randomized controlled trial. *J Med Internet Res* 2013;15:e233.
- Irvine AB, Gelatt VA, Seeley JR, et al. Web-Based intervention to promote physical activity by sedentary older adults: randomized controlled trial. *J Med Internet Res* 2013;15:e19.
- Ammann R, Vandelandotte C, de Vries H, et al. Can a website-delivered computer-tailored physical activity intervention be acceptable, usable, and effective for older people? *Health Educ Behav* 2013;40:160–70.
- Valenzuela T, Okubo Y, Woodbury A, et al. Adherence to technology-based exercise programs in older adults: a systematic review. *J Geriatr Phys Ther* 2018;41:49–61.
- Schoene D, Lord SR, Delbaere K, et al. A randomized controlled pilot study of home-based step training in older people using videogame technology. *PLoS One* 2013;8:e57734.
- Chao Y-Y, Scherer YK, Montgomery CA. Effects of using Nintendo Wii™ exergames in older adults: a review of the literature. *J Aging Health* 2015;27:379–402.
- Skjøret N, Nawaz A, Morat T, et al. Exercise and rehabilitation delivered through exergames in older adults: an integrative review of technologies, safety and efficacy. *Int J Med Inform* 2016;85:1–16.
- Pirovano M, Surer E, Mainetti R, et al. Exergaming and rehabilitation: a methodology for the design of effective and safe therapeutic exergames. *Entertain Comput* 2016;14:55–65.
- Holden MK, Todorov E. Use of virtual environments in motor learning and rehabilitation. In: Stanley KM, ed. *Handbook of virtual environments: design, implementation, and applications*. New Jersey: Lawrence Erlbaum Associates, 2002.
- Donath L, Rössler R, Faude O. Effects of virtual reality training (exergaming) compared to alternative exercise training and passive control on standing balance and functional mobility in healthy community-dwelling seniors: a meta-analytical review. *Sports Med* 2016;46:1293–309.
- Neri SG, Cardoso JR, Cruz L, et al. Do virtual reality games improve mobility skills and balance measurements in community-dwelling older adults? systematic review and meta-analysis. *Clin Rehabil* 2017;31:1292–304.
- Pietrzak E, Cotea C, Pullman S. Using commercial video games for falls prevention in older adults: the way for the future? *J Geriatr Phys Ther* 2014;37:166–77.
- Laufer Y, Dar G, Kodesh E. Does a Wii-based exercise program enhance balance control of independently functioning older adults? A systematic review. *Clin Interv Aging* 2014;9:9.
- Liberati A, Altman DG, Tetzlaff J, et al. The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate healthcare interventions: explanation and elaboration. *BMJ* 2009;339:b2700.
- Ambrens M, Tiedemann A, Delbaere K, et al. The effect of eHealth-based falls prevention programmes on balance in people aged 65 years and over living in the community: protocol for a systematic review of randomised controlled trials. *BMJ Open* 2020;10:e031200.
- Howe TE, Rochester L, Neil F, et al. Exercise for improving balance in older people. *Cochrane Database Syst Rev* 2011:CD004963.
- Winter DA. *A.B.C. (anatomy, biomechanics and control) of balance during standing and walking*. Waterloo Biomechanics, 1995.
- Physiotherapy Evidence Database [PEDro]. PEDro scale, 2020. Available: <https://pedro.org.au/english/resources/pedro-scale/2020>
- Maher CG, Sherrington C, Herbert RD, et al. Reliability of the PEDro scale for rating quality of randomized controlled trials. *Phys Ther* 2003;83:713–21.
- Cochrane Collaboration. The Cochrane Handbook for systematic reviews of interventions, 2020. Available: <https://methods.cochrane.org/bias/resources/cochrane-handbook> [Accessed cited 2020 November 20, 2020November 20, 2020].
- Hillsdale NJ, Cohen J. *Statistical power analysis for the behavioral sciences*. 2nd ed. New York, New York: Lawrence Erlbaum Associates, 1988.



- 35 Schoene D, Valenzuela T, Toson B, *et al.* Interactive cognitive-motor step training improves cognitive risk factors of falling in older adults - A randomized controlled trial. *PLoS One* 2015;10:e0145161.
- 36 Gschwind YJ, Schoene D, Lord SR, *et al.* The effect of sensor-based exercise at home on functional performance associated with fall risk in older people - a comparison of two exergame interventions. *Eur Rev Aging Phys Act* 2015;12:11.
- 37 Lee Y, Choi W, Lee K, *et al.* Virtual reality training with three-dimensional video games improves postural balance and lower extremity strength in community-dwelling older adults. *J Aging Phys Act* 2017;25:621-7.
- 38 Gschwind YJ, Eichberg S, Ejupi A, *et al.* ICT-based system to predict and prevent falls (iStoppFalls): results from an international multicenter randomized controlled trial. *Eur Rev Aging Phys Act* 2015;12.
- 39 Delbaere K, Valenzuela T, Lord SR, *et al.* E-Health StandingTall balance exercise for fall prevention in older people: results of a two year randomised controlled trial. *BMJ* 2021;373:n740.
- 40 Whyatt C, Merriman NA, Young WR, *et al.* A Wii bit of fun: a novel platform to deliver effective balance training to older adults. *Games Health J* 2015;4:423-33.
- 41 Hong J, Kong H-J, Yoon H-J. Web-Based telepresence exercise program for community-dwelling elderly women with a high risk of falling: randomized controlled trial. *JMIR Mhealth Uhealth* 2018;6:e132.
- 42 Bieryla KA, Dold NM. Feasibility of Wii fit training to improve clinical measures of balance in older adults. *Clin Interv Aging* 2013;8:775-81.
- 43 Phirom K, Kamnardsiri T, Sungkarat S. Beneficial effects of interactive Physical-Cognitive Game-Based training on fall risk and cognitive performance of older adults. *Int J Environ Res Public Health* 2020;17. doi:10.3390/ijerph17176079. [Epub ahead of print: 21 08 2020].
- 44 Sato K, Kuroki K, Saiki S, *et al.* Improving walking, muscle strength, and balance in the elderly with an exergame using Kinect: a randomized controlled trial. *Games Health J* 2015;4:161-7.
- 45 Jung D-I, Ko D-S, Jeong M-A. Kinematic effect of Nintendo Wii(TM) sports program exercise on obstacle gait in elderly women with falling risk. *J Phys Ther Sci* 2015;27:1397-400.
- 46 Chow DHK, Mann SKF. Effect of cyber-golfing on balance amongst the elderly in Hong Kong: a pilot randomised trial. *Hong Kong J Occup Ther* 2015;26:9-13.
- 47 Lai C-H, Peng C-W, Chen Y-L, *et al.* Effects of interactive video-game based system exercise on the balance of the elderly. *Gait Posture* 2013;37:511-5.
- 48 Sadeghi H, Jehu DA, Daneshjoo A, *et al.* Effects of 8 weeks of balance training, virtual reality training, and combined exercise on lower limb muscle strength, balance, and functional mobility among older men: a randomized controlled trial. *Sports Health* 2021;13:606-12.
- 49 Sherrington C, Michaleff ZA, Fairhall N, *et al.* Exercise to prevent falls in older adults: an updated systematic review and meta-analysis. *Br J Sports Med* 2017;51:1750-8.
- 50 Lin M-R, Hwang H-F, Hu M-H, *et al.* Psychometric comparisons of the timed up and go, one-leg stand, functional reach, and Tinetti balance measures in community-dwelling older people. *J Am Geriatr Soc* 2004;52:1343-8.