



ORIGINAL ARTICLE

A clinical, histopathological and laboratory study of 19 consecutive Italian paediatric patients with chilblain-like lesions: lights and shadows on the relationship with COVID-19 infection

M. El Hachem,¹ A. Diociaiuti,^{1,*,†}  C. Concato,^{2,*†} R. Carsetti,³ C. Carnevale,¹  M. Ciofi Degli Atti,⁴ L. Giovannelli,⁵ E. Latella,¹ O. Porzio,^{6,7} S. Rossi,⁸ A. Stracuzzi,⁸ S. Zaffina,⁹ A. Onetti Muda,¹⁰ G. Zambruno,¹¹ R. Alaggio⁸

¹Dermatology Unit, Bambino Gesù Children's Hospital, IRCCS, Rome, Italy

²Virology Unit, Bambino Gesù Children's Hospital, IRCCS, Rome, Italy

³Diagnostic Immunology Unit, Department of Laboratories, B Cell Pathophysiology Unit, Immunology Research Area, Bambino Gesù Children's Hospital, IRCCS, Rome, Italy

⁴Clinical Epidemiology Unit, Bambino Gesù Children's Hospital, IRCCS, Rome, Italy

⁵Medical Laboratory Unit, Bambino Gesù Children's Hospital, IRCCS, Rome, Italy

⁶Medical Laboratory Unit, Bambino Gesù Children's Hospital, IRCCS, Rome, Italy

⁷Department of Experimental Medicine, University of Rome Tor Vergata, Rome, Italy

⁸Pathology Unit, Bambino Gesù Children's Hospital, IRCCS, Rome, Italy

⁹Occupational Medicine, Bambino Gesù Children's Hospital, IRCCS, Rome, Italy

¹⁰Department of Laboratories, Bambino Gesù Children's Hospital, IRCCS, Rome, Italy

¹¹Genetics and Rare Diseases Research Division, Bambino Gesù Children's Hospital, IRCCS, Rome, Italy

*Correspondence: A. Diociaiuti. E-mail: andrea.diociaiuti@opbg.net

Abstract

Background Acral chilblain-like lesions are being increasingly reported during COVID-19 pandemic. However, only few patients proved positivity for SARS-CoV-2 infection. The relationship between this skin manifestation and COVID-19 infection has not been clarified yet.

Objective To thoroughly characterize a prospective group of patients with chilblain-like lesions and to investigate the possible relationship with SARS-CoV-2 infection.

Methods Following informed consent, patients underwent (i) clinical evaluation, (ii) RT-PCR and serology testing for SARS-CoV-2, (iii) digital videocapillaroscopy of finger and toe nailfolds, (iv) blood testing to screen for autoimmune diseases and coagulation anomalies, and (v) skin biopsy for histopathology, direct immunofluorescence and, in selected cases, electron microscopy.

Results Nineteen patients, all adolescents (mean age: 14 years), were recruited. 11/19 (58%) of them and/or their cohabitants reported flu-like symptoms one to two months prior to skin manifestation onset. Lesions were localized to toes and also heels and soles. Videocapillaroscopy showed pericapillary oedema, dilated and abnormal capillaries, and microhaemorrhages both in finger and toe in the majority of patients. Major pathological findings included epidermal basal layer vacuolation, papillary dermis oedema and erythrocyte extravasation, perivascular and perieccrine dermal lymphocytic infiltrate, and mucin deposition in the dermis and hypodermis; dermal vessel thrombi were observed in two cases. Blood examinations were normal. Nasopharyngeal swab for SARS-CoV-2 and IgG serology for SARS-CoV-2 nucleocapsid protein were negative. Importantly, IgA serology for S1 domain of SARS-CoV-2 spike protein was positive in 6 patients and borderline in 3.

Conclusions Chilblain-like lesions during COVID-19 pandemic have specific epidemiologic, clinical, capillaroscopic and histopathological characteristics, which distinguish them from idiopathic perniosis. Though we could not formally prove SARS-CoV-2 infection in our patients, history data and the detection of anti-SARS-CoV-2 IgA strongly suggest a relationship between skin lesions and COVID-19. Further investigations on the mechanisms of SARS-CoV-2 infection in children and pathogenesis of chilblain-like lesions are warranted.

Received: 11 May 2020; Accepted: 18 May 2020

[†]Equal contribution.

Conflict of interest

None to be reported for all authors.

Funding sources

None declared.

Introduction

Chilblains, also called pernio, are an inflammatory skin manifestation characterized by erythema and swelling of the extremities, lasting more than 24 h.^{1,2} The disease presents following exposure to cold humid weather, usually during winter or early spring. It affects more frequently young and middle-aged women. Chilblains may be idiopathic or, less commonly, occur in association with an underlying autoimmune connective tissue disease, in particular lupus erythematosus, haematologic disorders, viral hepatitis and malignancy. Chilblain lesions are oedematous, erythematous to violaceous macules, papules, plaques, or nodules on the dorsal fingers or toes, and less frequently, on the nose, ears and foot soles. Erosions or ulcerations may develop. Symptoms comprise burning, tenderness or pruritus. Idiopathic chilblains usually run a benign course with spontaneous resolution in 1–3 weeks; however, annual recurrences are frequent.

COVID-19 pandemic is caused by a novel coronavirus, SARS-CoV-2, first isolated in Chinese patients with severe pneumonia and rapidly spreading all over the world.^{3,4} COVID-19 infection manifests with fever, headache, dry cough, fatigue, dysgeusia, anosmia and breath shortness, which can progress to acute respiratory distress syndrome due to severe bilateral pneumonia. A certain proportion of patients, in particular children, are asymptomatic.^{5,6} Recently, skin manifestations have been described in COVID-19 patients.^{7–9} In addition to vesicular or maculopapular skin rashes, livedoid/necrotic lesions and urticaria, chilblains-like lesions are increasingly reported in particular in adolescents and young adults.^{8–22} We also observed from last March an unusual number of patients, mainly teenagers, affected with chilblains-like lesions. Thus, we decided to investigate the aetiology of these lesions and possible relationship with COVID-19 infection in our patients.

We report the clinical, capillaroscopic, histopathological, ultrastructural and laboratory findings in a prospective paediatric series of patients presenting with chilblains-like lesions during the COVID-19 pandemic.

Materials and methods

Study design and population

This study has been designed as a monocentre prospective study of paediatric patients presenting with chilblains-like lesions.

Patients were screened and recruited in the second half of April at the Dermatological Unit of Bambino Gesù Children's Hospital, Rome, Italy.

Consecutive patients, with a clinical diagnosis of chilblain-like lesions established by a dermatologist, and giving specific informed consent, were included in the study. Exclusion criteria were (i) patients aged less than 6 years and (ii) patients affected with a diagnosed autoimmune disease, in addition to (iii) patients or their legal guardians who refused to give the consent to participate to the study. The study was approved by our Institutional Ethical Committee and conducted in accordance with the Declaration of Helsinki.

Clinical evaluation

A standardized case report form was created. Information on patient demographics, past and recent family and personal history, clinical features (lesion localization and type), symptoms and follow-up was collected. In addition, results of different investigations performed (as detailed below) were recorded. Finally, clinical pictures were taken in all patients.

Study procedures

Patients recruited in the study underwent (i) testing for SARS-CoV-2 on nasopharyngeal swab and, in selected cases, skin biopsy, (ii) digital videocapillaroscopy of finger and toe nail-folds, (iii) blood testing including serology for SARS-CoV-2 and (iv) lesional skin biopsy for histopathology, direct immunofluorescence and, in 4 patients, transmission electron microscopy examination.

SARS-CoV-2 testing

'STARMag 96 × 4 Universal Cartridge kit' (Seegene Inc., South Korea) was used for extraction and isolation of nucleic acids from nasopharyngeal swabs and biopsy samples, followed by multiplex real-time reverse transcriptase (RT)-PCR assay for simultaneous detection of 3 target genes of SARS-CoV-2, RdRP and N genes specific for SARS-CoV-2, and E gene for all of sarbecovirus including SARS-CoV-2 ('Allplex™ 2019-nCoV Assay', Seegene Inc.). In addition, serology for SARS-CoV-2 was performed using a chemiluminescent microparticle immunoassay (CMIA) for detection of IgG antibodies directed against the nucleocapsid protein of SARS-CoV-2 (Abbott Laboratories, USA). In parallel, all sera were tested using an anti-SARS-CoV-2

ELISA (Euroimmun, Germany) for the detection of IgG and IgA antibodies against the S1 domain of the spike protein including the immunologically relevant receptor binding domain.

Digital videocapillaroscopy

It was performed on all recruited patients with VideoCAP® 3.0 (DS Medica, s.r.l., Italy), according to standard procedures.²³ Images were collected at 100× and 200× magnification. The following characteristics were evaluated: capillary density and architecture, pericapillary oedema, capillary morphology, capillary dimension (width of the apical limb of the capillary) and presence of avascular areas and microhaemorrhages.²⁴ Capillaroscopy images were independently reviewed by three dermatologists.

Blood testing

All patients underwent the following investigations: real-time PCR for parvovirus B19, complete blood cell count, erythrocyte sedimentation rate, liver enzymes, creatinine, cryoglobulins, ANCA, antinuclear, anti-dsDNA, antiphospholipid and anticardiolipin, ENA and ribonucleic protein antibodies, C3, C4, immunoglobulins (IgG, IgM, IgA) and coagulation screening (D-dimers, prothrombin time, activated partial thromboplastin time, fibrinogen).

Histopathology, immunohistochemistry and direct immunofluorescence

Haematoxylin–eosin-stained sections of formalin-fixed paraffin-embedded tissue were reviewed under light microscopy. Three pathologists analysed all cases independently. Histopathological features assessed in all cases included: apoptotic keratinocytes, spongiosis, exocytosis, vacuolar changes in the basal layer, inflammatory infiltrate (cell type, intensity and location), papillary dermal oedema, vascular damage, and dermal fibrin and mucin deposition. The intensity of the lymphocytic infiltrate was graded 1–3, according to previous studies.²⁵ Immunostains for CD3 (polyclonal, Dako Agilent), CD20 (clone L26, Dako Agilent), CD123 (clone BR4MS, Leica Novocastra) and Alcian blue stain were performed according to standard methods. Direct immunofluorescence (DIF) was performed using fluorescein isothiocyanate-conjugated anti-human IgG, IgM, IgA, fibrinogen, and complement fractions, C1q, C3c and C4c (all from Dako).

Transmission electron microscopy (TEM)

Skin biopsy specimens were fixed in 2% glutaraldehyde, post-fixed in 1% osmium tetroxide, dehydrated in graded alcohols, embedded in EPON resin and sectioned on an ultramicrotome (Ultracut S, Reichert-Jung, Austria). Ultrathin sections were stained with uranyl acetate and lead citrate and observed with a transmission electron microscope (JEM-1400, JEOL, Japan).

Results

Clinical features

The parents of nineteen out of twenty-one patients screened for chilblain-like lesions signed informed consent agreeing to the participation of their child in the study. Patient demographics, history and clinical findings are listed in Table 1. All patients were Italian and lived in Lazio Region (Central Italy). There was a predominance of male patients (M:F = 2.8: 1); the mean age was 14.0 years (range, 11–17). Two patients were siblings. Patient history was negative for perniosis, autoimmune, neoplastic or coagulation disorders in all cases. Five patients (26%) reported fever and cough 1–2 months prior to development of skin lesions, and one patient presented diarrhoea one week after skin manifestation onset. In addition, two patients recently stayed in areas with high prevalence of COVID-19 infection or hosted guests from these areas. In nine cases (47%), family history was positive for fever, headache, cough or gastrointestinal symptoms in one or more family members, 1–2 months prior to skin lesion onset. All patients were in good general health at the time of the first consultation, and they had respected home confinement since 9 March, according to the Italian COVID-19 pandemic regulations. Interestingly, 16 patients (84%) reported barefoot walking during confinement (5 with socks). Skin lesions had appeared 12–40 days (mean, 22.2 days) before screening visit, presenting at first as toe swelling and erythema. On physical examination, lesions were localized exclusively to feet in all cases, they were bilateral and predominantly distributed to the distal toes, followed by heels and lateral foot margin, and rarely spread to the soles (Fig. 1 and 2). All patients had toe swelling and erythema (Figs 1a,b,d and 2a), accompanied by erythematous-violaceous roundish macules and purpuric lesions in 12 cases (63%) (Figs 1c and 2b–d). In addition, some patients presented also pustules (Fig. 2a), and erosions covered by crusts (Figs 1d and 2a,c). Eleven patients (57.8%) reported pain and/or itching concomitant to lesion onset and resolving spontaneously within 7–10 days. No one required analgesic or specific therapy. At follow-up (14 days after screening), patients still presented asymptomatic minimal erythema, swelling, brownish macules and/or crusts.

Digital videocapillaroscopy findings

Capillaroscopy was performed on both finger and toe nailfolds (Table 2). On fingers, the capillary density and architecture were normal in all patients. Pericapillary oedema was present in 11 out of 19 cases (Fig. 3a,c). Capillary dimension was abnormal (>20 µm) in 7 patients (Fig. 3c). Abnormal capillary shape was the most frequent feature (14 cases, 73.7%; Fig. 3c). Microhaemorrhages were detected in 5 patients (Fig. 3a). Given the particular anatomical nailfold conformation on feet, the first and second toes were mainly examined. Capillary density was normal

Table 1 Demographics, history and clinical features in 19 patients with chilblain-like lesions

Pt.	Age (years)	Sex	Recent family history [†]	Recent personal history [†]	Lesion localization	Lesion type	Symptoms
1	13	F	Parents and 2 siblings: headache, fever, abdominal pain, one month before	Fever, headache, sore throat, one month before	Toes	Erythema, swelling [‡]	Intense pain
2	13	F	Negative	Negative	Toes	Erythema, swelling, purpuric macules, crusts	Intense pain
3	12	M	Parents: sore throat and fever, two months before	Fever, two months before	Toes, heels	Erythema, swelling, purpuric macules	Intense itching
4	15	M	Mother: cough and fatigue, one month before	Negative	Toes, heels, soles	Erythema, swelling, violaceous macules	No
5	14	M	Negative	Negative	Toes	Erythema, swelling	Itching
6	14	M	Negative	Negative	Toes, heels, soles	Erythema, purpuric macules, papules, crusts	Itching
7	13	F	Negative	Negative	Toes	Erythema, swelling, purpuric macules, crusts	Intense itching
8	15	M	Grandfather [§] fever and cough, one month before	Negative; travel to Milan, 1.5 month before	Toes, heel, soles	Erythema, swelling, purpuric macules, pustules, crusts	Pain, itching
9	17	M	Negative	Negative, hosting a friend from Northern Italy, 2 months before	Toes, heels, soles	Erythema, crusts	Pain, itching
10 [¶]	13	M	Negative	sore throat, fever, diarrhoea, 1.5 month before	Toes, heels	Erythema, purpuric macules, pustule	Burning, itching
11 [¶]	16	M	Brother: sore throat, fever, diarrhoea, 1.5 month before	Negative	Toes	Erythema, swelling	Asymptomatic
12	15	M	Negative	Negative	Toes, heels, soles	Erythema, swelling, purpuric macules, crusts	Asymptomatic
13	17	M	Negative	Negative	Toes, heels, soles	Erythema, swelling, purpuric macules, pustule, crusts	Asymptomatic
14	17	M	Father and his partner: long-lasting fever and cough, one month before	Negative	Toes, heels, soles	Erythema, swelling, purpuric macules, crusts	Asymptomatic
15	12	F	Father: fever and cough started 1.5 month before	Negative	Toes	Erythema, swelling, purpuric macules, crusts	Intense pain
16	12	M	Negative	Negative	Toes	Erythema, swelling	Asymptomatic
17	11	F	Negative	Fever, 2 months before	Toes	Swelling, purpuric macules, crusts	Asymptomatic
18	14	M	Negative	Diarrhoea one week after skin lesion onset	Toes	Erythema, swelling, crusts	Burning
19	14	M	Negative	Fever and cough, two months before	Toes, heel, soles	Erythema, swelling, purpuric macules, crusts	Asymptomatic

[†]Family and personal history timing are referred to the onset of skin manifestation.

[‡]Swelling was limited to toes in all patients.

[§]Patient cohabitant.

[¶]Patients 10 and 11 are brothers.

in almost all analysed toes. Pericapillary oedema was less frequently detectable in toes than in fingers (5 vs. 11 patients) (Fig. 3b,d), while toenail capillary dilation was present in 12 patients as observed in fingers (Fig. 3d). About half of the patients showed abnormal morphology of capillaries (Fig. 3b,d). The most striking capillaroscopic sign on toes were microhaemorrhages, detected in 7 out of 19 patients (Fig. 3d).

Histopathology, direct immunofluorescence and ultrastructural findings

Results from histopathological analysis of skin biopsies obtained from 18 patients are summarized in Table 3.

Oedema of the papillary dermis was observed in 12 cases, being more prominent in 3 (Fig. 4c). Red cell extravasation was present in almost all patients (15, 88.2%). In all cases, a mild



Figure 1 Clinical features. Erythema and swelling of right foot toes (a); erythema and swelling, particularly marked on the fourth right toe (b); purpuric macules, more evident on the second toe of both feet (c); erythema, mild swelling and multiple crusts on left toes (d).



Figure 2 Clinical features. Diffuse swelling, erythema and crusts on the left foot, and two pustules (arrows) on the big toes (a); purpuric macules on the lateral aspect of the right heel (b); confluent purpuric macules mostly covered by crusts on both heels and posterior ankles (c); brownish purpuric macules on the soles (d).

(12/18 cases) to moderate/intense (6 cases) perivascular lymphocytic infiltrate was seen in the superficial and deep dermis where it also localized around the eccrine glands (Fig. 4a–e). A perivascular infiltrate was detected also in the hypodermis in all biopsies containing subcutaneous tissue (11/11) (Fig. 4a,e). Small vessels showed swollen endothelia in 15 cases. Intriguingly, signs of lymphocytic vasculitis with a prominent lymphocytic infiltrate and isolated neutrophils were observed in 3 cases, with involvement of a deep arteriole (Fig. 4b) and a venule in one

Table 2 Capillaroscopic features in 19 patients with chilblain-like lesions

Capillaroscopic characteristics	Patient number (%)	
	Fingers	Toes
Capillary density		
Normal	19 (100%)	18 (94.7%)
Reduced (<7 capillaries/mm)	0 (0%)	1 (5.3%)
Pericapillary oedema		
Absent	8 (42.1%)	14 (73.7%)
Present	11 (57.9%)	5 (26.3%)
Capillary dimension		
Normal (<20 µm)	12 (63.2%)	12 (63.2%)
Abnormal (≥20 µm)	7 (36.8%)	7 (36.8%)
Morphology		
Normal	5 (26.3%)	9 (47.4%)
Abnormal	14 (73.7%)	10 (52.6%)
Microhaemorrhages		
Absent	15 (79%)	12 (63.2%)
Present	4 (21%)	7 (36.8%)




Figure 3 Videocapillaroscopy findings. Multiple and synchronous microhaemorrhages (asterisks), pericapillary oedema (arrows) on the finger nailfold (a); abnormal morphology of the capillaries (Black circle) and pericapillary oedema (arrows) on the toe nailfold of the same patient (b). Pericapillary oedema (arrows), dilated capillaries (Black triangle) and abnormal capillary morphology (Black circle) on the finger nailfold (c); numerous and prominent microhaemorrhages (asterisks), marked abnormal capillary morphology (circles) and pericapillary oedema (arrows) on the toe nailfold of the same patient (d).

case each. Fibrin thrombi were found in 2 patients (Fig. 4d). There was no evidence of fibrinoid necrosis in the vascular walls. Epidermis showed mild changes characterized by spongiosis (13

Table 3 Pathological features in 19 patients with chilblain-like lesions

Pathological features	Patient N (%)
Papillary dermis oedema	12/18 (67%)
Extravasated erythrocytes	15/18 (83%)
Perivascular and perieccrine lymphocytic infiltrate	
Mild	12/18 (67%)
Moderate/marked	6/18 (33%)
Involving subcutis	11/11 (100%)
Lymphocytic vasculitis	3/18 (17%)
Fibrin thrombi	2/18 (11%)
Spongiosis	13/18 (72%)
Basal epidermal layer vacuolation	14/18 (78%)
Exocytosis	6/18 (33%)
Apoptotic keratinocytes	4/18 (22%)
Mucin	
Minimal	10/18 (55%)
Prominent	7/18 (39%)

N, number.

cases), slight vacuolation of basal layer (14 cases; Fig. 4c), with lymphocyte exocytosis in 6 cases, and isolated apoptotic keratinocytes in 4. Alcian stain highlighted the presence of minimal mucin deposits mostly in the deep dermis in 10 cases, while in 7 it was more prominent with a diffuse involvement of subcutis (Fig. 4f). Immunostains showed a uniform phenotype in all cases, with a predominance of T-lymphocytes, CD3 positive (more than 95%), associated with scattered CD123 positive macrophages. Direct immunofluorescence performed on biopsies from 11 patients showed focal granular deposition of C3 in the walls of a few isolated dermal vessels (9 cases; Fig. 5a,b). In addition, colloid bodies along the dermal–epidermal junction stained positive for IgM (4 cases; Fig. 5c) and IgG (2 cases). TEM examination of lesional skin showed similar findings in the four patients examined. Several extravasated erythrocytes were present in the papillary dermis, which appeared diffusely oedematous (Fig. 6a). A dermal infiltrate chiefly composed of small to medium size lymphocytes with a few admixed macrophages was observed in a perivascular location but also scattered throughout the dermis (Fig. 6b). Dermal vessels frequently showed protruding endothelial cells with prominent intraluminal nuclei. In addition, some endothelial cells presented degenerative changes, with nuclear and cytoplasmic condensation and areas of denuded or interrupted basement membrane were observed (Fig. 6c,d).

SARS-CoV-2 testing results

Real-time PCR for SARS-CoV-2 on nasopharyngeal swabs from 19 patients and on skin biopsies from 3 patients proved always negative. In a similar way, serology for IgG antibodies directed against the nucleocapsid protein of SARS-CoV-2 was negative in all patients. Of note, serology for the detection of IgG and IgA




Figure 4 Histopathological findings. Representative low-power magnification of a punch skin biopsy showing perivascular inflammatory infiltrate in the superficial and deep dermis and subcutaneous tissue (a); higher magnification of the arteriole indicated with an arrow in (a) shows an intramural lymphocytic infiltrate (b); a scale-crust, epidermal spongiosis, basal layer smudging (arrow) and oedema of papillary dermis with extravasated erythrocytes are evident in (c); a dermal capillary blood vessel with intraluminal thrombus is visible in (d); dense perieccrine and perineural dermal inflammatory infiltrate associated with mucin deposits (e) highlighted by Alcian blue staining shown in (f). a–e: Haematoxylin–eosin staining; bars: 500 μ m in a; 50 μ m in b and d; 100 μ m in c, e and f.

antibodies against the S1 domain of the spike protein revealed: (i) positive IgG in one patient (n. 7 of Table 1), and borderline in 3 patients (n. 10, 13 and 14), and (ii) positive IgA in 6 patients (n. 5, 9, 12, 13, 14 and 16), and borderline in 3 patients (n. 8, 10 and 19). Finally, all other blood tests performed were within normal limits in all patients.

Discussion

Chilblain-like lesions are being increasingly reported in different European and non-European countries during COVID-19 pandemic.^{8–22} Indeed, first case series on skin manifestations during COVID-19 outbreak indicate that acral lesions are quite frequent: they were present in 106 of 277 dermatological outpatients (38.3%) in a French retrospective study²² and in 79 of 375 cases (21%) in a Spanish prospective survey.⁹ Therefore, we decided to undertake a systematic study of clinical,




Figure 5 Direct immunofluorescence findings. Granular deposits of C3 in the wall of vessels (asterisk) in the papillary and deep dermis (a and b, respectively) and focal deposits along the dermal–epidermal junction; colloid bodies at the dermal–epidermal junction staining positive for IgM (c). Bars: 50 µm in a and b; 100 µm in c.

capillaroscopic, histopathological and SARS-CoV-2 infection in paediatric patients referred to us for this peculiar skin manifestation, in order to investigate a possible relationship with COVID-19 infection.

We recruited 19 consecutive cases during a two-week recruitment period in the second half of April. Interestingly, all our patients were adolescents (mean age: 14 years). Most previously published case series comprised both paediatric and adult patients making difficult a direct comparison.^{9,14,15,17} Nevertheless, these studies reported a patient mean age ranging from 16 to 32.5 years. In addition, the first Italian case series included 63 patients, all adolescents, and a Spanish study on 27 paediatric patients reported a mean age of 14.44 years.^{12,19} Altogether, these data indicate a prevalence of this peculiar manifestation in teenagers. In our series, males were more numerous than females (2.8 : 1), which was not confirmed in other case series showing about 1 : 1 sex ratio^{12,14,15,19,22} and female predominance in a single study.⁹ In our patients, lesions were exclusively localized to feet, which represents the most frequently affected site also in previous reports, with 108 of 132 (81.8%), 37 of 41 (90%), 21 of 27 (78%) and 58 of 63 cases (92.7%) in different series.^{12,14,15,19} In a minority of patients, hands alone were affected. Interestingly, 9 of our patients also had lesions on soles and heels, in addition to toes. Similar to previous reports, clinical manifestations in our patients comprised swelling, erythema, purpuric macules and papules, pustules and crusts. In our cases, cutaneous manifestations lasted more than one month and no relapses were observed. Although shorter disease duration has been reported in some cases series,^{9,15} all reports agree on spontaneous resolution without sequelae. Overall, the demographic (similar sex ratio), epidemiological (onset with cool to warm temperatures and recent great increase in incidence) and clinical (involvement of soles and heels) features of chilblain-like lesions during the COVID pandemic distinguish them from idiopathic perniosis. Furthermore, in keeping with literature data, laboratory findings, including autoimmunity and coagulation screening, were negative or within norm in our patients, excluding connective tissue diseases.

We performed for the first time a videocapillaroscopy characterization of chilblain-like lesions showing the presence of pericapillary oedema, ectatic and abnormal capillaries, and microhaemorrhages in a significant proportion of cases. Of note, our patients presented anomalies both on hands and feet, even though skin lesions were limited to feet in all cases. Thus, this finding suggests that chilblain-like lesions represent




Figure 6 Ultrastructural features. Extravasated red blood cells (R) are visible in an oedematous (asterisks) papillary dermis (a, 'F' denotes two fibroblasts); a dermal infiltrate chiefly composed of small to medium size lymphocytes (L) is observed in the papillary dermis (b); protruding endothelial cells with prominent intraluminal nuclei, one with condensed chromatin (asterisk in c), line two dermal vessels (c and d), one showing partly denuded and interrupted basement membrane (asterisk in d). Bars: 10 μ m in (a) and (b), 5 μ m in (c) and (d).

manifestations of a systemic process. Although limited data are available about the physiopathologic morphology of toe capillaries, the values of the major capillaroscopic parameters, such as capillary size and density do not differ significantly from fingers.²⁶ In our series, the same percentage of patients (36.8%) presented dilated capillaries on hands and feet, while pericapillary oedema and microhaemorrhages were more common in toe than finger nailfolds (14 vs. 8 and 7 vs. 4 cases, respectively). Altogether, these features appear more severe than those described in idiopathic pernio where microhaemorrhages were not detected.²⁷ Indeed, our capillaroscopy findings well match the clinical features of erythema, oedema, and purpuric lesions and the histological ones of dilated capillaries, superficial dermal oedema, with extravasated red blood cells observed in our patients. Histopathological examination showed a perivascular, superficial and deep T-lymphocytic infiltrate and a combination

of other findings considered more characteristic of either idiopathic chilblains or chilblain lupus erythematosus. The morphologic overlap between these two entities has been extensively investigated.^{25,28,29} Interface dermatitis with basal layer vacuolation and necrotic keratinocytes together with a perivascular dermal infiltrate without periecrine distribution has been considered markers of chilblain lupus. By contrast, periecrine inflammatory infiltrate and papillary dermis oedema/spongiosis are frequent findings in chilblains.^{25,28} Our cases showed hybrid features: (i) a periecrine infiltrate in all biopsies and papillary oedema/spongiosis in about 70% of the cases, more in keeping with chilblains, and (ii) basal membrane smudging and mucin deposition, both detected in the large majority of the cases, are more reminiscent of lupus.^{25,27,28} Very few chilblain-like lesions with potential COVID-19 infection association have been histologically investigated to date.^{18,20–22} Similar to our findings, a

lymphocytic perivascular and periadnexal infiltrate, basal layer vacuolation and, in 3 biopsies, microthrombi were observed.^{18,20,22}

Despite increasing reports on chilblain-like lesions during COVID-19 pandemic, the possible relationship between this skin manifestation and SARS-CoV-2 infection remains matter of debate. All our patients were in good general health at the time of first examination and, except for an episode of diarrhoea in a single patient, did not develop any extracutaneous symptom during the follow-up period. However, the majority of them and/or their cohabitants (11/19, 58%) reported flu-like symptoms one to two months prior to skin lesion onset. Previous reports varied widely as to the presence of suspected and/or confirmed symptomatic COVID-19 infection in patients with chilblain-like lesions. Galván Casas *et al.* reported chilblain-like lesions in 71 patients with either confirmed COVID-19 infection (41%) or suggestive symptoms (59%).⁹ However, the percentage of patients with extracutaneous symptoms was much lower in other series ranging from 0 % to 44.4%.^{14,15,17,19,22} To date, a limited number of patients underwent SARS-CoV-2 testing.^{9,11–22} Interestingly in 3 reported patients with confirmed COVID-19 infection, cutaneous manifestations started 17 to 28 days after COVID-19 symptoms,²¹ indicating that chilblain-like lesions can appear late in disease course, as also suggested by our patient history. SARS-Cov2 testing by real-time RT-PCR on nasopharyngeal swabs and, in three cases, on skin biopsy proved negative in all our patients. However, the virus becomes undetectable in the upper respiratory airways usually after 6 to 11 days.³⁰ Indirectly, the infection can be measured by detecting SARS-COV-2-specific antibodies in the serum.³¹ SARS-COV-2 serology has been performed to date in very few patients with chilblain-like lesions and found positive in two patients with positive swab.^{12,19} IgG and IgA antibodies are secreted in patients with COVID-19.³² Most of the IgA is localized at mucosal sites for protection against pathogens and control of the microbiota. IgA is produced in response to respiratory infections in the nasal and bronchial-associated lymphoid tissue.³³ Besides its protective role, it has recently been demonstrated that IgA may also trigger inflammation at mucosal and non-mucosal sites.³⁴ However, many questions are still open regarding serology in COVID-19 and most of the published data is on adults.³¹ The tests current available vary because of the different techniques used and the antigen against which the specificity is measured.³⁴ We used two different methods: the first measures IgG antibodies against the SARS-COV-2 nucleoprotein (Abbott), and was negative in all patients, whereas the second (Euroimmun) detects IgG and IgA specific for the S1 domain of the spike protein.³⁵ Importantly, 6 patients (31.6%) were positive for IgA, and 3 (16%) borderline. The observation that children have serum antibodies against SARS-COV-2 of IgA isotype suggests that their immune response is based on strong mucosal protection that might even impair the triggering of an IgG response.

In conclusion, our study shows that chilblain-like lesions during COVID pandemic have several epidemiologic, clinical, capillaroscopic and histological characteristics different from idiopathic chilblains, making high unlikely that patients are affected with idiopathic perniosis. Though we could not formally prove SARS-CoV-2 infection in our patients, history data and presence of anti-SARS-COV-2 IgA strongly suggest a relationship between skin lesions and COVID-19. Importantly, most patients developing chilblain-like lesions have mild COVID-19 disease or are completely asymptomatic, and no fatal cases have been reported to date, suggesting that chilblain-like manifestations predominantly occur in COVID patients with a favourable disease course. Finally, our findings highlight the need for further investigations on the mechanisms of SARS-CoV-2 infection in children and pathogenesis of chilblain-like lesions.

Acknowledgements

We thank the patient's parents and patients themselves for accepting to participate in this study. The patients in this manuscript have given written informed consent to publication of their case details. We thank Mrs. Orietta Fiorani, nurse, for her support in patients' care, Mr. Mariani Riccardo for skilful technical assistance in electron microscopy and Mr. Gabriele Bacile for iconography preparation. Three of the authors (MEH, AD and GZ) of this publication are members of the European Reference Network ERN-SKIN and two of them (MEH and AD) are also members of vascular anomalies working group (VASCA WG) of the ERN for rare multisystemic vascular diseases (VAS-CERN).

References

- Nyssen A, Benhadou F, Magnée M, André J, Koopmansch C, Wautrecht JC. Chilblains. *Vasa* 2020; **49**: 133–140.
- Cappel JA Wetter DA. Clinical characteristics, etiologic associations, laboratory findings, treatment, and proposal of diagnostic criteria of pernio (chilblains) in a series of 104 patients at Mayo Clinic, 2000 to 2011. *Mayo Clin Proc* 2014; **89**: 207–215.
- Li Q, Guan X, Wu P *et al.* Early transmission dynamics in Wuhan, China, of novel coronavirus-infected pneumonia. *N Engl J Med* 2020; **382**: 1199–1207.
- Rodriguez-Morales AJ, Cardona-Ospina JA, Gutiérrez-Ocampo E *et al.* Clinical, laboratory and imaging features of COVID-19: A systematic review and meta-analysis. *Travel Med Infect Dis* 2020; **34**: 101623.
- Hu Z, Song C, Xu C *et al.* Clinical characteristics of 24 asymptomatic infections with COVID-19 screened among close contacts in Nanjing, China. *Sci China Life Sci* 2020; **63**: 706–711.
- Dong Y, Mo X, Hu Y *et al.* Epidemiology of COVID-19 among children in China. *Pediatrics* 2020; **145**: e20200702.
- Recalcati S. Cutaneous manifestations in COVID-19: a first perspective. *J Eur Acad Dermatol Venereol* 2020; **34**: e212–e213. <https://doi.org/10.1111/jdv.16387>.
- Bouaziz JD, Duong T, Jachiet M *et al.* Vascular skin symptoms in COVID-19: a French observational study. *J Eur Acad Dermatol Venereol* 2020. <https://doi.org/10.1111/jdv.16544>
- Galván Casas C, Català A, Carretero Hernández G *et al.* Classification of the cutaneous manifestations of COVID-19: a rapid prospective

- nationwide consensus study in Spain with 375 cases. *Br J Dermatol* 2020. <https://doi.org/10.1111/bjd.19163>
- 10 Alramthan A Aldaraji W. Two cases of COVID-19 presenting in clinical picture resembling chilblains disease. First report from the Middle East. *Clin Exp Dermatol* 2020. <https://doi.org/10.1111/ced.14243>
 - 11 Torres-Navarro I, Abril-Pérez C, Roca-Ginés J, Sánchez-Arráez J, Botella-Estrada R, Évole-Buselli M. Comment on 'Two cases of COVID-19 presenting with a clinical picture resembling chilblains: first report from the Middle East': pernio unrelated to COVID-19. *Clin Exp Dermatol* 2020. <https://doi.org/10.1111/ced.14255>
 - 12 Piccolo V, Neri I, Filippeschi C *et al*. Chilblain-like lesions during COVID-19 epidemic: a preliminary study on 63 patients. *J Eur Acad Dermatol Venereol* 2020. <https://doi.org/10.1111/jdv.16526>
 - 13 Kluger N, Scrivener Y. The use of Google trends for acral symptoms during COVID-19 outbreak in France. *J Eur Acad Dermatol Venereol* 2020. <https://doi.org/10.1111/jdv.16572>
 - 14 López-Robles J, de la Hera I, Pardo J, Martínez J, Cutillas-Marco E. Chilblain-like lesions: a case series of 41 patients during the COVID-19 pandemic. *Clin Exp Dermatol* 2020. <https://doi.org/10.1111/ced.14275>
 - 15 Fernandez-Nieto D, Jimenez-Cauhe J, Suarez-Valle A *et al*. Characterization of acute acro-ischemic lesions in non-hospitalized patients: a case series of 132 patients during the COVID-19 outbreak. *J Am Acad Dermatol* 2020. <https://doi.org/10.1016/j.jaad.2020.04.093>
 - 16 Landa N, Mendieta-Eckert M, Fonda-Pascual P, Aguirre T. Chilblain-like lesions on feet and hands during the COVID-19 pandemic. *Int J Dermatol* 2020; **59**: 739–743.
 - 17 Romani J, Baselga E, Mitja O *et al*. Chilblain and acral purpuric lesions in Spain during COVID confinement: Retrospective analysis of 12 cases. *Actas Dermosifiliogr* 2020. <https://doi.org/10.1016/j.ad.2020.04.002>
 - 18 Kolivras A, Dehavay F, Delplace D *et al*. Coronavirus (COVID-19) infection-induced chilblains: A case report with histopathologic findings. *JAAD Case Rep* 2020; **6**: 489–492.
 - 19 Garcia-Lara G, Linares-González L, Ródenas-Herranz T, Ruiz-Villaverde R. Chilblain-like lesions in pediatrics dermatological outpatients during the COVID-19 outbreak [published online ahead of print]. *Dermatol Ther* 2020; e13516. <https://doi.org/10.1111/dth.13516>
 - 20 Colonna C, Monzani NA, Rocchi A, Gianotti R, Boggio F, Gelmetti C. Chilblains-like lesions in children following suspected Covid-19 infection. *Pediatr Dermatol* 2020. <https://doi.org/10.1111/pde.14210>
 - 21 Suarez-Valle A, Fernandez-Nieto D, Diaz-Guimaraens B, Dominguez-Santas M, Carretero I, Perez-Garcia B. Acro-ischemia in hospitalized COVID-19 patients. *J Eur Acad Dermatol Venereol* 2020. <https://doi.org/10.1111/jdv.16592>
 - 22 de Masson A, Bouaziz JD, Sulimovic L *et al*. Chilblains are a common cutaneous finding during the COVID-19 pandemic: a retrospective nationwide study from France. *J Am Acad Dermatol* 2020. <https://doi.org/10.1016/j.jaad.2020.04.161>
 - 23 Chojnowski MM, Felis-Giemza A, Olesińska M. Capillaroscopy - a role in modern rheumatology. *Reumatologia* 2016; **54**: 67–72.
 - 24 Smith V, Herrick AL, Ingegnoli F *et al*. Standardisation of nailfold capillaroscopy for the assessment of patients with Raynaud's phenomenon and systemic sclerosis. *Autoimmun Rev* 2020; **19**(3): 102458.
 - 25 Boada A, Bielsa I, Fernández-Figueras MT, Ferrándiz C. Perniosis: clinical and histopathological analysis. *Am J Dermatopathol* 2010; **32**: 19–23.
 - 26 Lambova SN, Muller-Ladner U. Nailfold capillaroscopy of fingers and toes - variations of normal. *Curr Rheumatol Rev* 2018; **14**: 28–35.
 - 27 Ozmen M, Kurtoglu V, Can G, Tarhan EF, Soysal D, Aslan SL. The capillaroscopic findings in idiopathic pernio: is it a microvascular disease? *Mod Rheumatol* 2013; **23**: 897–903.
 - 28 Cribier B, Djeridi N, Peltre B, Grosshans E. A histologic and immunohistochemical study of chilblains. *J Am Acad Dermatol* 2001; **45**: 924–929.
 - 29 Wang ML, Chan MP. Comparative analysis of chilblain lupus erythematosus and idiopathic perniosis: histopathologic features and immunohistochemistry for CD123 and CD30. *Am J Dermatopathol* 2018; **40**: 265–271.
 - 30 Ling Y, Xu SB, Lin YX *et al*. Persistence and clearance of viral RNA in 2019 novel coronavirus disease rehabilitation patients. *Chin Med J (Engl)* 2020; **133**: 1039–1043.
 - 31 Sethuraman N, Jeremiah SS, Ryo A. Interpreting diagnostic tests for SARS-CoV-2. *JAMA* 2020. <https://doi.org/10.1001/jama.2020.8259>
 - 32 Padoan A, Sciacovelli L, Basso D *et al*. IgA-Ab response to spike glycoprotein of SARS-CoV-2 in patients with COVID-19: a longitudinal study. *Clin Chim Acta* 2020; **507**: 164–166.
 - 33 Macpherson AJ, McCoy KD, Johansen FE, Brandtzaeg P. The immune geography of IgA induction and function. *Mucosal Immunol* 2008; **1**: 11–22.
 - 34 Meyer B, Drosten C, Müller MA. Serological assays for emerging coronaviruses: challenges and pitfalls. *Virus Res* 2014; **194**: 175–183.
 - 35 Okba NMA, Müller MA, Li W *et al*. Severe acute respiratory syndrome coronavirus 2-specific antibody responses in coronavirus disease 2019 patients [published online ahead of print]. *Emerg Infect Dis* 2020; **26**(7). <https://doi.org/10.3201/eid2607.200841>.