

RESEARCH

Open Access

# Global expression studies in baker's yeast reveal target genes for the improvement of industrially-relevant traits: the cases of *CAF16* and *ORC2*

Roberto Pérez-Torrado, Joaquín Panadero, María José Hernández-López, José Antonio Prieto and Francisca Randez-Gil\*

## Abstract

**Background:** Recent years have seen a huge growth in the market of industrial yeasts with the need for strains affording better performance or to be used in new applications. Stress tolerance of commercial *Saccharomyces cerevisiae* yeasts is, without doubt, a trait that needs improving. Such trait is, however, complex, and therefore only in-depth knowledge of their biochemical, physiological and genetic principles can help us to define improvement strategies and to identify the key factors for strain selection.

**Results:** We have determined the transcriptional response of commercial baker's yeast cells to both high-sucrose and lean dough by using DNA macroarrays and liquid dough (LD) model system. Cells from compressed yeast blocks display a reciprocal transcription program to that commonly reported for laboratory strains exposed to osmotic stress. This discrepancy likely reflects differences in strain background and/or experimental design. Quite remarkably, we also found that the transcriptional response of starved baker's yeast cells was qualitatively similar in the presence or absence of sucrose in the LD. Nevertheless, there was a set of differentially regulated genes, which might be relevant for cells to adapt to high osmolarity. Consistent with this, overexpression of *CAF16* or *ORC2*, two transcriptional factor-encoding genes included in this group, had positive effects on leavening activity of baker's yeast. Moreover, these effects were more pronounced during freezing and frozen storage of high-sucrose LD.

**Conclusions:** Engineering of differentially regulated genes opens the possibility to improve the physiological behavior of baker's yeast cells under stress conditions like those encountered in downstream applications.

## Background

Variation in the osmotic pressure surrounding yeast occurs constantly in almost all steps from biomass production to bread making [1-3]. This is especially true for sweet bread products, which contain high amounts of sucrose or glucose/fructose syrup added as sweetener. Sweet dough exerts strong osmotic stress on yeasts, seriously affecting their fermentative capacity. Consequently, proofing times are longer for sweet bakery loaves and yield low volume products. To face these problems, manufacturers use greater amounts of yeast in the dough for-

mulation; however, this is expensive and final taste and texture are suboptimal. Therefore, there is a great interest in developing new baker's yeast strains with improved osmotic resistance.

The osmotic response of *Saccharomyces cerevisiae* has been well characterized in laboratory strains [4]. Exposure of yeast cells to highly osmotic environments provoke the up-regulation of ca. 400 genes, covering a wide variety of physiological functions, including carbon and amino acid metabolism, redox balance, anti-oxidant protection, ATPases, membrane proteins, chaperones, cytoskeletal and cell wall adaptations [5-7]. Such information has helped identify target genes, regulators and pathways involved in osmotic response [3]. However, given the special characteristics of these strains, it is

\* Correspondence: randez@iata.csic.es

<sup>1</sup> Department of Biotechnology, Instituto de Agroquímica y Tecnología de los Alimentos (CSIC), P.O. Box 73, E-46100-Burjassot Valencia, Spain  
Full list of author information is available at the end of the article

questionable whether such data can be used to develop molecular strategies to improve osmotic stress resistance in industrial yeasts.

Commercial baker's yeasts are domesticated strains of *S. cerevisiae*, selected and optimized for baking applications. Most of them are homothallic, with a high and irregular degree of ploidy and low sporulation ability [8]. Moreover, they exhibit chromosomal-length polymorphisms and rearranged chromosomes with multiple translocations. Thus, baker's yeast strains differ genomically from other *S. cerevisiae* strains in their adaptation to industrial conditions [3]. Their ability to ferment bread dough efficiently and behavior under stress conditions could also reflect specific expression patterns, as has been documented for wine strains [9,10]. Therefore, applying functional genomics to commercial strains under industrial conditions is a clue to understanding their osmotic adaptation mechanisms, thus enabling the design of new approaches to genetic improvement.

In this work, we have used compressed yeast blocks and flour-free liquid dough (LD) model system [11], to study the transcriptional response of industrial yeast to high-sucrose in bread dough. The aim was to reproduce as much as possible the conditions in which bakers employ commercial yeast, quite often as compressed yeast, in order to identify the set of genes regulated under these conditions. Overall, our work provides new insights into the links between technological traits and genetic determinants, showing the potential of this approach for genetic engineering of bakers' yeast.

## Results

We compared the genome-wide transcription pattern of starved cells from compressed yeast blocks with that of cells cultured for 60 min in high-sugar liquid dough (LD). In previous studies, we demonstrated that this flour-free model system mimics the nutritional and stressful environment encountered by baker's yeast cells in bread dough [11]. Furthermore, data obtained from the microarray analysis agreed very well with Northern blot data for five marker genes (additional file 1): two induced, *PIS1* (2.7-fold induction on gene filter and 1.6-fold induction in Northern) and *PHO3* (7.3/4.0), and three repressed, *OLE1* (2.7-fold repression on gene filter and 1.7-fold repression in Northern), *HSP12* (4.7/4.1) and *HSP26* (4.0/3.1). Hence, the combination of LD model system and gene filters is an appropriate and simple way to perform genome-wide transcriptional analysis under commercially important, but experimentally intractable, conditions.

### Global expression after transfer to high-sucrose LD: overview

Microarray data revealed that the shift of starved cells to high-sucrose LD greatly affected the gene expression pro-

gram of baker's yeast. The list of genes induced or repressed 4.0-fold or more ( $\log_2$  ratio of  $\pm 2$ ) is available as additional file 2. Significant variations in the mRNA levels of 1,029 genes, 423 induced and 606 repressed, were observed. The list of regulated genes was screened for enrichment of specific functional categories (Table 1), from which several general conclusions can be drawn. First, the observed expression changes mainly reflect the suppression of starvation conditions in compressed baker's yeast cells, rather than exposure to a hostile medium. Thus, a significant number of genes encoding ribosomal proteins or involved in protein biosynthesis and ribosomal processing, functions known to be transiently repressed by stress conditions [5,6], were significantly up-regulated (Table 1). On the contrary, genes associated with GO biological processes as response to stress, i.e. *TPS1*, *TSL1*, *CTT1*, *GLO1*, *HSP12*, *HSP26*, *HSP30* or *GRE1*, that are normally induced by global stress responses [5,6], tended to be repressed (Table 1). Such reciprocal expression pattern has been reported previously when yeast cells were subjected to opposite stresses [6]. This view was confirmed by comparing the dataset of genes differentially regulated in our study, with the list of ESR (Environmental Stress Response) genes, a set of genes showing a common transcriptional response to different stress conditions [6]. As can be seen, only 4 genes showed a common response (Figure 1A). On the contrary, 239 and 137 genes, induced and repressed in high-sucrose LD, respectively, displayed an opposite transcriptional response in the ESR (Figure 1B). Second, enriched GO terms are mainly associated with energy generation and metabolic functions. Indeed, aerobic respiration, tricarboxylic acid cycle, oxidative phosphorylation or fatty acid oxidation were strongly overrepresented among the genes that were repressed (Table 1), reflecting a shift from carbon starvation conditions to active fermentation. Third, the transcriptional response after 60 min of inoculation of baker's yeast cells also reflects the variation in environmental conditions and/or nutrient concentration during the fermentation process. For instance, we noted a significant induction of *THI6*, *THI7*, *THI21*, *THI22*, *THI80* and *PHO3*, all of them involved in thiamine biosynthesis or transport under anaerobic conditions (additional file 2). On the contrary, *THI5*, *THI11* and *THI12*, encoding biosynthetic enzymes involved in the production of thiamine under aerobic conditions, were repressed. This clearly reflects the microaerophilic conditions found by yeast cells in regular bread dough.

### Differences in the global transcriptional response between lean and high-sucrose dough

The transcriptional response of commercial baker's yeast cells in lean and high-sucrose LD were compared. Additional file 2 provides the complete list of genes showing at

**Table 1: Enriched GO categories for up- and down-regulated genes at 60 min after onset of fermentation of baker's yeast cells in high-sucrose LD.**

Functional Group	n	Functional Group	n
<b>Up (423 genes)</b>		<b>Down (606 genes)</b>	
Ribosome (SCE03010 p = 1.5E-59)	78	Cellular carbohydrate metabolic process (GO0044262 p = 2.1E-12)	57
Translation (GO0006412 p = 6.2E-09)	113	Aerobic respiration (GO0009060 p = 9.5E-07)	25
Ribonucleoside monophosphate metabolic process (GO 0009161) p = 8.1E-05)	10	Response to stress (GO006950 p = 9.8E-05)	16
tRNA metabolic process (GO0006399 p = 2.1E-03)	21	Starch and sucrose metabolism (Kegg pathway p = 7.7E-05)	18
Nuclear transport (GO0051169 p = 1.2E-03)	20	Oxidative phosphorylation (GO0006119 p = 2.9E-04)	16
RNA helicase activity (GO003724 p = 1.2E-03)	10	Tricarboxylic acid cycle (GO0006099 p = 1.8E-04)	10
		Coenzyme metabolic process (GO0006732 p = 1.0E-04)	32
		Response to water deprivation (GO0009414 p = 9.0E-03)	4
		Water soluble vitamin metabolic process (GO0006767 p = 3.7E-03)	18
		Carbohydrate transport (GO0008643 p = 3.1E-03)	12
		Fatty acid oxidation (GO0019305 p = 1.1E-03)	6

n, Number of genes found in the indicated category

least 4-fold up- or down-regulation ( $\log_2$  ratio of  $\pm 2$ ) at 60 min after the transfer of starved cells from compressed yeast blocks to lean LD. A total of 1,105 ORFs showed significant expression changes, being 438 induced and 667 repressed. Again, comparison of regulated genes in lean LD with the list of ERS genes revealed the lack of common responses (Figure 1). Moreover, the list of enriched GO terms (Table 2) was similar to that observed in cells transferred by 60 min to high-sucrose LD (Table 1). Only a few functional categories, comprising a small number of genes were differentially over-represented in lean LD (Table 2). Consistent with this, direct comparison of the set of regulated genes in lean and high-sucrose LD, showed again a high similarity. In this analysis, genes with no reliable data in any of the two conditions were excluded. Of 423 and 606 genes up- and down-regulated in high-sucrose, 288 (68%) and 461 (76%) genes, followed the same change in lean LD. Moreover, there was none ORF showing reverse regulation. The complete list of commonly regulated genes is also shown in additional file 2.


#### Genes differentially regulated in high-sucrose

We looked more in detail at those genes that were differentially regulated by the presence of sucrose in the LD system. A total of 135 and 145 ORFs, showing no response in lean LD were significantly induced and repressed, respectively (additional file 2). There were no enriched GO terms among the down-regulated genes. However, the list of induced genes showed prevalent

functional categories including ribosome biogenesis and assembly (GO0005840 p = 6.7E-12 n = 32), tRNA metabolic process (GO0006399 p = 4.8E-03 n = 10) and snRNA modification (GO00040031 p = 6.7E-03 n = 3). Furthermore, the list of up-regulated genes included six genes encoding proteins with regulatory functions on transcription and/or translation, *CAF16*, *HAL9*, *MED4*, *SWC4*, *ORC2* and *TAF1*. All of them have been reported to be induced by osmotic stress conditions [5], and two, *CAF16* and *ORC2* have been additionally identified as cold-regulated [12]. These observations suggest the need for a remodeling of the transcriptional and translational machinery under extreme osmolarity conditions.

#### LD-induced genes versus HST genes

We next compared the list of genes specifically or commonly up-regulated in LD under lean and high-sucrose conditions with the set of genes identified by Ando *et al.* [13], concretely 273 genes, as required for tolerance to high sucrose (HST) in a genome-wide screening of *S. cerevisiae* deletion strains (Figure 2). The reason for do this was that in this study, as much as 269 from 273 genes showed cross-sensitivities to sorbitol and NaCl [13]. Consequently, the expression of these essential genes might be expected to be affected in either high-sucrose or lean dough. Seven genes were found in common exclusively between high-sugar LD and HST set of genes (Figure 2). We next compared the list of HST genes with that of 150 genes specifically up-regulated in lean LD and 288 genes that were commonly induced in both high-sugar and lean



LD (additional file 2). The comparison revealed a further overlap of 8 and 33 genes among those specifically induced in lean LD and co-induced in both LD systems, respectively (Figure 2). Of these 48 genes found in any condition, 24 genes (50%) encode proteins involved in translation (GO0006412  $p = 5.8E-11$ ). Hence, our results make again emphasis in the importance of modulating the transcript levels for genes encoding the translation apparatus and its regulators.

#### Overexpression of *CAF16* and *ORC2* enhances baking performance of yeast cells exposed to osmotic and freeze stress

We addressed the question of whether gene filter data might reveal target genes for strain selection. We rationalized that products of genes differentially regulated in the LD system (additional file 2) could be important for yeast cells to adapt to extremely low water activity conditions. Eleven genes displaying a significant induction at

60 min after inoculation of commercial baker's yeast cells in LD were tested. Of them, 6 genes (*CAF130*, *CDC10*, *FUR1*, *SEC14*, *YVH1* and *ZUO1*) showed to be commonly up-regulated in lean and high-sucrose LD, while 5 genes (*CAF16*, *MFT1*, *NMT1*, *ORC2* and *SSF2*) were specifically induced in high-sucrose (additional file 2).

The selected genes were cloned into the shuttle vector YEplac195 and the resulting plasmids were used to transform the industrial HS13 baker's yeast strain ( $Ura^-$ ). Transformants were selected by auxotrophic complementation and the functionality of the recombinant strains analyzed for testing their ability to produce  $CO_2$  in high-sucrose LD. However, only overexpression of *CAF16* and *ORC2*, two of the six transcriptional factor-encoding genes [14-16], identified as specifically induced in high-sucrose LD (additional file 2), had significant positive effects on leavening activity of baker's yeast cells (Table 3 and Figure 3, control, 0 days). Because of this, only strains overexpressing these two genes were further characterized.

Thus, we then inspected the behavior of the recombinant strains in frozen products. Freezing is a complex stress in which cellular damage by osmotic shrinkage is one of the main causes of death [17,18]. As can be seen in Figure 3A,  $CO_2$  production during frozen storage of lean LD was almost similar for all the strains. However, overexpression of *CAF16* or *ORC2* appeared to play a stress-protective role in high-sucrose LD (Figure 3B). Indeed,  $CO_2$  production attained by overexpression of either of these genes in 45-day pre-frozen high-sucrose LD was about 2-fold higher than that observed with the control strain (Figure 3B).

Finally, YE $pCAF16$  and YE $pORC2$  transformants were analyzed for growth in molasses medium or in high-sucrose LD (Table 4). A good behavior in these media is critical for the technological applicability of industrial strains [1-3]. As can be seen, neither high copy number of *CAF16* nor *ORC2* altered the specific growth rate under the assay conditions (Table 4).

#### Discussion

Here we show relevant information concerning the transcriptional response of commercial baker's yeast in high-sucrose and lean dough. We demonstrated that cells from compressed yeast blocks, the main source of fresh baker's yeast, display a reciprocal transcription program to that commonly reported for laboratory strains exposed to high-osmotic conditions [5,6]. This discrepancy likely reflects differences in strain background and/or experimental design. Indeed, most or all commercial baker's yeasts are polyploid or aneuploid strains with approximately  $3n-4n$  DNA content [19,20]. Moreover, we used starved yeast cells, as do the bakers in the bread-making process, in contrast to the YPD-grown early-log phase

**Table 2: Enriched GO categories for up- and down-regulated genes at 60 min after onset of fermentation of baker's yeast cells in lean LD.**


Functional Group	n	Functional Group	n
<b>Up (438 genes)</b>		<b>Down (667 genes)</b>	
Ribosome (SCE03010 p = 3.8E-64)	81	Aerobic respiration (GO0009060 p = 2.1E-06)	26
Translation (GO0006412 p = 1.5E-47)	124	Oxidative phosphorylation (GO0006119 p = 9.9E-04)	16
Ribonucleoside monophosphate metabolic process (GO 0009161) p = 2.8E-03)	8	Tricarboxylic acid cycle (GO0006099 p = 4.4E-04)	10
tRNA metabolic process (GO0006399 p = 3.9E-04)	23	Starch and sucrose metabolism (Kegg path. p = 1.4E-04)	17
Nuclear transport (GO0051169 p = 1.8E-04)	22	Response to external stimulus (GO0009605 p = 6.1E-03)	10
RNA helicase activity (GO003724 p = 5.1E-03)	9	Carbohydrate transport (GO0008643 p = 2.5E-03)	13
snRNA modification (GO00 p = 2.9E-04)	5	Coenzyme catabolic process (GO0009109 p = 1.7E-03)	33

n, Number of genes found in the indicated category

cells of laboratory strains examined in previous works [5-7]. Overall, these data further support the idea that studies involving industrial yeast should be performed under those particular conditions encountered in downstream applications, as it has been previously suggested [1-3,20].

We found that the transcriptional response of starved baker's yeast cells was qualitatively similar in the presence or absence of sucrose in the LD model system. This was, in some way, not surprising since a previous study by Causton *et al.* [5] had shown that the genetic response to high salt and to sorbitol are remarkably similar to each other. Cells inoculated in either lean or high-sucrose LD are indeed exposed to the same nutrient environment, except for sucrose. As a result, water activity is low in both systems ( $a_w = 0.97$  and  $0.95$ , respectively). Moreover,

a recent survey for sucrose-sensitivity mutants revealed that 269 of 273 genes identified as required under these conditions (High Sucrose Tolerance, HST genes), showed cross-sensitivities to sorbitol and NaCl [13]. Thus, decreased water activity appears to be a main drive force of the transcriptional program triggered by yeast cells in bread dough. In this line, our study has identified genes that might be important to cope with high-osmolarity. Indeed, 48 genes found as specifically or commonly induced in high-sucrose and lean LD overlap with genes reported as essential in the HST analysis [13]. Of them, 50% encode proteins involved in or related with translation. Osmotic stress is generally considered to be a growth-restrictive condition and regulation of genes involved in protein biosynthesis constitutes an important



**Table 3: CO<sub>2</sub> production by different HS13 transformants<sup>a</sup>**

Plasmid	ml CO <sub>2</sub> /mg yeast (d.w.) ± SD (P value)	
	20% Sucrose	30% Sucrose
YEplac195	0.401 ± 0.005	0.255 ± 0.003
YE <sub>p</sub> CAF130	0.385 ± 0.015 (0.218)	nd
YE <sub>p</sub> CDC10	0.425 ± 0.017 (0.140)	nd
YE <sub>p</sub> FUR1	0.401 ± 0.009 (0.137)	nd
YE <sub>p</sub> SEC14	0.422 ± 0.022 (0.112)	nd
YE <sub>p</sub> YVH1	0.383 ± 0.013 (0.095)	nd
YE <sub>p</sub> ZUO1	0.435 ± 0.019 (0.110)	nd
YE <sub>p</sub> CAF16	0.462 ± 0.012 (0.017)*	0.400 ± 0.030 (0.003)*
YE <sub>p</sub> MFT1	0.440 ± 0.030 (0.155)	nd
YE <sub>p</sub> NMT1	0.426 ± 0.022 (0.204)	nd
YE <sub>p</sub> ORC2	0.446 ± 0.018 (0.022)*	0.402 ± 0.036 (0.012)*
YE <sub>p</sub> SSF2	0.385 ± 0.015 (0.175)	nd

<sup>a</sup>Cells were grown on molasses plates and tested for gas production as described in the Materials and Methods section. Values represent the mean of at least three independent experiments. SD, standard deviation. nd, non-determined. \*, *P* < 0.05 for gas production of the overexpressing strain compared to gas production of the control strain (YEplac195).


part of the overall adaptive response in *S. cerevisiae* [7,21].

Our study also revealed the existence of genes differentially regulated in high-sucrose LD, whose activity might be important for the adaptive response of yeast cells to severe osmotic stress. In particular, engineering of genes with regulatory functions has the potential to affect simultaneously the activity of many cellular functions. In this line, previous work had shown that overexpression of the calcineurin-target *CRZ1* and unrelated genes encoding Cys<sub>2</sub>/His<sub>2</sub>-type zinc finger proteins confers freeze tol-

**Table 4: Specific growth rate of HS13 transformants**

Plasmid	μ <sub>max</sub> (h <sup>-1</sup> ) <sup>a</sup> ± SD	
	HSLD	Molasses
YEplac195	0.21 ± 0.02	0.51 ± 0.03
YE <sub>p</sub> CAF16	0.19 ± 0.02	0.50 ± 0.02
YE <sub>p</sub> ORC2	0.22 ± 0.04	0.50 ± 0.03

<sup>a</sup>Cells were grown in high-sucrose liquid dough (HSLD) or molasses medium. Values represent the mean of at least three independent experiments. SD, standard deviation. μ<sub>max</sub> was calculated as described in the Materials and Methods section.



erance and enhances fermentative capacity of baker's yeast [22].

Six genes with these characteristics, *CAF16*, *HAL9*, *MED4*, *SWC4*, *ORC2* and *TAF1*, were found from the list of 135 genes specifically up-regulated in high-sucrose LD. Two of them, *CAF16* and *ORC2* were tested together with other nine genes for their ability to provide enhanced fermentative performance to baker's yeast cells. Quite remarkably, both *CAF16* and *ORC2* have been reported to be induced by either, osmotic [5] and cold stress [12]. In agreement with this, we found that transformants of the industrial HS13 strain, in which the *CAF16* or *ORC2* genes had been introduced in a high-copy number, exhibited increased metabolic activity in LD. This was especially true in frozen high-sucrose LD, where the presence of salt (around 2%, flour basis), combined with sucrose and ice-crystal formation exposes yeast cells to both ionic and osmotic stress [22,23]. Indeed, ionic imbalance caused by ice-crystal formation is an important factor

determining freeze injury in all living organisms [24]. Thus, enhanced expression of *CAF16* and *ORC2* would allow yeast cells to alleviate the harmful effects of ionic stress during freezing. *CAF16* is a member of the non-transporter group of the ATP-binding cassette (ABC) superfamily [25], and component of the Ccr4-Not transcriptional regulatory complex [14]. In the case of environmental stress, it has been described that this complex affects the Msn2p-dependent transcriptional activation [26,27], and the stress-specific response modulated by the transcription factor Skn7p [28]. On the other hand, Orc2p is a subunit of the origin recognition complex (ORC) that functions in pre-replication complex formation [29] and in chromatin silencing at telomere [16]. Interestingly, Orc2p also functions in the transcriptional regulation of stress-responsive genes, acting as a repressor [30,31] or an activator able to induce expression of highly transcribed genes positioned nearby ORCs [15]. Nevertheless, more work is needed to establish the functional connections between overexpression of *CAF16* and *ORC2* and ion and freeze tolerance in high-sucrose dough.

## Conclusions

Global transcriptional approach is a powerful tool to overcome yeast response studies under complex conditions. Such information is clearly necessary in establishing the relationship between genetic determinants and industrial traits and in defining targets for strain selection and improvement. Thus, the finding that overexpression of *CAF16* and *ORC2* helps yeast cells to face with osmotic and freeze stress, underlies the importance of this strategy and opens the possibility to new advances. Indeed, other genes determining the same phenotypic character might be also found from the list of regulated genes. If confirmed, the use of engineered strains for these genes may be a way to optimize gassing rate in both fresh and frozen high-sucrose dough, leading to bakery products with lower cost and better organoleptic properties.

## Materials and methods

### Strains, culture media and general methods

The baker's yeast strains L'Hirondelle and HS13 were used throughout this work. HS13 is a uracil auxotrophic non-commercial strain (Lesaffre International, Lille, France), while L'Hirondelle is a commercial strain produced by the Lesaffre Group and usually employed for general baking. The *E. coli* strain DH10B was used as the host for plasmid construction. Yeast cells were cultured at 30°C in defined media, SD (0.2% yeast nitrogen base without amino acids [DIFCO], 0.5% (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>, 2% glucose). *E. coli* was grown in Luria Bertani (LB) medium (1% peptone, 0.5% yeast extract, 0.5% NaCl) supple-

mented with ampicillin (50 mg/l). Yeast cells were transformed by the lithium acetate method [32], and transformants were selected by auxotrophic complementation in SD plates. *E. coli* was transformed by electroporation following the manufacturer's instructions (Eppendorf AG, Hamburg, Germany).

### Yeast biomass preparation

Compressed yeast packs were acquired from a local distributor, maintained at 4°C, and used no longer than 5 days after the production date. Weighed samples were resuspended in 4°C distilled water containing 27 g/l NaCl, vortexed, and the OD<sub>600</sub> of the resulting suspension was measured. Final yeast concentration was adjusted to approximately 15 mg (dry weight) per ml. To do this, cell mass was related to optical density measurements, OD<sub>600</sub> = 1 equals 0.35 mg cells dry weight/ml [11].

Yeast biomass from HS13 transformants was prepared by cultivating cells (7.6 units of OD<sub>600</sub>) on molasses (5.0 g beet molasses [49% sucrose], 0.5 g (NH<sub>4</sub>)<sub>2</sub>HPO<sub>4</sub>, 26.0 g agar and 20 µg biotin per liter; pH 5.0) plates (140-mm diameter) for 20 h at 30°C. Then, yeast cells were recovered by washing the plate surface with 2 × 10 ml of distilled water and the yeast suspension was poured into a tube. After centrifugation, the yeast cake was washed twice with distilled water (4°C), resuspended in saline solution and the final yeast concentration was adjusted as above for further analysis of CO<sub>2</sub> production.

### Liquid dough (LD) model system

High-sucrose and lean LD solutions were prepared as previously described [11]. Briefly, a 5 × concentrated nutrient solution, containing 5 g MgSO<sub>4</sub>·7H<sub>2</sub>O, 2 g KCl, 11.75 g (NH<sub>4</sub>)<sub>2</sub>HPO<sub>4</sub>, 4 mg thiamine, 4 mg pyridoxine, and 40 mg nicotinic acid in a final volume of 250 ml of 0.75 M citrate buffer (pH 5.5), was prepared. Twenty ml of the concentrated nutrient solution was added to a tube containing 0.5 g yeast extract, 3 g glucose, 9 g maltose, 12 g sorbitol and 50 g sucrose, and the mixture dissolved by sonication. Distilled water was added to a final volume of 100 ml, and the solution filter-sterilized. Lean LD was prepared as above, except that no sucrose was added.

### Gas production measurements

Fifteen ml of yeast suspension was poured into a 250-ml screw cap graduated bottle, placed in a 30°C water bath and gently shaken (80 rpm). After 15 min, 15 ml of 30°C pre-warmed LD was added and the amount of CO<sub>2</sub> evolved recorded in a Fermograph II (ATTO Co., Ltd., Tokyo, Japan). Samples for freezing were kept at -80°C for 1 h and then stored at -20°C. At different times, they were thawed at 30°C for 30 min before measuring gassing power. In all cases, CO<sub>2</sub> production was recorded for 180

min. Values are expressed as ml of CO<sub>2</sub> per mg of yeast cells, dry weight.

#### Growth rate estimation

Culture growth was followed by measuring OD<sub>600</sub> with a Polarstar plate spectrophotometer in six replicates of 225 µl volumes for high-sucrose LD and molasses medium. Maximum specific growth rate was calculated from each condition by directly fitting OD<sub>600</sub>/ml versus time to the reparametrized Gompertz equation proposed by Zwietering *et al.* [33]:  $y = D \cdot \exp\{-\exp[(\mu_{\max} \cdot e)/D] \cdot (\lambda - t) + 1\}$  where  $y = \ln(N_t/N_0)$ ,  $N_0$  is the initial population (OD<sub>600</sub>/ml) and  $N_t$  is the population at time  $t$ ;  $D = \ln(N_{\infty}/N_0)$  is the maximum population value reached with  $N_{\infty}$  as the asymptotic maximum,  $\mu_{\max}$  is the maximum specific growth rate ( $h^{-1}$ ), and  $\lambda$  the lag phase period (h). Growth data from each treatment and yeast were fitted by a non-linear regression procedure, minimizing the sum of squares of the difference between experimental data and the fitted model, i.e., loss function (observed-predicted). This task was accomplished using the non-linear module of the Statistica 6.0 software package and its Quasi-Newton option.

#### Plasmids

PCR-amplified fragments containing the whole sequence of *CAF130*, *CDC10*, *FUR1*, *SEC14*, *YVH1*, *ZUO1*, *CAF16*, *MFT1*, *NMT1*, *ORC2* and *SSF2* gene, including its own promoter and terminator were obtained with specific synthetic oligonucleotides (additional file 3). The corresponding fragments were digested with the appropriate set of enzymes, XbaI/PstI (*CAF16*, *CDC10*, *NMT1*, *ORC2*, *SEC14*, *ZUO1*), SalI/HindIII (*CAF130*, *SSF2*, *YVH1*), EcoRI/PstI (*FUR1*) or XbaI/HindIII (*MFT1*) and cloned into the plasmid YEplac195 digested with the same set of enzymes [34].

#### RNA purification and Northern blot analysis

To prepare control RNA from baker's yeast cells, 0.2 g pieces of compressed yeast blocks were homogenized with 10 ml of ice-cold LETS buffer (200 mM LiCl, 20 mM EDTA, 20 mM Tris-HCl [pH 8.0], 0.4% SDS). Aliquots of 0.5 ml of the cell suspension were transferred to screw-cap microcentrifuge tubes containing 0.5 ml of phenol and 0.5 ml of glass beads (acid-washed, 0.4-mm diameter). Then, the suspension was mixed vigorously 2 times for 45 s each time in a FastPrep device (Bio101). RNA was purified as previously described [11]. Cells from LD were harvested by centrifugation, resuspended in 0.5 ml of LETS buffer and treated as above.

Equal amounts of RNA (30 µg) were separated in 1% (w/v) agarose gels, containing formaldehyde (2.5% v/v), transferred to a Nylon membrane and hybridized with nonradioactive DIG labeled probes containing sequences

of *PIS1* (+41 to +571), *PHO3* (+1 to +1,381), *OLE1* (+112 to +1,106), *HSP12* (+1 to +311) and *HSP26* (+51 to +508). Oligonucleotides for synthesis of probes are reported in additional file 3. DNA sequences were obtained from the MIPS database (available at <http://mips.gsf.de>). PCR labeling of DNA probes, membrane pre-hybridizations and hybridizations were performed with the PCR DIG Probe Synthesis Kit and DIG Easy Hyb solution of Roche (Roche Diagnostics GmbH, Mannheim, Germany). After stringency washes, the blots were subjected to immunological detection using anti-digoxigenin antibody conjugated to alkaline phosphatase (Roche), followed by CDP-Star detection (Roche). Images were captured with the Las-1000 Plus imaging system (Fuji, Kyoto, Japan). Spot intensities were quantified with the Image Gauge software version 3.12 (Fuji). Values of spot intensity were evaluated with respect to the rRNA level and represented as the folds of induction/repression of each mRNA.

#### Synthesis of target cDNA and DNA filter hybridization

Labeling by random priming using [ $\alpha$ -<sup>33</sup>P]dCTP (3,000 Ci/mmol; 10 µCi/µl) was performed as described [35]. The labeled cDNAs were purified by using a MicroSpin S-300 column (Amersham Biosciences). Between  $3 \times 10^6$  and  $5 \times 10^6$  dpm/ml of labeled cDNA was used for filter hybridization. The purified target cDNA was kept at 4°C until used. Prehybridization, hybridization, and washing were carried out according to published protocols [35]. Replicate samples were prepared from three independent experiments of both high-sucrose and lean LD fermentation.

#### Data generation, correction, and normalization

Yeast macroarrays containing the whole genome (6,052 ORFs) of yeast strain FY1679 on  $7.5 \times 11$ -cm nylon filters, were provided by the DNA chips laboratory of the University of Valencia (SCSIE, Valencia, Spain, <http://scsie.uv.es/chipsdna>). The macroarrays have been constructed and the protocols for their use have been evaluated as previously described [35]. A total of 3 different nylon filters were used (one for each experimental replicate). cDNA labeled samples belonging to the same experimental replicate were successively hybridized against the same filter. Therefore, each determination is obtained from the averaged results of three independent filters. Previous studies have shown that the same membrane can be used up to 12 times with satisfactory results [35]. Hybridization signals were quantified using Array-Vision 7.0 software (Imaging Research, Inc., St. Catharines, Ontario, Canada), taking the artifact-removed median density (with the corresponding subtracted background) as signal. Poor or inconsistent signals were not considered for further analysis. The use of the same DNA chip for successive cDNA hybridization improved the



comparisons between values for each gene. cDNA hybridizations were normalized within each experiment replicate by the global mean procedure. Reproducibility of the replicates was tested by ArrayStat software (Imaging Research, Inc.), considering the data as independent and allowing the program to take a minimum number of two valid replicates in order to calculate the mean and standard deviation values for every gene (only one of the three replicates was allowed to be a removable outlier). To detect differentially expressed genes, a z-score was obtained for every gene. A *p*-value of 0.05 and a median false discovery rate of 0.5% were used. The change in expression level of a transcript is considered when the  $\log_2$  ratio is equal to  $\pm 2$  (fold change of 4).

### Functional category searches

Statistical assessment of overrepresentation of GO biological processes categories [36]<http://www.geneontology.org/> among significantly changed sets of transcripts was achieved using the Database for Annotation, Visualization and Integrated Discovery (DAVID) 2006 [37], which is available at <http://david.niaid.nih.gov>. The *Saccharomyces* Genome Database <http://www.yeastgenome.org> and the MIPS Comprehensive Yeast Genome Database <http://mips.helmholtz-muenchen.de/genre/proj/yeast/> were used to retrieve information about specific gene function and biological process. The list of *S. cerevisiae* ESR genes was obtained from the Web site of Gasch *et al.* [6], at [http://www-genome.stanford.edu/yeast\\_stress](http://www-genome.stanford.edu/yeast_stress). The data of deletions resulting in high-sucrose sensitivity was obtained from Ando *et al.* [13].

### Additional material

**Additional file 1 Comparison of Northern blots versus gene filter data.** The fold-change in expression level of five marker genes, *PI51*, *PHO3*, *OLE1*, *HSP12* and *HSP26*, as obtained by Northern blot (white bars) or gene filters (black bars) was compared. Total RNA extraction from cells of the L'Hirondelle strain and global gene expression analysis were performed as described in the Materials and Methods section. Cells from compressed yeast blocks (time zero) were used as control.

**Additional file 2 List of regulated genes after 60 min of transfer of baker's yeast cells to lean or high-sucrose liquid dough.** Genes specifically up- or down-regulated in lean liquid dough (LLD) and high-sucrose liquid dough (HSLD), and those commonly up- or down-regulated are shown. Expression values showing more than 4-fold induction or repression ( $\log_2$  ratio of  $\pm 2$ ) are colored in red and green, respectively. Each value indicates the ratio of the level of expression in liquid dough-transferred cells at 60 min to that in control cells from compressed yeast blocks.

**Additional file 3 Oligonucleotides used in this study.** Sequences of forward and reverse primers employed to amplify by PCR the mentioned genes.

### Competing interests

The authors declare that they have no competing interests.

### Authors' contributions

RPT assisted with data analysis and interpretation, statistical analysis and manuscript writing. JP carried transcriptomic experiments. MJHL performed the

Northern blot experiments. JAP conceived the study, participated in its design, and contributed to the writing of the manuscript. FRG constructed the strains, performed fermentative capacity experiments, conceived the study, participated in its design, and writing of the manuscript. All authors read and approved the final manuscript.

### Acknowledgements

We thank A. Blasco for technical assistance. We also thank Lesaffre International for providing us with the H513 strain. This research was funded by the Comisión Interministerial de Ciencia y Tecnología (projects AGL2004-00462 and AGL2007-65498-C02-01) from the Ministry of Science and Technology of Spain. J.P. was supported by a F.P.I. fellowship.

### Author Details

Department of Biotechnology, Instituto de Agroquímica y Tecnología de los Alimentos (CSIC), P.O. Box 73, E-46100-Burjassot Valencia, Spain

Received: 14 June 2010 Accepted: 13 July 2010

Published: 13 July 2010

### References

1. Attfeld PV: **Stress tolerance: the key to effective strains of industrial baker's yeast.** *Nature Biotechnol* 1997, **15**:1351-1357.
2. Rande-Gil F, Sanz P, Prieto JA: **Engineering baker's yeast: room for improvement.** *Trends Biotechnol* 1999, **17**:237-244.
3. Rande-Gil F, Aguilera J, Codón A, Rincón AM, Estruch F, Prieto JA: **Baker's yeast: challenges and future prospects.** In *Functional Genetics of Industrial Yeasts* Edited by: de Winde JH. Heidelberg: Springer-Verlag; 2003:57-97.
4. Hohmann S: **Osmotic stress signaling and osmoadaptation in yeasts.** *Microbiol Mol Biol Rev* 2002, **66**:300-372.
5. Causton HC, Ren B, Koh SS, Harbison CT, Kanin E, Jennings EG, Lee TI, True HL, Lander ES, Young RA: **Remodeling of yeast genome expression in response to environmental changes.** *Mol Biol Cell* 2001, **12**:323-337.
6. Gasch AP, Spellman PT, Kao CM, Carmel-Harel O, Eisen MB, Storz G, Botstein D, Brown PO: **Genomic expression programs in the response of yeast cells to environmental changes.** *Mol Biol Cell* 2000, **11**:4241-4257.
7. Rep M, Krantz M, Thevelein JM, Hohmann S: **The transcriptional response of *Saccharomyces cerevisiae* to osmotic shock. Hot1p and Msn2p/Msn4p are required for the induction of subsets of high osmolarity glycerol pathway-dependent genes.** *J Biol Chem* 2000, **275**:8290-8300.
8. Codon AC, Benitez T, Korhola M: **Chromosomal polymorphism and adaptation to specific industrial environments of *Saccharomyces* strains.** *Appl Microbiol Biotechnol* 1998, **49**:154-63.
9. Perez-Ortin JE, Garcia-Martinez J, Alberola TM: **DNA chips for yeast biotechnology. The case of wine yeasts.** *J Biotechnol* 2002, **98**:227-241.
10. Bisson LF, Karpel JE, Ramakrishnan V, Joseph L: **Functional genomics of wine yeast *Saccharomyces cerevisiae*.** *Adv Food Nutr Res* 2007, **53**:65-121.
11. Panadero J, Rande-Gil F, Prieto JA: **Validation of a flour-free model dough system for throughput studies of baker's yeast.** *Appl Environ Microbiol* 2005, **71**:1142-1147.
12. Sahara T, Goda T, Ohgiya S: **Comprehensive expression analysis of time-dependent genetic responses in yeast cells to low temperature.** *J Biol Chem* 2002, **277**:50015-50021.
13. Ando A, Tanaka F, Murata Y, Takagi H, Shima J: **Identification and classification of genes required for tolerance to high-sucrose stress revealed by genome-wide screening of *Saccharomyces cerevisiae*.** *FEMS Yeast Res* 2006, **6**:249-267.
14. Liu HY, Chiang YC, Pan J, Chen J, Salvatore C, Audino DC, Badarinarayana V, Palaniswamy V, Anderson B, Denis CL: **Characterization of *CAF4* and *CAF16* reveals a functional connection between the CCR4-NOT complex and a subset of SRB proteins of the RNA polymerase II holoenzyme.** *J Biol Chem* 2001, **276**:7541-7548.
15. Shor E, Warren CL, Tietjen J, Hou Z, Müller U, Alborelli I, Gohard FH, Yemm AI, Borisov L, Broach JR, Weinreich M, Nieduszynski CA, Ansari AZ, Fox CA: **The origin recognition complex interacts with a subset of metabolic genes tightly linked to origins of replication.** *PLoS Genet* 2009, **5**:e1000755.
16. Ozaydin B, Rine J: **Expanded roles of the origin recognition complex in the architecture and function of silenced chromatin in *Saccharomyces cerevisiae*.** *Mol Cell Biol* 2010, **30**:626-639.

17. Tanghe A, van Dijck P, Thevelein JM: **Determinants of freeze tolerance in microorganisms, physiological importance, and biotechnological applications.** *Adv Appl Microbiol* 2003, **53**:129-176.
18. Aguilera J, Ranz-Gil F, Prieto JA: **Cold response in *Saccharomyces cerevisiae*: new functions for old mechanisms.** *FEMS Microbiol Rev* 2007, **31**:327-341.
19. Higgins VJ, Bell PJ, Dawes IW, Attfield PV: **Generation of a novel *Saccharomyces cerevisiae* strain that exhibits strong maltose utilization and hyperosmotic resistance using nonrecombinant techniques.** *Appl Environ Microbiol* 2001, **67**:4346-4348.
20. Aguilera J, Andreu P, Ranz-Gil F, Prieto JA: **Adaptive evolution of baker's yeast in a dough-like environment enhances freeze and salinity tolerance.** *Microb Biotechnol* 2010, **3**:210-221.
21. Uesono Y, Toh-E A: **Transient inhibition of translation initiation by osmotic stress.** *J Biol Chem* 2002, **277**:13848-13855.
22. Panadero J, Hernández-López MJ, Prieto JA, Ranz-Gil F: **Overexpression of the calcineurin target *CRZ1* provides freeze tolerance and enhances the fermentative capacity of baker's yeast.** *Appl Environ Microbiol* 2007, **73**:4824-31.
23. Hernández-López MJ, Prieto JA, Ranz-Gil F: **Osmotolerance and leavening ability in sweet and frozen sweet dough. Comparative analysis between *Torulopsis delbrueckii* and *Saccharomyces cerevisiae* baker's yeast strains.** *Antonie Van Leeuwenhoek* 2003, **84**:125-134.
24. Wolfe J, Bryant G: **Freezing, crying, and/or vitrification of membrane-solute-water systems.** *Cryobiology* 1999, **39**:103-129.
25. Decottignies A, Goffeau A: **Complete inventory of the yeast ABC proteins.** *Nat Genet* 1997, **15**:137-145.
26. Lenssen E, James N, Pedruzzi I, Dubouloz F, Camerani E, Bisig R, Mailet L, Werner M, Roosen J, Petrovic K, Winderickx J, Collart MA, De Virgilio C: **The Ccr4-Not complex independently controls both Msn2-dependent transcriptional activation--via a newly identified Glc7/Bud14 type I protein phosphatase module--and TFIID promoter distribution.** *Mol Cell Biol* 2005, **25**:488-498.
27. Martínez-Pastor MT, Marchler G, Schüller C, Marchler-Bauer A, Ruis H, Estruch F: **The *Saccharomyces cerevisiae* zinc finger proteins Msn2p and Msn4p are required for transcriptional induction through the stress response element (STRE).** *EMBO J* 1996, **15**:2227-2235.
28. Lenssen E, Azzouz N, Michel A, Landrieux E, Collart MA: **The Ccr4-not complex regulates Skn7 through Srb10 kinase.** *Eukaryot Cell* 2007, **6**:2251-2259.
29. Lee DG, Bell SP: **Architecture of the yeast origin recognition complex bound to origins of DNA replication.** *Mol Cell Biol* 1997, **17**:7159-7168.
30. Ramachandran L, Burhans DT, Laun P, Wang J, Liang P, Weinberger M, Wissing S, Jarolim S, Suter B, Madeo F, Breitenbach M, Burhans WC: **Evidence for ORC-dependent repression of budding yeast genes induced by starvation and other stresses.** *FEMS Yeast Res* 2006, **6**:763-776.
31. Malavé TM, Dent SY: **Transcriptional repression by Tup1-Ssn6.** *Biochem Cell Biol* 2006, **84**:437-443.
32. Ito H, Jukuda K, Murata K, Kimura A: **Transformation of intact yeast cells treated with alkali cations.** *J Bacteriol* 1983, **153**:163-168.
33. Zwietering MH, Jongenburger I, Rombouts FM, van 't Riet K: **Modeling of the Bacterial Growth Curve.** *Appl Environ Microbiol* 1990, **56**:1875-1881.
34. Gietz RD, Sugino A: **New yeast-*Escherichia coli* shuttle vectors constructed with *in vivo* mutagenized yeast genes lacking six-base pair restriction enzymes.** *Gene* 1988, **74**:527-534.
35. Alberola TM, García-Martínez J, Antúnez O, Viladevall L, Barceló A, Ariño J, Pérez-Ortín JE: **A new set of DNA microchips for the yeast *Saccharomyces cerevisiae*: features and uses.** *Int Microbiol* 2004, **7**:199-206.
36. Eilbeck K, Lewis SE, Mungall CJ, Yandell M, Stein L, Durbin R, Ashburner M: **The Sequence Ontology: a tool for the unification of genome annotations.** *Genome Biol* 2005, **6**:R44.
37. Huang DW, Sherman BT, Tan Q, Kir J, Liu D, Bryant D, Guo Y, Stephens R, Baseler MW, Lane HC, Lempicki RA: **DAVID Bioinformatics Resources: Expanded annotation database and novel algorithms to better extract biology from large gene lists.** *Nucleic Acids Res* 2007, **35**:W169-175.

doi: 10.1186/1475-2859-9-56

**Cite this article as:** Pérez-Torrado *et al.*, Global expression studies in baker's yeast reveal target genes for the improvement of industrially-relevant traits: the cases of *CAF16* and *ORC2* *Microbial Cell Factories* 2010, **9**:56

**Submit your next manuscript to BioMed Central and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)

