

CORRECTION

Differences in color between early gastric cancer and cancer-suspected non-cancerous mucosa on linked color imaging

Hirimitsu Kanzaki, Yoshiro Kawahara, Takuya Satomi et al.


Endoscopy International Open 2023; 11: E90–E96.

DOI: 10.1055/a-1989-6643

published online: 2023-01-19

In the above-mentioned article ► **Fig. 3** was corrected.

This was corrected in the online version on 2 February 2023.



► **Fig. 3** Plot on the ΔL and Δb dimension. **a** Plot of all cases on ΔL and Δb dimensions. Many of the cases in the area of the first quadrant, which had positive scores for both ΔL and Δb , were EGC. **b** The accuracy, sensitivity, specificity, PPV and NPV for EGC in the 1st quadrant, which had both positive ΔL and Δb values, were 54.7%, 39.6%, 88.2%, 88.2%, and 39.6%, respectively. In contrast, many of the cases in the area of the third quadrant, which had negative scores for both ΔL and Δb , were CSM. The accuracy, sensitivity, specificity, PPV and NPV for CSM were 71.8%, 23.7%, 84.1%, 64.3%, and 73.3%, respectively. EGC, early gastric cancer; CSM, cancer-suspected non-cancerous mucosa; PPV, positive predictive value; NPV, negative predictive value.