

STUDY PROTOCOL

Open Access



Photobiomodulation versus light-emitting diode (LED) therapy in the treatment of temporomandibular disorder: study protocol for a randomized, controlled clinical trial

Luciana G. Langella, Paula F. C. Silva, Larissa Costa-Santos, Marcela L. L. Gonçalves, Lara J. Motta, Alessandro M. Deana, Kristianne P. S. Fernandes, Raquel A. Mesquita-Ferrari and Sandra Kalil Bussadori*

Abstract

Background: Temporomandibular disorder (TMD) is described as a subgroup of orofacial pain with a set of signs and symptoms that involve the temporomandibular joint, masticatory muscles, ears, and neck. TMD can occur unilaterally or bilaterally and approximately 70% of the population is affected with at least one sign. The disorder progresses with orofacial pain, muscle pain involving the masticatory and cervical muscles, joint noises (clicks and pops), joint block, mandibular dysfunction, and headache. The etiology can be abnormal occlusion and/or posture, trauma involving local tissues, repetitive microtrauma, parafunctional habits, and an increase in emotional stress. Studies have demonstrated that phototherapy is an efficient option for the treatment of TMD, leading to improvements in pain and orofacial function.

Methods: The aim of the proposed study is to compare the effects of two sources of photobiomodulation in individuals with TMD. A randomized, controlled, double-blind, clinical trial is proposed, which will involve 80 individuals aged 18–65 years allocated to either a laser group or light-emitting diode (LED) group submitted to 12 sessions of phototherapy. The Research Diagnostic Criteria for TMDs will be used to evaluate all participants. Pain will be measured using the visual analog scale and maximum vertical mandibular movement will be determined with the aid of digital calipers.

Discussion: This study compares the effects of two modalities of laser therapy on the pain and orofacial function of patients with TMD dysfunction. Photobiomodulation and LED therapy are treatment options for reducing the inflammatory process and pain as well as inducing the regeneration of the target tissue.

Trial registration: ClinicalTrials.gov, NCT03257748. Registered on 8 August 2017.

Keywords: Temporomandibular disorder, Photobiomodulation, Light-emitting diode (LED)

* Correspondence: sandra.skb@gmail.com
Nove de Julho University, 235/249 Vergueiro Street, Liberdade, São Paulo
01504-001, Brazil

Background

The temporomandibular joints link the lower jaw (mandible) to the temporal bone of the skull, which is immediately in front of the ear on each side of the head. The flexibility of the joints allows the mandible to move up and down as well as from side to side [1, 2].

Temporomandibular disorder (TMD) is described as a subgroup of orofacial pain with a set of signs and symptoms that involve the temporomandibular joint, masticatory muscles, ears, and neck either unilaterally or bilaterally [2–6]. The signs and symptoms include orofacial pain [2, 8], muscle pain involving the masticatory and cervical muscles, joint noises (clicks and pops), joint block, mandibular dysfunction [8–11], and headache [2, 7, 8]. The etiology can be abnormal occlusion and/or posture, trauma involving local tissues, repetitive microtrauma, parafunctional habits, and an increase in emotional stress [2, 7, 8, 10, 12]. With regard to prevalence, studies report that approximately 70% of the population can have at least one sign of TMD [13–15], but only 10% seek treatment [13, 15]. Pain is the most common symptom and can affect quality of life due to limitations regarding actions such as laughing, eating, speaking, and yawning [3, 10, 14].

Due to the pain and functional limitations inherent to TMD, phototherapy has been investigated and has proven to be an important treatment option for this disorder [1, 4, 5, 7–9, 11, 13, 16, 17]. Photobiomodulation and light-emitting diode (LED) therapy are treatment options for reducing pain and inflammatory processes as well as inducing the regeneration of the target tissue [18–20]. Both types of phototherapy have been suggested for the treatment of inflammatory joint conditions due to the absence of side effects, which are common with the use of non-steroidal anti-inflammatory drugs (NSAIDs) [20]. Studies have reported that lasers and LEDs operating with similar parameters produce equivalent effects [21, 22]. Different theories have been proposed to explain the analgesic action of phototherapy, but no consensus has been reached. Some authors report effects such as biomodulation, the capacity to stimulate cell division, analgesia, modulation of the production of β -endorphins, as well as increases in cortisol and protein synthesis [9, 16–18, 23–27].

Drugs are widely employed for the treatment of pain, but are expensive, which generates high costs in the public health realm, and have limited effectiveness as well as potentially serious side effects, especially when administered for prolonged periods of time. Thus, non-pharmacological, non-invasive resources for pain relief are of the utmost importance and have been widely investigated in the scientific literature.

Hypothesis

Based on previous studies conducted by different research groups, the proposed study will test the hypothesis that photobiomodulation in the form of low intensity laser

compared to LED decreases pain and improves temporomandibular joint function, thereby accelerating the return to activities of daily living.

Objective

The aim of the proposed study is to analyze the effect of laser therapy and LED therapy on pain and function in patients with TMD and its association with neck pain.

Expected results

At the end of the study, we expect to find a reduction in pain in the temporomandibular joint and temporal muscle region, and improvement in the function of the temporomandibular joint through photobiomodulation and LED therapy in patients with TMD, with a consequent improvement in their quality of life.

Methods / design

Study design


This protocol is presented in accordance with the 2013 SPIRIT (Standard Protocol Items: Recommendations for Interventional Trials) Statement (see Additional file 1 for the populated SPIRIT Checklist and Fig. 1 for the trial schedule of enrollment, interventions, and assessments in accordance with recommended SPIRIT figure), which was developed to provide guidance in the form of a checklist of recommended items to include in a clinical trial protocol to help improve its content and quality. A randomized, controlled, clinical trial will be conducted at the dentistry and physical therapy clinics of University Nove de Julho (Brazil) involving individuals with TMD distributed between two groups. This protocol study received approval from the Human Research Ethics Committee of University Nove de Julho (São Paulo, Brazil) under process number 1.706.160. All potential participants will receive clarifications regarding the objectives and procedures and those who agree to participate voluntarily will sign a statement of informed consent, as stipulated in Resolution 466/2012 of the Brazilian National Board of Health.

The patients will be randomly allocated to either the photobiomodulation or LED group using a set of opaque, sealed, numbered envelopes to ensure concealment. Each envelope will contain a card stipulating to which group the volunteer will be allocated. The participants will then be sent for evaluations (Fig. 2).

Participants

Male and female individuals aged 18–65 years with a clinical diagnosis of TMD based on the Research Diagnostic Criteria for Temporomandibular Disorders (RDC/TMD) will be selected to participate in the study. No restrictions will be imposed with regard to race.

The volunteers will be informed of all procedures and sign a statement of informed consent before the onset of the study.



Inclusion criteria

- Clinical diagnosis of TMD, separated by the degrees of this function based on the Research Diagnosis Criteria for Temporomandibular Disorders, as shown in Table 1;
- Age 18–65 years;
- Presence of at least 20 functional teeth, at least three being molars.

Exclusion criteria

- Dentofacial anomalies;
- Currently in orthodontic or orthopedic treatment for the jaw;
- Currently undergoing psychological treatment;
- Use of muscle relaxant or anti-inflammatory agent.

Sample composition

The sample size was calculated based on studies found in the literature using maximum vertical

mandibular movement as the basis, with a standard deviation of 9.82, a 5% significance level ($p < 0.05$), and 80% test power. The sample was determined to be 80 individuals, who will be allocated to two groups.

Step 1 – Recruitment for the selection of patients who will participate in the study. The patients will receive clarifications regarding the objectives and procedures and those who agree to participate voluntarily will sign a statement of informed consent.


Step 2 – The selected patients will receive an envelope containing a card stipulating to which group they will be allocated. The patients will not be aware of the content of the envelope.

Interventions

Phototherapy

The volunteers will be allocated to two groups:

The sessions will be held in a reserved, noise-free room. The volunteer will be positioned with the



Frankfurt plane parallel to the ground. The devices will be covered in disposable plastic wrap to avoid cross-contamination and for hygiene reasons. The operator will wear suitable protective clothing. The skin at the site to be irradiated will have previously been cleaned with 70% alcohol. Both the operator and volunteer will use protective eyewear.

Both groups (laser and LED) will be submitted to eight sessions of phototherapy held twice a week for four weeks, during which only the researcher in charge of programming the phototherapy device will be aware of which treatment is being employed. However, the programmer will not participate in the execution of the treatments, evaluations, or data analysis.

Table 1 Diagnostic subgroups according to RDC/TMD

Group I Muscle diagnoses	
Ia	– myofascial pain
Ib	– myofascial pain with limited opening
Group II Disk displacement	
IIa	– disk displacement with reduction
IIb	– disk displacement without reduction with limited opening
IIc	– disk displacement without reduction without limited opening
Group III Arthralgia, osteoarthritis, and osteoarthritis	
IIIa	– arthralgia
IIIb	– osteoarthritis of temporomandibular joint
IIIc	– osteoarthritis of temporomandibular joint

Phototherapy will be applied:

- Temporomandibular joint region: around the temporomandibular joint;
- Masseter and temporal muscles: on the masseter and the anterior temporal.

The number of irradiated points will depend on the treatment. Patients in the laser group will have three points irradiated, while those in the LED group will have 36 points irradiated. The parameters that shall be used can be found in Table 2.

Outcome measures**Reliability of examiner**

The initial evaluations will be performed by an examiner who will be duly trained by a “gold standard” evaluator following an evaluation protocol. To confer greater reliability to the results, the examiner will be blinded to the allocation of the participants to the different groups. A second examiner will perform the re-evaluations following the treatment protocols and will also be blinded to the allocation of

the participants. This examiner will also be duly trained following the evaluation protocol.

Evaluations

- Research Diagnostic Criteria for Temporomandibular Disorders (RDC/TMD): Potentially eligible patients will be diagnosed using the RDC/TMD, which consists of a clinical exam involving the palpation of the temporomandibular joints an analysis of the mandible with the aid of digital calipers to measure vertical and horizontal movements and a stethoscope to evaluate joint sounds. Symptoms such as headache, fatigue and difficulty when chewing, bruxism, psychological state, and parafunctional habits will then be investigated.
- Visual analog scale: This scale will be used for the assessment of pain and consists of a 10-cm line with 0 (absence of pain) printed at one end and 10 (debilitating pain) printed at the other end. The participants will be asked to mark a place on the line that represents their current pain intensity. The researcher will subsequently use a ruler to register the distance from zero to obtain a numeric representation of the pain level. These procedures will be performed before and after treatment.
- Maximum vertical mandibular movement: The volunteer will be instructed to open his/her mouth as wide as possible. Maximum vertical mandibular movement will be measured as the distance between the maxillary and mandibular central incisors determined with the aid of digital calipers. The volunteer will then be instructed to exert pressure on the mandibular teeth with the mouth open and move the mandible to the right and left for the determination of excursion (distance between upper and lower mid points). These procedures will be performed before and after treatment.

Table 2 Parameters of phototherapy

Parameters	Laser group	LED group
Wavelength (nm)	780	780
Target area (cm ²)	130	130
Spot area (cm ²)	43	3.58
Power (mW)	60	5
Irradiation time (s)	600	600
Points irradiated	3	36
Energy per point (J)	36	3
Total energy (J)	108	108
Radiant exposure (J/cm ²)	0.8	0.8
Irradiance (mW/cm ²)	1.38	1.38

Statistical analysis

The results will be tested for normality using the Kolmogorov–Smirnov test. Parametric data will be submitted to analysis of variance (ANOVA) with the Tukey–Kramer post hoc test. Non-parametric data will be submitted to Friedman’s test. The level of significance will be set to 5% ($p < 0.05$).

Monitoring

Phototherapy has been employed for 30 years and there are no reports of side effects in the literature.

The following are the expected benefits: reductions in signs of inflammation, pain and muscle tension of the masticatory, and cervical muscles as well as an improvement

in orofacial function, enabling greater mobility and functionality of the temporomandibular joints.

Discussion

Pain and functional limitations are inherent to temporomandibular dysfunction, and studies indicate that 70% of the population is affected with at least one TMD signal [14–16]. Medications are commonly used to treat the pain, but they have a high cost and potential side effects, especially in the administration of NSAIDs [20] and long-term use [27, 28]. Therefore, there is a need for studies that evaluate the use of non-pharmacological methods for the treatment of this dysfunction.

Phototherapy has the potential to reduce pain and improve temporomandibular joint function in TMD patients, with consequent improvement in their quality of life. Photobiomodulation and LED therapy are treatment options for reduction of the inflammatory process and pain, besides inducing the regeneration of the target tissue [18–20].

The efficacy of phototherapy for the treatment of TMD has been proven by several authors [1, 4, 9, 11, 17], but there are few controlled clinical studies comparing photobiomodulation performance with LED therapy in patients with TMDs.

Trials status

This study is currently recruiting participants.

Additional file

Additional file 1: SPIRIT 2013 Checklist: Recommended items to address in a clinical trial protocol and related documents. (DOC 121 kb)

Abbreviations

ANOVA: Analysis of variance; LED: Light-emitting diode; RDC/TMD: Research Diagnostic Criteria for Temporomandibular Disorders; TMD: Temporomandibular disorder

Acknowledgements

The authors gratefully thank Nove de Julho University for the technological support it will provide.

Funding

This research received no specific funding agency in the public, commercial, or not-for-profit sectors.

Availability of data and materials

Not applicable.

Authors' contributions

LGL developed the design and methodology, developed the analysis plan, and reviewed and commented on drafts of the protocol of the paper. PFCS, LC-S, and MLLG reviewed and commented on drafts of the protocol of the paper. AMD, LJM, KPS, and RAM-F contributed to drafting the article, and read and approved the final manuscript. SKB developed the original concept of the trial, drafted the original protocol, and adapted the trial proposal as a protocol paper. All authors read and approved the final manuscript.

Ethics approval and consent to participate

This protocol study received approval from the Human Research Ethics Committee of University Nove de Julho (São Paulo, Brazil) under process

number 1.706.160. All potential participants will receive clarifications regarding the objectives and procedures and those who agree to participate voluntarily will sign a statement of informed consent, as stipulated in Resolution 196/96 of the Brazilian National Board of Health.

Consent for publication

Written informed consent for publication of their clinical details and/or clinical images will be obtained from the patient/parent/guardian/relative of the patient. A copy of the consent form will be available for review by the Editor of this journal.

Competing interests

The authors declare that they have no competing interests.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Received: 18 October 2017 Accepted: 2 January 2018

Published online: 26 January 2018

References

- Dostolová T, Hlinakova P, Kasparova M, Rehacek A, Vavrichova L, Navratil L. Effectiveness of physiotherapy disorders and GaAlAs laser in the management of temporomandibular joint. *Photomed Laser Surg.* 2012;30(5):275–80.
- Biasotto-Gonzalez DA. Abordageminterdisciplinar das disfunçõestemporomandibulares. Barueri: Manole; 2005.
- de Godoy CHL, Silva PFC, Araujo DS, Motta LJ, Biasotto-Gonzalez DA, Politti F, et al. Evaluation of effect of low-level laser therapy on adolescents with temporomandibular disorder: study protocol for a randomized controlled trial. *Trials.* 2013;14:229–34.
- Pereira TS, Flecha OD, Guimarães RC, de Oliveira D, Botelho AM, Glória JCR, et al. Efficacy of red and infrared lasers in treatment of temporomandibular disorders—a double-blind, randomized, parallel clinical trial. *Cranio.* 2014;32(1):51–6.
- Salmos-Brito JAL, Menezes RF, Teixeira CEC, Gonzaga RKM, Rodrigues BHM, Braz R, et al. Evaluation of low-level laser therapy in patients with acute and chronic temporomandibular disorders. *Lasers Med Sci.* 2013;28:57–64.
- IASP. Classification of chronic pain: descriptors of chronic pain syndromes and definitions of pain terms. 2nd ed. Seattle: IASP; 1994.
- Mazzetto MO, Carrasco TG, Bidinelo EF, de Andrade Pizzo RC, Mazzetto RG. Low intensity laser application in temporomandibular disorders: a phase I double-blind study. *Cranio.* 2007;25:186–92.
- Andrade TNC, Frare JC. Estudocomparativo entre osefeitos de técnicas de terapia manual isoladas e associadas à laserterapia de baixapotênciasobre a dorempacientes com disfunção temporomandibular. *Rev Bras Fisioter.* 2008; 56(3):287–95.
- Carrasco TG, Mazzetto MO, Mazatto RG, Mestriner Jr W. Low intensity laser therapy in temporomandibular disorder: a phase II double-blind study. *Cranio.* 2008;26(4):274–81.
- Gonçalves DA, Dal Fabbro AL, Campos JA, Bigal ME, Speciali JG. Symptoms of temporomandibular disorders in the population: an epidemiological study. *J Orofac Pain.* 2010;24:270–78.
- Mazzetto MO, Hotta TH, de Andrade Pizzo RC. Measurements of jaw movements and TMJ pain intensity in patients treated with GaAlAs laser. *Braz Dent J.* 2010;21(4):356–60.
- Sarlani E. Diagnosis and treatment of orofacial pain. *Braz J Oral Sci.* 2003;2:283–90.
- Kulekioglu S, Sivrioglu K, Ozcan O, Parlak M. Effectiveness of low-level laser therapy in temporomandibular disorder. *Scand J Rheumatol.* 2003; 32:114–18.
- Dworkin SF, Huggins KH, LeResche L, Von Korff M, Howard J, Truelove E, et al. Epidemiology of signs and symptoms in temporomandibular disorders: clinical signs in cases and controls. *J Am Dent Assoc.* 1990;120(3):273–81.
- Nassif NJ, Al-Salleeh F, Al-Admawi M. The prevalence and treatment needs of symptoms and signs of temporomandibular disorders among Young adult males. *J Oral Rehabil.* 2003;30:944–50.
- Frare JC, Nicolau RA. Análiseclínica do efeito da fotobiomodulação laser (GaAs-904 nm) sobre a disfunção temporomandibular. *Rev Bras Fisioter.* 2008;12(1):37–42.
- Petrucci A, Sgolastra F, Gatto R, Mattei A, Monaco A. Effectiveness of low level laser therapy in temporomandibular disorders: a systematic review and meta-analysis. *J Orofac Pain.* 2011;25(4):298–307.

18. Gynther GW, Dijkgraaf LC, Reinholt FP, Holmlund AB, Liem RS, de Bont LG. Synovial inflammation in arthroscopically obtained biopsy specimens from the temporomandibular joint: A review of the literature and a proposed histologic grading system. *J Oral Maxillofac Surg.* 1998;56:1281–6.
19. Pallotta RC, Bjordal JM, Frigo L, Leal Junior EC, Teixeira S, Marcos RL, et al. Infrared (810-nm) low-level laser therapy on rat experimental knee inflammation. *Lasers Med Sci.* 2011;27:71–8.
20. Panhoca VH, Lizarelli Rde F, Nunez SC, Pizzo RC, Grecco C, Paolillo FR, et al. Comparative clinical study of light analgesic effect on temporomandibular disorder (TMD) using red and infrared led therapy. *Lasers Med Sci.* 2015;30(2):815–22.
21. Peter-Szabo M, Kekesi G, Nagy E, Sziver E, Benedek G, Horvath G. Quantitative characterization of a repeated acute joint inflammation model in rats. *Clin Exp Pharmacol Physiol.* 2007;34:520–26.
22. Yamazaki Y, Ren K, Shimada M, Iwata K. Modulation of paratrigeminal nociceptive neurons following temporomandibular joint inflammation in rats. *Exp Neurol.* 2008;214:209–18.
23. Laakso E, Cramond T, Richardson C, Galligan J. Plasma ACTH and beta-endorphin levels in response to low-level laser therapy (LLLT) for myofascial trigger points. *Laser Ther.* 1994;6:133–42.
24. Vinck EM, Cagnie BJ, Cornelissen MJ, Declercq HA, Cambier DC. Increased fibroblast proliferation induced by light emitting diode and low power laser irradiation. *Lasers Med Sci.* 2003;18:89–94.
25. Laakso E, Cabot PJ. Nociceptive scores and endorphin-containing cells reduced by low-level laser therapy (LLLT) in inflamed paws of Wistar rat. *Photomed Laser Surg.* 2005;23:32–5.
26. Brugnera JRA. Biomodulatory effect of laser therapy-clinical indications. *Dentistry Braz Dent J.* 2005;15(Suppl):60.
27. Bjordal JM, Ljunggren AE, Klovning A, Slordal L. Non-steroidal anti-inflammatory drugs, including cyclo-oxygenase-2 inhibitors, in osteoarthritic knee pain: meta-analysis of randomised placebo-controlled trials. *BMJ.* 2004;329:1317.
28. Bjordal J, Johnson MI, Lopes-Martins RA, Bogen B, Chow R, Ljunggren AE. Short-term efficacy of physical interventions in osteoarthritic knee pain: A systematic review and meta-analysis of randomised placebo controlled trials. *BMC MusculoskeletDisord.* 2007;8:51.

Submit your next manuscript to BioMed Central and we will help you at every step:

- We accept pre-submission inquiries
- Our selector tool helps you to find the most relevant journal
- We provide round the clock customer support
- Convenient online submission
- Thorough peer review
- Inclusion in PubMed and all major indexing services
- Maximum visibility for your research

Submit your manuscript at
www.biomedcentral.com/submit

