

Short report

Open Access

Mental health measures of anxiety and depression in patients with retinal detachment

Maneli Mozaffarieh¹, Stefan Sacu¹, Thomas Benesch² and Andreas Wedrich^{*3}

Address: ¹Department of Ophthalmology, Medical University of Vienna, Austria, ²Institute of Medical Statistics, Medical University of Vienna, Austria and ³Department of Ophthalmology, Medical University of Graz, Austria

Email: Maneli Mozaffarieh - mmozaffarieh@uhbs.ch; Stefan Sacu - stefan.sacu@meduniwien.ac.at; Thomas Benesch - thomas.benesch@meduniwien.ac.at; Andreas Wedrich* - andreas.wedrich@meduni-graz.at

* Corresponding author

Published: 19 July 2007

Received: 14 March 2007

Accepted: 19 July 2007

Clinical Practice and Epidemiology in Mental Health 2007, 3:10 doi:10.1186/1745-0179-3-10

This article is available from: <http://www.cpementalhealth.com/content/3/1/10>

© 2007 Mozaffarieh et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

In this study, the researchers examined anxious and depressive symptoms of patients with rhegmatogenous retinal detachment (RRD) prior to and up to year after retinal detachment surgery. One hundred and thirteen (113) patients with RRD took part in this prospective longitudinal study. Anxiety and depression were evaluated using the Hospital Anxiety and Depression Scale (HADS). Visual acuity (VA) results and HADS scores of all participants were recorded prior to and 3, 6 and 12 months after retinal detachment surgery. Pearson correlation analysis showed a significant association between the patients' VA and HADS psychological scores both prior to and three months after surgery, regardless of the type of surgery performed. Psychological distress is a significant problem associated with retinal detachments that requires more attention.

Background

Retinal detachment (RD) is a condition which affects approximately one in 300 patients in the course of a lifetime and more often requires urgent surgical repair [1]. Despite the knowledge on visual outcomes of the patients with retinal detachment, and the multiple reports on outcomes after various surgical interventions [1-6], little is known on the effect of retinal detachment on the patients' psychological status.

It is recognized that psychological symptoms accompany visual loss [7,8]. However, most research in this domain has focused on blind individuals and only little information is available concerning psychological distress among visually impaired (but not blind individuals). In a study done on pre-school children treatment for unilateral vis-

ual impairment (amblyopia) was shown to be associated with some degree of psychological distress [9]. Reports suggest that a large number of people with advanced age-related macular degeneration have depressive disorders leading to serious consequences for their quality of life [10-12]. The impact of ocular surgery on patient's psychological reactions is also limited to certain surgical interventions. Patients for cataract surgery, for example, suffer from psychological disorders of anxiety and depression after surgery, in particular if their vision has not improved significantly [13]. A study on patients with uveal melanoma reported a prevalence of mental disorders of anxiety and depression after eye surgery [14]. Unfortunately, to date, there have been no studies investigating whether RD has any effect on the patients' psychological status. The lack of familiarity with the psychological con-

sequences of acute visual loss secondary to RD prevents most physicians from supplying these patients with optimal additional treatment and support. As a result, the psychological symptoms which accompany RD are overlooked in many patients.

The purpose of this article is to examine the psychological reactions of patients with RD prior to and after the successful surgical intervention. To our knowledge this is the first study seeking to find the impact of RD on the patients' psychological status.

Methods

Participants

One hundred and thirteen patients (113 eyes) with RD were included in this prospective longitudinal clinical study. The study was performed at the Department of Ophthalmology at the Vienna General Hospital (Medical University of Vienna) between 2000 and 2003. The inclusion criteria included the identification of rhegmatogenous detachment by a vitreoretinal specialist. Patients with glaucoma, uveitis or any other coexisting visually limiting condition (e.g. exudative age-related macular degeneration) or RRD with a proliferative vitreoretinopathy stage 3 or 4 were excluded from the study. Additionally, patients with potentially confounding events (e.g. death of family member) that may have influenced overall psychological stress were excluded from the study (1 patient). The operation was considered anatomically successful if the retina was still re-attached at least six months after the operation. Patients with re-detachment during the study period for second surgical repair were excluded from the study. Written informed consent was obtained from all patients after their involvement in the study had been explained in detail.

Surgical Procedure

All surgery was performed by an experienced vitreoretinal surgeon. The surgeon (AW) involved in the study decided the surgical approach in each case according to his experience. Patients underwent either a vitrectomy with fluid-gas exchange or scleral buckling surgery under general anaesthesia.

Visual Acuity

Visual acuity was recorded as the best corrected visual acuity in the study eye and the fellow eye using a Snellen chart prior to and three, six and twelve months after surgery. The Snellen fractions were converted to a log scale (Log MAR) as previously described [15].

The Hospital Anxiety and Depression Scale (HADS)

The hospital anxiety and depression scale was developed to measure anxiety and depressive symptoms among patients with somatic diseases [16]. The reliability and

validity of the HADS have been tested in a number of international studies [17]. The HADS contains 14 items and consists of two subscales: 7 items for anxiety and 7 for depression. Each item is rated on a four-point scale (scored 0–3), giving maximum scores of 21 for anxiety and depression. This questionnaire was administered to the patients prior to, and 3, 6 and 12 months after surgery.

Statistical Analysis

Using SPSS 12.0 for Windows, the descriptive and inferential data analyses were performed. Mean visual acuity and HADS psychological scores were evaluated using a paired t-test for change from baseline. Analysis of variance (ANOVA) was used for between types of surgical treatments evaluation (vitrectomy/buckle surgery) and for subgroup analysis based on gender of patients (female/male). For nonparametric data, the Mann-Whitney U test was applied. Univariate associations between mean visual acuity and HADS psychological scores were evaluated using the Pearson correlation test. The probabilities were all corrected for multiple testing, applying the Bonferroni-Holm step-down test ($\alpha = 0.05$).

Results

Descriptive data of the study population

A total of 113 patients with RD were eligible according to our inclusion criteria. Of these ten patients were not included (5 did not complete questionnaires, 4 did not attend their follow-up's, and 1 lived abroad). Follow-up data were available for the remaining 103 patients who were studied prospectively. The mean age of the patients 56.5 years (SD, ± 11.6 years; range: 27–81). Fifty-two (50%) of the patients were male and 51 (50%) were female. 65 of the patients (63%) had vitrectomy with fluid-gas exchange and 38 (37%) had buckle surgery.

Changes from baseline for all parameters

The mean best corrected visual acuity (Log MAR) of the patients prior to and 3, 6 and 12 months after surgery, was 1.3 ± 0.2 , 0.62 ± 0.06 , 0.43 ± 0.04 and 0.26 ± 0.06 , respectively ($p < 0.05$ to baseline, table 1). The mean best corrected visual acuity of the patients fellow-eye prior to and 3, 6 and 12 months after surgery, was 0.21 ± 0.09 , 0.19 ± 0.07 , 0.20 ± 0.06 , and 0.19 ± 0.06 , respectively ($p > 0.05$ to baseline).

The cut-off values for anxious and depressive moods are depicted in table 2. Patients were significantly more psychologically affected prior to than after detachment surgery (HADS anxiety and depression, $p < 0.01$). More than half of the patients (65%) experienced 'probable' levels of anxiety, prior to surgery. This value decreased to 59.2%, 12 months after RRD surgery. Percent of patients psychologically affected and mean HADS scores are presented in table 2.

Table 1: Visual acuity (Log MAR) and hospital anxiety and depression scores (HADS)

	Baseline	3 months after surgery		6 months after surgery		12 months after surgery	
	Mean ± SD	Mean ± SD	P*	Mean ± SD	P*	Mean ± SD	P*
Study eye Log MAR	1.3 ± 0.2	0.62 ± 0.06	< 0.05	0.43 ± 0.04	< 0.01	0.26 ± 0.06	< 0.01
Fellow-eye Log MAR	0.21 ± 0.09	0.19 ± 0.07	> 0.2	0.2 ± 0.06	> 0.2	0.19 ± 0.06	> 0.2
Anxiety	18 ± 3	17 ± 3	< 0.01	16 ± 3	< 0.01	16 ± 3	< 0.01
Depression	16 ±	13 ± 2	< 0.01	8 ± 3	> 0.1	5 ± 2	> 0.1

* for change from baseline

Means and standard deviations (SD) of visual acuity and Hospital Anxiety and Depression Scores (HADS) at the 4 points of assessment.

Subgroup analysis

No statistically significant difference was found between the type of surgery performed (vitrectomy or buckle) and HADS psychological scores at any follow-up time (p > 0.05). The patients' gender had no statistically significant effect on HADS psychological scores (p = 0.31).

Association between the visual acuity and the psychological reactions

Study eye

There were strong correlations between the visual acuity and HADS psychological scores for depression and anxiety 3 months after surgery ($r_{3A} = 0.68$, $r_{3D} = 0.74$, $p < 0.01$). Moderate correlations were found between the visual acuity and HADS psychological scores for anxiety both six and twelve after surgery ($r_{6A} = 0.45$, $r_{12A} = 0.39$, $p < 0.01$). No significant association was found between the visual acuity and HADS depression scores of the patients' either at 6 or 12 months ($r_{6D} = 0.35$, $r_{12D} = 0.26$ $p > 0.05$).

Fellow-eye

Interestingly, there was a significant association between the patients' fellow-eye visual acuity and HADS anxiety scores both 6 and 12 months after surgery ($r_6 = 0.86$, $r_{12} = 0.74$, $p < 0.01$). No significant association was found

between patients' fellow-eye visual acuity and HADS depression scores either 6 or 12 months after surgery ($r_6 = 0.32$, $r_{12} = 0.21$, $p > 0.1$).

Discussion

This investigation reports the long-term impact of psychological affects in patients with retinal detachment. The assessment and follow-up of the patients' psychological characteristics were measured by means of the reliable and valid HADS questionnaire. The reliability of this measure has been reported in numerous previous studies making its use internationally applicable [16,17]. We then further sought to observe the relationships between the patients' psychological status and visual acuity prior to and after surgery.

RRD is usually an acute condition for which surgery is generally considered mandatory in order to prevent severe and permanent visual loss. We found that more than half of our patients (65%) experienced probable levels of anxiety, prior to surgery. This value decreased gradually to 59%, 12 months after surgery. Patients were most probably relieved to have had surgery. Still, the prevalence of anxiety, demands our attention as to why and what we could possibly do to alleviate this symptom.

Table 2: Hospital anxiety and depression scores (HADS) of patients prior to and after retinal detachment surgery

	Prior to surgical treatment N (%)	3 months after treatment N (%)	6 months after treatment N (%)	12 months after treatment N (%)
Anxiety				
0-7	6 (5.8)	9 (8.7)	11 (10.6)	13 (12.6)
8-10	67 (65.0)	60 (58.2)	70 (67.9)	61 (59.2)
> 10	30 (29.1)	29 (28.1)	27 (26.2)	29 (28.1)
Depression				
0-7	31 (30.1)	57 (55.3)	72 (69.9)	82 (79.6)
8-10	52 (50.5)	31 (30.1)	22 (21.3)	16 (15.5)
> 10	20 (19.4)	15 (14.5)	9 (8.7)	5 (4.8)

Number (%) of patients scoring in categories 'No case' (< 7), 'Possible clinical case' (8-10) and 'Clinical case' (> 10) on the Hospital Anxiety and Depression Subscales. A score of 0 to 7 for either subscale of depression or anxiety is regarded as being in the normal range, a score of 8 to 10 indicates probable presence of the mood disorder, and a score larger of 10 or larger is suggestive of the clinical presence of the mood disorder.

Retinal detachment is accompanied by a certain degree of functional loss which is regained gradually [18]; this may explain the reason for the high HADS anxiety scores. Ophthalmologists may support patients by acting as facilitators providing information about RD and the various surgical interventions, and by suggesting ways to ameliorate anxious symptoms. Patients may, for example, find it alleviating to share their experiences with others who were similarly affected or they may just simply need more reassurance to relieve their concerns.

Anxiety is a multidimensional response to stimuli in the person's environment, or a response to an internal stimulus resulting from a combination of general biological and individual psychological processes. Physicians should be aware that there may be a certain degree of anxiety associated with the fear of similar RRD occurrence in the fellow-eye which may also have substantive negative effects in the patients' overall status. Data from other studies suggest that giving patients' choices about, information, and responsibility for certain aspects of care has important implications for the quality of care and health outcomes [19-21]. Especially patients with RD should be adequately informed on all the consequences of surgery, the status of their fellow-eye and possibly be advised that they are capable of, and may take over daily routine activities.

Unfortunately obstacles such as resource and time constraints or the primary lack of knowledge that indeed patients with RD are psychologically affected, prevent us in our efforts to optimally treat patients. Evidence of interactions between physician and patient suggests that as the pressures to contain costs increase, physicians respond by increasing the volume of their practices with a corresponding decrease in time spent per patient [22]. Busy hospital practices and shorter office visits have been empirically linked with less effective patient behaviour, whereas more time with the physician has been associated with more favourable ratings of the clinical encounter. Our data suggest that more time with patients after surgery may be required to adequately discuss visual and psychological outcomes of vitreoretinal surgery.

A limitation in this study is the fact that this study includes data from patients operated on by one vitreoretinal surgeon at one center. Data from this patient pool may not reflect the outcomes of persons undergoing vitreoretinal surgical procedures by other surgeons at other centers. Another limitation is that although the HADS instrument has been reported to be a very reliable instrument, its reliability was not measured in the current study. HADS is a very useful first step of a screening which must be multiphasic. More data is therefore required to back-up our results firmly. An extensive clinical interview, by pro-

fessionals, would be an important and necessary second step. In our investigation we evaluated psychological responses of patients with uncomplicated cases of RD; cases requiring additional surgery, which may have been exposed to much more psychological stress, were not included in this study.

Further research should evaluate various psychosocial aspects of RRD and provide patients with suitable therapeutic interventions. If data from future research supports our results, then more attention must be paid to the psychological manifestations of acute visual impairment and to approaches which may improve psychological management to provide optimal care for patients.

Authors' contributions

MM carried out the study.

SS participated in the design of the study, drafted the study and contributed in the statistical analysis.

TB performed the statistical analysis.

AW conceived of the study, and participated in its design and coordination.

References

1. Alhassan MB, Rabi MM, Olongusua Y, Ahmed A: **Outcome of scleral buckling for primary rhegmatogenous retinal detachment in Nigeria.** *Med Sci Monit* 2005, **11**:CR589-CR593.
2. Martinez-Castillo V, Verdugo A, Boixadera A, Garcia-Arumi J, Corcostegui B: **Management of inferior breaks in pseudophakic rhegmatogenous retinal detachment with pars plana vitrectomy and air.** *Arch Ophthalmol* 2005, **123**:1078-1081.
3. Salam GA, Greene JM, Deramo VA, Tibrewala RK, Ferrone PJ, Fastenberg DM: **Retinal tears and retinal detachment as factors affecting visual outcome after cataract extraction complicated by posteriorly dislocated lens material.** *Retina* 2005, **25**:570-575.
4. Heimann H, Zou X, Jandek C, Kellner U, Bechrakis NE, Kreusel KM, Helbig H, Krause L, Schuler A, Bornfeld N, Foerster MH: **Primary vitrectomy for rhegmatogenous retinal detachment: an analysis of 512 cases.** *Graefes Arch Clin Exp Ophthalmol* 2006, **244**:69-78.
5. Sharma YR, Karunanithi S, Azad RV, Vohra R, Pal N, Singh DV, Chandra P: **Functional and anatomic outcome of scleral buckling versus primary vitrectomy in pseudophakic retinal detachment.** *Acta Ophthalmol Scand* 2005, **83**:293-297.
6. Halberstadt M, Chatterjee-Sanz N, Brandenberg L, Koerner-Stiefbold U, Koerner F, Garweg JG: **Primary retinal reattachment surgery: anatomical and functional outcome in phakic and pseudophakic eyes.** *Eye* 2005, **19**:891-898.
7. CHOLDEN L: **Some psychiatric problems in the rehabilitation of the blind.** *Bull Menninger Clin* 1954, **18**:107-112.
8. Adams GL, Pearlman JT, Sloan SH: **Guidelines for the psychiatric referral of visually handicapped patients.** *Ann Ophthalmol* 1971, **3**:72-73.
9. Hrisos S, Clarke MP, Wright CM: **The emotional impact of amblyopia treatment in preschool children: randomized controlled trial.** *Ophthalmology* 2004, **111**:1550-1556.
10. Brody BL, Roch-Leveq AC, Kaplan RM, Moutier CY, Brown SI: **Age-related macular degeneration: self-management and reduction of depressive symptoms in a randomized, controlled study.** *J Am Geriatr Soc* 2006, **54**:1557-1562.
11. Berman K, Brodaty H: **Psychosocial effects of age-related macular degeneration.** *Int Psychogeriatr* 2006, **18**:415-428.

12. Tolman J, Hill RD, Kleinschmidt JJ, Gregg CH: **Psychosocial adaptation to visual impairment and its relationship to depressive affect in older adults with age-related macular degeneration.** *Gerontologist* 2005, **45**:747-753.
13. Mitsonis CI, Mitropoulos PA, Dimopoulos NP, Mitsonis MI, Andriotis NM, Gitsa OE, Mitsonis IM: **Anxiety and depression in cataract surgery: a pilot study in the elderly.** *Psychol Rep* 2006, **99**:257-265.
14. Brandberg Y, Kock E, Oskar K, af TE, Seregard S: **Psychological reactions and quality of life in patients with posterior uveal melanoma treated with ruthenium plaque therapy or enucleation: a one year follow-up study.** *Eye* 2000, **14**:839-846.
15. Holladay JT, Prager TC: **Mean visual acuity.** *Am J Ophthalmol* 1991, **111**:372-374.
16. Zigmond AS, Snaith RP: **The hospital anxiety and depression scale.** *Acta Psychiatr Scand* 1983, **67**:361-370.
17. Poole NA, Morgan JF: **Validity and reliability of the Hospital Anxiety and Depression Scale in a hypertrophic cardiomyopathy clinic: the HADS in a cardiomyopathy population.** *Gen Hosp Psychiatry* 2006, Jan-Feb; **28**(1):55-58.
18. Mozaffarieh M, Krepler K, Heinzl H, Sacu S, Wedrich A: **Visual function, quality of life and patient satisfaction after ophthalmic surgery: a comparative study.** *Ophthalmologica* 2004, **218**:26-30.
19. Woolley FR, Kane RL, Hughes CC, Wright DD: **The effects of doctor--patient communication on satisfaction and outcome of care.** *Soc Sci Med* 1978, **12**:123-128.
20. Starfield B, Wray C, Hess K, Gross R, Birk PS, D'Lugoff BC: **The influence of patient-practitioner agreement on outcome of care.** *Am J Public Health* 1981, **71**:127-131.
21. Stewart MA, McWhinney IR, Buck CW: **The doctor/patient relationship and its effect upon outcome.** *J R Coll Gen Pract* 1979, **29**:77-81.
22. Radecki SE, Kane RL, Solomon DH, Mendenhall RC, Beck JC: **Do physicians spend less time with older patients?** *J Am Geriatr Soc* 1988, **36**:713-718.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

