

E46K-like α -synuclein mutants increase lipid interactions and disrupt membrane selectivity

Received for publication, November 9, 2018, and in revised form, May 1, 2019. Published, Papers in Press, May 2, 2019, DOI 10.1074/jbc.RA118.006551

Matteo Rovere[‡], Alex E. Powers[‡], Haiyang Jiang^{‡,1}, Julia C. Pitino^{‡,1}, Luis Fonseca-Ornelas[‡], Dushyant S. Patel[‡], Alessandro Achille[§], Ralf Langen[¶], Jobin Varkey[¶], and Tim Bartels^{‡,||,2}

From the [‡]Ann Romney Center for Neurologic Diseases, Brigham and Women's Hospital and Harvard Medical School, Boston, Massachusetts 02115, the [§]Department of Computer Science, University of California, Los Angeles, California 90095, the [¶]Zilkha Neurogenetic Institute, University of Southern California, Los Angeles, California 90033, and the ^{||}Dementia Research Institute, University College London, London WC1E 6BT, United Kingdom

Edited by Paul E. Fraser


Parkinson's disease (PD) is one of the most common neurodegenerative disorders, and both genetic and histopathological evidence have implicated the ubiquitous presynaptic protein α -synuclein (α Syn) in its pathogenesis. Recent work has investigated how disrupting α Syn's interaction with membranes triggers trafficking defects, cellular stress, and apoptosis. Special interest has been devoted to a series of mutants exacerbating the effects of the E46K mutation (associated with autosomal dominant PD) through homologous Glu-to-Lys substitutions in α Syn's N-terminal region (*i.e.* E35K and E61K). Such E46K-like mutants have been shown to cause dopaminergic neuron loss and severe but L-DOPA-responsive motor defects in mouse overexpression models, presenting enormous translational potential for PD and other "synucleinopathies." In this work, using a variety of biophysical techniques, we characterize the molecular pathology of E46K-like α Syn mutants by studying their structure and membrane-binding and remodeling abilities. We find that, although a slight increase in the mutants' avidity for synaptic vesicle-like membranes can be detected, most of their deleterious effects are connected to their complete disruption of α Syn's curvature selectivity. Indiscriminate binding can shift α Syn's subcellular localization away from its physiological interactants at the synaptic bouton toward trafficking vesicles and organelles, as observed in E46K-like cellular and murine models, as well as in human pathology. In conclusion, our findings suggest that a loss of curvature selectivity, rather than increased membrane affinity, could be the critical dyshomeostasis in synucleinopathies.

Parkinson's disease (PD)³ is a progressive neurodegenerative disorder characterized in its early stages by resting tremor, bradykinesia, and rigidity because of loss of dopaminergic neurons in the substantia nigra (1). Its prevalence in industrialized countries is estimated at 1% in people over 60 years of age, making it the second most common neurodegenerative disorder after Alzheimer's disease (2). α -Synuclein (α Syn), a small presynaptic protein ubiquitously expressed in nervous tissue (3), has long been considered a key player in the pathogenesis of PD, as supported by both genetic and histopathologic evidence (4). Its role in neuronal physiology, although still debated, has been tied to the maintenance of synaptic homeostasis (5, 6). These mechanisms are especially important in dopaminergic neurons, which exhibit high-frequency activity bursts (7).

One of the recurring themes of α Syn biology is its affinity for lipid interfaces and how such interplay shapes its function and dysfunction. Although α Syn has been classically considered an intrinsically disordered protein (8, 9), its N-terminal region contains seven imperfect 11-mer repeats (with the consensus XKTKEGVXXXX), homologous to those of apolipoproteins (10–12). In fact, upon interaction with liposomes and micelles, α Syn undergoes a coil-to-helix transition and binds robustly to curved, negatively charged interfaces (13, 14). In particular, its specificity for lipid bilayers similar in charge and curvature to those of synaptic vesicles (15) is attained by virtue of the periodic arrangement of mildly hydrophobic and hydrophilic amino acid side chains (16). In addition, the numerous lysines in the N-terminal region mediate the formation of ionic bridges with the charged heads of negative membranes (17). α Syn's N terminus has also been shown to be critical for its subcellular localization to the presynaptic terminals (18) and its partitioning between the cytosol and membranes, regulated by electrical activity (19).

Along with its role in cellular physiology, the lipid-binding ability of α Syn's N-terminal region has also been implicated in the pathogenesis of neurodegeneration (20). Overexpression of

This work was supported by NINDS, National Institutes of Health Grant R21NS107950 (to T. B.), American Parkinson Disease Association Grant PF-JFA-1884 (to T. B.), National Institutes of Health Grants NS084345 and GM115736 (to R. L.), and Office of Naval Research Grant N00014-13-1-0563 (to A. A.) and Parkinson's Disease Foundation Grant PF-JFA-1884 (to T. B.). The authors declare that they have no conflicts of interest with the contents of this article. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

 Author's Choice—Final version open access under the terms of the Creative Commons CC-BY license.

This article contains **Figures S1–S6 and Tables S1–S3**.

¹ Both authors contributed equally to this work.

² To whom correspondence should be addressed: Dementia Research Institute, University College London, Gower St., Cruciform Bldg., Wing 2.4, London WC1E 6BT, UK. Tel.: 44-20-31086814; E-mail: t.bartels@ucl.ac.uk.

³ The abbreviations used are: PD, Parkinson's disease; α Syn, α -synuclein; fPD, familial Parkinson's disease; HSQC, heteronuclear single-quantum coherence; SUV, small unilamellar vesicle; LUV, large unilamellar vesicle; POPC, 1-palmitoyl-2-oleoyl-glycero-3-phosphocholine; DOPS, 1,2-dioleoyl-*sn*-glycero-3-phospho-L-serine; ITC, isothermal titration calorimetry; MLV, multilamellar vesicle; POPG, 1-palmitoyl-2-oleoyl-*sn*-glycero-3-phospho-(1'-*rac*-glycerol).

E46K-like α -synuclein mutants disrupt membrane selectivity




Figure 1. Structural features of the E46K-like α Syn mutants in the lipid-unbound state. A, primary sequence of the N-terminal region (1–95) of WT, K, 2K, and 3K α Syn. Glu-to-Lys mutation sites are shown in red. The non-amyloid- β component (NAC) domain (61–95) and the seven 11-mer repeats XKTKEGVXXXX are annotated on the sequence. B, far-UV CD spectra of 10 μ M WT, K, 2K, and 3K α Syn in 10 mM NH_4Ac (pH 7.4), measured at 25 $^\circ\text{C}$. The single minimum around 200 nm is characteristic of a predominantly unfolded (random coil) secondary structure. C, size-exclusion chromatography (Superdex 200 10/300 GL) of the E46K-like α Syn mutants, monitored by 280-nm absorbance (the absorbance signal shown in the chromatogram was normalized to equalize the peak heights). As a reference, an ovalbumin standard (44 kDa) elutes on the same column at 15.375 ml.

α Syn and its mutants associated with autosomal dominant PD (fPD) in yeast and mammalian cell models severely compromises vesicle trafficking and protein clearance through abnormal membrane association (21–23). In addition, vesicle and organelle clusters have been recognized as prominent features of Lewy pathology (24).

Recent work has shown that modifications in α Syn's repeats that enhance the amphipathic nature of its membrane-binding helix modulate toxicity upon overexpression (25, 26). Inserting either more hydrophobic residues on the largely apolar face of the helix or charged residues on the cytosol-exposed side appears to increase the ability of α Syn mutants to induce toxicity and cluster a variety of intracellular vesicles and organelles into α Syn-rich inclusions. Among these mutants, Glu-to-Lys mutations homologous to the fPD-associated E46K mutation (27) (e.g. E35K and E61K) are of particular interest because of their direct connection to disease and their robust toxicity in cellular models (28). Readouts of cellular toxicity (e.g. viability and apoptotic markers) and their ability to cluster vesicles correlate with the number of Glu-to-Lys substitutions, suggesting a common molecular mechanism between the E46K mutation and these “exacerbated” E46K-like mutants. In addition, such mutants have been shown to markedly decrease the ratio between aggregation-resistant α Syn tetramers (29) and aggregation-prone monomers, whose imbalance has been proposed as one of the molecular mechanisms underlying disease inception (28, 30). Lastly, heterozygous transgenic mice overexpressing E46K-like α Syn mutants present a robust and progressive PD-like phenotype with severe tremor and gait deficits (par-

tially rescued by L-DOPA administration), loss of dopaminergic neurons, and formation of protease-resistant, α Syn-rich inclusions in neuronal bodies (31).

In this work we characterize the lipid-binding and lipid-remodeling abilities of three of these Glu-to-Lys mutants (E46K, “K”; E35K,E46K, “2K”; and E35K,E46K,E61K, “3K”; Fig. 1) using a variety of biophysical methods. Through comparisons between multiple E46K-like substitutions in α Syn's N-terminal region, we draw insights into the molecular pathology of E46K (and E46K-like) α Syn mutants and the structure of membrane-bound α Syn.

Results

E46K-like mutants increase N-to-C interactions and coil compactness

To characterize whether the Glu-to-Lys mutations induced any change in the structure of the lipid-unbound protein, CD spectra of the recombinant mutants in buffer were measured. Recombinant WT α Syn has long been known to exist in a predominantly unstructured “random coil” conformation (8), and the fPD mutant E46K and the 2K and 3K artificial mutants do not appear to affect the protein secondary structure enough to cause a change in the far-UV CD spectrum (Fig. 1B). Analogously, the weak near-UV CD signal of the protein is minimally affected by the mutations, if at all (Fig. S1). Nevertheless, size-exclusion chromatography analysis of the mutants' hydrodynamic radii shows a marked decrease moving from the WT to E46K (as reported previously (32)) and from E46K to 2K and 3K




Figure 2. NMR spectroscopy of the E46K-like α Syn mutants in the lipid-unbound state. A, overlaid ^1H - ^{15}N HSQC spectra of 120 μM WT, K, 2K, and 3K α Syn in 50 mM phosphate buffer and 10% D_2O (pH 6.8), measured at 15 $^\circ\text{C}$ and realigned using Ala-140. B, histogram of the mean weighted chemical shift perturbations (calculated as $[(\Delta\delta^1\text{H})^2 + (0.15 \cdot \Delta\delta^{15}\text{N})^2]^{1/2}$) of each mutant versus WT α Syn, plotted against the corresponding residue number. In addition to the N-terminal region, the C terminus is also affected and shows a clear correlation between the entity of the perturbations and the number of Glu-to-Lys mutations. C, charge (and charge density, *i.e.* absolute charge value) distribution of WT α Syn and its mutants, averaged over a window of nine amino acids. The transition from the N-terminal amphipathic region to the negatively charged C terminus can be observed around residue 100. Glu-to-Lys mutations, although not changing the charge density distribution, increase the charge polarization, making electrostatic N-to-C interactions more favorable.

(Fig. 1C). Although the elution volumes still indicate that all mutants have an extended conformation (because their hydrodynamic radii are unusually high for 14.5-kDa proteins (8)), the addition of lysines induces increasingly more compact structures.

Because CD cannot document small changes in the proteins' conformational equilibria upon Glu-to-Lys substitution, ^1H - ^{15}N HSQC spectra of the ^{15}N -labeled proteins in buffer (at 15 $^\circ\text{C}$) were also recorded and compared (Fig. 2A and Table S1). In accord with the CD data, the overall spread of the amide peaks of the E46K-like mutants reflects their disordered structure. However, numerous shifts in the resonances can be clearly observed, and some of these changes (especially in the case of C-terminal residues) correlate well with the increment in Glu-to-Lys mutations. After assignment of the amide peaks, relying on deposited datasets of the ^1H - ^{15}N HSQC of WT α Syn (33, 34), a histogram of the absolute weighted chemical shift perturbations throughout the protein sequence was obtained, highlighting the regions that are mostly affected by the sequential Glu-to-Lys substitutions (Fig. 2B). Compared with WT α Syn, all mutations show changes in the N terminus, not only around the site of the Glu-to-Lys substitutions but also mild-, medium-, and long-range effects. Interestingly, the C-terminal region is also markedly affected by the mutations and in a much clearer incremental fashion (*i.e.* correlating with the insertion of more and more lysines) than the N terminus. WT α Syn is known to exist in a variety of unstructured conformational states, many of which are stabilized by long-range N-to-C interactions, probably electrostatic in nature (35, 36). E46K α Syn has also been shown, in both experiments and simulations, to have

increased long-range contacts, most likely because of the substitution of the Glu-46 negative charge with a positively charged lysine, making transient bonding with the negative C-terminal region more favorable (37, 38). We can imagine similar (and much more marked, at least according to the magnitude of the perturbations) effects playing a role in this case, explaining the decrease in hydrodynamic radius as well (Fig. 1C). Indeed, analysis of the charge distribution of the N terminus of WT α Syn and comparison with its K, 2K, and 3K mutants confirms that, although the overall charge density is unperturbed by the mutations, the Glu-to-Lys substitutions increase the overall charge polarization (Fig. 2C). If we take the segment 1–95 as the N terminus of the protein, we observe a net +6 charge increase going from WT α Syn (which has a mild imbalance of +3.1 charges) to 3K α Syn.

E35K and E46K most affect the micelle-bound "broken helix"

Extensive biophysical work has concluded that a single continuous helix is the preferred conformation of α Syn upon interaction with curved lipid bilayers (*e.g.* small unilamellar vesicles (SUVs) and large unilamellar vesicles (LUVs)) similar in morphology and composition to the physiological membranous interactants of α Syn (17, 39). Nevertheless, the best-characterized helical structure of α Syn is that classically described as a broken helix (33) and observed mainly in lipoprotein nanoparticles (12) and on detergent micelles (33, 40). We therefore decided to explore how the Glu-to-Lys mutations affected α Syn's broken-helix structure by recording ^1H - ^{15}N HSQC spectra of the recombinant ^{15}N -labeled mutants in the presence of SDS micelles (Fig. 3A and Table S1). The distribution of

E46K-like α -synuclein mutants disrupt membrane selectivity




Figure 3. NMR spectroscopy of the E46K-like α Syn mutants in the SDS-bound state. *A*, overlaid ^1H - ^{15}N HSQC spectra of 120 μM WT, K, 2K, and 3K α Syn in 15 mM SDS, 50 mM phosphate buffer, and 10% D_2O (pH 6.8), measured at 37 $^\circ\text{C}$ and realigned using Ala-140. *B*, histogram of the mean weighted chemical shift perturbations (calculated as $[(\Delta\delta^1\text{H})^2 + (0.15 \cdot \Delta\delta^{15}\text{N})^2]^{1/2}$) of each mutant versus WT α Syn, plotted against the corresponding residue number. *C*, histogram of the mean weighted chemical shift perturbations of E46K versus WT (E46K), 2K versus E46K (E35K), and 3K versus 2K (E61K), plotted against the corresponding residue number. Although the E35K and E46K mutations cause wide-ranging perturbations in the N-terminal region of SDS-bound α Syn, the effects of E61K are mostly clustered around the site of the mutation. This is easily understood upon examination of the deposited structure of SDS-bound α Syn (inset, PDB code 1XQ8, showing the position and orientation of the glutamates in the sites of Glu-to-Lys mutation).

the amide resonances of all mutants is consistent with a folded structure, and the Glu-to-Lys mutations seem to cause relatively few perturbations in the chemical shifts, which are mainly clustered in the N terminus of the protein, as is clearest after plotting the chemical shift perturbations of each mutant compared with WT α Syn (Fig. 3*B*). By comparing each of the K mutants with its homolog (with one less Glu-to-Lys substitution) we can also probe how each site affects the structure of the broken helix (although in an imperfect fashion, as the perturbations could well not be perfectly additive). These differences are shown in Fig. 3*C*. Although both E46K and E35K have robust, long-ranging effects on α Syn's resonances, E61K appears to act prevalently on a local scale, with only minor "ripples" in the residues away from the substitution site. Inspection of the deposited structure of micelle-bound α Syn (PDB code 1XQ8) allows us to draw structural correlations that confirm our NMR findings. Although Glu-35 and Glu-46 are localized close to the "ordered loop" region and oriented in a way that facilitates electrostatic interactions with the negatively charged heads of the detergent, Glu-61 points directly opposite of the hydrophobic side of the second amphipathic helix and can therefore be expected to only marginally affect the structure of α Syn upon charge inversion (Fig. 3*C*, inset). These results contradict previous *in vitro* and *in vivo* work with the E46K-like mutants, which showed a stepwise increase in toxicity with further Glu-to-Lys mutations and, if anything, a more prominent effect of the E61K mutation (28, 31). A structure other than the broken helix could be predominant in cellular and murine models and would explain such inconsistencies.

Glu-to-Lys mutations mildly increase the "avidity" for curved membranes

To further explore the molecular events leading to the vesicle clustering and toxicity caused by overexpression of the E46K-like mutants (26), we studied the binding of WT α Syn and its Glu-to-Lys mutants to negatively charged SUVs. Upon interaction with curved membranes, such as when titrating the recombinant protein with SUVs, α Syn undergoes a coil-to-helix transition that can be monitored by CD (13). SUVs were prepared by sonication (obtaining a dispersion of vesicles ~ 33 nm in diameter; Table S2) of a 70:30 hydrated mixture of zwitterionic 1-palmitoyl-2-oleoylphosphatidylcholine (POPC) and anionic 1,2-dioleoyl-*sn*-glycero-3-phospho-L-serine (DOPS). α Syn, as expected, bound avidly (Fig. 4, *A* and *B*) to these membranes; the titration curves obtained were analyzed to derive the thermodynamic and stoichiometric parameters of the binding event. An *N*-independent binding site model was used to describe the binding events (see "Experimental procedures"). Because the molar lipid concentration was used as the ligand concentration, the *N* value, rather than the number of binding sites, measures the average number of lipid molecules that are necessary to form a lipid binding site for α Syn. As the number of lipid molecules constituting a vesicle can be considered constant, *N* is then inversely proportional to the number of binding sites on a given vesicle. A decrease in *N* therefore reflects a higher number of saturable binding sites on the vesicles' surface and vice versa. We observe a mild decreasing trend in the *N* values obtained by fitting the mutants' titration curves; this represents an increase in the number of α Syn binding sites,




Figure 4. Glu-to-Lys mutations result in mildly increased α Syn binding to curved charged membranes. *A*, representative far-UV CD spectra from a titration of 10 μ M WT α Syn (in 10 mM NH₄Ac (pH 7.4)) with 6 mM 70:30 POPC:DOPS SUVs, measured at 25 °C. There is a clear transition from the (almost completely) unstructured lipid-unbound protein to the robustly helical endpoints of the titration (e.g. lipid:protein molar ratio 270:1), reflecting the binding and folding of α Syn. *B*, titration curves obtained from raw CD data after extracting the molar ellipticity at 222 nm and graphing it against the lipid:protein molar ratio at each point of the titration. Datapoints are shown along with their standard errors (from $n = 3$ independent titrations) and their best fit with an N independent binding sites model (see “Experimental procedures”). *C*, histogram of the N values (along with their standard errors, $n = 3$) obtained from the best fit of the titration curves with an N -independent binding sites model. Decreasing N s indicate increased binding (or avidity, see “Results”). *D*, histogram of the 222-nm plateau molar ellipticity values (along with their standard errors, $n = 3$) obtained from the end-point CD spectra of each mutant after renormalization using the WT α Syn isodichroic point. n.s., $p > 0.05$; **, $p \leq 0.01$.

although this result does not reach significance in the cases of K and 2K α Syn (Fig. 4C and Table S3). Although E46K has been reported previously to have a higher affinity for negatively charged, curved lipid interfaces (41, 42), more recent work has not been able to detect differences with WT α Syn, either when using recombinant E46K α Syn and artificial liposomes or measuring synaptic localization in cultured primary neurons (43). It must also be noted that E61K has the highest effect on N of all Glu-to-Lys mutations. Finally, the increase in binding does not correspond to significant changes in the plateau ellipticity values associated with the membrane-bound helical structures, suggesting only minor (if any) conformational rearrangements. Similar CD-monitored lipid-into-protein titrations were also performed in the presence of 150 mM NaCl to test the effect of a higher ionic strength on α Syn’s binding behavior and confirmed the results obtained under low-salt conditions (Fig. S2 and Table S3). As expected, the end-point ellipticities and the k_B values are smaller, probably because of the electrostatic masking of the charged moieties in the lipid heads and in some of α Syn’s residues, decreasing the strength of ionic interactions at the vesicular interface.

Knowing the critical role of cholesterol-rich lipid rafts in α Syn’s physiology (18) and the high content of cholesterol in the membranes of synaptic vesicles (15), we also studied whether the rigidity of the double layer affected the binding of

the Glu-to-Lys mutants. CD-monitored titrations were performed in the same way as those with 70:30 POPC:DOPS, but a 52.5:17.5:30 POPC:DOPS:cholesterol lipid dispersion was sonicated to generate SUVs (~36 nm in diameter). The results (both N and the plateau ellipticity values) are similar to those obtained with POPC:DOPS SUVs (Fig. 5), but, in this case, there is a much clearer distinction between the binding behavior of the 3K and that of the other mutants, which are practically indistinguishable and show no trends in their N values (Fig. 5B). Again, E61K appears to be the mutation site that most significantly affects the ability of α Syn to associate with membranes.

Although CD-monitored lipid-into-protein titrations can provide accurate estimates of the number of lipids per binding site (N) and macroscopic structural information regarding the coil-to-helix transition, their estimates of the apparent microscopic binding constants (k_B) are poor. For this reason, and to confirm the previous results by means of an independent technique, we also measured the binding curves of WT α Syn and its Glu-to-Lys mutants to SUVs by isothermal titration calorimetry (ITC). The results for 70:30 POPC:DOPS are analogous to those found by CD-monitored titrations (Fig. S3, A and C, and Table S3). In addition to N , ITC provides thermodynamic information about the heat released upon binding (Fig. S3D), but the interpretation of these data, although marked differences are observed between the mutants, is made challenging by the fact

E46K-like α -synuclein mutants disrupt membrane selectivity




Figure 5. E61K alone induces mildly increased binding to curved cholesterol-rich membranes. *A*, titration curves of 10 μ M WT, K, 2K, and 3K α Syn (in 10 mM NH_4Ac (pH 7.4)) with 6 mM 52.5:17.5:30 POPC:DOPS:cholesterol SUVs (measured at 25 $^\circ\text{C}$). Plots were obtained from raw CD data after extracting the molar ellipticity at 222 nm and graphing it against the lipid:protein molar ratio at each point of the titration. Datapoints are shown along with their standard errors (from $n = 3$ independent titrations) and their best fit with an N -independent binding site model (see “Experimental procedures”). *B*, histogram of the N values (along with their standard errors, $n = 3$) obtained from the best fit of the titration curves with an N -independent binding site model. Decreasing N s indicate increased binding (or avidity, see “Results”). *C*, histogram of the 222-nm plateau molar ellipticity values (along with their standard errors, $n = 3$) obtained from the end-point CD spectra of each mutant after renormalization using the WT α Syn isodichroic point. *n.s.*, $p > 0.05$; **, $p \leq 0.01$; ***, $p \leq 0.001$.

that the ΔH values are a combination, among others, of the heat of binding of α Syn to the membranes, the heat exchanged because of the coil-to-helix transition, and the heat contributions of the lipids themselves (e.g. tail hydration, bilayer rearrangement, and remodeling). ITC, as expected, provides much more accurate estimates of the apparent binding constants, which, however, do not appear to change significantly between mutants (Fig. S3E and Table S3). ITC of 52.5:17.5:30 POPC:DOPS:cholesterol SUVs did not detect any binding processes in the cases of WT, K, and 2K α Syn, possibly because the heat exchanges of bilayer reorganization processes masked the heat from the protein’s binding and folding. The only mutant that showed a clear sigmoidal profile by ITC was 3K α Syn, again in accord with the CD evidence that the E61K mutation alone increases binding to more rigid membranes (Fig. S3B and Table S3).

In conclusion, it is important to consider how, based on the combined CD and ITC data, the α Syn mutants (3K in particular) do not necessarily appear to have a higher “affinity”, in its

classic interpretation (higher k_B), for curved charged membranes. Rather, the more Glu-to-Lys mutations are present, the more α Syn molecules seem to be able to bind to a single vesicle, which, to borrow from the nomenclature of polyvalent interactions, corresponds to a higher avidity of lipid bilayers for the E46K-like mutants (44).

Glu-to-Lys mutations cause a progressive loss of curvature selectivity

One rarely explored aspect of the molecular pathology of fPD-associated α Syn mutants is how their binding selectivity is affected by mutations. Because α Syn has long been known to bind preferentially to small vesicles (13), and its dynamic association with synaptic vesicles upon exo- and endocytosis is believed to be central to its function (19, 45), we chose to explore how Glu-to-Lys mutations modify α Syn’s selectivity for certain curvature radii. We performed CD-monitored titrations of WT α Syn and its Glu-to-Lys mutants with LUVs prepared by extrusion of a 70:30 POPC:DOPS dispersion using 0.1- μ m polycarbonate membranes, obtaining vesicles ~ 100 nm in diameter (Table S2). As expected, the binding propensity of both WT α Syn and its mutants to charged LUVs is markedly lower than that observed with SUVs (Fig. 6, A and B). The curvature of the titration curves is much shallower than the sharp bends of SUVs titrations, signaling that the k_B values are orders of magnitude lower (Table S3). The ellipticities of the helical conformers of LUV-bound α Syn are also roughly half of those recorded before (Figs. 6A and 4A), possibly because of a more dynamic equilibrium between partly folded and partly unfolded conformations, which has been proposed to be one of the curvature-sensing mechanisms adopted by α Syn (46). Furthermore, although the effects of Glu-to-Lys mutations are generally mild in the SUV titrations (possibly because of the overall high affinities), robust changes are evident in the LUV titrations, especially after curve fitting with our N -independent binding site model (Fig. 6, C and D). Going from WT α Syn to 3K, we see a decrease in N (number of lipids per binding site) that is truly stepwise incremental, as are the effects observed in cellular models (28). In addition, although we could not detect significant differences between the avidity (nor the affinity) of E46K and WT α Syn using SUVs, here we observe a clear loss of curvature selectivity with the introduction of the fPD-causing mutation, which could be one of the factors underlying its propensity for pathology. In addition to the increased avidity of negatively charged LUVs for E46K-like mutants, a slight increase in plateau ellipticities is also detected, but, although significant, the effects are as minor as those described previously in the case of SUVs (Fig. 6D).

As with the SUVs titrations, the experiments were repeated in the presence of 150 mM NaCl to explore the effect of increased ionic strength (Fig. S4). However, given the already weak affinity of α Syn for the mildly charged LUVs, the presence of a higher salt concentration almost completely abolished binding for all mutants except 3K (confirming its much stronger ability to bind bilayers in all of the conditions studied). To be able to quantitatively compare the different Glu-to-Lys mutants, we used the slope of the binding curve in its early phase as a measure of the strength of the affinity (Fig. S4C).




Figure 6. α Syn's Glu-to-Lys mutations cause a progressive loss of curvature selectivity. *A*, representative far-UV CD spectra from a titration of 10 μ M WT α Syn (in 10 mM NH_4Ac (pH 7.4)) with 10 mM 70:30 POPC:DOPS LUVs, measured at 25 $^\circ\text{C}$. Although a coil-to-helix transition is also observed in the presence of LUVs, both its sharpness and the entity of the conformational rearrangement are much less than those observed with SUVs (Fig. 4). *B*, titration curves obtained from raw CD data after extracting the molar ellipticity at 222 nm and graphing it against the lipid:protein molar ratio at each point of the titration. Datapoints are shown along with their standard errors (from $n = 3$ independent titrations) and their best fit with an N -independent binding site model (see "Experimental procedures"). *C*, histogram of the N values (along with their standard errors, $n = 3$) obtained from the best fit of the titration curves with an N -independent binding site model. Decreasing N s indicate increased binding (or avidity, see "Results"). *D*, histogram of the 222-nm plateau molar ellipticity values (along with their standard errors, $n = 3$) obtained from the best fit of the titration curves after renormalization using the WT α Syn isodichroic point. *n.s.*, $p > 0.05$; *, $p \leq 0.05$; **, $p \leq 0.01$; ****, $p \leq 0.0001$.

Although not as mechanistically valuable as the N -independent binding site model, comparison of the binding slopes among the different mutants still shows a much stronger progression (WT to 3K) than the one observed with SUVs (Fig. S4D), confirming previous conclusions.

E46K-like mutants have increasing membrane remodeling abilities

One further aspect of α Syn's interplay with membranes was then studied; namely, its ability to remodel lipid bilayers into bilayer tubes, micellar tubes, and lipoprotein nanoparticles (12, 47, 48). A dispersion of multilamellar vesicles (MLVs) was prepared by hydration of a 90:10 mixture of 1-palmitoyl-2-oleoyl-*sn*-glycero-3-phospho-(1'-*rac*-glycerol) (16:0–18:1 PG or POPG) and POPC. The content of charged lipids was chosen to magnify differences in tubulation rates arising from progressive Glu-to-Lys mutations. Upon addition of α Syn, the robust scattering signal from the heterogeneous dispersion of large ($d > 1 \mu\text{m}$) MLVs is drastically decreased as α Syn induces the formation of tubules, vesicular structures, and lipoprotein nanoparticles, which all exhibit reduced scatter. Remodeling rates can thus be studied by light scattering (*e.g.* absorbance at 500 nm) and also by CD because it is the association of α Syn with the

vesicles (and its consequent folding) that drives the membrane remodeling (Fig. 7, A and B). The rates of vesicle clearance correlate well with α Syn's increasing avidity for LUVs, with a stepwise increase going from WT to 2K α Syn; however, they plateau at 3K, showing little difference upon introduction of E61K. Fitting of the clearance curves using a two-phase exponential decay confirms qualitative observations (the decay rates of the "slow" phase were compared, because the early "fast" phase has too few points to provide accurate parameters). Half-lives decrease upon introduction of the first two Glu-to-Lys mutations (E46K and E35K), and no differences can be detected between 2K and 3K (both in absorbance-monitored and CD-monitored experiments; Fig. 7C). Electron micrographs obtained after mixing WT α Syn and its Glu-to-Lys mutants with MLVs also reveal robust membrane remodeling in all cases. However, the introduction of successive Glu-to-Lys mutations leads to the more frequent appearances of narrower ($d \sim 5\text{--}6 \text{ nm}$) membrane tubes (Fig. 7, D–G). Previous work (47, 48) has shown how such narrow tubes are micellar in nature and are increasingly generated at higher protein-to-lipid ratios. This phenomenon reflects the increased avidity of Glu-to-Lys mutants toward large curved membranes (Fig. 6), result-

E46K-like α -synuclein mutants disrupt membrane selectivity




Figure 7. α Syn's Glu-to-Lys mutations result in increased membrane remodeling. *A*, vesicle clearance experiments, monitored by absorbance at 500 nm of 400 mM 90:10 POPG:POPC MLV dispersions by 40 μ M WT α Syn and its E46K-like mutants (lipid:protein molar ratio 10:1). Best fits were obtained using a two-phase exponential decay model (Table S3) and are shown in *black*. *B*, vesicle clearance experiments, monitored by CD (222-nm ellipticity), of 400 mM 90:10 POPG:POPC MLV dispersions by 40 μ M WT α Syn and its E46K-like mutants (lipid:protein molar ratio 10:1). Best fits were obtained as in *A* and are shown in *black* (Table S3). *C*, histogram of the (slow-phase) half-life values (along with their 95% confidence interval) obtained from the best fit of the clearance kinetics in *A* and *B* with a two-phase exponential decay model. As more Glu-to-Lys mutations are introduced, the clearance rate increases (and, thus, half-lives decrease), but the trend plateaus at 3K α Syn. *D–G*, electron micrographs, negatively stained, of lipid tubule dispersions generated by mixing WT (*D*), E46K (*E*), 2K (*F*), and 3K (*G*) α Syn to 90:10 POPG:POPC MLV preparations. Scale bars = 200 nm. Although all mutants present the ability to robustly tubulate membranes, with progressive Glu-to-Lys mutations, the number of large, vesicular-like “cisternae” (e.g. *D*, left) and bilayer tubes (black arrowheads) formed upon incubation with α Syn decreases, and thinner, micellar (e.g. *G*, right) tubes can be observed (white arrowheads). *abs*, absorbance.

ing in a greater number of bound α Syn molecules and more profound bilayer remodeling.

Discussion

In the search for the molecular determinants underlying the inception of PD and other synucleinopathies, a great deal of

attention has been devoted to the biophysics and biochemistry of fPD-associated α Syn mutants. Historically, most of these mutants were characterized in their ability to generate toxic oligomers upon aggregation (49, 50). However, after the discovery of the robust and specific lipid-binding ability of WT α Syn (13), several lines of evidence have shown how disruption of

such binding equilibria can also lead to cellular toxicity and disease (20, 24, 51).

In this work, we focused on the molecular mechanism behind the lipid-mediated toxicity of E46K α Syn and its exacerbated analogs (2K and 3K α Syn, Fig. 1A), which have been shown, in both cellular and murine models, to cause trafficking defects, α Syn-rich vesicular inclusions, and a PD-like motor phenotype (26, 28, 31). We described how E46K, 2K, and 3K α Syn, in accord with previous reports (26, 31), exhibit a stepwise increase in membrane avidity (Figs. 4–6) and in their membrane-remodeling abilities (Fig. 7). Although the effects of E46K and E46K-like mutations on α Syn's binding to small curved vesicles were relatively minor (Figs. 4 and 5), these mutants caused a much more profound disruption in the protein's curvature selectivity, which was almost completely abolished in 3K α Syn (Fig. 6). Although some reports have previously described E46K α Syn as binding more avidly to small, charged vesicles (41, 42), our tightly controlled CD-monitored titration experiments, combined with an independent technique such as ITC, failed to reproduce these results, in agreement with more recent work (43). 2K and 3K α Syn also displayed only small (or even undetectable) effects when SUVs were used as their binding partners (Figs. 4 and 5) but exhibited much more severe changes in their LUV-binding ability (Fig. 6), strengthening the case that they behave as exacerbated E46K mutants. Finally, membrane tubulation assays, performed using heterogeneous dispersions of large MLVs, confirmed such trends (Fig. 7). Successive Glu-to-Lys mutations appear to transform α Syn from being predominantly a "curvature sensor" (WT) to acting as a robust "curvature inducer" (2K and 3K), although it is likely that these aspects are ultimately linked and can both be traced back to the loss of curvature selectivity.

We hypothesize that behind the pathogenesis of both E46K-associated fPD and 2K/3K murine models of synucleinopathy there could be a redistribution of the membrane-bound α Syn population to larger intracellular vesicles (such as those involved in secretory or endosomal pathways) rather than an increased avidity for synaptic vesicles, their natural binding partners. Such a shift could then act as a combined toxic gain of function, with α Syn disrupting vesicular trafficking through its abnormal binding and damaging loss of function because of α Syn being less available at the synaptic bouton, where it would normally regulate synaptic vesicle fusion and recycling (6).

In addition to characterizing the molecular pathology of E46K-like mutants, the sequential insertion of Glu-to-Lys mutations at specific points in the 11-mer repeat region allows us to rationalize the observed phenomena using structural determinants. α Syn's primary structure, upon inspection of its sequence hydrophobicity (Fig. S5A), is clearly split between a neutral (neither robustly polar nor apolar) N terminus (1–95) and a negatively charged C terminus, consistently characterized as disordered. Also, either by plotting Eisenberg's hydrophobic moment (measuring the degree of polarity segregation in a helical structure) *versus* the helix's periodicity (52, 53) or by calculating the fast Fourier transform of the vector of the side-chain hydrophobicities in segment 1–95 (54, 55), we can see how the N terminus can be arranged into robustly amphiphilic helical structures with periodicities around 3.6 amino acids per




Figure 8. α Syn's N-terminal region has an intrinsic propensity toward an 11/3 helix. Shown is the hydrophobic moment, according to Eisenberg's definition (52), of the N-terminal segment of α Syn (1–95), calculated for different helix periodicities (using the hydrophathy index values of Kyte and Doolittle (64)) and graphed against the angle between consecutive amino acids. The angles corresponding to an ideal α -helix (100°) and an ideal 11/3 helix (98.18°) are marked and labeled, showing how the amino acid arrangement in the N terminus of α Syn leans toward an 11/3 arrangement if amphiphilicity (*i.e.* segregation of polar and apolar residues) is to be maximized.

turn (Fig. 8 and S5B). Depending on the amphiphilic interactant, α Syn's membrane-bound (or detergent-bound) structure has been either described as a broken helix, with two α -helical segments (3.6 amino acids per turn) connected by an ordered loop (33, 40), or as a single continuous 11/3 helix (\sim 3.67 amino acids per turn), spanning throughout the whole 1–95 region (17, 39). Subsequent work has described the possibility of an interchange between these two forms, quantifying the equilibrium as shifted toward the 11/3 helix when α Syn is bound to charged SUVs rather than detergent micelles (12, 56).

If we plot a Schiffer–Edmundson helical wheel diagram (57) with an 11/3 periodicity and its helical net (58), spanning α Syn's N-terminal region (1–95), and highlight the positions of the three Glu residues progressively substituted by lysines, we can hypothesize how their arrangement could increase their avidity for bilayers with a progression comparable with that observed in our experiments (Fig. S6). The proximity of Glu-35, Glu-46, and Glu-61 to the negatively charged surface of the vesicle's membrane could allow the formation of ionic bridges between the positively charged lysines and the phosphate heads, favoring the binding and stabilizing the 11/3 helix. Although all three residues are in very similar positions, the presence of the 53–56 (ATVA, Fig. 1A) break in the 11-mer periodicity region shifts Glu-61 slightly closer to the membrane surface. While this small difference might be masked in 3K α Syn by the dynamic nature of the 11/3 helix (17, 46, 59), especially in fluid POPC/DOPS SUVs (Fig. 4), it could become more important in the case of rigid membranes, rich in cholesterol, also explaining the lone increased avidity of 3K for such interfaces (Fig. 5).

In conclusion, our biophysical characterization recapitulates all relevant features of cellular and murine models of the E46K-like mutants and, at the same time, allows us to advance the possibility of a loss of curvature selectivity, rather than increased membrane affinity, as the critical dyshomeostasis in synucleinopathies. In light of these results, there appears to be an even greater translational potential in therapies modulating the cellular lipid composition to compensate for α Syn defects, which has recently started to be explored (60, 61).

E46K-like α -synuclein mutants disrupt membrane selectivity

Experimental procedures

Materials

1-palmitoyl-2-oleoyl-glycero-3-phosphocholine (16:0–18:1 PC or POPC), 1,2-dioleoyl-*sn*-glycero-3-phospho-L-serine (18:1 PS or DOPS), and 1-palmitoyl-2-oleoyl-*sn*-glycero-3-phospho-(1'-*rac*-glycerol) (16:0–18:1 PG or POPG) were purchased from Avanti Polar Lipids (Alabaster, AL) as chloroform solutions. All other chemicals were obtained from Sigma-Aldrich (St. Louis, MO) unless otherwise noted.

Molecular cloning

pET21a- α -synuclein, encoding codon-optimized untagged human α Syn, was a gift from the Michael J. Fox Foundation (Addgene plasmid 51486). Fragments encoding E46K (K), E35K,E46K (2K), and E35K,E46K,E61K (3K) α Syn were amplified with PfuUltra II Fusion HS (Agilent Technologies, Santa Clara, CA) using their constructs in pcDNA4 (28) as templates and the primers α Syn_InFu_FWD 5'-AAGGAGATATACATATGGATGTATTCATGAAAGGACTTTCAAAGG-3' and α Syn_InFu_REV 5'-TGCTCGAGTGC GGCCGCTCAGGCTCAGGTCGTAGTCTTGATACC-3'. The amplified fragments were then inserted into pET21a (linearized with NdeI/NotI) using the In-Fusion HD Cloning Kit (Takara Bio, Mountain View, CA), following the manufacturer's protocol. Cloning of pET21a-K- α Syn, pET21a-2K- α Syn, and pET21a-3K- α Syn was confirmed by DNA sequencing and restriction analysis.

Protein expression and purification

BL21(DE3) *Escherichia coli* (New England Biolabs, Ipswich, MA) were transformed with the pET21a-based constructs, and single colonies were inoculated in LB + ampicillin. Cultures were induced at an OD₆₀₀ of ~0.5 with 1 mM isopropyl 1-thio- β -D-galactopyranoside (IPTG) for 4 h. The cell pellet, after harvesting, was resuspended in 20 mM Tris buffer and 25 mM NaCl (pH 8) and lysed by boiling for 15 min. The supernatant of a 20-min., 20,000 \times g spin of the lysate was then processed further. The sample was loaded on two 5-ml (tandem) HiTrap Q HP anion exchange columns (GE Healthcare) equilibrated with 20 mM Tris buffer and 25 mM NaCl (pH 8). α Syn was eluted from the columns with a 25–1000 mM NaCl gradient of 20 mM Tris buffer and 1 M NaCl (pH 8). Peak fractions were pooled and further purified via gel filtration on a HiPrep Sephacryl S-200 HR 16/60 gel filtration column (GE Healthcare) using 50 mM NH₄Ac (pH 7.4) as running buffer. Peak fractions (>95% pure, by Coomassie Brilliant Blue–stained SDS-PAGE) were pooled, aliquoted, lyophilized, and stored at –20 °C. Purified samples were also analyzed by MALDI-TOF MS and by trypsin digestion followed by MALDI-TOF MS to confirm the intact mass and sequence of the proteins (Molecular Biology Core Facilities, Dana-Farber Cancer Institute). Lyophilized proteins were reconstituted fresh, before experiments, in 10 mM NH₄Ac (pH 7.4) and not reused (relyophilized and reconstituted) more than twice. Protein solutions were spun down every time after resuspension, before spectroscopic quantitation of the protein concentration, at 21,130 \times g for 20 min at 4 °C to remove any large

α Syn aggregates. Concentration was quantified by measuring the absorbance at 280 nm ($\epsilon = 0.412 \text{ mg}\cdot\text{ml}^{-1}\cdot\text{cm}^{-1}$).

¹⁵N-labeled protein expression

For the expression of ¹⁵N-labeled α Syn, the bacteria were grown in M9 complete medium + ampicillin supplemented with [¹⁵N]H₄Cl. M9 complete medium was prepared fresh from a 10 \times M9 medium stock (60 g·L⁻¹ Na₂HPO₄, 30 g·L⁻¹ KH₂PO₄, 5 g·L⁻¹ NaCl), 50 \times 20% D-glucose, 1000 \times 1 M MgSO₄, 5000 \times 1 M CaCl₂, 1000 \times 1 mg·ml⁻¹ thiamine and biotin, 100 \times trace element solution, 1000 \times vitamin solution, and 1 g·L⁻¹ [¹⁵N]H₄Cl. The 100 \times trace element solution was prepared by dissolving (in 1 liter of MilliQ water) 5 g of EDTA, 0.50 g of FeCl₃, 84 mg of ZnCl₂, 12 mg of CuSO₄, 10 mg of CoCl₂·6H₂O, 10 mg of H₃BO₃, 1.6 mg of MnCl₂·4H₂O, and 1.6 mg of Na₂MoO₄·2H₂O and adjusting the pH to 7.0 (it was then filter-sterilized and stored at room temperature). The 1000 \times vitamin solution was prepared by dissolving (in 1 liter of 95% MilliQ water and 5% acetonitrile) 50 mg of riboflavin, 0.5 g of niacinamide, 0.5 g of pyridoxine hydrochloride, and 0.5 g of thiamine. After filtration with a 0.22- μ m cutoff (to sterilize and remove the undissolved riboflavin), the solution was stored at 4 °C protected from light. BL21(DE3) *E. coli* (New England Biolabs) were transformed with the pET21a-based constructs and single colonies inoculated in 2 \times YT + ampicillin. Cultures were grown for 6–8 h. at 37 °C under shaking and then used to inoculate an overnight preculture in M9 complete medium + ampicillin, which, on the following day, was diluted again 1:10 in M9 complete medium + ampicillin. The expression and purification of ¹⁵N-labeled proteins were, from this point on, performed as described under “Protein expression and purification.”

Size-exclusion chromatography

Samples were injected on a Superdex 200 (10/300 GL) column (GE Healthcare) at room temperature and eluted with 50 mM NH₄Ac (pH 7.4) while measuring (in-line) the conductivity and the 280-nm absorption of the eluate. For size estimation, a gel filtration standard (catalog no. 151-1901, Bio-Rad) was run on the column, and the calibration curve was obtained by semi-logarithmic plotting of molecular weight *versus* the elution volume divided by the void volume.

Liposome preparation

Aliquots from chloroform stock solutions of POPC, DOPS, and cholesterol were mixed thoroughly in the desired proportions and dried under a stream of argon. The dried lipid film was then kept under a vacuum overnight to remove traces of organic solvent. For both SUVs and LUVs, lipid solutions were freshly prepared by resuspending the dried lipids in 10 mM NH₄Ac (pH 7.4) to their final concentrations and hydrating them for 1 h at 37 °C while nutating. SUVs were prepared by pulse-sonicating the phospholipid suspensions for 10–20 min at room temperature with a microtip sonicator (Misonix S-4000, Qsonica, Newtown, CT). The SUV suspensions were then spun down at 21,130 \times g for 10 min to remove any metal particles and larger lipid aggregates. LUVs were prepared by manual extrusion (25 passes) using an Avanti Mini-Extruder

(Avanti Polar Lipids) and 0.1- μ m polycarbonate membranes, following the manufacturer's protocol. For MLVs, aliquots from the lipid stock solutions in chloroform were mixed thoroughly in the desired proportions and dried under a stream of nitrogen. The dried lipid film was then kept under vacuum for more than 12 h to remove traces of organic solvent. MLV dispersions were prepared by resuspending the dried lipids in the buffer of choice and vortexing.

CD

Spectra were recorded with a Jasco J-815 spectropolarimeter (Jasco, Easton, MD) at 25 °C using a 1-mm path length cuvette for far-UV CD (a 1-cm path length cuvette was used for near-UV CD measurements). Temperature control with an accuracy of ± 0.1 °C was achieved with a heating/cooling accessory equipped with a Peltier element (PFD-425S) connected to a water thermostatic bath. When possible, buffer spectra were recorded and subtracted. In titration experiments, spectra were corrected for the dilution effect from the addition of titrant; control lipid-into-buffer titrations were also recorded to confirm the negligibility of the liposome scattering signal. Three independent titrations (although with the same vesicle preparation) were performed for each condition. In "high-salt" (150 mM NaCl) titrations, NaCl was spiked into protein and lipid samples right before the experiments using a 5 M stock.

Stoichiometric and thermodynamic parameters (N , Θ , k_B) were determined by fitting the CD titration curves according to an N -independent binding site model. The expression of the fractional saturation ϑ (ratio of occupied sites to total sites) as a function of P_t and L_t (the total concentrations of protein and lipid), k_B (the apparent microscopic binding constant), and N (the number of saturable binding sites on the surface of the vesicles) was taken from the classic derivation of the Wiseman isotherm for an N -independent binding site model (62).

$$\theta = \frac{1}{2} \left[1 + \frac{L_t}{NP_t} + \frac{1}{Nk_B P_t} - \sqrt{\left(1 + \frac{L_t}{NP_t} + \frac{1}{Nk_B P_t} \right)^2 - \frac{4L_t}{NP_t}} \right] \quad (\text{Eq. 1})$$

In CD-monitored titrations, the 222-nm signal comes from the combination of the random coil signal from unbound α Syn and the helical signal of lipid-bound α Syn, each proportional to the molar ratios of the two species (Θ is the molar, or residue, ellipticity at each point of the titration).

$$[\theta]_i^{222} = \frac{[PL_n]}{P_t} [\theta]_{helix}^{222} + \frac{[P]}{P_t} [\theta]_{coil}^{222} \quad (\text{Eq. 2})$$

ϑ (at each point of the titration) can then be expressed as

$$\frac{[PL_n]}{P_t} = \theta = \frac{[\theta]_i^{222} - [\theta]_{coil}^{222}}{[\theta]_{helix}^{222} - [\theta]_{coil}^{222}} \quad (\text{Eq. 3})$$

Combining Equations 1 and 3, we can write Θ as a function of the lipid:protein molar ratio X , obtaining the final form of the fitting equation.

$$[\theta]_i^{222} = [\theta]_{coil}^{222} + \frac{[\theta]_{helix}^{222} - [\theta]_{coil}^{222}}{2} \left[1 + \frac{X}{N} + \frac{1}{Nk_B P_t} - \sqrt{\left(1 + \frac{X}{N} + \frac{1}{Nk_B P_t} \right)^2 - \frac{4X}{N}} \right] \quad (\text{Eq. 4})$$

Because the molar lipid concentration in the cuvette is used as L_p , the N value, rather than the number of binding sites, measures the average number of lipid molecules forming one lipid binding site.

ITC

ITC measurements were performed on an iTC200 instrument (Malvern Instruments, Westborough, MA). Typically, 18 fresh SUV dispersion injections of 2 μ l each (+1 0.4- μ l preinjection) were titrated in the calorimeter chamber. Both linear baselines and control titrations of vesicles into buffer alone were used for baseline subtraction. The data were processed using the MicroCal ORIGIN 7.0 software (Malvern Instruments, OriginLab, Northampton, MA). After integration of the differential power signal, stoichiometric and thermodynamic parameters (N , ΔH , and k_B) were determined by fitting the sigmoidal titration curve using an N -independent binding site model (with the sites assumed to localize on the vesicles' outer surface). The apparent (given the approximations used) microscopic binding constant was labeled k_B . The total lipid concentration in the syringe was taken as the ligand concentration; thus, the N value measures the average number of lipid molecules forming a binding site. In all experiments, the stoichiometric and thermodynamic parameters were obtained from three independent titrations (although performed using the same vesicle preparation). ΔS was obtained from $\Delta G = -RT \ln k_B$ and ΔH .

Dynamic light scattering

The hydrodynamic radius of the liposomes (5 mM) was measured using a DynaPro dynamic light scattering instrument (Wyatt Technology Corp., Santa Barbara, CA). The dynamic light-scattering cell was thermostated at 20.0 °C; the hydrodynamic radii were obtained by averaging three independent samples.

Phospholipid vesicle clearance assay

The ability of α Syn and its mutants to remodel phospholipid vesicles was studied by measuring light scattering as a function of time using a Jasco V-550 UV/VIS spectrophotometer as described previously (47). Briefly, light scattering was monitored at 500 nm using a 2-nm slit width and a "medium" response time. MLVs were prepared in 20 mM HEPES and 100 mM NaCl (pH 7.4). No changes in light scattering of the MLV dispersion were observed in the absence of protein. CD-monitored vesicle clearance experiments were performed at 25 °C using a Jasco J-810 spectropolarimeter in a 1-mm quartz cell. Blanks were recorded under similar conditions and subtracted.

E46K-like α -synuclein mutants disrupt membrane selectivity

Transmission electron microscopy

Transmission electron microscopy specimens were prepared on carbon-coated formvar grids (Electron Microscopy Sciences, Hatfield, PA). Samples were adsorbed on the grids for 5 min and then negatively stained with 1% (w/v) aqueous uranyl acetate. Images were taken on a JEOL 1400 transmission electron microscope (JEOL USA Inc., Peabody, MA) at an accelerating voltage of 100 kV.

NMR

^1H - ^{15}N HSQC spectra of ^{15}N -labeled αSyn were recorded in 50 mM phosphate buffer (pH 6.8) and 10% D_2O . Spectra were recorded on a 600 MHz Bruker AVANCE II spectrometer (Bruker, Billerica, MA) equipped with a Prodigy CryoProbe at either 15 °C (120 μM αSyn) or 37 °C (120 μM αSyn and 15 mM SDS). Resonances were assigned, where possible, by inspection and comparison with the previously assigned spectra of WT αSyn (BMRB 5744, 6968) (33, 34). Data processing and analysis were performed with Bruker TopSpin software and CcpNmr Analysis (63). Perturbations in the chemical shift values for ^1H and ^{15}N were calculated as $[(\Delta\delta^1\text{H})^2 + (0.15 \cdot \Delta\delta^{15}\text{N})^2]^{1/2}$.

Data analysis and primary sequence analysis

Data analysis was performed using GraphPad Prism 7 (GraphPad Software, La Jolla, CA). Statistical significance was determined by ordinary one-way analysis of variance followed by Sidak's multiple comparisons test, with a single pooled variance: n.s., $p > 0.05$; *, $p \leq 0.05$; **, $p \leq 0.01$; ***, $p \leq 0.001$; ****, $p \leq 0.0001$.

Primary sequence analyses were performed with Microsoft Excel (Microsoft, Redmond, WA) and Wolfram Mathematica 10 (Wolfram Research, Champaign, IL), using the hydropathy index values of Kyte and Doolittle (64). Helical wheel and net diagrams were plotted using in-house Python code, modeled after the WHEEL and HELNET programs (65).

Author contributions—M. R., A. E. P., H. J., J. C. P., L. F.-O., and D. S. P. data curation; M. R., A. E. P., H. J., J. C. P., L. F.-O., A. A., R. L., J. V., and T. B. formal analysis; M. R., A. E. P., H. J., J. C. P., L. F.-O., and D. S. P. investigation; M. R., A. E. P., H. J., J. C. P., L. F.-O., and A. A. visualization; M. R., J. V., and T. B. writing-original draft; M. R., A. E. P., H. J., J. C. P., L. F.-O., D. S. P., A. A., R. L., J. V., and T. B. writing-review and editing; D. S. P. validation; D. S. P. methodology; A. A. software; R. L., J. V., and T. B. conceptualization; R. L., J. V., and T. B. supervision; R. L., J. V., and T. B. funding acquisition.

Acknowledgments—We thank Kelly Arnett and the Center for Macromolecular Interactions at the Harvard Medical School Department of Biological Chemistry and Molecular Pharmacology for assistance with CD and ITC measurements and many helpful discussions; Simon Jenni, Yoana Dimitrova, and the Harrison laboratory at the Harvard Medical School Department of Biological Chemistry and Molecular Pharmacology for the use of instruments and assistance with dynamic light scattering measurements; and Jim Lee and the Molecular Biology Core Facilities of the Dana-Farber Cancer Institute for assistance with MS. Finally, we also thank our colleagues at the Ann Romney Center for Neurologic Diseases for many helpful discussions and Martina Astrid Rodda for editing and revising the manuscript.

References

- Poewe, W., Seppi, K., Tanner, C. M., Halliday, G. M., Brundin, P., Volkmann, J., Schrag, A.-E., and Lang, A. E. (2017) Parkinson disease. *Nat. Rev. Dis. Prim.* **3**, 17013 [CrossRef Medline](#)
- de Lau, L. M., and Breteler, M. M. (2006) Epidemiology of Parkinson's disease. *Lancet Neurol.* **5**, 525–535 [CrossRef Medline](#)
- Iwai, A., Masliah, E., Yoshimoto, M., Ge, N., Flanagan, L., de Silva, H. A., Kittel, A., and Saitoh, T. (1995) The precursor protein of non-A β component of Alzheimer's disease amyloid is a presynaptic protein of the central nervous system. *Neuron* **14**, 467–475 [CrossRef Medline](#)
- Shulman, J. M., De Jager, P. L., and Feany, M. B. (2011) Parkinson's disease: genetics and pathogenesis. *Annu. Rev. Pathol.* **6**, 193–222 [CrossRef Medline](#)
- Bendor, J. T., Logan, T. P., and Edwards, R. H. (2013) The function of α -synuclein. *Neuron* **79**, 1044–1066 [CrossRef Medline](#)
- Lautenschläger, J., Kaminski, C. F., and Kaminski Schierle, G. S. (2017) α -Synuclein: regulator of exocytosis, endocytosis, or both? *Trends Cell Biol.* **27**, 468–479 [CrossRef Medline](#)
- Grace, A. A., and Bunney, B. S. (1984) The control of firing pattern in nigral dopamine neurons: burst firing. *J. Neurosci.* **4**, 2877–2890 [CrossRef Medline](#)
- Weinreb, P. H., Zhen, W., Poon, A. W., Conway, K. A., and Lansbury, P. T. (1996) NACP, a protein implicated in Alzheimer's disease and learning, is natively unfolded. *Biochemistry* **35**, 13709–13715 [CrossRef Medline](#)
- Nielsen, J. T., and Mulder, F. A. (2016) There is diversity in disorder—"in all chaos there is a cosmos, in all disorder a secret order". *Front. Mol. Biosci.* **3**, 4 [Medline](#)
- George, J. M., Jin, H., Woods, W. S., and Clayton, D. F. (1995) Characterization of a novel protein regulated during the critical period for song learning in the zebra finch. *Neuron* **15**, 361–372 [CrossRef Medline](#)
- Bussell, R., Jr., and Eliezer, D. (2003) A structural and functional role for 11-mer repeats in α -synuclein and other exchangeable lipid binding proteins. *J. Mol. Biol.* **329**, 763–778 [CrossRef Medline](#)
- Varkey, J., Mizuno, N., Hegde, B. G., Cheng, N., Steven, A. C., and Langen, R. (2013) α -Synuclein oligomers with broken helical conformation form lipoprotein nanoparticles. *J. Biol. Chem.* **288**, 17620–17630 [CrossRef Medline](#)
- Davidson, W. S., Jonas, A., Clayton, D. F., and George, J. M. (1998) Stabilization of α -synuclein secondary structure upon binding to synthetic membranes. *J. Biol. Chem.* **273**, 9443–9449 [CrossRef Medline](#)
- Eliezer, D., Kutluay, E., Bussell, R., Jr., and Browne, G. (2001) Conformational properties of α -synuclein in its free and lipid-associated states. *J. Mol. Biol.* **307**, 1061–1073 [CrossRef Medline](#)
- Takamori, S., Holt, M., Stenius, K., Lemke, E. A., Grønborg, M., Riedel, D., Urlaub, H., Schenck, S., Brügger, B., Ringler, P., Müller, S. A., Rammner, B., Gräter, F., Hub, J. S., De Groot, B. L., et al. (2006) Molecular anatomy of a trafficking organelle. *Cell* **127**, 831–846 [CrossRef Medline](#)
- Pranke, I. M., Morello, V., Bigay, J., Gibson, K., Verbavatz, J.-M., Antonny, B., and Jackson, C. L. (2011) α -Synuclein and ALPS motifs are membrane curvature sensors whose contrasting chemistry mediates selective vesicle binding. *J. Cell Biol.* **194**, 89–103 [CrossRef Medline](#)
- Jao, C. C., Hegde, B. G., Chen, J., Haworth, I. S., and Langen, R. (2008) Structure of membrane-bound α -synuclein from site-directed spin labeling and computational refinement. *Proc. Natl. Acad. Sci. U.S.A.* **105**, 19666–19671 [CrossRef Medline](#)
- Fortin, D. L., Troyer, M. D., Nakamura, K., Kubo, S., Anthony, M. D., and Edwards, R. H. (2004) Lipid rafts mediate the synaptic localization of α -synuclein. *J. Neurosci.* **24**, 6715–6723 [CrossRef Medline](#)
- Fortin, D. L., Nemani, V. M., Voglmaier, S. M., Anthony, M. D., Ryan, T. A., and Edwards, R. H. (2005) Neural activity controls the synaptic accumulation of α -synuclein. *J. Neurosci.* **25**, 10913–10921 [CrossRef Medline](#)
- Auluck, P. K., Caraveo, G., and Lindquist, S. (2010) α -Synuclein: membrane interactions and toxicity in Parkinson's disease. *Annu. Rev. Cell Dev. Biol.* **26**, 211–233 [CrossRef Medline](#)
- Cooper, A. A., Gitler, A. D., Cashikar, A., Haynes, C. M., Hill, K. J., Bhullar, B., Liu, K., Xu, K., Strathearn, K. E., Liu, F., Cao, S., Caldwell, K. A.,

- Caldwell, G. A., Marsischky, G., Kolodner, R. D., *et al.* (2006) α -Synuclein blocks ER-Golgi traffic and Rab1 rescues neuron loss in Parkinson's models. *Science* **313**, 324–328 [CrossRef Medline](#)
22. Gitler, A. D., Bevis, B. J., Shorter, J., Strathearn, K. E., Hamamichi, S., Su, L. J., Caldwell, K. A., Caldwell, G. A., Rochet, J.-C., McCaffery, J. M., Barlowe, C., and Lindquist, S. (2008) The Parkinson's disease protein α -synuclein disrupts cellular Rab homeostasis. *Proc. Natl. Acad. Sci. U.S.A.* **105**, 145–150 [CrossRef Medline](#)
 23. Thayanidhi, N., Helm, J. R., Nycz, D. C., Bentley, M., Liang, Y., and Hay, J. C. (2010) α -Synuclein delays endoplasmic reticulum (ER)-to-Golgi transport in mammalian cells by antagonizing ER/Golgi SNAREs. *Mol. Biol. Cell* **21**, 1850–1863 [CrossRef Medline](#)
 24. Shahmoradian, S. H., Lewis, A. J., Genoud, C., Graff-Meyer, A., Hench, J., Moors, T., Schweighauser, G., Wang, J., Goldie, K. N., Suetterlin, R., Castano-Diez, D., Perez-Navarro, P., Huisman, E., Ipsen, S., Ingrassia, A., *et al.* (2018) Lewy pathology in Parkinson's disease consists of a crowded organellar, membranous medley. *bioRxiv* [CrossRef](#)
 25. Dettmer, U., Newman, A. J., von Saucken, V. E., Bartels, T., and Selkoe, D. (2015) KTKEGV repeat motifs are key mediators of normal α -synuclein tetramerization: their mutation causes excess monomers and neurotoxicity. *Proc. Natl. Acad. Sci. U.S.A.* **112**, 9596–9601 [CrossRef Medline](#)
 26. Dettmer, U., Ramalingam, N., von Saucken, V. E., Kim, T.-E., Newman, A. J., Terry-Kantor, E., Nuber, S., Ericsson, M., Fanning, S., Bartels, T., Lindquist, S., Levy, O. A., and Selkoe, D. (2017) Loss of native α -synuclein multimerization by strategically mutating its amphipathic helix causes abnormal vesicle interactions in neuronal cells. *Hum. Mol. Genet.* **26**, 3466–3481 [CrossRef Medline](#)
 27. Zarranz, J. J., Alegre, J., Gómez-Esteban, J. C., Lezcano, E., Ros, R., Ampuero, I., Vidal, L., Hoenicka, J., Rodriguez, O., Atarés, B., Llorens, V., Gomez Tortosa, E., del Ser, T., Muñoz, D. G., and de Yebenes, J. G. (2004) The new mutation, E46K, of α -synuclein causes Parkinson and Lewy body dementia. *Ann. Neurol.* **55**, 164–173 [CrossRef Medline](#)
 28. Dettmer, U., Newman, A. J., Soldner, F., Luth, E. S., Kim, N. C., von Saucken, V. E., Sanderson, J. B., Jaenisch, R., Bartels, T., and Selkoe, D. (2015) Parkinson-causing α -synuclein missense mutations shift native tetramers to monomers as a mechanism for disease initiation. *Nat. Commun.* **6**, 7314 [CrossRef Medline](#)
 29. Bartels, T., Choi, J. G., and Selkoe, D. J. (2011) α -Synuclein occurs physiologically as a helically folded tetramer that resists aggregation. *Nature* **477**, 107–110 [CrossRef Medline](#)
 30. Dettmer, U., Selkoe, D., and Bartels, T. (2016) New insights into cellular α -synuclein homeostasis in health and disease. *Curr. Opin. Neurobiol.* **36**, 15–22 [CrossRef Medline](#)
 31. Nuber, S., Rajsombath, M., Minakaki, G., Winkler, J., Müller, C. P., Ericsson, M., Caldarone, B., Dettmer, U., and Selkoe, D. J. (2018) Abrogating native α -synuclein tetramers in mice causes a L-DOPA-responsive motor syndrome closely resembling Parkinson's disease. *Neuron* **100**, 75–90.e5 [CrossRef Medline](#)
 32. Fredenburgh, R. A., Rospigliosi, C., Meray, R. K., Kessler, J. C., Lashuel, H. A., Eliezer, D., and Lansbury, P. T. (2007) The impact of the E46K mutation on the properties of α -synuclein in its monomeric and oligomeric states. *Biochemistry* **46**, 7107–7118 [CrossRef Medline](#)
 33. Chandra, S., Chen, X., Rizo, J., Jahn, R., and Südhof, T. C. (2003) A broken α -helix in folded α -Synuclein. *J. Biol. Chem.* **278**, 15313–15318 [CrossRef Medline](#)
 34. Bermel, W., Bertini, I., Felli, I. C., Lee, Y.-M., Luchinat, C., and Pierattelli, R. (2006) Protonless NMR experiments for sequence-specific assignment of backbone nuclei in unfolded proteins. *J. Am. Chem. Soc.* **128**, 3918–3919 [CrossRef Medline](#)
 35. Bertoncini, C. W., Jung, Y.-S., Fernandez, C. O., Hoyer, W., Griesinger, C., Jovin, T. M., and Zweckstetter, M. (2005) Release of long-range tertiary interactions potentiates aggregation of natively unstructured α -synuclein. *Proc. Natl. Acad. Sci. U.S.A.* **102**, 1430–1435 [CrossRef Medline](#)
 36. Croke, R. L., Sallum, C. O., Watson, E., Watt, E. D., and Alexandrescu, A. T. (2008) Hydrogen exchange of monomeric alpha-synuclein shows unfolded structure persists at physiological temperature and is independent of molecular crowding in *Escherichia coli*. *Protein Sci.* **17**, 1434–1445 [CrossRef Medline](#)
 37. Rospigliosi, C. C., McClendon, S., Schmid, A. W., Ramlall, T. F., Barré, P., Lashuel, H. A., and Eliezer, D. (2009) E46K Parkinson's-linked mutation enhances C-terminal-to-N-terminal contacts in α -synuclein. *J. Mol. Biol.* **388**, 1022–1032 [CrossRef Medline](#)
 38. Wise-Scira, O., Dunn, A., Aloglu, A. K., Sakallioğlu, I. T., and Coskuner, O. (2013) Structures of the E46K mutant-type α -synuclein protein and impact of E46K mutation on the structures of the wild-type α -synuclein protein. *ACS Chem. Neurosci.* **4**, 498–508 [CrossRef Medline](#)
 39. Jao, C. C., Der-Sarkissian, A., Chen, J., and Langen, R. (2004) Structure of membrane-bound α -synuclein studied by site-directed spin labeling. *Proc. Natl. Acad. Sci. U.S.A.* **101**, 8331–8336 [CrossRef Medline](#)
 40. Ulmer, T. S., Bax, A., Cole, N. B., and Nussbaum, R. L. (2005) Structure and dynamics of micelle-bound human α -synuclein. *J. Biol. Chem.* **280**, 9595–9603 [CrossRef Medline](#)
 41. Choi, W., Zibae, S., Jakes, R., Serpell, L. C., Davletov, B., Crowther, R. A., and Goedert, M. (2004) Mutation E46K increases phospholipid binding and assembly into filaments of human α -synuclein. *FEBS Lett.* **576**, 363–368 [CrossRef Medline](#)
 42. Bodner, C. R., Maltsev, A. S., Dobson, C. M., and Bax, A. (2010) Differential phospholipid binding of α -synuclein variants implicated in Parkinson's disease revealed by solution NMR spectroscopy. *Biochemistry* **49**, 862–871 [CrossRef Medline](#)
 43. Burré, J., Sharma, M., and Südhof, T. C. (2012) Systematic mutagenesis of α -synuclein reveals distinct sequence requirements for physiological and pathological activities. *J. Neurosci.* **32**, 15227–15242 [CrossRef Medline](#)
 44. Mammen, M., Choi, S.-K., and Whitesides, G. M. (1998) Polyvalent interactions in biological systems: implications for design and use of multivalent ligands and inhibitors. *Angew. Chem. Int. Ed. Engl.* **37**, 2754–2794 [CrossRef Medline](#)
 45. Bartels, T., Ahlstrom, L. S., Leftin, A., Kamp, F., Haass, C., Brown, M. F., and Beyer, K. (2010) The N-terminus of the intrinsically disordered protein α -synuclein triggers membrane binding and helix folding. *Biophys. J.* **99**, 2116–2124 [CrossRef Medline](#)
 46. Fusco, G., De Simone, A., Gopinath, T., Vostrikov, V., Vendruscolo, M., Dobson, C. M., and Veglia, G. (2014) Direct observation of the three regions in α -synuclein that determine its membrane-bound behaviour. *Nat. Commun.* **5**, 3827 [CrossRef Medline](#)
 47. Varkey, J., Isas, J. M., Mizuno, N., Jensen, M. B., Bhatia, V. K., Jao, C. C., Petrlova, J., Voss, J. C., Stamou, D. G., Steven, A. C., and Langen, R. (2010) Membrane curvature induction and tubulation are common features of synucleins and apolipoproteins. *J. Biol. Chem.* **285**, 32486–32493 [CrossRef Medline](#)
 48. Mizuno, N., Varkey, J., Kegulian, N. C., Hegde, B. G., Cheng, N., Langen, R., and Steven, A. C. (2012) Remodeling of lipid vesicles into cylindrical micelles by α -synuclein in an extended α -helical conformation. *J. Biol. Chem.* **287**, 29301–29311 [CrossRef Medline](#)
 49. Conway, K. A., Lee, S. J., Rochet, J. C., Ding, T. T., Williamson, R. E., and Lansbury, P. T. (2000) Acceleration of oligomerization, not fibrillization, is a shared property of both α -synuclein mutations linked to early-onset Parkinson's disease: implications for pathogenesis and therapy. *Proc. Natl. Acad. Sci. U.S.A.* **97**, 571–576 [CrossRef Medline](#)
 50. Lashuel, H. A., Petre, B. M., Wall, J., Simon, M., Nowak, R. J., Walz, T., and Lansbury, P. T. (2002) Alpha-synuclein, especially the Parkinson's disease-associated mutants, forms pore-like annular and tubular protofibrils. *J. Mol. Biol.* **322**, 1089–1102 [CrossRef Medline](#)
 51. Dettmer, U. (2018) Rationally designed variants of α -synuclein illuminate its *in vivo* structural properties in health and disease. *Front. Neurosci.* **12**, 623 [CrossRef Medline](#)
 52. Eisenberg, D., Weiss, R. M., and Terwilliger, T. C. (1982) The helical hydrophobic moment: a measure of the amphiphilicity of a helix. *Nature* **299**, 371–374 [CrossRef Medline](#)
 53. Eisenberg, D., Weiss, R. M., and Terwilliger, T. C. (1984) The hydrophobic moment detects periodicity in protein hydrophobicity. *Proc. Natl. Acad. Sci. U.S.A.* **81**, 140–144 [CrossRef Medline](#)
 54. McLachlan, A. D., and Stewart, M. (1976) The 14-fold periodicity in α -tropomyosin and the interaction with actin. *J. Mol. Biol.* **103**, 271–298 [CrossRef Medline](#)

E46K-like α -synuclein mutants disrupt membrane selectivity

55. Finer-Moore, J., and Stroud, R. M. (1984) Amphipathic analysis and possible formation of the ion channel in an acetylcholine receptor. *Proc. Natl. Acad. Sci. U.S.A.* **81**, 155–159 [CrossRef Medline](#)
56. Lokappa, S. B., and Ulmer, T. S. (2011) Alpha-synuclein populates both elongated and broken helix states on small unilamellar vesicles. *J. Biol. Chem.* **286**, 21450–21457 [CrossRef Medline](#)
57. Schiffer, M., and Edmundson, A. B. (1967) Use of helical wheels to represent the structures of proteins and to identify segments with helical potential. *Biophys. J.* **7**, 121–135 [CrossRef Medline](#)
58. Lim, V. I. (1978) Polypeptide chain folding through a highly helical intermediate as a general principle of globular protein structure formation. *FEBS Lett.* **89**, 10–14 [CrossRef Medline](#)
59. Jiang, Z., Hess, S. K., Heinrich, F., and Lee, J. C. (2015) Molecular details of α -synuclein membrane association revealed by neutrons and photons. *J. Phys. Chem. B.* **119**, 4812–4823 [CrossRef Medline](#)
60. Fanning, S., Haque, A., Imberdis, T., Baru, V., Barrasa, M. I., Nuber, S., Termine, D., Ramalingam, N., Ho, G. P. H., Noble, T., Sandoe, J., Lou, Y., Landgraf, D., Freyzon, Y., Newby, G., *et al.* (2018) Lipidomic analysis of α -synuclein neurotoxicity identifies stearoyl CoA desaturase as a target for Parkinson treatment. *Mol. Cell* **73**, 1001–1014.e8 [CrossRef Medline](#)
61. Vincent, B. M., Tardiff, D. F., Piotrowski, J. S., Aron, R., Lucas, M. C., Chung, C. Y., Bacherman, H., Chen, Y., Pires, M., Subramaniam, R., Doshi, D. B., Sadlish, H., Raja, W. K., Solís, E. J., Khurana, V., *et al.* (2018) Inhibiting stearoyl-CoA desaturase ameliorates α -synuclein cytotoxicity. *Cell Rep.* **25**, 2742–2754.e31 [CrossRef Medline](#)
62. Indyk, L., and Fisher, H. F. (1998) Theoretical aspects of isothermal titration calorimetry. *Methods Enzymol.* **295**, 350–364 [CrossRef Medline](#)
63. Vranken, W. F., Boucher, W., Stevens, T. J., Fogh, R. H., Pajon, A., Llinas, M., Ulrich, E. L., Markley, J. L., Ionides, J., and Laue, E. D. (2005) The CCPN data model for NMR spectroscopy: development of a software pipeline. *Proteins* **59**, 687–696 [CrossRef Medline](#)
64. Kyte, J., and Doolittle, R. F. (1982) A simple method for displaying the hydropathic character of a protein. *J. Mol. Biol.* **157**, 105–132 [CrossRef Medline](#)
65. Jones, M. K., Anantharamaiah, G. M., and Segrest, J. P. (1992) Computer programs to identify and classify amphipathic α helical domains. *J. Lipid Res.* **33**, 287–296 [Medline](#)