

# Cyanidin and Cyanidin-3-Glucoside Alleviate Peptic Ulcer Disease: Insights from in vitro, and in vivo Studies

Deshanda Kurniawan Prayoga<sup>1,\*</sup>, Diah Lia Aulifa<sup>2,\*</sup>, Arif Budiman<sup>3,\*</sup>, Jutti Levita<sup>4,\*</sup> , Supat Jiranusornkul<sup>5,\*</sup>

<sup>1</sup>Doctoral Program in Pharmacy, Faculty of Pharmacy, Padjadjaran University, Sumedang, West Java, Indonesia; <sup>2</sup>Department of Pharmaceutical Analysis and Medicinal Chemistry, Faculty of Pharmacy, Padjadjaran University, Sumedang, Indonesia; <sup>3</sup>Department of Pharmaceutics and Pharmaceutical Technology, Faculty of Pharmacy, Padjadjaran University, Sumedang, Indonesia; <sup>4</sup>Department of Pharmacology and Clinical Pharmacy, Faculty of Pharmacy, Padjadjaran University, Sumedang, Indonesia; <sup>5</sup>Department of Pharmaceutical Science, Faculty of Pharmacy, Chiang Mai University, Chiang Mai, Thailand

\*These authors contributed equally to this work

Correspondence: Jutti Levita, Department of Pharmacology and Clinical Pharmacy, Faculty of Pharmacy, Padjadjaran University, Sumedang, West Java, 46363, Indonesia, Tel +6222 84288888 Ext 3510, Email [jutti.levita@unpad.ac.id](mailto:jutti.levita@unpad.ac.id)

**Abstract:** Peptic ulcer disease (PUD) remains a significant global health issue, affecting millions despite a decrease in overall prevalence. However, complications continue to persist, with substantial mortality rates in regions like India and China. Current treatments, though effective, have limitations, driving interest in plant-derived therapy. Anthocyanins, including cyanidin and cyanidin-3-glucoside (C3G), are known for their antioxidant and anti-inflammatory properties. This study aims to explore the potential of cyanidin and C3G in alleviating PUD, focusing on their mechanisms of action and therapeutic efficacy in preclinical studies. Articles were searched in Scopus and PubMed databases and filtered for publication from 2014 to 2024, resulting in 89 articles from Scopus and 11 articles from PubMed. The articles were further screened by title, abstract, and full text, resulting in 6 articles. Cyanidin and C3G were described to be able to alleviate PUD by inhibiting the cytokine pro-inflammatory, reducing inflammation in gastric mucosa, and reducing lipid peroxidation in the gastric mucosa. These compounds have proven effective in managing other health problems, including peptic ulcers, but more in-depth exploration in clinical settings is required to confirm therapeutic potential in humans. It is necessary to validate the therapeutic efficacy and safety in human populations. This review provides an overview of preclinical studies of cyanidin and C3G, such as in vitro and in vivo, focusing on mechanism of action or their effectiveness in alleviating peptic ulcers.



**Keywords:** anthocyanins, anti-inflammatory, antioxidant, gastroprotective, preclinical studies

## Introduction

Peptic ulcer disease (PUD) affects millions globally, with an estimated lifetime prevalence of 5–10% in the general population.<sup>1–3</sup> Although the overall prevalence of PUD has decreased over recent decades, the incidence of complications remains steady.<sup>4,5</sup> In 2019, there were approximately 8.09 million cases worldwide, marking a 25.82% increase since 1990. However, despite this rise in case numbers, the age-standardized prevalence rate dropped from 143.47 per 100,000 population in 1990 to 99.50 per 100,000 population in 2019.<sup>6,7</sup> Based on world life expectancy data ([www.worldlifeexpectancy.com](http://www.worldlifeexpectancy.com)) and the World Health Organization's Mortality Database ([www.platfrom.who.int/mortality](http://www.platfrom.who.int/mortality)), India (68,108 deaths) and China (37,723 deaths) report the highest number of ulcer-related deaths, followed by several other countries with significant mortality rates. Regions like Australia, Canada, and Europe show a moderate impact, while some countries, including New Zealand and certain African nations, report fewer than 100 deaths, indicating a lower effect (Figure 1).

The geographical variation in ulcer-related mortality is influenced by factors such as *Helicobacter pylori* infections, prolonged non-steroidal anti-inflammatory drugs (NSAIDs) use, and lifestyle choices that may weaken the stomach's protective barrier,<sup>8</sup> leading to ulcers and severe complications like perforation or obstruction and disrupting physiological

## Graphical Abstract



**Figure 1** The global number of deaths due to PUD, reported by countries to WHO, covers the period from 2010 to 2022 and is compiled by World Life Expectancy (This figure was created using [mapchart.net](https://www.mapchart.net/)).

functions.<sup>9–11</sup> Peptic ulcers can also result from excessive alcohol consumption, elevated stress, and bile salt reflux.<sup>12</sup> Current treatments for PUD, including proton pump inhibitors (PPIs), histamine-2 receptor antagonists (H2RAs), antacids, and sucralfate, operate by suppressing gastric acid secretion, neutralizing stomach acid, and protecting the gastric mucosa. Additionally, the development and refinement of synthetic medications have significantly decreased the incidence of complications related to PUD, including bleeding and perforations.<sup>13,14</sup> Notably, PPIs are exceptionally effective in controlling acid production and fostering an environment conducive to ulcer healing, and their usage has increased substantially over the past decade.<sup>15</sup> However, prolonged use of these drugs raises side effects and financial costs. Sudden discontinuation of PPI may cause recurrence, thus secure discontinuation strategy is by tapering the dose.<sup>16–18</sup>

Ulcers pose a significant clinical challenge, often leading to weak mucosal regeneration and increased vulnerability to further damage. As the first line, PPIs were thought to have a low incidence of side effects. Still, evidence has cumulated to indicate that long-term use might not be as harmless as first considered, with concerns about possible adverse side effects, such as hypomagnesemia and drug interactions, raised.<sup>15</sup> Nonetheless, the discontinuation of PPIs can trigger a withdrawal, such as rebound acid hypersecretion (RAHS), where the stomach produces excessive acid levels beyond pre-treatment amounts.<sup>19,20</sup> Due to these challenges, there is increasing interest in alternative treatments from natural sources, as reported by the WHO, that 80–85% of the global population relies on natural products for disease management.<sup>21,22</sup>


Anthocyanins, eg, cyanidin and cyanidin-3-glucoside (C3G), are plant-derived secondary metabolites belonging to the monomeric anthocyanin class within the flavonoid family.<sup>23–25</sup> These water-soluble compounds are widely found in fruits, vegetables, and various plant parts, including leaves, petals, flowers, and red-colored fruits.<sup>26</sup> They are traditionally used as medicinal treatments in many countries and have been extensively reported to offer therapeutic potential against various diseases.<sup>27</sup> Notably, cyanidin is recognized for its significant anti-inflammatory and antioxidant properties, suggesting potential therapeutic applications for conditions like asthma,<sup>28,29</sup> cancer,<sup>30–32</sup> diabetes,<sup>33–35</sup> and inflammatory bowel disease.<sup>36,37</sup> Additionally, cyanidins exhibit antiproliferative effects and promote apoptosis, contributing to their role in cancer prevention. Recent studies have highlighted the potential of anthocyanins in modulating key receptors such as tumor necrosis factor receptor 1 (TNFR1) and toll-like receptor 1–9 (TLR1-9), which can reduce inflammation and oxidative stress.<sup>38,39</sup> Furthermore, these compounds enhance mucosal defense by neutralizing reactive oxygen species (ROS) and strengthening endogenous antioxidant systems, including catalase (CAT) and glutathione (GSH).<sup>40</sup> This presents a promising avenue for developing safer and more effective treatment strategies for PUD.

Considering all, this review aims to provide a comprehensive overview of the role of cyanidin and C3G in alleviating PUD. The review synthesizes from *in vitro* studies, animal models, and human clinical trials to present an understanding of the mechanisms through which these plant-derived metabolites exert gastroprotective effects. This review thoroughly analyzes preclinical studies investigating the critical areas of cyanidin and C3G potential in alleviating PUD, focusing on their efficacy and safety across various study models. Special emphasis is placed on elucidating the underlying mechanisms of action, including their role in enhancing gastric mucosal protection, modulating oxidative stress, and reducing inflammation. This review will provide a nuanced understanding and may contribute to improved therapeutic outcomes in PUD management.

## Methods

### Search Strategy

Briefly, a literature search using the (1) PubMed database using the keywords cyanidin OR cyanidin-3-glucoside AND gastroprotective searched within Abstract, free full text, in the last 10 years, and resulted in *n* = 11; (2) in the Scopus database using the keywords cyanidin OR cyanidin-3-glucoside AND gastroprotective searched within All fields, resulted in *n* = 89, with a total of all publications found through these searches were reviewed. The authors retrieved and evaluated all documents that met the inclusion criteria, such as using cyanidin and cyanidin-3-glucoside in treating peptic ulcers. Finally, 6 articles were selected for review, as depicted in Figure 2. However, an additional search was conducted to support the discussion during the process.



**Figure 2** The study design of the article review.

## Data Extraction

Each investigator independently evaluated and selected articles that were reasonable or relevant to the topic. Key data extracted from each article includes the medicinal plant isolated, collection location and year, plant part used, extraction method and solvents, other metabolites aside from cyanidin and cyanidin-3-glucoside, animal models employed, study controls, inducement method dosage and duration, reported findings, and interpretations of the results.

## Results

Cyanidin and C3G may have an essential role in human health. Studies considered these metabolites to work synergistically as a gastroprotective by inhibiting pro-inflammatory cytokines, reducing inflammation, and antioxidants. Of the 6 articles, 3 discussed in vitro studies, 1 discussed in vivo studies, and 2 discussed both in vitro and in vivo studies. For example, 3 articles described the mechanism for alleviating PUD as an antioxidant that decreases or inhibits ROS productions; 2 articles described the mechanism for alleviating PUD as an anti-inflammatory that inhibits the production of pro-inflammatory cytokines, including IL-1 $\beta$ , IL-6, IL-8, IL-10, IL-12, IL-18, and TNF- $\alpha$ .<sup>41–43</sup> These are the key cytokines that contribute to the development and progression of the inflammatory pathways within the gastric mucosa.<sup>44</sup> Of these studies, three articles confirmed the effects of gastroprotective using in vivo studies.

## Chemical Structure and Properties of Cyanidin and Cyanidin 3- Glucoside

Cyanidins are natural, organic pigments that belong to the anthocyanins group.<sup>27</sup> They are the most represented pigments in plants,<sup>45,46</sup> including leaves, stems, and flowers, and in fruits and vegetables such as cranberry, blackberry, cherry, grapes, raspberry, apples, plums, peaches, beans, and onions.<sup>26,47–50</sup> Anthocyanins have been categorized to belong to the flavonoid family and possess a C6-C3-C6 molecular structure, 15 carbons constituted by a common skeleton of phenyl benzo- $\gamma$ -pyran, composed of two phenyl rings (A and B) and a heterocyclic ring (pyran, C).<sup>51</sup> Cyanidin or cyanidol (3,5,7,3',4'-pentahydroxy flavylum) is a relatively unstable molecule. It is present rarely in free form in plant tissues.<sup>52</sup> Cyanidin predominantly exists as glycosides, where monosaccharides or disaccharides are linked to anthocyanidin analogs through glycosidic bonds.<sup>53</sup> A secondary sugar residue is often attached to the aglycone structure.<sup>54</sup> Its glycosylation gives it better stability and water solubility.<sup>52</sup> Cyanidin is characterized by its high hydrophilicity, with a solubility of 0.049 mg/mL, a Log P of 3.05, a molecular surface area of 114.3 Å<sup>2</sup>, and molecular weight by mass spectroscopy in tandem with mass spectroscopy (MS/MS) at an m/z of 287.<sup>55</sup> It has a distinctive red-orange color, although this can change with pH, indeed, its solutions are red at acidic pH of < 3, purple/blue/violet at pH 3–10, and green to yellow at basic pH of > 10.<sup>52</sup> However, at very high pH, this compound loses its color due to its molecular degeneration.<sup>27</sup> Moreover, very stable salt formation with heavy metal cations may lead to different colors in plant tissues, in particular when the 3' and 4' positions contain free phenolic hydroxyls.<sup>56</sup>

Cyanidin-3-glucoside (C3G) and cyanidin-3-galactoside (C3Gal) were identified as the most widespread cyanidin glycosides in the plant kingdom.<sup>55</sup> Additionally, other glycosides, such as cyanidin-3-arabinoside, cyanidin-3-xyloside, cyanidin-3-sambubioside, cyanidin-3-sophoroside, cyanidin-3-rutinoside, were also found. These derivatives, belonging to the anthocyanin family, provide a basis for detailed exploration of the anthocyanin composition across various plant species.<sup>25</sup>

It has long been assumed that anthocyanidins only, ie, the aglycones of anthocyanins, were absorbed by intestinal cells and entered the bloodstream due to the absence of a bound sugar residue. Previously, it was presumed that anthocyanins were poorly absorbed because specific enzymes to hydrolyze glycosidic bonds were unknown.<sup>25</sup> However, recent research has demonstrated the absorption of glycosidic flavonoids, particularly C3G, an anthocyanin derived from the aglycone cyanidin. C3G is a standard red pigment found naturally in various vegetables and fruits.<sup>57</sup> C3G is a water-soluble secondary metabolite of the monomeric anthocyanin class within the flavonoid family, widely utilized in food products such as yogurt and porridge attributed to their dual role as natural pigments,<sup>58–60</sup> and their potential health benefits from regular consumption.<sup>57</sup> C3G is characterized by its high hydrophilicity, with a solubility of 0.6 mg/mL, a Log P of 0.39, a molecular surface area of 193.4 Å<sup>2</sup>, and molecular weight by mass spectroscopy in tandem with mass spectroscopy (MS/MS) at an m/z of 449.<sup>61</sup> Furthermore, the UV-visible spectrum analysis of C3G across different pH levels (pH 3–8) revealed that C3G exhibits pH-dependent behavior.<sup>62</sup> Specifically, the results indicate that at pH 3, the absorbance is dominated by the flavylum cation, but this intensity diminishes at pH 4–5. At pH 5, the quinoidal species becomes predominant, and at pH 6–8, C3G transitions into quinoidal and chalcone forms.<sup>62</sup>



## Metabolism and Absorption of Cyanidin and Cyanidin 3-Glucoside


In the gastrointestinal tract, most anthocyanins, including C3G, remain stable, particularly in the stomach and upper intestine, where they are primarily absorbed.<sup>57,63,64</sup> As the pH increases in the intestine, anthocyanins lose their color and change form.<sup>53</sup> Although a significant amount (85%) is found in the distal intestine,<sup>65</sup> the stomach is still considered a key site for anthocyanin absorption. The absorption process is accompanied by extensive first-pass metabolism, allowing these compounds to enter the systemic circulation as metabolites.<sup>66</sup> C3G catabolism predominantly occurs in the distal small intestine (the ileum) and the upper large intestine (the colon), where the gut microbiota plays a crucial role in breaking it down.<sup>67–69</sup> Enzymes in the small intestine hydrolyze C3G into aglycones, which are further degraded into phenolic compounds. The gut microbiota cleaves the heterocyclic flavylium ring of C3G, leading to dehydroxylation or decarboxylation. Following these processes, Phase II and multistage metabolites, including bacterial metabolites, enter the liver and kidneys, where they undergo further modifications such as methylation, glucuronidation, and sulfation through enterohepatic and blood circulations.<sup>57,70</sup>

At the oral pH of approximately 6.8, C3G predominantly exists in its quinoidal form, with the glucose moiety enhancing solubility,<sup>71</sup> and nearly 50% of C3G undergoes biotransformation into metabolites like protocatechuic acid (PCA), facilitated by microbial B-glycosidase, which may contribute to its preventive effects against oral diseases,<sup>68,69,72–75</sup> and in the stomach it is rapidly absorbed by gastric epithelial cells (0.25 to 2 h),<sup>76–78</sup> after intake through active transport mechanisms,<sup>79</sup> including bili translocase, SGLT1 (sodium-glucose transporter 1), GLUT1 (glucose-transporter 1), GLUT3 (glucose-transporter 3), and mono-carboxylated transporter 1 (MCT1).<sup>77,80–82</sup> Despite efficient absorption, extensive first-pass metabolism significantly reduces the bioavailability of intact C3G, although both the intact compound and its metabolites are detectable in plasma shortly after ingestion.<sup>83–86</sup> Upon reaching the small intestine, C3G bioavailability is further diminished by 40–50% due to the intestinal environment and has been confirmed by studies conducted in humans,<sup>87</sup> mice,<sup>88</sup> and rats,<sup>89</sup> where factors such as pH, enzymatic activity, and the food matrix play crucial roles.<sup>90</sup> Although cleavage of the glucose moiety is not required for C3G metabolism, it does facilitate trans-epithelial transport.<sup>76,91</sup> Subsequent metabolism in the small intestine produces Phase I enzymes that metabolize anthocyanins into hippuric acid (HA), vanillic acid (VA), or isovanillic acid (IVA). Subsequently, C3G, cyanidin, PCA, VA, and IVA undergo further metabolism by phase II enzymes, forming their respective glucuronide or sulfate conjugates.<sup>61,67</sup> In the large intestine, C3G and its remaining metabolites are released from fibrous food matrices and undergo further transformation by the gut microbiome.<sup>67</sup> These metabolites are predominantly excreted after additional phase II conjugation, with less than 0.005% C3G remaining intact.<sup>87</sup> The slightly basic pH of the large intestine supports these transformations, leading to the production of various phenolic acids and other metabolites.<sup>71,92,93</sup>

In general, C3G is considered non-toxic within the range of regular dietary consumption. Safe intake levels (LD50) have been established in mice and rats at 25,000 mg/kg and 20,000 mg/kg, respectively, with no observed adverse effects; studies in rabbits administered anthocyanins orally (6 g/kg BW) revealed no changes in blood pressure.<sup>94</sup> C3G is considered safe and anti-mutagenic.<sup>95</sup> Despite C3G's chemical instability and low bioavailability, foods rich in anthocyanin (Figure 3) have shown biological value in treating many acute and chronic human disorders.<sup>96–99</sup> Multiple studies have also confirmed that cyanidin and C3G are essential in human health.

## In vitro Studies

The protective effect of these compounds in alleviating PUD is primarily based on their antioxidant ability in vitro studies (Table 1). However, cyanidin, C3G, and their bioactive phenolic metabolites, such as PCA, VA, and FA, can enhance the antioxidant enzyme system by increasing the activities of manganese-dependent superoxide dismutase (MnSOD) and glutathione (GSH).<sup>100,101</sup> Additionally, these compounds down-regulate pro-oxidant system by reducing cyclooxygenase-2 (COX-2), inducible nitric oxide synthase (iNOS),<sup>37,101</sup> as well as lowering levels of free radicals, including reactive oxygen species (ROS) such as superoxide anion ( $O_2^{\cdot-}$ ), superoxide hydroxy radicals ( $HO^{\cdot}$ ), alkyl peroxy radicals ( $ROO^{\cdot}$ ),<sup>102,103</sup> and reactive nitrogen species (RNS).<sup>37</sup> Cyanidin and its derivatives exhibit antioxidant effects primarily through their ability to donate hydrogen atoms or electrons, thereby neutralizing free radicals. The diphenyl structure in the B ring plays a key role in this activity. Additionally, the hydroxyl groups at specific positions (eg, 3' and 4') enhance the antioxidant capacity by stabilizing free radicals.<sup>104–108</sup>



**Figure 3** The role of cyanidin and C3G in alleviating PUD.

The in vitro studies also demonstrate that cyanidin and C3G exhibit significant anti-inflammatory effects in addition to their antioxidant activity (Table 1). These compounds inhibit the production of key pro-inflammatory cytokines such as IL-6, IL-10, and TNF- $\alpha$ , which are crucial in the development and progression of gastric ulcers.<sup>41–43</sup> Cyanidin and its derivatives' anti-inflammatory properties extend to gastric ulcers, where they effectively inhibit cytokines secreted by macrophages.<sup>113</sup> Specifically, C3G has been shown to inhibit TNF- $\alpha$  production by 50% and IL-6 production by 30% in human neutrophils,<sup>112</sup> Thus highlighting its potential therapeutic role in managing gastric inflammation.

## In vivo Studies

In vivo studies further confirm the anti-inflammatory potential of cyanidin and C3G (Table 2). These compounds significantly reduce the inflammation induced by ethanol and hydrochloric acid, which mimics gastric ulcers.<sup>111</sup> Results from various studies indicate that alcohol consumption is strongly linked to the development of gastric mucosal

**Table 1** In vitro Studies Approach in Cyanidin and Cyanidin 3-O-Glucoside

Medicinal Plant (Family), Collected In, Year	Plant Part Used/ Compound Used	Solvent	Class of Metabolites	Type of Study		%Inhibition or Effective Concentration		Interpretation	Ref.
				Chemical Reaction (Redox) Scavenging Activity Using	Determination of pro-Inflammatory Cytokines in Response to				
<i>Quercus crassifolia</i> Humb. and Bonpl. (Beech). Michoacan. 2018.	Barks	Ethanol	<b>Cyanidin chloride</b> , Gallic Acid	NBT Sodium nitroprusside ORAC HClO Fenton Reaction	N/A	EC <sub>50</sub>	40.9 µg/mL 653 µg/mL 873 µg/mL 1747 µg/mL 2024 µg/mL	Moderate Activity	[103]
<i>Callistemon citrinus</i> (Myrtaceae), Messina, 2020	Flowers	Acetic acid: methanol: water	<b>Cyanidin 3-Glucoside</b> , Cyanidin 3-5-diglucoside, Cyanidin-coumaroylglucoside-pyruvic acid, Peonidin-3,5-diglucoside	DPPH TEAC ORAC	N/A	IC <sub>50</sub> :	2.20 µg/mL 1.64 µg/mL 5.66 µg/mL	Strong Activity	[109]
<i>Euterpe oleracea</i> Mart. (Arecaceae), Sao Paulo, 2020	Fruits	Methanol, HCl	<b>Cyanidin-3-Glucoside</b> , Cyanidin-3,5-hexoside-pentoside, Pelargonin-3-rutinoside, Pelargonidin-3-glucoside, Peonidin-3-glucoside, Peonidin-3-rutinoside	DPPH	N/A	IC <sub>50</sub>	90%	Strong Activity	[110]
<i>Prunus cerasus</i> (Rosaceae). N/a	Fruits	Ethyl Acetate	<b>Cyanidin-3-Glucoside</b>	N/A	TNF- $\alpha$ IL-6 IL-10	IC <sub>50</sub>	1.75 µg/mL 1.04 µg/mL 1.99 µg/mL	Strong Activity	[111]
<i>Prunus spinosa</i> L. (Rosaceae). Krasnobród, 2018	Fruits	Methanol: water	<b>Cyanidin-3-Glucoside</b> , Cyanidin-3-rutinoside, Peonidin-3-glucoside	Luminol-dependent chemiluminescence	IL-8 TNF- $\alpha$	IC <sub>50</sub>	90% 50% 30%	Strong Activity	[112]

**Abbreviations:** DPPH, 2,2-diphenyl-1-picrylhydrazyl; EC, Effective Concentration; HClO, Hypochlorous Acid; IC, Inhibition Concentration; IL-6, Interleukin-6; IL-8, Interleukin-8; IL-10, Interleukin-10; NBT, Nitroblue Tetrazolium; ORAC, Oxygen Radical Absorbance Capacity; TEAC, Trolox Equivalent Antioxidant Capacity; TNF- $\alpha$ , Tumor Necrosis Factor-alpha.



**Table 2** In vivo Studies Approach in Cyanidin and Cyanidin-3-Glucoside

Isolated Medicinal Plants Authority (Family), Collected In, Year	Plant part, and Extraction Method (Solvent Used)	Class of Metabolites	Animal used in the Preclinical Study (n)	Control (Negative and Positive)	Inducement Method and Dose	Duration (Days)	Results	Interpretation	Ref.
<i>Euterpe oleracea</i> Mart. (Arecaceae), Sao Paulo, 2020	Fruits, Fractionation (Methanol, HCl)	<b>Cyanidin-3-Glucoside</b> , Cyanidin-3,5-hexoside-pentoside, Pelargonin-3-rutinoside, Pelargonidin-3-glucoside, Peonidin-3-glucoside, Peonidin-3-rutinoside	Female Wistar rats ( <i>Rattus norvegicus</i> ) (200–250 g) (n= 6)	Vehicle (Water) as control, and Omeprazole (30 mg/kg) as positive control	Ethanol (98% ethanol)	I	The plants that contained anthocyanidin including C3G showed doses 30 and 100 mg/kg were effective as gastroprotective which were reduced by 83% and 67% ulcer area	Strong as Gastroprotective	[110]
N/a	N/a	Purified <b>Cyanidin Chloride</b>	Male and female Swiss mice ( <i>Mus musculus</i> ) (n=20)	Vehicle (0.9% saline) as negative control and Lansoprazole (30 mg/kg) as Positive Control	Ethanol (Absolute Ethanol)	56	The purified cyanidin showed all doses (5; 10; 20 mg/kg) were effective and indicated a high level of gastroprotection. Male mice at doses 20 mg/kg achieved a 93.4% reduction in lesions (p<0.01) and female mice at the same doses achieved an 80.7% reduction in lesions;	Strong as Gastroprotective	[116]
<i>Prunus cerasus</i> (Rosaceae). N/a	Fruits, Sonicated (Ethyl acetate)	Purified <b>Cyanidin-3-Glucoside</b>	Male mice <i>Swiss-albino</i> ( <i>Mus musculus</i> ) (20–35 g) (n=28)	Vehicle (Saline) as negative control and Ranitidine HCl (50 mg/kg) as Positive Control	Ethanol (60% ethanol) and hydrochloride Acid (60%)	I	The purified C3G showed all doses (10; 15; 20 mg/kg) were effective as gastroprotective indicated 68.5% gastro-lesion reduction in 20 mg/kg doses, respectively	Strong as Gastroprotective	[111]
		Isolated <b>Cyanidin-3-Glucoside</b>					The isolated C3G from plants showed all doses (100; 150; 200 mg/kg) were effective as gastroprotection indicated 74.9% gastro-lesion reduction in 200 mg/kg doses, respectively		

lesions. Chronic ethanol intake has been shown to promote gastric ulceration by decreasing mucus production and reducing blood flow, leading to microvascular injury and triggering inflammatory processes.<sup>114</sup> The damage induced by ethanol is primarily attributed to tissue lipid peroxidation caused by the accumulation of ROS and inflammation.<sup>115</sup> Upon induction of inflammation, both cyanidin and C3G have shown considerable promise in counteracting these ethanol-induced effects by enhancing the integrity of the gastric mucosal barrier.

Based on *in vivo* studies, cyanidin has demonstrated efficiency as gastroprotection varied between males and females, with males showing reduced myeloperoxidase (MPO) activity. In contrast, females, both supplemented and non-supplemented hormones, exhibited reduced MPO activity alongside increased glutathione (GSH) levels, thus suggesting that the antioxidant systems play a more significant role in females treated with cyanidin compared to males and indicating that MPO and GSH are crucial in reducing inflammation caused by ethanol. MPO, an enzyme found in neutrophils, is closely associated with the body's inflammatory response. Elevated MPO levels typically correlate with increased oxidative stress and inflammation. However, Cyanidin and its derivatives have shown that a reduction in MPO activity leads to decreased oxidative stress and inflammation, thereby contributing to the protective effects against ethanol-induced gastric damage.<sup>116</sup>

## Discussion

This study unveils the remarkable potential of cyanidin and C3G as powerful agents in alleviating PUD, combining insights from preclinical studies such as *in vitro* and *in vivo* studies. The findings reveal that these compounds offer a dual-action approach to ulcer management, effectively targeting the underlying inflammation and the oxidative damage that exacerbates gastric injuries. C3G, a glycosylated form of cyanidin, enhances water solubility, stability, and bioactivity. C3G is hydrolyzed by  $\beta$ -glucosidase in the gastrointestinal tract upon ingestion, releasing cyanidin. Both cyanidin and its metabolites are then absorbed and undergo first-pass metabolism in the liver, which involves phase I oxidation followed by phase II conjugation.<sup>36</sup> Hence, these compounds isolated from plants are paving the way for developing targeted and efficacious treatments for PUD. Cyanidin and C3G are metabolized into active phenolic compounds, including PA, VA, and FA, through pathways involving gut microbiota. These stable metabolites enhance antioxidant activity by neutralizing reactive oxygen species (ROS) and modulating pro-inflammatory cytokines such as TNF- $\alpha$  and IL-6. Additionally, the gut microbiota's role in producing low-molecular-weight phenolics further improves their absorption and protective effects on the gastric mucosa.<sup>49,65</sup> Assessing the anti-ulcer potential of cyanidin and C3G is crucial for validating their therapeutic effectiveness and understanding the broader role that plant-based therapies can play in managing complex gastrointestinal conditions (Figure 3).

Oxidative stress plays a pivotal role in the inflammation of the digestive tract, primarily driven by the excessive production of ROS during the oxidative burst by neutrophils.<sup>117</sup> This overproduction leads to progressive intestinal mucosa damage and exacerbates tissue inflammation, prolonged neutrophil activation further accelerates the recruitment of monocytes, which differentiate into macrophages at the site of injury, thereby perpetuating the inflammatory response.<sup>118</sup> The antioxidant properties of polyphenols may counteract this process either by directly scavenging ROS or through indirect pathways, including the modulation of transcription factors like NF-kappaB and the secretion of regulatory cytokines, such as TNF- $\alpha$ , which primes neutrophils for oxidative burst.<sup>119</sup> Cyanidin and C3G, with their potent antioxidant properties, have been shown to effectively scavenge ROS, thereby mitigating oxidative stress and its associated inflammatory pathways.<sup>120</sup> This is particularly important as regulating transcription factors, such as the secretion of pro-inflammatory cytokines like TNF- $\alpha$ , is crucial in controlling the oxidative burst that drives the severity of gastric ulcers.<sup>111,121</sup>

The anti-inflammatory effects of these compounds are further highlighted by their ability to inhibit key pro-inflammatory cytokines, such as TNF- $\alpha$  and IL-6, which play a role in the development of gastric ulcers. TNF- $\alpha$ , secreted progressively by macrophages during ulcer formation,<sup>113</sup> and IL-6, a cytokine involved in acute inflammation and immune regulation,<sup>122</sup> Both significantly amplify the oxidative pathways that lead to local tissue damage. Elevated IL-6 levels stimulate the activation of lymphocytes, neutrophils, and macrophages at the site of inflammation, thereby intensifying oxidative stress driven by mitochondrial ROS generation.<sup>123</sup> The regulation of these pro-inflammatory cytokines is essential in determining the severity of gastric ulcers as they contribute to oxidative stress-induced tissue

damage.<sup>124</sup> Furthermore, the anti-inflammatory cytokine IL-10 counteracts this inflammatory response by inhibiting TNF- $\alpha$  production, thereby reducing the overall impact of oxidative stress on gastric tissue.<sup>113</sup> These cytokines exacerbate local tissue damage by activating oxidative pathways and contribute to the overall severity of gastric ulcers.

Furthermore, the study demonstrated that ethanol-induced neutrophil infiltration in the gastric mucosa significantly contributes to lesion formation.<sup>125</sup> This process, characterized by elevated MPO activity (a key marker of neutrophil infiltration), is closely linked to acute gastric injury.<sup>126</sup> The increased presence of neutrophils not only intensifies mucosal damage but also initiates the release of additional pro-inflammatory mediators, thereby escalating the inflammatory response and worsening gastric tissue damage.<sup>127</sup> In the context of antioxidant defense, catalase (CAT) is vital for neutralizing ROS by catalyzing the decomposition of hydrogen peroxide into water and oxygen.<sup>128</sup> GSH, an endogenous antioxidant with a thiol group in its structure, also plays a crucial role in scavenging free radicals or acting as a cofactor for antioxidant enzymes.<sup>129,130</sup>

Ethanol administration can increase ROS levels while depleting GSH levels in the gastric mucosa.<sup>131</sup> However, treatment with cyanidin demonstrated a beneficial effect, with a dose of 5 mg/kg enhancing catalase activity in male rats and a dose of 20 mg/kg elevating GSH levels in ovariectomized female rats, compared to the control group. Cyanidin and C3G exhibited gastroprotective effects through different mechanisms in this model. Additionally, estrogen appears to be directly linked to the antioxidant defense, as it activates antioxidant enzymes and reduces ROS levels.<sup>132</sup> Based on a study, men over 70 years old and postmenopausal women had a higher incidence of peptic ulcers, likely due to decreased serum estrogen levels leading to reduced mucus production in the stomach.<sup>133</sup> Estrogen is also known for its anti-inflammatory properties and ability to delay gastroesophageal reflux development.<sup>134</sup> Furthermore, research has shown that estrogen possesses anti-ulcer activity by maintaining the gastric mucus layer and providing protection through its antioxidant properties.<sup>135</sup>

These findings highlight the gastroprotective effects of cyanidin and C3G against ethanol-induced damage, mediated through their influence on antioxidant and anti-inflammatory pathways (Figure 3). These compounds significantly suggested mechanisms of action, including cytoprotective effects, such as increasing gastric mucus production, and antioxidant activities that prevent and reduce lipid peroxidation and myeloperoxidase activity in the gastric mucosa. Additionally, their anti-inflammatory properties are evident in their ability to inhibit pro-inflammatory cytokines responsible for ulceration. Furthermore, the small size of these metabolites allows them to efficiently pass through the mucus layer, facilitating interactions with colonic cells and initiating an immune response.

## Conclusion

This review elaborates on an overview of the complex and multifaceted processes involved in the absorption, metabolism, and safety of anthocyanins, particularly cyanidin and C3G, as promising agents for mitigating PUD from preclinical studies. Cyanidin and C3G, derived from various plants, exhibit a dual-action approach, addressing inflammation and oxidative stress (key contributors to the progression of gastric injuries). Both anthocyanins are effective in scavenging ROS, thereby reducing oxidative stress and inflammation, which are critical in the development and severity of gastric ulcers. Cyanidin and C3G inhibit key pro-inflammatory cytokines like TNF- $\alpha$  and IL-6, and thus can be developed as novel therapies for alleviating PUD. While the study provides valuable insights into the potential therapeutic effects of cyanidin and C3G, we encountered some limitations, such as the fact that all the articles are primarily based on preclinical studies, which may not fully translate to human physiology. Moreover, there are varying effects observed between genders, as well as the influence of hormonal factors like estrogen, and the mechanisms by which these compounds exert their protective effects require more in-depth exploration in clinical settings to confirm their therapeutic effects in humans. From a future perspective, clinical trials are necessary to validate the therapeutic efficacy and safety of cyanidin and C3G in humans, these studies should focus on different demographic groups, considering ethnicity, age, gender, BMI, and hormonal status, to understand better how these variables influence the outcomes. However, this review provides scientific evidence supporting the utilization of anthocyanins as a potential approach for treating PUD.

## Abbreviations

C3G, cyanidin-3-glucoside; CAT, catalase; COX-2, cyclooxygenase-2; FA, ferulic acid; GLUT1, glucose-transporter 1; GLUT3, glucose-transporter 3; GSH, glutathione; H2RA, histamine-2 receptor antagonist; HA, hippuric acid; HO, superoxide hydroxy radicals; IL-10, interleukin-10; IL-12, interleukin-12; IL-18, interleukin-18; IL-1 $\beta$ , interleukin-1 $\beta$ ; IL-6, interleukin-6; IL-8,

interleukin-8; iNOS, inducible nitric oxide synthase; IVA, iso-vanillic acid; MnSOD, manganese-dependent superoxide dismutase; MPO, myeloperoxidase; NSAID, non-steroidal anti-inflammatory drugs; O<sub>2</sub><sup>-</sup>, Superoxide anion; PCA, protocatechuic acid; PPI, Proton Pump Inhibitor; PUD, peptic ulcer disease; RAHS, rebound acid hypersecretion; RNS, reactive nitrogen species; ROO, alkyl peroxy radicals; ROS, reactive oxygen species; SGLT1, sodium-glucose transporter 1; TNF- $\alpha$ , tumor necrosis factor- $\alpha$ ; TNFR1, tumor necrosis factor receptor 1; VA, vanillic acid.

## Acknowledgments

The authors thank the Rector of Padjadjaran University via the Directorate of Research and Community Engagement for facilitating the APC. This review study is within the framework of the doctoral research of the first author at the Doctoral Program in Pharmacy, Faculty of Pharmacy, Padjadjaran University, West Java, Indonesia, funded by the Academic-Leadership Grant of Prof. Dr. Jutti Levita (No. 1479/UN6.3.1/PT.00/2024).

## Disclosure

The authors declared no potential conflicts of interest to the research, authorship, or publication of this article.

## References

1. Zelickson MS, Bronder CM, Johnson BL, et al. Helicobacter pylori is not the predominant etiology for peptic ulcers requiring operation. *Am Surg*. 2011;77(8):1054–1060. doi:10.1177/000313481107700827
2. Kruk ME, Gage AD, Joseph NT, Danaei G, García-Saisó S, Salomon JA. Mortality due to low-quality health systems in the universal health coverage era: a systematic analysis of amenable deaths in 137 countries. *Lancet*. 2018;392(10160):2203–2212. doi:10.1016/S0140-6736(18)31668-4
3. Lanás A, Chan FKL. Peptic ulcer disease. *Lancet*. 2017;390(10094):613–624. doi:10.1016/S0140-6736(16)32404-7
4. Kavitt RT, Lipowska AM, Anyane-Yeboah A, Gralnek IM. Diagnosis and treatment of peptic ulcer disease. *Am J Med*. 2019;132(4):447–456. doi:10.1016/j.amjmed.2018.12.009
5. Sung JY, Kuipers EJ, El-Serag HB. Systematic review: the global incidence and prevalence of peptic ulcer disease. *Aliment Pharmacol Ther*. 2009;29(9):938–946. doi:10.1111/j.1365-2036.2009.03960.x
6. Abbasi-Kangevari M, Ahmadi N, Fattahi N, et al. Quality of care of peptic ulcer disease worldwide: a systematic analysis for the global burden of disease study 1990–2019. *PLoS One*. 2022;17(8):e0271284. doi:10.1371/journal.pone.0271284
7. Xie X, Ren K, Zhou Z, Dang C, Zhang H. The global, regional and national burden of peptic ulcer disease from 1990 to 2019: a population-based study. *BMC Gastroenterol*. 2022;22(1):58. doi:10.1186/s12876-022-02130-2
8. Jones MP. The role of psychosocial factors in peptic ulcer disease: beyond Helicobacter pylori and NSAIDs. *J Psychosom Res*. 2006;60(4):407–412. doi:10.1016/j.jpsychores.2005.08.009
9. Malik T, Singh K, Gnanapandithan K. *Peptic Ulcer Disease*. StatPearls Publishing; 2023. <https://www.ncbi.nlm.nih.gov/books/NBK534792/>.
10. Al-garni AA, Khalifa FK, Zeyadi MA. Comparative study of the efficacy of prebiotics and probiotics as dietary supplements in rats with gastric ulcer. *J Pharm Res Int*. 2021;137–145. doi:10.9734/jpri/2021/v33i28A31518
11. Hosseini E, Poursina F, van de Wiele T, Safaei HG, Adibi P. Helicobacter pylori in Iran: a systematic review on the association of genotypes and gastroduodenal diseases. *J Res Med Sci*. 2012;17(3):280.
12. Taha MME, Salga MS, Ali HM, Abdulla MA, Abdelwahab SI, Hadi AHA. Gastroprotective activities of Turnera diffusa Willd. ex Schult. revisited: role of arbutin. *J Ethnopharmacol*. 2012;141(1):273–281. doi:10.1016/j.jep.2012.02.030
13. Begg M, Tarhuni M, Fotso MN, et al. Comparing the Safety and Efficacy of Proton Pump Inhibitors and Histamine-2 receptor antagonists in the management of patients with peptic ulcer disease: a systematic review. *Cureus*. 2023. doi:10.7759/cureus.44341
14. Prayoga D, Aulifa D, Budiman A, Levita J. Plants with anti-ulcer activity and mechanism: a review of preclinical and clinical studies. *Drug Des Devel Ther*. 2024;18:193–213. doi:10.2147/DDDT.S446949
15. Hastrup P, Paulsen MS, Begtrup LM, Hansen JM, Jarbol DE. Strategies for discontinuation of proton pump inhibitors: a systematic review. *Fam Pract*. 2014;31(6):625–630. doi:10.1093/fampra/cmu050
16. Shaw DH. Drugs acting on the gastrointestinal tract. In: *Pharmacology and Therapeutics for Dentistry*. Elsevier; 2017:404–416. doi:10.1016/B978-0-323-39307-2.00028-X
17. Molnar T. Pathogenesis of Ulcerative Colitis and Crohn's Disease: similarities, differences and a lot of things we do not know yet. *J Clin Cell Immunol*. 2014;05(04). doi:10.4172/2155-9899.1000253
18. Probert CSJ, Dignass AU, Lindgren S, Oudkerk Pool M, Marteau P. Combined oral and rectal mesalazine for the treatment of mild-to-moderately active ulcerative colitis: rapid symptom resolution and improvements in quality of life. *J Crohn's Colitis*. 2014;8(3):200–207. doi:10.1016/j.crohns.2013.08.007
19. Wang WX, Li RJ, Li XF. Efficacy and safety of potassium-competitive acid blockers vs proton pump inhibitors for peptic ulcer disease or postprocedural artificial ulcers: a systematic review and meta-analysis. *Clin Transl Gastroenterol*. 2024;15(9):e1. doi:10.14309/ctg.0000000000000754
20. Reimer C, Søndergaard B, Hilsted L, Bytzer P. Proton-pump inhibitor therapy induces acid-related symptoms in healthy volunteers after withdrawal of therapy. *Gastroenterology*. 2009;137(1):80–87.e1. doi:10.1053/j.gastro.2009.03.058
21. Gupta M, Kapoor B, Gupta R, Singh N. Plants and phytochemicals for treatment of peptic ulcer: an overview. *South Afr J Bot*. 2021;138:105–114. doi:10.1016/j.sajb.2020.11.030
22. Kuna L, Jakab J, Smolic R, Raguz-Lucic N, Vcev A, Smolic M. Peptic ulcer disease: a brief review of conventional therapy and herbal treatment options. *J Clin Med*. 2019;8(2):179. doi:10.3390/jcm8020179

23. Cao G, Muccitelli HU, Sánchez-Moreno C, Prior RL. Anthocyanins are absorbed in glycosylated forms in elderly women: a pharmacokinetic study. *Am J Clin Nutr.* 2001;73(5):920–926. doi:10.1093/ajcn/73.5.920
24. Hertog MGL, Hollman PCH, Katan MB, Kromhout D. Intake of potentially anticarcinogenic flavonoids and their determinants in adults in the Netherlands. *Nutr Cancer.* 1993;20(1):21–29. doi:10.1080/01635589309514267
25. Galvano F, La Fauci L, Lazzarino G, et al. Cyanidins: metabolism and biological properties. *J Nutr Biochem.* 2004;15(1):2–11. doi:10.1016/j.jnutbio.2003.07.004
26. de Araújo FF, de Paulo Farias D, Neri-Numa IA, Pastore GM. Polyphenols and their applications: an approach in food chemistry and innovation potential. *Food Chem.* 2021;338:127535. doi:10.1016/j.foodchem.2020.127535
27. Posadino AM, Giordo R, Ramli I, et al. An updated overview of cyanidins for chemoprevention and cancer therapy. *Biomed Pharmacother.* 2023;163:114783. doi:10.1016/j.biopha.2023.114783
28. Min SW, Ryu SN, Kim DH. Anti-inflammatory effects of black rice, cyanidin-3-O- $\beta$ -D-glucoside, and its metabolites, cyanidin and protocatechuic acid. *Int Immunopharmacol.* 2010;10(8):959–966. doi:10.1016/j.intimp.2010.05.009
29. Fu Y, Zhou E, Wei Z, et al. Cyanidin-3-O- $\beta$ -glucoside ameliorates lipopolysaccharide-induced acute lung injury by reducing TLR4 recruitment into lipid rafts. *Biochem Pharmacol.* 2014;90(2):126–134. doi:10.1016/j.bcp.2014.05.004
30. Chen PN, Chu SC, Chiou HL, Kuo WH, Chiang CL, Hsieh YS. Mulberry anthocyanins, cyanidin 3-rutinoside and cyanidin 3-glucoside, exhibited an inhibitory effect on the migration and invasion of a human lung cancer cell line. *Cancer Lett.* 2006;235(2):248–259. doi:10.1016/j.canlet.2005.04.033
31. Cho E, Chung EY, Jang HY, et al. Anti-cancer effect of Cyanidin-3-glucoside from Mulberry via Caspase-3 Cleavage and DNA Fragmentation in vitro and in vivo. *Anticancer Agents Med Chem.* 2017;17(11). doi:10.2174/1871520617666170327152026
32. Chen PN, Chu SC, Chiou HL, Chiang CL, Yang SF, Hsieh YS. Cyanidin 3-Glucoside and Peonidin 3-Glucoside inhibit tumor cell growth and induce apoptosis in vitro and suppress tumor growth in vivo. *Nutr Cancer.* 2005;53(2):232–243. doi:10.1207/s15327914nc5302\_12
33. Cásedas G, Les F, González-Burgos E, Gómez-Serranillos MP, Smith C, López V. Cyanidin-3-O-glucoside inhibits different enzymes involved in central nervous system pathologies and type-2 diabetes. *South Afr J Bot.* 2019;120:241–246. doi:10.1016/j.sajb.2018.07.001
34. Zheng HX, Qi SS, He J, et al. Cyanidin-3-glucoside from Black Rice Ameliorates diabetic nephropathy via reducing blood glucose, suppressing oxidative stress and inflammation, and regulating transforming growth factor  $\beta$ 1/Smad expression. *J Agric Food Chem.* 2020;68(15):4399–4410. doi:10.1021/acs.jafc.0c00680
35. Akkarachiyasit S, Charoenlertkul P, Yibchok-anun S, Adisakwattana S. Inhibitory activities of cyanidin and its glycosides and synergistic effect with Acarbose against Intestinal  $\alpha$ -Glucosidase and Pancreatic  $\alpha$ -Amylase. *Int J mol Sci.* 2010;11(9):3387–3396. doi:10.3390/ijms11093387
36. Gan Y, Fu Y, Yang L, Chen J, Lei H, Liu Q. Cyanidin-3- O -Glucoside and Cyanidin protect against intestinal barrier damage and 2,4,6-Trinitrobenzenesulfonic Acid-Induced Colitis. *J Med Food.* 2020;23(1):90–99. doi:10.1089/jmf.2019.4524
37. Serra D, Paixão J, Nunes C, Dinis TCP, Almeida LM. Cyanidin-3-Glucoside Suppresses Cytokine-Induced Inflammatory Response in Human Intestinal Cells: comparison with 5-Aminosalicylic Acid. *PLoS One.* 2013;8(9):e73001. doi:10.1371/journal.pone.0073001
38. Ferrari D, Speciale A, Cristani M, et al. Cyanidin-3- O -glucoside inhibits NF- $\kappa$ B signalling in intestinal epithelial cells exposed to TNF- $\alpha$  and exerts protective effects via Nrf2 pathway activation. *Toxicol Lett.* 2016;264:51–58. doi:10.1016/j.toxlet.2016.10.014
39. Fagundes FL, Pereira QC, Zarricueta ML, Dos Santos RDC. Malvidin Protects against and Repairs Peptic Ulcers in Mice by Alleviating Oxidative Stress and Inflammation. *Nutrients.* 2021;13(10):3312. doi:10.3390/nu13103312
40. Shih PH, Yeh CT, Yen GC. Anthocyanins Induce the Activation of Phase II Enzymes through the Antioxidant Response Element Pathway against Oxidative Stress-Induced Apoptosis. *J Agric Food Chem.* 2007;55(23):9427–9435. doi:10.1021/jf071933i
41. Yu B, Xiang L, Peppelenbosch MP, Fuhler GM. Overlapping cytokines in H. pylori infection and gastric cancer: a tandem meta-analysis. *Front Immunol.* 2023;14. doi:10.3389/fimmu.2023.1125658
42. Mosiychuk LM, Tatarchuk OM, Petishko OP. Gender features of the cytokine profile in patients with chronic atrophic gastritis. *Gastroenterology.* 2022;55(4):217–222. doi:10.22141/2308-2097.55.4.2021.247911
43. Esedov EM, Akbieva DS. Proinflammatory cytokines in the gastric juice in acid-related diseases before and after standard therapy. *Russ Clin Lab Diagnostics.* 2019;64(8):484–489. doi:10.18821/0869-2084-2019-64-8-484-489
44. Yang XT, Niu PQ, Li XF, et al. Differential cytokine expression in gastric tissues highlights helicobacter pylori's role in gastritis. *Sci Rep.* 2024;14(1):7683. doi:10.1038/s41598-024-58407-x
45. Akashi T, Saito N, Hirota H, Ayabe SI. Anthocyanin-producing dandelion callus as a chalcone synthase source in recombinant polyketide reductase assay. *Phytochemistry.* 1997;46(2):283–287. doi:10.1016/S0031-9422(97)00298-7
46. Ashton OBO, Wong M, McGhie TK, et al. Pigments in Avocado Tissue and Oil. *J Agric Food Chem.* 2006;54(26):10151–10158. doi:10.1021/jf061809j
47. Kamsteeg J, van Brederode J, van Nigtevecht G. Identification and properties of UDP-glucose: cyanidin-3-O-glucosyltransferase isolated from petals of the red campion (*Silene dioica*). *Biochem Genet.* 1978;16(11–12):1045–1058. doi:10.1007/BF00484525
48. Rau D, Forkmann G. Anthocyanin synthesis in tissue cultures of *Callistephus chinensis* (China aster). *Plant Cell Rep.* 1986;5(6):435–438. doi:10.1007/BF00269635
49. Wang H, Nair MG, Strasburg GM, et al. Antioxidant and Antiinflammatory Activities of Anthocyanins and Their Aglycon, Cyanidin, from Tart Cherries. *J Nat Prod.* 1999;62(2):294–296. doi:10.1021/np980501m
50. Hosseinian FS, Beta T. Saskatoon and Wild Blueberries Have Higher Anthocyanin Contents than Other Manitoba Berries. *J Agric Food Chem.* 2007;55(26):10832–10838. doi:10.1021/jf072529m
51. Kozłowska A, Dzierżanowski T. Targeting Inflammation by Anthocyanins as the Novel Therapeutic Potential for Chronic Diseases: an Update. *Molecules.* 2021;26(14):4380. doi:10.3390/molecules26144380
52. Tang B, He Y, Liu J, et al. Kinetic investigation into pH-dependent color of anthocyanin and its sensing performance. *Dye Pigment.* 2019;170:107643. doi:10.1016/j.dyepig.2019.107643
53. Castañeda-Ovando A, Pacheco-Hernández MDL, Páez-Hernández ME, Rodríguez JA, Galán-Vidal CA. Chemical studies of anthocyanins: a review. *Food Chem.* 2009;113(4):859–871. doi:10.1016/j.foodchem.2008.09.001



54. Barnes JS, Schug KA. Structural characterization of cyanidin-3,5-diglucoside and pelargonidin-3,5-diglucoside anthocyanins: multi-dimensional fragmentation pathways using high performance liquid chromatography-electrospray ionization-ion trap-time of flight mass spectrometry. *Int J Mass Spectrom.* **2011**;308(1):71–80. doi:10.1016/j.ijms.2011.07.026
55. Duchnowicz P, Broncel M, Podsek A, Koter-Michalak M. Hypolipidemic and antioxidant effects of hydroxycinnamic acids, quercetin, and cyanidin 3-glucoside in hypercholesterolemic erythrocytes (in vitro study). *Eur J Nutr.* **2012**;51(4):435–443. doi:10.1007/s00394-011-0227-y
56. Jiang T, Mao Y, Sui L, et al. Degradation of anthocyanins and polymeric color formation during heat treatment of purple sweet potato extract at different pH. *Food Chem.* **2019**;274:460–470. doi:10.1016/j.foodchem.2018.07.141
57. Tan J, Li Y, Hou DX, Wu S. The Effects and Mechanisms of Cyanidin-3-Glucoside and Its Phenolic Metabolites in Maintaining Intestinal Integrity. *Antioxidants.* **2019**;8(10):479. doi:10.3390/antiox8100479
58. Akogou FU, Kayodé AP, den Besten HM, Linnemann AR. Extraction methods and food uses of a natural red colorant from dye sorghum. *J Sci Food Agric.* **2018**;98(1):361–368. doi:10.1002/jsfa.8479
59. Dias S, Castanheira EMS, Fortes AG, Pereira DM, Gonçalves MST. Natural Pigments of Anthocyanin and Betalain for Coloring Soy-Based Yogurt Alternative. *Foods.* **2020**;9(6):771. doi:10.3390/foods9060771
60. Saini P, Kumar N, Kumar S, et al. Bioactive compounds, nutritional benefits and food applications of colored wheat: a comprehensive review. *Crit Rev Food Sci Nutr.* **2021**;61(19):3197–3210. doi:10.1080/10408398.2020.1793727
61. Olivas-Aguirre F, Rodrigo-García J, Martínez-Ruiz N, et al. Cyanidin-3-O-glucoside: physical-Chemistry, Foodomics and Health Effects. *Molecules.* **2016**;21(9):1264. doi:10.3390/molecules21091264
62. Tirupula KC, Balem F, Yanamala N, Klein-Seetharaman J. pH-dependent Interaction of Rhodopsin with Cyanidin-3-glucoside. 2. Functional Aspects †. *Photochem Photobiol.* **2009**;85(2):463–470. doi:10.1111/j.1751-1097.2008.00533.x
63. Esposito D, Damsud T, Wilson M, et al. Black Currant Anthocyanins Attenuate Weight Gain and Improve Glucose Metabolism in Diet-Induced Obese Mice with Intact, but Not Disrupted, Gut Microbiome. *J Agric Food Chem.* **2015**;63(27):6172–6180. doi:10.1021/acs.jafc.5b00963
64. Yang P, Yuan C, Wang H, et al. Stability of Anthocyanins and Their Degradation Products from Cabernet Sauvignon Red Wine under Gastrointestinal pH and Temperature Conditions. *Molecules.* **2018**;23(2):354. doi:10.3390/molecules23020354
65. Morais CA, de Rosso VV, Estadella D, Pisani LP. Anthocyanins as inflammatory modulators and the role of the gut microbiota. *J Nutr Biochem.* **2016**;33:1–7. doi:10.1016/j.jnutbio.2015.11.008
66. Fang J. Some Anthocyanins Could Be Efficiently Absorbed across the Gastrointestinal Mucosa: extensive Presystemic Metabolism Reduces Apparent Bioavailability. *J Agric Food Chem.* **2014**;62(18):3904–3911. doi:10.1021/jf405356b
67. Hanske L, Engst W, Loh G, Sczesny S, Blaut M, Braune A. Contribution of gut bacteria to the metabolism of cyanidin 3-glucoside in human microbiota-associated rats. *Br J Nutr.* **2013**;109(8):1433–1441. doi:10.1017/S0007114512003376
68. Aura AM, Martín-López P, O’Leary KA, et al. In vitro metabolism of anthocyanins by human gut microflora. *Eur J Nutr.* **2005**;44(3):133–142. doi:10.1007/s00394-004-0502-2
69. Cai H, Thomasset SC, Berry DP, et al. Determination of anthocyanins in the urine of patients with colorectal liver metastases after administration of bilberry extract. *Biomed Chromatogr.* **2011**;25(6):660–663. doi:10.1002/bmc.1499
70. Zhang X, Sandhu A, Edirisinghe I, Burton-Freeman B. An exploratory study of red raspberry (*Rubus idaeus* L.) (poly)phenols/metabolites in human biological samples. *Food Funct.* **2018**;9(2):806–818. doi:10.1039/C7FO00893G
71. Kopf-Bolanz KA, Schwander F, Gijs M, Vergères G, Portmann R, Egger L. Validation of an In Vitro Digestive System for Studying Macronutrient Decomposition in Humans3. *J Nutr.* **2012**;142(2):245–250. doi:10.3945/jn.111.148635
72. Mallery SR, Budendorf DE, Larsen MP, et al. Effects of Human Oral Mucosal Tissue, Saliva, and Oral Microflora on Intraoral Metabolism and Bioactivation of Black Raspberry Anthocyanins. *Cancer Prev Res.* **2011**;4(8):1209–1221. doi:10.1158/1940-6207.CAPR-11-0040
73. Carpenter GH. The Secretion, Components, and Properties of Saliva. *Annu Rev Food Sci Technol.* **2013**;4(1):267–276. doi:10.1146/annurev-food-030212-182700
74. Chen J. Food oral processing: some important underpinning principles of eating and sensory perception. *Food Struct.* **2014**;1(2):91–105. doi:10.1016/j.foosr.2014.03.001
75. Knobloch TJ, Uhrig LK, Pearl DK, et al. Suppression of Proinflammatory and Prosurvival Biomarkers in Oral Cancer Patients Consuming a Black Raspberry Phytochemical-Rich Troche. *Cancer Prev Res.* **2016**;9(2):159–171. doi:10.1158/1940-6207.CAPR-15-0187
76. Miyazawa T, Nakagawa K, Kudo M, Muraishi K, Someya K. Direct Intestinal Absorption of Red Fruit Anthocyanins, Cyanidin-3-glucoside and Cyanidin-3,5-diglucoside, into Rats and Humans. *J Agric Food Chem.* **1999**;47(3):1083–1091. doi:10.1021/jf9809582
77. McGhie TK, Walton MC. The bioavailability and absorption of anthocyanins: towards a better understanding. *mol Nutr Food Res.* **2007**;51(6):702–713. doi:10.1002/mnfr.200700092
78. Czank C, Cassidy A, Zhang Q, et al. Human metabolism and elimination of the anthocyanin, cyanidin-3-glucoside: a 13C-tracer study. *Am J Clin Nutr.* **2013**;97(5):995–1003. doi:10.3945/ajcn.112.049247
79. Hassimotto NMA, Genovese MI, Lajolo FM. Absorption and metabolism of cyanidin-3-glucoside and cyanidin-3-rutinoside extracted from wild mulberry (*Morus nigra* L.) in rats. *Nutr Res.* **2008**;28(3):198–207. doi:10.1016/j.nutres.2007.12.012
80. Oliveira H, Fernandes I, Brás NF, et al. Experimental and Theoretical Data on the Mechanism by Which Red Wine Anthocyanins Are Transported through a Human MKN-28 Gastric Cell Model. *J Agric Food Chem.* **2015**;63(35):7685–7692. doi:10.1021/acs.jafc.5b00412
81. Del Bo’ C, Riso P, Brambilla A, et al. Blanching Improves Anthocyanin Absorption from Highbush Blueberry (*Vaccinium corymbosum* L.) Purée in Healthy Human Volunteers: a Pilot Study. *J Agric Food Chem.* **2012**;60(36):9298–9304. doi:10.1021/jf3021333
82. Passamonti S, Terdoslavich M, Franca R, et al. Bioavailability of Flavonoids: a Review of Their Membrane Transport and the Function of Bilitranslocase in Animal and Plant Organisms. *Curr Drug Metab.* **2009**;10(4):369–394. doi:10.2174/138920009788498950
83. Fernandes I, Marques F, de Freitas V, Mateus N. Antioxidant and antiproliferative properties of methylated metabolites of anthocyanins. *Food Chem.* **2013**;141(3):2923–2933. doi:10.1016/j.foodchem.2013.05.033
84. Shih PH, Yeh CT, Yen GC. Effects of anthocyanidin on the inhibition of proliferation and induction of apoptosis in human gastric adenocarcinoma cells. *Food Chem Toxicol.* **2005**;43(10):1557–1566. doi:10.1016/j.fct.2005.05.001
85. Kim SH, Woo H, Park M, et al. Cyanidin 3-O-glucoside reduces *Helicobacter pylori* VacA-induced cell death of gastric KATO III cells through inhibition of the SecA pathway. *Int J Med Sci.* **2014**;11(7):742–747. doi:10.7150/ijms.7167

86. Phan ADT, Netzel G, Wang D, Flanagan BM, D'Arcy BR, Gidley MJ. Binding of dietary polyphenols to cellulose: structural and nutritional aspects. *Food Chem.* **2015**;171:388–396. doi:10.1016/j.foodchem.2014.08.118
87. de Ferrars RM, Czank C, Zhang Q, et al. The pharmacokinetics of anthocyanins and their metabolites in humans. *Br J Pharmacol.* **2014**;171(13):3268–3282. doi:10.1111/bph.12676
88. Felgines C, Krisa S, Mauray A, et al. Radiolabelled cyanidin 3- O -glucoside is poorly absorbed in the mouse. *Br J Nutr.* **2010**;103(12):1738–1745. doi:10.1017/S0007114510000061
89. Felgines C, Talavéra S, Texier O, et al. Absorption and metabolism of red Orange juice anthocyanins in rats. *Br J Nutr.* **2006**;95(5):898–904. doi:10.1079/BJN20061728
90. Fang J. Bioavailability of anthocyanins. *Drug Metab Rev.* **2014**;46(4):508–520. doi:10.3109/03602532.2014.978080
91. Woodward GM, Needs PW, Kay CD. Anthocyanin-derived phenolic acids form glucuronides following simulated gastrointestinal digestion and microsomal glucuronidation. *mol Nutr Food Res.* **2011**;55(3):378–386. doi:10.1002/mnfr.201000355
92. Ozdal T, Sela DA, Xiao J, Boyacioglu D, Chen F, Capanoglu E. The Reciprocal Interactions between Polyphenols and Gut Microbiota and Effects on Bioaccessibility. *Nutrients.* **2016**;8(2):78. doi:10.3390/nu8020078
93. Valdés L, Cuervo A, Salazar N, Ruas-Madiedo P, Gueimonde M, González S. The relationship between phenolic compounds from diet and microbiota: impact on human health. *Food Funct.* **2015**;6(8):2424–2439. doi:10.1039/C5FO00322A
94. Aguilar F, Crebelli R, Dusemund B, et al. Scientific Opinion on the re-evaluation of anthocyanins (E 163) as a food additive. *EFSA J.* **2013**;11(4). doi:10.2903/j.efsa.2013.3145
95. Charoensin S, Taya S, Wongpoomchai S, Wongpoomchai R. Assessment of genotoxicity and antigenotoxicity of an aqueous extract of *Cleistocalyx nervosum* var. *paniala* in vitro and in vivo models. *Interdiscip Toxicol.* **2012**;5(4):201–206. doi:10.2478/v10102-012-0033-2
96. Li D, Wang P, Luo Y, Zhao M, Chen F. Health benefits of anthocyanins and molecular mechanisms: update from recent decade. *Crit Rev Food Sci Nutr.* **2017**;57(8):1729–1741. doi:10.1080/10408398.2015.1030064
97. Hassellund SS, Flaa A, Kjeldsen SE, et al. Effects of anthocyanins on cardiovascular risk factors and inflammation in pre-hypertensive men: a double-blind randomized placebo-controlled crossover study. *J Hum Hypertens.* **2013**;27(2):100–106. doi:10.1038/jhh.2012.4
98. Mehta AJ, Cassidy A, Litonjua AA, Sparrow D, Vokonas P, Schwartz J. Dietary anthocyanin intake and age-related decline in lung function: longitudinal findings from the VA Normative Aging Study. *Am J Clin Nutr.* **2016**;103(2):542–550. doi:10.3945/ajcn.115.121467
99. Cassidy A, Rogers G, Peterson JJ, Dwyer JT, Lin H, Jacques PF. Higher dietary anthocyanin and flavonol intakes are associated with anti-inflammatory effects in a population of US adults. *Am J Clin Nutr.* **2015**;102(1):172–181. doi:10.3945/ajcn.115.108555
100. Ma L, Wang G, Chen Z, et al. Modulating the p66shc Signaling Pathway with Protocatechuic Acid Protects the Intestine from Ischemia-Reperfusion Injury and Alleviates Secondary Liver Damage. *Sci World J.* **2014**;2014:1–11. doi:10.1155/2014/387640
101. Pereira SR, Pereira R, Figueiredo I, Freitas V, Dinis TCP, Almeida LM. Comparison of anti-inflammatory activities of an anthocyanin-rich fraction from Portuguese blueberries (*Vaccinium corymbosum* L.) and 5-aminosalicylic acid in a TNBS-induced colitis rat model. *PLoS One.* **2017**;12(3):e0174116. doi:10.1371/journal.pone.0174116
102. Jiménez S, Gascón S, Luquin A, Laguna M, Ancin-Azpilicueta C, Rodríguez-Yoldi MJ. Rosa canina Extracts Have Antiproliferative and Antioxidant Effects on Caco-2 human Colon Cancer. *PLoS One.* **2016**;11(7):e0159136. doi:10.1371/journal.pone.0159136
103. Valencia-Avilés E, García-Pérez ME, Garnica-Romo MG, et al. Antioxidant Properties of Polyphenolic Extracts from *Quercus Laurina*, *Quercus Crassifolia*, and *Quercus Scytophylla* Bark. *Antioxidants.* **2018**;7(7):81. doi:10.3390/antiox7070081
104. Barreca D, Gattuso G, Laganà G, Leuzzi U, Bellocco E. C- and O-glycosyl flavonoids in Sanguinello and Tarocco blood Orange (*Citrus sinensis* (L.) Osbeck) juice: identification and influence on antioxidant properties and acetylcholinesterase activity. *Food Chem.* **2016**;196:619–627. doi:10.1016/J.FOODCHEM.2015.09.098
105. Pesca MS, Dal Piaz F, Sanogo R, et al. Bioassay-guided isolation of proanthocyanidins with antiangiogenic activities. *J Nat Prod.* **2013**;76(1):29–35. doi:10.1021/NP300614U
106. Kähkönen MP, Heinonen M. Antioxidant activity of anthocyanins and their aglycons. *J Agric Food Chem.* **2003**;51(3):628–633. doi:10.1021/JF025551I
107. Wang H, Cao G, Prior RL. Oxygen Radical Absorbing Capacity of Anthocyanins. *J Agric Food Chem.* **1997**;45(2):304–309. doi:10.1021/JF960421T
108. Rice-Evans CA, Miller NJ, Paganga G. Structure-antioxidant activity relationships of flavonoids and phenolic acids. *Free Radic Biol Med.* **1996**;20(7):933–956. doi:10.1016/0891-5849(95)02227-9
109. Laganà G, Barreca D, Smeriglio A, et al. Evaluation of Anthocyanin Profile, Antioxidant, Cytoprotective, and Anti-Angiogenic Properties of *Callistemon citrinus* Flowers. *Plants.* **2020**;9(8):1045. doi:10.3390/plants9081045
110. Cury BJ, Boeing T, Somensi LB, et al. Açai berries (*Euterpe oleracea* Mart.) dried extract improves ethanol-induced ulcer in rats. *J Pharm Pharmacol.* **2020**;72(9):1239–1244. doi:10.1111/jphp.13290
111. Raafat K, El-Darra N, Saleh FA. Gastroprotective and anti-inflammatory effects of *Prunus cerasus* phytochemicals and their possible mechanisms of action. *J Tradit Complement Med.* **2020**;10(4):345–353. doi:10.1016/j.jtcme.2019.06.001
112. Magiera A, Czerwińska ME, Owczarek A, Marchelak A, Granica S, Olszewska MA. Polyphenol-Enriched Extracts of *Prunus spinosa* Fruits: anti-Inflammatory and Antioxidant Effects in Human Immune Cells Ex Vivo in Relation to Phytochemical Profile. *Molecules.* **2022**;27(5):1691. doi:10.3390/molecules27051691
113. Rozza AL, Meira de Faria F, Souza Brito AR, Pellizzon CH. The Gastroprotective Effect of Menthol: involvement of Anti-Apoptotic, Antioxidant and Anti-Inflammatory Activities. *PLoS One.* **2014**;9(1):e86686. doi:10.1371/journal.pone.0086686
114. Mishra AP, Bajpai A, Chandra S. A Comprehensive Review on the Screening Models for the Pharmacological Assessment of Antiulcer Drugs. *Curr Clin Pharmacol.* **2019**;14(3):175–196. doi:10.2174/1574884714666190312143846
115. Chen H, Liao H, Liu Y, et al. Protective effects of pogostone from *Pogostemonis* Herba against ethanol-induced gastric ulcer in rats. *Fitoterapia.* **2015**;100:110–117. doi:10.1016/j.fitote.2014.11.017
116. Zarricueta ML, Fagundes FL, Pereira QC, Pantaleão SQ, Dos Santos RDC. Relationship between Hormonal Modulation and Gastroprotective Activity of Malvidin and Cyanidin Chloride: in Vivo and In Silico Approach. *Pharmaceutics.* **2022**;14(3):565. doi:10.3390/pharmaceutics14030565

117. Bourgonje AR, Feelisch M, Faber KN, Pasch A, Dijkstra G, van Goor H. Oxidative Stress and Redox-Modulating Therapeutics in Inflammatory Bowel Disease. *Trends Mol Med*. 2020;26(11):1034–1046. doi:10.1016/j.molmed.2020.06.006
118. Prame Kumar K, Nicholls AJ, Wong CHY. Partners in crime: neutrophils and monocytes/macrophages in inflammation and disease. *Cell Tissue Res*. 2018;371(3):551–565. doi:10.1007/s00441-017-2753-2
119. Michel P, Granica S, Rosińska K, Rojek J, Poraj L, Olszewska MA. Biological and chemical insight into Gaultheria procumbens fruits: a rich source of anti-inflammatory and antioxidant salicylate glycosides and procyanidins for food and functional application. *Food Funct*. 2020;11(9):7532–7544. doi:10.1039/D0FO01750G
120. Gao S, Chen T, Choi MY, Liang Y, Xue J, Wong YS. Cyanidin reverses cisplatin-induced apoptosis in HK-2 proximal tubular cells through inhibition of ROS-mediated DNA damage and modulation of the ERK and AKT pathways. *Cancer Lett*. 2013;333(1):36–46. doi:10.1016/j.canlet.2012.12.029
121. Speciale A, Canali R, Chirafisi J, Saija A, Virgili F, Cimino F. Cyanidin-3- O -glucoside Protection against TNF- $\alpha$ -Induced Endothelial Dysfunction: involvement of Nuclear Factor- $\kappa$ B Signaling. *J Agric Food Chem*. 2010;58(22):12048–12054. doi:10.1021/jf1029515
122. Kishimoto T. INTERLEUKIN-6: from Basic Science to Medicine—40 Years in Immunology. *Annu Rev Immunol*. 2005;23(1):1–21. doi:10.1146/annurev.immunol.23.021704.115806
123. Mei X, Xu D, Xu S, Zheng Y, Xu S. Novel role of Zn(II)-curcumin in enhancing cell proliferation and adjusting proinflammatory cytokine-mediated oxidative damage of ethanol-induced acute gastric ulcers. *Chem Biol Interact*. 2012;197(1):31–39. doi:10.1016/j.cbi.2012.03.006
124. Augusto AC, Miguel F, Mendonça S, Pedrazzoli J, Gurgueira SA. Oxidative stress expression status associated to Helicobacter pylori virulence in gastric diseases. *Clin Biochem*. 2007;40(9–10):615–622. doi:10.1016/j.clinbiochem.2007.03.014
125. La Casa C, Villegas I, Alarcón de la Lastra C, Motilva V, Martín Calero M. Evidence for protective and antioxidant properties of rutin, a natural flavone, against ethanol induced gastric lesions. *J Ethnopharmacol*. 2000;71(1–2):45–53. doi:10.1016/S0378-8741(99)00174-9
126. Takeuchi K, Ueshima K, Hironaka Y, Fujioka Y, Matsumoto J, Okabe S. Oxygen free radicals and lipid peroxidation in the pathogenesis of gastric mucosal lesions induced by indomethacin in rats. *Digestion*. 1991;49(3):175–184. doi:10.1159/000200718
127. Wallace JL, Ma L. Inflammatory mediators in gastrointestinal defense and injury. *Exp Biol Med*. 2001;226(11):1003–1015. doi:10.1177/153537020122601107
128. Suo H, Zhao X, Qian Y, et al. Lactobacillus fermentum Suo Attenuates HCl/Ethanol Induced Gastric Injury in Mice through Its Antioxidant Effects. *Nutrients*. 2016;8(3):155. doi:10.3390/nu8030155
129. Glavin GB, Szabo S. Experimental gastric mucosal injury: laboratory models reveal mechanisms of pathogenesis and new therapeutic strategies. *FASEB J*. 1992;6(3):825–831. doi:10.1096/fasebj.6.3.1740232
130. Ibrahim AA, Ameen Abdulla M, Hajrezaie M, et al. The gastroprotective effects of hydroalcoholic extract of Monolluma quadrangula against ethanol-induced gastric mucosal injuries in Sprague Dawley rats. *Drug Des Devel Ther*. 2015;93. doi:10.2147/DDDT.S91247.
131. YT S, Li J, Chen Y, et al. [Effects of glutathione on plasma heat shock protein 70 of acute gastric mucosal injury in rats exposed to positive acceleration]. *Zhonghua Yi Xue Za Zhi*. 2013;93(46):3708–3710. Danish
132. Speir E, Yu ZX, Takeda K, Ferrans VJ, Cannon RO. Antioxidant Effect of Estrogen on Cytomegalovirus-Induced Gene Expression in Coronary Artery Smooth Muscle Cells. *Circulation*. 2000;102(24):2990–2996. doi:10.1161/01.CIR.102.24.2990
133. Okada K. Gender differences of low-dose aspirin-associated gastroduodenal ulcer in Japanese patients. *World J Gastroenterol*. 2010;16(15):1896. doi:10.3748/wjg.v16.i15.1896
134. Iijima K, Shimosegawa T. Involvement of luminal nitric oxide in the pathogenesis of the gastroesophageal reflux disease spectrum. *J Gastroenterol Hepatol*. 2014;29(5):898–905. doi:10.1111/jgh.12548
135. Boeckstaens G, El-Serag HB, Smout AJPM, Kahrilas PJ. Symptomatic reflux disease: the present, the past and the future. *Gut*. 2014;63(7):1185–1193. doi:10.1136/gutjnl-2013-306393

## Drug Design, Development and Therapy

### Publish your work in this journal

Drug Design, Development and Therapy is an international, peer-reviewed open-access journal that spans the spectrum of drug design and development through to clinical applications. Clinical outcomes, patient safety, and programs for the development and effective, safe, and sustained use of medicines are a feature of the journal, which has also been accepted for indexing on PubMed Central. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/drug-design-development-and-therapy-journal>

**Dovepress**  
Taylor & Francis Group