

MEETING ABSTRACT

Open Access

Diet, nutrition and osteoarthritis

Margaret P Rayman

From Musculoskeletal Health in the 21st Century
Guildford, UK. 30 June - 1 July 2015

Osteoarthritis (OA) is the fastest growing cause of disability worldwide[1]. In the absence of effective therapies, patients may wish to take some control of their own condition by making dietary changes that have the potential to ameliorate symptoms or reduce disease progression.

A number of dietary factors have been associated with OA symptoms or progression. Most notably, in those overweight, weight reduction of $\geq 10\%$ has the potential to lead to important changes in pain and function[2]. Losing weight also reduces pain-associated inflammation[3]. Weight loss combined with physical activity has an even greater capacity to improve pain and function[4].

It has been suggested that OA is a metabolic disease in which lipids essentially contribute to the pathophysiology of cartilage degradation[5]. Dietary long-chain ω -3 PUFA may affect articular cartilage composition and appear to have beneficial effects in OA[5]. In a US cohort of individuals with, or at high risk of, knee OA, there was a significant inverse relationship between total n-3 PUFAs and patella-femoral cartilage loss[6].

A positive association has been shown between elevated serum cholesterol and OA; hypercholesterolemia (OR 1.61; 95% CI 1.06-2.47) and high serum cholesterol (3rd vs. 1st tertile: OR 1.73; 95% CI 1.02-2.92) were independently associated with generalized OA in the Ulm study [7]. Hence there may be a potential benefit in adopting dietary cholesterol-lowering strategies (such as consumption of sterol/stanol spreads/drinks).

Vitamin D affects the state of multiple articular structures. The evidence for association between the vitamin D biomarker, serum 25(OH)D, and OA was assessed in a systematic review. For knee radiographic OA progression and cartilage loss, there was strong evidence for an association with low 25(OH)D[8].

Vitamin K is important in cartilage metabolism as an inhibitor of extracellular matrix calcification and a

promoter of cell survival/proliferation. In the US MOST study, vitamin K deficiency was associated with incident radiographic knee OA and MRI-based cartilage lesions (RR 2.39; 95% CI, 1.05-5.40) compared with no deficiency[9].

Hence, dietary recommendations are the following:

- lose weight, if overweight, preferably combined with exercise;
- reduce plasma cholesterol by dietary means;
- at least for a trial period, increase intake of long-chain n-3 fatty acids preferably by eating oily fish twice a week;
- aim for a safe level of sun exposure, eat rich vitamin-D dietary sources or take vitamin D supplements, $\leq 25 \mu\text{g}/\text{d}$;
- increase vitamin K intake by eating green leafy vegetables.

Published: 1 December 2015

References

1. Conaghan PG, Porcheret M, Kingsbury SR, Gammon A, Soni A, Hurley M, Rayman MP, Barlow J, Hull RG, Cumming J, Llewelyn K, Moscovici F, Lyons J, Birrell F: **Impact and therapy of osteoarthritis: the Arthritis Care OA Nation 2012 survey.** *Clin Rheumatol* 2014, Jun 3. [Epub ahead of print].
2. Riddle DL, Stratford PW: **Body weight changes and corresponding changes in pain and function in persons with symptomatic knee osteoarthritis: a cohort study.** *Arthritis Care Res (Hoboken)* 2013, **65**:15-22.
3. Hauner H: **Secretory factors from human adipose tissue and their functional role.** *Proc Nutr Soc* 2005, **64**:163-9.
4. Brosseau L, Wells GA, Tugwell P, Egan M, Dubouloz CJ, Casimiro L, Bugnariu N, Welch VA, De Angelis G, Francoeur L, Milne S, Loew L, McEwan J, Messier SP, Doucet E, Kenny GP, Prud'homme D, Lineker S, Bell M, Poitras S, Li JX, Finestone HM, Laferrrière L, Haines-Wangda A, Russell-Doreleyers M, Lambert K, Marshall AD, Cartizzone M, Teav A: **Ottawa Panel. Ottawa Panel evidence-based clinical practice guidelines for the management of osteoarthritis in adults who are obese or overweight.** *Phys Ther* 2011, **91**:843-61.
5. Masuko K, Murata M, Suematsu N, Okamoto K, Yudoh K, Nakamura H, Kato T: **A metabolic aspect of osteoarthritis: lipid as a possible contributor to the pathogenesis of cartilage degradation.** *Clin Exp Rheumatol* 2009, **27**:347-53.
6. Baker KR, Matthan NR, Lichtenstein AH, Niu J, Guermazi A, Roemer F, Grainger A, Nevitt MC, Clancy M, Lewis CE, Torner JC, Felson DT: **Association of plasma n-6 and n-3 polyunsaturated fatty acids with synovitis in the knee: the MOST study.** *Osteoarthritis Cartilage* 2012, **20**:382-7.

Correspondence: m.rayman@surrey.ac.uk
Department of Nutritional Sciences, Faculty of Health and Medical Sciences,
University of Surrey, Guildford, GU2 7XH, UK

7. Stürmer T, Sun Y, Sauerland S, Zeissig I, Günther KP, Puhl W, Brenner H: **Serum cholesterol and osteoarthritis. The baseline examination of the Ulm Osteoarthritis Study.** *J Rheumatol* 1998, **25**:1827-32.
8. Cao Y, Winzenberg T, Nguo K, Lin J, Jones G, Ding C: **Association between serum levels of 25-hydroxyvitamin D and osteoarthritis: a systematic review.** *Rheumatology (Oxford)* 2013, **52**:1323-34.
9. Misra D, Booth SL, Tolstykh I, Felson DT, Nevitt MC, Lewis CE, Torner J, Neogi T: **Vitamin K deficiency is associated with incident knee osteoarthritis.** *Am J Med* 2013, **126**(3):243-8.

doi:10.1186/1471-2474-16-S1-S7

Cite this article as: Rayman: Diet, nutrition and osteoarthritis. *BMC Musculoskeletal Disorders* 2015 **16**(Suppl 1):S7.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

