

Outcomes of laparoscopic common bile duct exploration (LCBDE) after failed endoscopic retrograde cholangiopancreatography versus primary LCBDE for managing cholecystocholedocholithiasis

Journal of International Medical Research

48(10) 1–9

© The Author(s) 2020

Article reuse guidelines:

sagepub.com/journals-permissions

DOI: 10.1177/0300060520957560

journals.sagepub.com/home/imr



Hanbaro Kim¹ , Suk Pyo Shin²,
Ji Woong Hwang³ and Jung Woo Lee⁴

Abstract

Objective: This study was performed to compare the outcomes of laparoscopic common bile duct exploration (LCBDE) after failed endoscopic retrograde cholangiopancreatography (ERCP) versus primary LCBDE for managing cholecystocholedocholithiasis.

Methods: We retrospectively analyzed data from 59 patients who underwent LCBDE during laparoscopic cholecystectomy (LC) for managing cholecystocholedocholithiasis from January 2013 to August 2019. The patients underwent either primary LCBDE plus LC (Group I) or LCBDE plus LC after failed ERCP (Group II). The demographics, reason for ERCP failure, peri-operative details, and postoperative outcomes were evaluated.

Results: CBD stone removal using preoperative ERCP failed in 31 patients (Group II) because of remaining stones after ERCP (n = 9), failed cannulation (n = 6), failed sedation (n = 6), a periam-pullary diverticulum (n = 5), previous Billroth II gastrectomy (n = 3), a huge stone (n = 1), and an impacted stone (n = 1). The CBD stone clearance rate was >96% in both groups. The mean

¹Department of Surgery, Hallym University College of Medicine, Chuncheon Sacred Heart Hospital, South Korea

²Department of Internal Medicine, Hallym University College of Medicine, Chuncheon Sacred Heart Hospital, South Korea

³Department of Surgery, Hallym University College of Medicine, Kangnam Sacred Heart Hospital, South Korea

⁴Department of Surgery, Hallym University College of Medicine, Hallym Sacred Heart Hospital, South Korea

Corresponding author:

Jung Woo Lee, Department of Surgery, Hallym University College of Medicine, Hallym Sacred Heart Hospital, 22 Gwanpyeong-ro 170 beon-gil, Dongan-gu, Anyang-si, Gyeonggi-do 14068, Republic of Korea.
Email: km98woo@hanmail.net



operative time, hospital stay, overall complication rate, and open conversion rate were not significantly different between the two groups.

Conclusions: When extraction of CBD stones by ERCP is likely to be difficult or fail, primary LCBDE is an acceptable alternative treatment for managing cholecystocholedocholithiasis.

Keywords

Common bile duct stone, exploration, endoscopic retrograde cholangiopancreatography failure, laparoscopic cholecystectomy, cholecystocholedocholithiasis, stone clearance rate

Date received: 3 May 2020; accepted: 17 August 2020

Introduction

Common bile duct (CBD) stones are found in approximately 5% to 20% of patients with symptomatic gallstones.¹⁻³ The optimal management of cholecystocholedocholithiasis is still under debate. Therapeutic strategies for CBD stones vary widely and include laparotomy, laparoscopic surgery, percutaneous intervention, and endoscopic retrograde cholangiopancreatography (ERCP). Among these, preoperative ERCP and stone extraction followed by laparoscopic cholecystectomy (LC) are preferred treatments; these are recommended in the European Association for the Study of the Liver clinical practice guidelines on the prevention, diagnosis, and treatment of gallstones.¹ However, potential ERCP-related adverse effects include pancreatitis, duodenal perforation, and bleeding.^{2,4} Moreover, the use of ERCP to extract CBD stones can fail for many reasons, including failed cannulation, previous Billroth II gastrectomy, large CBD stones, and high numbers of CBD stones.⁵ Single-stage laparoscopic CBD exploration (LCBDE) during LC is now primarily used to treat cholecystocholedocholithiasis, although advanced surgical skills are required to perform the procedure. As the number of surgeons with laparoscopic skill

and experience has increased, the number of centers that perform LCBDE has increased as well. A recent meta-analysis showed that LCBDE during LC is superior to preoperative ERCP and stone extraction followed by LC.⁶ However, few case series have been performed to compare the outcomes of LCBDE after failed preoperative ERCP versus the outcomes of primary LCBDE during LC for managing cholecystocholedocholithiasis. The present study was performed to analyze the reasons for ERCP failure when treating CBD stones and to evaluate our results of primary LCBDE compared with the results of LCBDE after failed ERCP for managing cholecystocholedocholithiasis.

Materials and methods

Data collection

We retrospectively analyzed the medical records of patients with cholecystocholedocholithiasis who underwent LCBDE and LC at the Hallym Sacred Heart Hospital and Chuncheon Sacred Heart Hospital from January 2013 to August 2019. We divided these patients into two groups: those who underwent primary LCBDE and LC (Group I) and those who

underwent LCBDE and LC after failed ERCP (Group II).

At our institution, the general protocol is that the endoscopic approach is initially used to treat CBD stones. The diagnosis of CBD stones was based on the results of radiologic investigations such as ultrasonography and magnetic resonance cholangiopancreatography. The main ERCP procedure involved endoscopic sphincterotomy, endoscopic papillary balloon dilation, and stone removal using a retrieval balloon or basket. We used mechanical lithotripsy for large CBD stones. If this failed, the physician decided whether to repeat the endoscopic procedure or perform a surgical procedure after consulting with the surgeon. Patients whose CBD stones were completely removed after repeating the ERCP procedure were excluded. Primary LCBDE without ERCP was performed if the physician determined that the endoscopic approach would be difficult because of the presence of huge stones, a history of Billroth II surgery, or a huge diverticulum or if the surgeon determined that surgical treatment would be more advantageous. Patients who had a non-dilated CBD with a diameter of <7 mm usually did not undergo LCBDE because of the risk of CBD stricture occurring as a long-term complication.

The following clinical characteristics were analyzed: age, sex, body mass index, American Society of Anesthesiologists physical status, and comorbidities. The following operative data were analyzed: operating time, postoperative complications, open conversion, start of oral intake, and length of hospital stay. The CBD diameter, CBD stone size, CBD stone clearance rate, and CBD stone recurrence rate were also analyzed. We did not perform intraoperative choledochoscopy to confirm CBD clearance. However, duct clearance was confirmed by performing intraoperative cholangiography, postoperative T-tube

cholangiography, or magnetic resonance cholangiopancreatography. The data associated with ERCP failure were analyzed. We also analyzed the degree of difficulty in performing ERCP according to the following grading system: grade 1, simple diagnostic ERCP; grade 2, simple therapeutic ERCP; grade 3, complex diagnostic ERCP; grade 4, complex therapeutic ERCP; and grade 5, very advanced ERCP in Group II.⁷ Readmission was defined as re-hospitalization within 30 days of discharge. The present study was approved by the Institutional Review Board of the Hallym Hospital. Consent was not required because this was a retrospective study performed using medical records of patients who received standard treatment.

Surgical procedure

The type of surgery (transductal or transcystic approach) was determined according to the surgeon's preference, and the surgical technique was not standardized. We used four trocars (two 12-mm trocars and two 5-mm trocars) for the transductal approach. The transductal approach was performed with various techniques involving primary closure, T-tube insertion, and internal stent insertion. First, choledochotomy was performed 1 to 2 cm vertically at the location of the anterior CBD. Normal saline was injected through a 7-Fr Nelaton catheter or 3-mm pediatric feeding tube to flush obstructive stones through the choledochotomy. Primary closure was performed for patients in whom all CBD stones had been removed, patients in whom the feeding tube passed through the ampulla of Vater (AoV), and patients for whom an intraoperative cholangiogram was available. Internal stent insertion was performed when all CBD stones were removed and the feeding tube passed through the AoV but no intraoperative cholangiogram was obtained. When it was uncertain whether all CBD

stones were removed, or when the feeding tube did not smoothly pass through the AoV, T-tube insertion was performed. During the transcystic approach, we used four trocars (one 12-mm trocar and three 5-mm trocars). The endoscopic basket was passed over a guide wire through the cystic duct to extract stones under fluoroscopic guidance.

Statistical analysis

Continuous variables were compared between the two groups using the χ^2 test and are expressed as median and range or mean \pm standard deviation. Categorical variables are reported as number and percentage. Statistical analyses were performed using IBM SPSS Statistics for Windows, Version 22.0 (IBM Corp., Armonk, NY, USA).

Results

Patient characteristics

Fifty-nine patients were included in this study (Group I, n = 28; Group II, n = 31). The patients' clinical characteristics are listed in Table 1. There were no significant differences in the patients' age, American Society of Anesthesiologists physical status, or history of an abdominal operation between the two groups. Symptoms and comorbidities were also not significantly different between the two groups. The causes of ERCP failure are listed in Table 2. The most common cause of incomplete CBD stone removal was remaining stones after ERCP (n = 9), followed by a huge stone (n = 1) and an impacted stone (n = 1). The reasons for failed cannulation were an abnormal position of the AoV (n = 6), a periampullary diverticulum

Table 1. Patient demographics.

	Group I (n = 28)	Group II (n = 31)	P
Age, years	70.1 \pm 12.3	65.0 \pm 19.2	0.235
Sex, male:female	13:15	15:16	0.880
BMI, kg/m ²	23.4 \pm 3.5	24.0 \pm 5.1	0.586
ASA physical status			0.576
1	1 (3.6)	2 (6.5)	
2	8 (28.6)	10 (32.3)	
3	14 (50.0)	17 (54.8)	
4	5 (17.9)	2 (6.5)	
Operation history	8 (28.6)	6 (19.4)	0.406
Symptoms			
Fever	9 (32.1)	8 (25.8)	0.592
Abdominal pain	28 (100.0)	28 (90.3)	0.091
Jaundice	15 (53.6)	22 (71.0)	0.168
Comorbidities			
Hypertension	10 (35.7)	12 (38.7)	0.812
Diabetes	6 (21.4)	7 (22.6)	0.915
Other cardiovascular disease	6 (21.4)	10 (32.2)	0.644
Other pulmonary disease	2 (7.1)	1 (3.2)	0.513
Other conditions	3 (10.7)	4 (12.9)	0.795

Data are presented as mean \pm standard deviation, n, or n (%).

BMI, body mass index; ASA, American Society of Anesthesiologists.

Table 2. Reasons for failed endoscopic stone extraction.

	n = 31
Incomplete CBD stone removal	
Remaining stones after ERCP	9
Huge stones	1
Impacted stone	1
Failed cannulation	
Abnormal position of ampulla of Vater	6
Periampullary diverticulum	5
Previous Billroth II gastrectomy	3
Failed sedation	6

ERCP, endoscopic retrograde cholangiopancreatography; CBD, common bile duct.

(n = 5), and previous Billroth II gastrectomy (n = 3). ERCP failed in six patients because of failed sedation. The degree of ERCP difficulty according to the grading scale in Group II was grade 2 (n = 11), grade 4 (n = 19), and grade 5 (n = 1).

Perioperative results

Stone extraction under direct laparoscopic choledochotomy was achieved in 28 patients (100%) in Group I and in 29 patients (93.5%) in Group II. The transcystic approach was performed in only two patients (6.5%) in Group II. There were no differences in the surgical type between the two groups. The mean operative time was also similar between Groups I and II (92.0 vs. 84.4 min, respectively). Two patients (8.0%) in Group I required conversion to open surgery because of adhesions caused by previous surgery. The mean time to oral diet commencement and the duration of the hospital stay did not differ between Groups I and II (1.5 vs. 1.3 days and 7.4 vs. 7.0 days, respectively). The postoperative complication rates in Groups I and II were 3.6% and 3.2%, respectively, with no significant difference. Bile leakage occurred in one patient in Group II. No postoperative mortality, reoperation, or

readmission was recorded in either group. The perioperative findings are summarized in Table 3.

CBD stone clearance rate and recurrence

The CBD stone size and diameter were not significantly different between Groups I and II (9.1 vs. 9.3 mm and 14.8 vs. 13.0 mm, respectively). The degree of ERCP difficulty according to the grading scale was also not significantly different between the two groups. The CBD stone clearance rate in Groups I and II was 96.4% and 100%, respectively. Additional procedures were needed in Group I, primarily involving subsequent stone extraction by ERCP. The median follow-up period was 26 months. Two patients (8.0%) in Group I and one patient (3.2%) in Group II had recurrent stones (Table 4).

Discussion

The first choice for CBD stone management is endoscopic treatment, which is now performed worldwide. Many reports have indicated high technical success rates with extraction of at least 90% of stones.^{8,9} However, the endoscopic approach occasionally fails when either removing CBD stones or cannulating the CBD. In the present study, the most common reason for incomplete CBD stone removal was the presence of remaining stones (9/31) followed by a huge stone (1/31) and an impacted stone (1/31). Extraction of CBD stones via an endoscopic approach may be difficult for several reasons. In most situations, the size, shape, and number of stones are the key determinants of whether extraction will be easy. A prospective study by Kim et al.¹⁰ showed that large CBD stones (>15 mm), impacted CBD stones, a shorter length of the distal CBD arm (<36 mm), and more acute distal CBD angulation were significant factors affecting the

Table 3. Overall comparison of perioperative results.

	Group I (n = 28)	Group II (n = 31)	p
Operation time, minutes	92.0 ± 38.6	84.4 ± 30.4	0.399
Commencement of water sipping, days	1.5 ± 0.6	1.3 ± 0.7	0.113
Commencement of oral diet, days	2.5 ± 0.8	2.2 ± 0.8	0.113
Hospital stay, days	7.4 ± 3.7	7.0 ± 2.3	0.591
Surgery type			0.140
Transductal approach	28 (100.0)	29 (93.5)	
T-tube insertion	7 (25.1)	14 (45.2)	
Primary closure	17 (60.7)	11 (35.5)	
Internal stent	4 (14.3)	4 (12.9)	
Transcystic approach	0 (0.0)	2 (6.5)	
Open conversion	2 (8.0)	0 (0.0)	0.115
All complications	1 (3.6)	1 (3.2)	0.942
Bile leakage	0 (0.0)	1 (3.2)	
Wound infection	1 (3.6)	0 (0.0)	
Reoperation	0 (0.0)	0 (0.0)	–
Readmission	0 (0.0)	0 (0.0)	–
Mortality	0 (0.0)	0 (0.0)	–

Data are presented as mean ± standard deviation or n (%).

Table 4. CBD stone clearance and recurrence.

	Group I (n = 28)	Group II (n = 31)	p
CBD stone size, mm	9.1 ± 6.7	9.3 ± 6.7	0.932
CBD diameter, mm	14.8 ± 5.2	13.0 ± 5.5	0.225
Multiple CBD stones	13 (46.4)	16 (51.6)	0.691
Gallstones	25 (89.3)	26 (83.9)	0.544
CBD stone clearance rate	27 (96.4)	34 (100.0)	0.272
Additional procedures	1 (3.6)	0 (0.0)	0.289
Recurrent stones	2 (8.0)	1 (3.2)	0.513

Data are presented as mean ± standard deviation or n (%).

CBD, common bile duct.

technical difficulty of CBD stone clearance. Failed cannulation is another reason for failed ERCP. The success rate of biliary cannulation is 84% to 94%.^{5,11} However, even the most skilled endoscopist will fail to achieve deep biliary cannulation in a minority of cases. In the present study, the reasons for failed cannulation included an abnormal position of the AoV (6/31), a periampullary diverticulum (5/31), and

previous Billroth II gastrectomy (3/31). Hong et al.¹² recently reported that the success rate of cannulation for 107 patients with an ectopic papilla of Vater was 83.2%. They reported that the ectopic opening was so obscure that it may have increased the difficulty of cannulation.¹² Sfarti et al.¹³ reported that a periampullary diverticulum can increase the rate of difficult or failed cannulation. Li et al.¹⁴

reported that the success rate of duodenal ampullary access and selective biliary cannulation was 84.1% (191/227) and 92.1% (176/191), respectively, in patients with Billroth II anatomy. Patient sedation is also important for ERCP success. Conscious sedation is routinely performed by gastroenterologists for endoscopic procedures; however, ERCP failure is often due to inadequate conscious sedation.¹⁵ Predicting the possibility of failure in advance is helpful because ERCP fails to remove CBD stones for many of the reasons mentioned above. Many studies have already used a grading system to assess the technical difficulty of ERCP.^{7,10,16,17} We also used a grading scale to analyze the degree of difficulty of ERCP in Group II. Grade 2 was seen in 11 patients, grade 4 in 19, and grade 5 in only 1. The success of ERCP is influenced by the technical difficulty of the procedure. Therefore, a more accurate grading system to predict failure of ERCP is needed.

When ERCP fails, surgery is needed for CBD stone extraction. Recently, LCBDE has been increasingly advocated in the primary management of CBD stones. A prospective study that compared single-stage LCBDE and LC as a two-stage endoscopic stone extraction procedure followed by LC for cholecystocholedocholithiasis illustrated that the former achieved a shorter hospital stay and comparable stone clearance rate compared with the latter.¹⁸ In addition, a recent meta-analysis showed that LCBDE produced superior results in terms of perioperative safety and short- and long-term postoperative efficacy compared with ERCP.⁶ The meta-analysis concluded that LCBDE should be considered as optimal treatment for cholecystocholedocholithiasis. In the present study, the success rate of laparoscopic management of CBD stones was 98.3%, with an average morbidity rate of 3.4% and mortality rate of 0.0%. We also confirmed that LCBDE

is safe and effective for managing cholecystocholedocholithiasis.

In some reports, LCBDE after failed ERCP was described as more technically challenging than primary LCBDE.^{18,19} Di Mauro et al.²⁰ reported that the mean operative time was longer in LCBDE after failed ERCP because of local fibrosis around Calot's triangle. They also reported that patients who underwent LCBDE after failed ERCP had a longer hospital stay because of the longer operative time and the use of a T-tube, which required time to manage.²⁰ In our study, however, LCBDE after failed ERCP yielded a similar mean operative time and hospital stay as primary LCBDE. In fact, the 14 patients who underwent failed CBD cannulation and 6 patients who did not remain sedated were the same patients as those who did not undergo ERCP. These may be the reasons why the mean operative time and hospital stay were similar between the two groups in our study. There were also no significant differences in the CBD stone clearance rate and stone recurrence rate between the two groups.

This study has limitations associated with its retrospective nature and small study population. A study with a larger sample size would be needed to reach more definitive conclusions.

In conclusion, our surgical outcomes indicate that LCBDE can be safely and effectively performed for managing cholecystocholedocholithiasis. The outcomes were similar between primary LCBDE and LCBDE after failed ERCP. If endoscopic stone removal is predicted to be difficult or to fail, primary LCBDE is an acceptable alternative treatment for managing cholecystocholedocholithiasis.

Declaration of conflicting interest

The authors declare that there is no conflict of interest.

Funding

This research was supported by Hallym University Research Fund (HURF-2019-55).

ORCID iD

Hanbaro Kim  <https://orcid.org/0000-0002-9835-7168>

References

1. EASL Clinical Practice Guidelines on the prevention, diagnosis and treatment of gallstones. *J Hepatol* 2016; 65: 146–181.
2. Gad EH, Zakaria H, Kamel Y, et al. Surgical (open and laparoscopic) management of large difficult CBD stones after different sessions of endoscopic failure: a retrospective cohort study. *Ann Med Surg (Lond)* 2019; 43: 52–63.
3. Tarantino G, Magistri P, Ballarin R, et al. Surgery in biliary lithiasis: from the traditional “open” approach to laparoscopy and the “rendezvous” technique. *Hepatobiliary Pancreat Dis Int* 2017; 16: 595–601.
4. Williams EJ, Taylor S, Fairclough P, et al. Are we meeting the standards set for endoscopy? Results of a large-scale prospective survey of endoscopic retrograde cholangiopancreatograph practice. *Gut* 2007; 56: 821–829.
5. Williams EJ, Ogollah R, Thomas P, et al. What predicts failed cannulation and therapy at ERCP? Results of a large-scale multicenter analysis. *Endoscopy* 2012; 44: 674–683.
6. Pan L, Chen M, Ji L, et al. The safety and efficacy of laparoscopic common bile duct exploration combined with cholecystectomy for the management of choledocholithiasis: an up-to-date meta-analysis. *Ann Surg* 2018; 268: 247–253.
7. Schutz SM and Abbott RM. Grading ERCPs by degree of difficulty: a new concept to produce more meaningful outcome data. *Gastrointest Endosc* 2000; 51: 535–539.
8. Binmoeller KF, Bruckner M, Thonke F, et al. Treatment of difficult bile duct stones using mechanical, electrohydraulic and extracorporeal shock wave lithotripsy. *Endoscopy* 1993; 25: 201–206.
9. Cipolletta L, Costamagna G, Bianco MA, et al. Endoscopic mechanical lithotripsy of difficult common bile duct stones. *Br J Surg* 1997; 84: 1407–1409.
10. Kim HJ, Choi HS, Park JH, et al. Factors influencing the technical difficulty of endoscopic clearance of bile duct stones. *Gastrointest Endosc* 2007; 66: 1154–1160.
11. Colton JB and Curran CC. Quality indicators, including complications, of ERCP in a community setting: a prospective study. *Gastrointest Endosc* 2009; 70: 457–467.
12. Hong J, Pan W, Zuo W, et al. Efficacy and safety of therapeutic ERCP in patients with ectopic papilla of Vater. *Medicine (Baltimore)* 2020; 99: e18536. Advance online publication.
13. Sfarti VC, Balan G Jr, Chiriac AS, et al. Endoscopic retrograde cholangiopancreatography (ERCP) in patients with periampullary diverticula. *Rom J Morphol Embryol* 2018; 59: 833–837.
14. Li JS, Zou DW, Jin ZD, et al. Predictive factors for extraction of common bile duct stones during endoscopic retrograde cholangiopancreatography in Billroth II anatomy patients. *Surg Endosc* 2020; 34: 2454–2459. Advance online publication.
15. Raymondos K, Panning B, Bachem I, et al. Evaluation of endoscopic retrograde cholangiopancreatography under conscious sedation and general anesthesia. *Endoscopy* 2002; 34: 721–726.
16. Rangunath K, Thomas LA, Cheung WY, et al. Objective evaluation of ERCP procedures: a simple grading scale for evaluating technical difficulty. *Postgrad Med J* 2003; 79: 467–470.
17. Sahar N, La Selva D, Gluck M, et al. The ASGE grading system for ERCP can predict success and complication rates in a tertiary referral hospital. *Surg Endosc* 2019; 33: 448–453.
18. Bansal VK, Misra MC, Rajan K, et al. Single-stage laparoscopic common bile duct exploration and cholecystectomy versus two-stage endoscopic stone extraction followed by laparoscopic cholecystectomy for patients with concomitant gallbladder stones and common bile duct stones: a

- randomized controlled trial. *Surg Endosc* 2014; 28: 875–885.
19. Karaliotas C, Sgourakis G, Goumas C, et al. Laparoscopic common bile duct exploration after failed endoscopic stone extraction. *Surg Endosc* 2008; 22: 1826–1831.
 20. Di Mauro D, Ricciardi E, Siragusa L, et al. Outcome of laparoscopic common bile duct exploration after failed endoscopic retrograde cholangiopancreatography: a comparative study. *J Laparoendosc Adv Surg Tech A* 2019; 29: 1391–1396.