

RESEARCH ARTICLE

Complex community-wide consequences of consumer sexual dimorphism

Stephen P. De Lisle¹  | Sebastian J. Schreiber²  | Daniel I. Bolnick¹ 

¹Department of Ecology & Evolutionary Biology, University of Connecticut, Storrs, CT, USA

²Department of Evolution and Ecology and Center for Population Biology, University of California, Davis, CA, USA

Correspondence

Stephen P. De Lisle

Email: stephen.de_lisle@biol.lu.se**Present address**

Stephen P. De Lisle, Evolutionary Ecology Unit, Department of Biology, Lund University, Lund, Sweden

Handling Editor: Greta Bocedi**Abstract**

1. Sexual dimorphism is a ubiquitous source of within-species variation, yet the community-level consequences of sex differences remain poorly understood.
2. Here, we analyse a bitrophic model of two competing resource species and a sexually reproducing consumer species.
3. We show that consumer sex differences in resource acquisition can have striking consequences for consumer-resource coexistence, abundance and dynamics.
4. Under both direct interspecific competition and apparent competition between two resource species, sexual dimorphism in consumers' attack rates can mediate coexistence of the resource species, while in other cases can lead to exclusion when stable coexistence is typically expected. Slight sex differences in total resource acquisition also can reverse competitive outcomes and lead to density cycles. These effects are expected whenever both consumer sexes require different amounts or types of resources to reproduce.
5. Our results suggest that consumer sexual dimorphism, which is common, has wide-reaching implications for the assembly and dynamics of natural communities.

KEYWORDS

apparent competition, community assembly, competitive exclusion, ecological sexual dimorphism, resource competition

1 | INTRODUCTION

Within-species variation is one key feature of natural populations that has emerged as a critical contributor to community ecology and species interactions (Hughes et al., 2008; Bolnick et al., 2011; Des Roches et al., 2018). Even in the simplest scenario of a single-species community with a single optimum phenotype, within-population variation is expected to reduce population growth rate of a well-adapted population via effects on mean fitness (Haldane, 1937). In more complex multispecies communities, phenotypic variation in

consumers or their resources can either promote or constrain coexistence between competing species, under many circumstances yielding different conclusions than would be reached from models ignoring within-species variation (Bolnick et al., 2011; Schreiber et al., 2011; Patel & Schreiber, 2015; Cortez & Patel, 2017). One emerging question (Bolnick et al., 2011) from this work is the degree to which understanding the specific source of phenotypic variation matters in community ecology.

Here, we explore the ecological consequences of sexual dimorphism, a central feature of metazoans and many plant populations

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited and is not used for commercial purposes.

© 2022 The Authors. *Journal of Animal Ecology* published by John Wiley & Sons Ltd on behalf of British Ecological Society.

that has not been fully incorporated into ecological theory. The gamete differences that define the sexes are expected to lead to divergence between males and females in a suite of life-history traits (Trivers, 1972; Shärer et al., 2012) and other phenotypic traits. This divergence is iconically manifested as striking sexual dimorphism in sexually selected traits such as body size or courtship displays (Darwin, 1871; Andersson, 1994). Although sexual selection and associated mate choice behaviours themselves may have relevance to interspecific ecological interactions (Gomez-Llano et al., 2018), the different life histories that define the sexes are often expected to lead to different nutritional and resource requirements for males and females (Maklakov et al., 2008). As a specific example, in insects, differential contributions of longevity and fecundity to male and female lifetime reproductive success result in different combinations of macronutrients that maximize male and female fitness (Maklakov et al., 2008; Reddix et al., 2013; Garlapow et al., 2015; Jensen et al., 2015; Camus et al., 2017). Similarly, nutritional models of optimal foraging have been proposed to explain sex differences in diets of moose (Belovksy, 1978). Consequently, males and females of many species have evolved divergent resource use, either as an indirect outcome of divergent reproductive roles or through other forms of sex-specific natural selection, resulting in sex differences in diet composition and often in corresponding trophic morphology (Slatkin, 1984; Temeles, 1985; Shine, 1989; Temeles et al., 2000; De Lisle & Rowe, 2015a; De Lisle, 2019).

Data suggest that these 'ecological' sexual dimorphisms can have substantial consequences for community structure (Fryxell et al., 2015; Pincheira-Donoso et al., 2018; Start & De Lisle, 2018; Tsuji & Fukami, 2018, 2020). More generally, sex differences may play an important role in the relationship between ecological and evolutionary dynamics (Giery & Layman, 2019; Svensson, 2019; Fryxell et al., 2019). Sexual dimorphism thus represents a key source of ecologically relevant variation within species. Within-species variation in resource specialization is commonplace (Bolnick et al., 2003), with important consequences for species interactions and community assembly (Hughes et al., 2008; Bolnick et al., 2011; Des Roches et al., 2018). Critically, as a source of intraspecific variation, sexual dimorphism may have different consequences from other types of variation. Unlike most phenotypes, whose relative abundances can evolve to reflect local ratios of alternative resources, the ratio of males to females (at birth) is expected to be maintained at 1:1 (Darwin, 1871; Fisher, 1930), though their phenotypes and demography may diverge to have differential ecological impacts. Thus, the rate at which males are born (which determines predation pressure on males' prey) is strongly dependent on females' foraging success. Perhaps to a lesser extent (depending on mating system), the rate of female offspring production can depend on male foraging success. Importantly, persistence of a sexually reproducing consumer depends on the persistence of both sexes. This coupling of the dynamics of two consumer phenotypes has unknown consequences for consumer-resource dynamics, consumer-mediated coexistence of the resource species and apparent competition between the resource species. Using a simple model of consumer-resource

dynamics, we show that consumer sexual dimorphism can influence (both positively and negatively) coexistence between resource species (competing directly or apparently) and species' abundance.

2 | MATERIALS AND METHODS

Our model represents a simple extension of a classic model of consumer-resource dynamics, in which a consumer species exploits two resources that themselves undergo density-dependent growth (Figure 1). Consumer growth rate is limited only by resource abundance. The original version of this model did not consider within-species phenotypic variation and can lead to exclusion of one resource ('apparent competition'; e.g. Holt, 1977). More recently, Schreiber et al. (2011) showed that quantitative trait variation in the predator, affecting attack rates on the two resources, can facilitate coexistence between the prey. Here, we instead allow for the possibility that male and female consumers differ in resource-specific attack rates. We describe the population dynamics of two resources (with densities R_1, R_2) and consumer males (with density M) and females (with density F) using the system of ordinary differential equations:

$$\frac{dR_j}{dt} = r_j R_j (1 - \alpha_{ij} R_i - \alpha_{jj} R_j) - a_{M,j} R_j M - a_{F,j} R_j F \text{ for } j \neq i, \quad (1a)$$

$$\frac{dM}{dt} = \frac{1}{2} B(M, F, R_1, R_2) - d_M M, \quad (1b)$$




FIGURE 1 Illustration of the model structure. Males (M) and females (F) of a consumer species exploit two resources (R_1 and R_2) that may or may not also compete directly (double arrow). Sex-specific attack rates (black arrows) generate sex differences in ecological niche such that males and females preferentially attack R_1 and R_2 , respectively, with varying degrees of overlap (grey arrows). Although males and females may consume different resources, fitness is equal across the sexes and so the dynamics of males, females, resource 1 and resource 2 are coupled

$$\frac{dF}{dt} = \frac{1}{2} B(M, F, R_1, R_2) - d_F F. \quad (1c)$$

Each resource undergoes density-dependent population growth with intrinsic per capita growth rates r , with competition represented by competition coefficients α . Resources are also regulated by consumers depending on sex-specific attack rates by males a_M and by females a_F . Consumer dynamics are governed by females' birth rate B (half of the offspring being female), and limited by sex-specific intrinsic mortality rates d_M and d_F .

An appropriate function describing birth rates has been a point of debate for demographers, with the general conclusion being that any function describing birth rates should capture the negative effects of extreme sex ratio skew on birth rates, with the extreme being that birth rates should be zero when one sex is absent (Caswell & Weeks, 1986), or unable to breed. Generally, treating birth rates proportional to the harmonic mean density of males and females is agreed as the best approach (Caswell & Weeks, 1986; Lindstöm & Kokko, 1998), and is also empirically supported (Miller & Inouye, 2011, 2011). Here, we extend this harmonic mean birth function to account for each sex's foraging success, on the logic that each sex must both be present and sufficiently well-fed to reproduce:

$$B(M, F, R_i, R_j) = 2 \frac{M I_M * F I_F}{M I_M + F I_F}, \quad (2)$$

where I is the energy intake of a given sex:

$$I_F = b_F (a_{F,1} R_1 + a_{F,2} R_2) \text{ and } I_M = b_M (a_{M,1} R_1 + a_{M,2} R_2), \quad (3)$$

which couples the dynamics of consumers and resources. In this model the rate of consumer births depends on not just the number of males and females but also on the abundance of both prey and the prey preference of each sex, where $M I_M$ is the number of males weighted by their energy to reproduce (similarly with females), b is a scaling constant reflecting the degree to which sex-specific resource acquisition influences birth rate and $a_{M,i}, a_{F,i}$ are the attack rates on the resource i . Although our birth function is phenomenological and we make no genetic assumptions, we note that our birth function, and thus our model, applies to sexually reproducing diploid consumers. As an alternative to the inclusion of constants b_M and b_F , Equation 2 can be equivalently expressed with the inclusion of a 'harem size' parameter, h (Caswell and Weeks 1986, Lindstöm and Kokko 1998; see supplemental material); changes in b_M relative to b_F alter the degree to which each sex contributes to birth rates in an equivalent manner to the effects of deviations of h from unity. We also note that although $h = 1$ is typically interpreted as a condition of monogamy, this in fact simply represents a condition where the average number of mating partners is equal for males and females, which corresponds to monogamy (relatively rare in nature) as well as polygynandry (common in nature). Thus, we focus on the parameterization in Equation 2 because it illustrates that the mathematical/demographic effects of 'harem size' can in fact be brought

about by any factor that alters the relative contribution of sex-specific density to birth rates.

Because we are specifically interested in understanding the effects of sexual dimorphism in prey preference, we relate male and female resource-specific attack rates via the degree of sexual dimorphism, β , such that

$$a_{M,1} = a_{M,\max} \left(\frac{1}{2} + \frac{\beta}{2} \right) \text{ and } a_{F,1} = a_{F,\max} \left(\frac{1}{2} - \frac{\beta}{2} \right), \quad (4a)$$

$$a_{M,2} = a_{M,\max} \left(\frac{1}{2} - \frac{\beta}{2} \right) \text{ and } a_{F,2} = a_{F,\max} \left(\frac{1}{2} + \frac{\beta}{2} \right), \quad (4b)$$

where β ranges from 0 to 1, with $\beta = 0$ representing complete sexual monomorphism if maximum attack rates are equal across the sexes (both sexes are generalists, attacking each prey at the rates, $a_{F,\max}/2$ and $a_{M,\max}/2$) and $\beta = 1$ representing complete sexual dimorphism such that males only attack R_1 and females only attack R_2 . Sex differences in a_{\max} reflect a situation where males and females intake different total amounts of resources, independent of any difference in resource preference. In our study, we explore both types of sex differences in resource acquisition; sexual dimorphism in resource preference captured by β , as well as sex differences in total resource acquisition captured by sex-specific a_{\max} .

By assuming that resource acquisition of males and females contributes to the consumer birth rate, we are not necessarily making any assumption about parental care or condition-dependent fecundity. Rather, this is an assumption that the fraction of each sex available for reproduction depends in part on resource acquisition, a realistic feature of many organisms. For instance, males' energetic costs of finding a mate, defending a territory, care for offspring or expressing a sexually selected trait are all expected to depend on the resource pool available to males (Rowe and Houle 1996). Females' ability to produce eggs, gestate and provision young (e.g. lactation in mammals) similarly depends on their energy intake. Parental care and other factors can also lead to associations between sex-specific resource abundance and consumer birth rates, beyond what would be expected beyond the costs of simply being available to mate. We also note that there are alternative ways in which resource acquisition could be parameterized to relate to consumer birth rates, namely through resource-dependent consumer mortality. We have focused on the scenario of a resource-dependent birth function, which is not exclusive of the operation of resource-dependent mortality which was not modelled here.

We analysed this model to find equilibria when one or two resources are present, as well as invasion criteria for a resource into a two-species community at equilibrium. Using the mathematical theory of permanence (Patel & Schreiber, 2018; Schreiber, 2000), we can use these invasion criteria to determine whether all three species coexist (in the sense of permanence), exhibit a bistability or competitive exclusion occurs. In addition to presenting our

analytical results below, we provide further mathematical details in Supplement A. We also numerically explored the behaviour of the model under a variety of scenarios using the packages `DESOLVE` v. 1.21 (Soetgart et al. 2010) and `caTools` v. 1.17.1.2 (Tuszynski 2019) in R v. 3.5.0 (R Core Team 2018). Importantly, using simulations allowed us to explore the behaviour of our model for cases where two-species consumer-resource equilibria do not exist. Because there are a large number of potential combinations of parameters that could be explored, we focused our simulations on three different biological scenarios: (a) completely symmetric male and female total attack rates and contributions to birth rate, $a_{F,\max} = a_{M,\max}$ and $b_M = b_F$, (b) asymmetric total attack rates across the sexes, $a_{F,\max} \neq a_{M,\max}$ and (c) asymmetric contribution of male and female resource acquisition to consumer birth rates, $b_M \neq b_F$. The first scenario applies most readily to organisms with biparental care and similar total caloric requirements across the sexes. The second scenario likely applies to many organisms where total lifetime resource acquisition is higher for one sex, which is the case for many organisms where the sexes differ, for example in body size. The third scenario is another realistic departure from the first, relevant for organisms where resource acquisition in one sex (e.g. females) has a greater influence on that sex's mating propensity, or cases with biased operational sex ratios due to polyandry or polygyny. For each scenario, we explore the consequences of consumer ecological sexual dimorphism (β) for consumer persistence and coexistence. For each set of parameter values, we simulated the model 1,000 time steps (to ensure the simulation reaches its equilibrium). For each run, we determined the equilibrium abundances of M , F , R_1 and R_2 at $t = 1,000$. We also calculated the standard deviation of species' abundances in the last 50 time steps. Although we present solutions simulated on the scale of $\alpha_{ij} = 0.1$, we obtained qualitatively equivalent conclusions rescaling under a wide range values of competition coefficients (see Table S1B, Supplement B). The R script to generate all figures and results presented is archived at <https://doi.org/10.5281/zenodo.6324965> (De Lisle et al. 2022).

3 | RESULTS

3.1 | Analytical results

Our analysis begins by considering a subsystem of a single resource species, say species i , and the consumer species. That is, we explore the conditions under which a sexually dimorphic consumer can persist on a single resource species. These two species can coexist if

$$H_i \frac{1}{\alpha_{ii}} > 2, \quad (5)$$

where

$$H_i := 2 \frac{\frac{b_M a_{iM}}{d_M} \frac{b_F a_{iF}}{d_F}}{\frac{b_M a_{iM}}{d_M} + \frac{b_F a_{iF}}{d_F}}, \quad (6)$$

is the harmonic mean of the lifetime per capita resource contributions of each sex to reproduction, and $\frac{1}{\alpha_{ii}}$ corresponds to the carrying capacity of resource species i . Intuitively, $H_i \frac{1}{\alpha_{ii}}$ corresponds to the average number of offspring produced by a mating pair during their lifetime when the resource species is at its carrying capacity. These harmonic means decrease to zero with the degree of sexual dimorphism (i.e. H_i is a decreasing function of β and $H_i = 0$ when $\beta = 1$). There always is a critical degree of sexual dimorphism above which the consumer cannot persist on a single resource species. Intuitively, this arises when one of the sexes specializes too much on the other (absent) resource species and, consequently, contributes too little to reproduction. When the coexistence condition holds, the consumer-resource species pair coexist at the following equilibrium densities:

$$R_i^* = 2 / H_i, \quad (7)$$

$$F_i^* = \frac{r_i \left(1 - \frac{2\alpha_{ii}}{H_i}\right)}{a_{iF} + \frac{a_{iM} d_F}{d_M}},$$

$$M_i^* = \frac{d_F}{d_M} F_i^*.$$

At these densities of the consumer and resource i , the second (rare) resource species j can invade if its per capita growth rate

$$I_j = r_j \left(1 - \frac{2\alpha_{jj}}{H_j}\right) - A_j F_i^*, \quad (8)$$

is positive. Here,

$$A_j = a_{Fj} + \frac{a_{Mj} d_F}{d_M}, \quad (9)$$

is the average attack rate on resource j when the males and females are at an equilibrium. Provided the consumer species can persist on each of the resource species individually, the invasion growth rates I_1 and I_2 determine the ecological outcomes. If both invasion growth rates are positive (mutual invasibility), all three species coexist. If both invasion rates are negative, there is a priority effect in which both single resource-consumer equilibria are stable. If one invasion growth rate is positive and the other is negative, this suggests (as confirmed by numerical simulations) that the resource species with the positive invasion growth rate excludes the other resource species. In Supplement A, we provide more details about this analysis and describe the invasion conditions for these outcomes or when the consumer only persists in the presence of both resource species.

As a step towards understanding the general conditions for coexistence and exclusion, we first examine two special cases corresponding to (a) resource competitive symmetry whereby $\alpha_{11} = \alpha_{12} = \alpha_{21} = \alpha_{22}$ and (b) pure apparent competition whereby $\alpha_{12} = \alpha_{21} = 0$. When there is competitive symmetry, the resource species with the larger value of r_i / A_i excludes the other resource species. The quantity r_i / A_i corresponds to the consumer density

supported by resource species i when intraspecific competition is very weak that is $\alpha_{ii} = 0$. Hence, the resource species that supports the highest consumer equilibrium density (in the absence of intraspecific competition) excludes the other. This is an analogue of the p^* rule (Holt 1977, Holt and Lawton 1993, Schreiber 2021). In this case, sexual dimorphism (as measured by β) influences outcomes only if there is an asymmetry in the maximal attack rates of the female and male (e.g. female masked boobies dive for prey more often than males; Weimerskirch et al. 2009). For example, if the female-preferred resource (R_1) has the higher intrinsic rate of growth ($r_1 > r_2$) and the female consumer has the higher maximal attack rate of the two sexes, then the female-preferred resource species excludes the other resource species when sexual dimorphism is low (i.e. $A_1 = A_2$ when $\beta = 0$). However, as the average attack rate A_1 on the female-preferred resource increases with the magnitude β of the sexual dimorphism, sexual dimorphism reverses the outcome of apparent competition whenever $r_1/a_{F,\max} < r_2/a_{M,\max}$ and the self-limitation in the female-preferred resource is weak.

Next, we consider the case of pure apparent competition in which there is no direct competition between the resources, that is $\alpha_{12} = \alpha_{21} = 0$. Then, coexistence occurs when

$$\frac{1}{1 - 2\alpha_{ii}/H_i} > \frac{r_i/A_i}{r_j/A_j} \text{ for } i \neq j. \quad (10)$$

The right-hand side of this condition determines the outcome of apparent competition in the case of competitive symmetry, that is the species with larger r_i/A_i wins. The left-hand side increases to infinity as the sexual dimorphism increases such that H_i approaches $2\alpha_{ii}$. This has two implications: (a) when r_1/A_1 and r_2/A_2 are equal, the resource species coexist at any level of sexual dimorphism, and (b) when r_1/A_1 and r_2/A_2 are unequal, sufficiently strong sexual dimorphisms (that still allow the consumer to persist, per Equation 5) will ensure coexistence of the two resource species by diluting the strength of apparent competition.

Finally, the general coexistence condition is

$$\frac{1 - 2\alpha_{ji}/H_j}{1 - 2\alpha_{ii}/H_i} > \frac{r_i/A_i}{r_j/A_j} \text{ for } i \neq j. \quad (11)$$

The effect of sexual dimorphism on the left-hand side term depends on the relative strengths of intra- and interspecific competition. When intraspecific competition is greater than interspecific competition (i.e. $\alpha_{ii} > \alpha_{ji}$), the left-hand side term increases to infinity with increasing sexual dimorphism. In contrast, when interspecific competition between resources is greater than intraspecific competition (i.e. $\alpha_{ji} > \alpha_{ii}$), the left-hand side term of the general coexistence condition decreases to zero with increasing levels of sexual dimorphism. The effect of sexual dimorphism on the right-hand side term of the general coexistence criterion is as discussed for the case of competitive symmetry. Together these observations imply that if resource 1 is the better direct competitor, that is $\alpha_{21} > \alpha_{11}$ and $\alpha_{21} < \alpha_{22}$, then coexistence requires that resource 2 is the better apparent competitor, that is

$r_2/A_2 > r_1/A_1$. In which case, a sexual dimorphism can help consumer-mediated coexistence.

3.2 | Numerical results: Symmetric resource acquisition across the sexes

Our numerical results, which show the effects of sex-specific resource preference in comparison to the case of both sexes being generalists, generally matched conclusion from our analytical solutions and are summarized in Table 1. We generally find that ecological sexual dimorphism leads to a reduction in consumer density and corresponding increase in resource density.

When competition is purely apparent and resource acquisition by the consumer is symmetric across the sexes, increasing consumer sexual dimorphism in resource preference (i.e. increasing β) leads to an increase in resource density (Figure 2a vs. 2b) and a reduction in consumer density (similar to the effects in a single resource community shown in Figure 3a). These effects result in an increase in the parameter space under which resources can coexist, consistent with our analytical results (Figure 4a). In particular, as sexual dimorphism approaches the extreme (β approaches unity and each sex uses a different resource exclusively), competitively inferior resource species can coexist with a superior competitor (Figure 4). This effect is due to the fact that consumer persistence depends on the density of both resources at this extreme. Sex differences in mortality alter the shape of the relationship between sexual dimorphism and resource coexistence (Figure 4c,d), although a pattern of expanded coexistence at the extreme of consumer sexual dimorphism β is maintained.

When resources compete directly with symmetric intraspecific competition, resource acquisition is symmetric across the sexes and intrinsic per capita growth rates are equal between the resources, consumer sexual dimorphism in resource preference (β) has little effect on invasion of a resource into a consumer-single-resource community at equilibrium (S1A-C, S4A). Yet even in the absence of effects on coexistence, consumer sexual dimorphism in resource preference (β) has strong effects on equilibrium resource abundance, both under analytical solutions of a two-species community and under simulations of three species communities (Figure 3). As predicted by our analytical conditions, when resource growth rates are unequal, sexual dimorphism in resource preference (β) changes

TABLE 1 Summary of the effects of sex differences in resource use. Arrows indicate effects on mean species density and the range of parameter space under which resource coexistence is observed

Density	Coexistence	
Consumer ↓	Apparent competition	↑
Resource ↑	Direct competition, equal resource growth rates, symmetric acquisition	0
	Direct competition, different resource growth rates, symmetric acquisition	↑
	Direct competition, asymmetric acquisition	↑↓




FIGURE 2 Coexistence mediated by sexual dimorphic consumers. When competition is completely apparent (a, b), the resource with superior growth rate (resource 2, red) excludes the inferior resource (resource 1, blue) when consumers are sexually monomorphic (a). Under the same parameterization but with sexually dimorphic consumers ($\beta = 0.9$; b), density of both resources is increased and they coexist. When resources compete directly, c, resource 2 excludes resource 1 and the sexually dimorphic consumer does not persist. However, adding consumer sex differences in total resource acquisition, under otherwise identical parameter values, leads to consumer persistence and coexistence of the resource species with cyclical dynamics (d). Parameter values: (a) $\alpha_{11} = 0.1$, $\alpha_{12} = 0$, $\alpha = 0.1$, $\alpha_{21} = 0$, $\beta = 0$, $d_M = d_F = 0.02$, $b_M = b_F = 0.1$, $a_{M,\max} = 1$, $a_{F,\max} = 1$, $r_1 = 1$ and $r_2 = 1.1$; (b) $\alpha_{11} = 0.1$, $\alpha_{12} = 0$, $\alpha_{22} = 0.1$, $\alpha_{21} = 0$, $\beta = 0.9$, $d_M = d_F = 0.02$, $b_M = b_F = 0.1$, $a_{M,\max} = 1$, $a_{F,\max} = 1$, $r_1 = 1$ and $r_2 = 1.1$; (c) $\alpha_{11} = 0.1$, $\alpha_{12} = 0.12$, $\alpha_{22} = 0.1$, $\alpha_{21} = 0.09$, $\beta = 1$, $d_M = d_F = 0.02$, $b_M = b_F = 0.1$, $a_{M,\max} = a_{F,\max} = 1$ and $r_1 = r_2 = 1$; (d) $\alpha_{11} = 0.1$, $\alpha_{12} = 0.12$, $\alpha_{22} = 0.1$, $\alpha_{21} = 0.09$, $\beta = 1$, $d_M = d_F = 0.02$, $b_M = b_F = 0.1$, $a_{M,\max} = 0.8$, $a_{F,\max} = 1.2$, $r_1 = 1$ and $r_2 = 1$

expected regions of coexistence of both resources and the consumer, leading to invasion and persistence of competitively inferior resource that is a superior apparent competitor in that it has a larger value of r_i/A_i (Figure 5b; Figure S4B; Figure S1).

3.3 | Numerical results: Asymmetric total resource acquisition

Introducing asymmetric total resource acquisition, represented as sex differences in the maximal attack rates $a_{F,\max}$ and $a_{M,\max}$, has complex effects on both population dynamics and coexistence between competing resources. Introducing asymmetric attack rates $a_{F,\max} \neq a_{M,\max}$ can promote coexistence (as predicted by the analytical results) while simultaneously creating cyclical resource dynamics (Figure 2c,d). Under pure apparent competition and asymmetric attack rates $a_{F,\max} \neq a_{M,\max}$, consumer sexual dimorphism in resource preference has some similar effects to the case of symmetry ($a_{F,\max} = a_{M,\max}$), mediating the coexistence of resources that differ in intrinsic per capita growth rates and leading to expanded regions of resource coexistence in comparison to a community with a sexually monomorphic consumer (Figure 4b).

Under direct interspecific competition, sexual dimorphism in resource preference (high value of β) has striking effects on resource persistence when the sexes differ in total resource acquisition ($a_{F,\max} \neq a_{M,\max}$), expanding regions of coexistence (Figure 5a,b) and in some

cases leading to persistence of a competitively inferior resource (Figure 5c,d; Supplement B Figure S4b,c). Moreover, sex differences in total resource acquisition ($a_{F,\max} \neq a_{M,\max}$) can reverse competitive outcomes when resources differ in their intrinsic per capita growth rates and the degree of resource specialization β is held constant (Figure 5c,d). The effect of sex differences in total resource acquisition ($a_{F,\max} \neq a_{M,\max}$) on patterns of prey persistence can be as striking as the effects of sex differences in resource preference (β ; Supplement B Figures S1 and S2), as can the effects of sex differences in morality (Figure 5e vs. 5f), which serve to couple consumer abundance and resource abundance asymmetrically across resources when sexual dimorphism is present.

3.4 | Asymmetric contribution of resource acquisition to birth rate

Introducing asymmetries in the contribution of sex-specific resource acquisition to birth rate (e.g. polygyny or polyandry, manifest as changes of the constant b_M) had little qualitative effect. This similarity is illustrated in Supplement B Figure S3 (compare to S1). Although changing these constants results in shifts of absolute regions of coexistence, the influence of consumer sex differences was similar. The lack of sensitivity in asymmetries in the contribution of resource acquisition to birth rates is consistent with our analytical results (Supplement A) that apply to all first-order homogenous




FIGURE 3 Consumer sexual dimorphism results in increased resource density and reduced consumer density in two and three species communities. Panel a: Equilibrium consumer female density decreases and resource 1 density increases with increased sexual dimorphism in a two-species (consumer-1 resource) community (Equation 6). Panel b: these results hold in simulations in three species communities, where resource 1 density is plotted against competition coefficients in both resource species under three levels of sexual dimorphism, increasing from left to right. Assuming symmetric total attack rates across the sexes, symmetric contribution of resource acquisition to consumer birth rates, and equal growth rates across resources

mating functions including geometric and arithmetic contributions to reproduction.

4 | DISCUSSION

Using a general model of consumer-resource dynamics, we show that consumer sexual dimorphism has substantial consequences for community assembly. Competitive exclusion via apparent competition is expected and observed when males and females are monomorphic generalists and resources differ substantially in their intrinsic per capita growth rates. However, when male and female consumers differ in their resource-specific attack rates, resource species that differ substantially in their intrinsic per capita growth rates can coexist. Similar effects of consumer sexual dimorphism are observed when resources compete directly, with sexual dimorphism in some cases permitting coexistence or persistence of a competitively inferior resource (summarized in Table 1). However, for both direct and apparent competition, consumer sexual dimorphism can also lead to competitive exclusion between

resources that would typically be expected to coexist. Thus, consumer sex differences result in fundamental changes in the types of competing resources that can establish during community assembly. These results also support conclusions from other food web models suggesting trophic position may impact the observed ecological effects of sexual reproduction (Kawatsu 2018). Moreover, equilibrium resource abundances and temporal dynamics are altered by consumer sexual dimorphism even when long-term ecological outcomes are unaffected. Our results in many ways echo recent work demonstrating that ontogenetic differences in resource acquisition in a species can have similar complex consequences for community assembly (de Roos 2020). However, a key difference is that in sexually reproducing species, we expect to observe these types of effects whenever consumer mating propensity depends, in part, on resource acquisition, a biological reality for many sexually reproducing consumers (e.g. widespread evidence of developmental thresholds for transitions to reproduction, reviewed in Wilbur and Collins 1973, Day and Rowe 2002, and resource-dependent sexual displays, Bonduriansky 2007), and whenever male abundance matters for population growth rates.




FIGURE 4 Sexual dimorphism in resource acquisition expands regions of coexistence between resources under apparent competition. Panels show relative density of resource 1 ($R_1/[R_1 + R_2]$) from numerical simulations with starting conditions of $R_1 = R_2 = 1$ and $M = F = 1$. Panel a shows the outcomes under symmetric maximal attack rates across the sexes ($a_{M,\max} = a_{F,\max} = 1$) and equal death rates $d_M = d_F = 0.02$. Panel b shows the outcomes assuming sex differences in maximal attack rate ($a_{M,\max} = 0.8$, $a_{F,\max} = 1.2$). Panel c shows outcomes under identical conditions to a, but with a sex difference in death rate of approximately 11% (reduced male mortality, $d_M = 0.018$, $d_F = 0.02$). Panel d shows the opposite sex difference (reduced female mortality, $d_M = 0.02$, $d_F = 0.018$). White dashed lines indicate analytical solutions (Equation 8) limited to the range of consumer coexistence $H_i \frac{1}{a_{ii}} > 2$, which corresponds to $\beta < 0.96$

Although our results suggest that the effects of consumer sexual dimorphism on community assembly can be complex, our analysis reveals some key predictions from our model. First, consumer sex differences in resource-specific attack rates result in an increase in the equilibrium density of each resource and a decrease in the abundance of the consumer species. Intuitively, sexual dimorphism can frequently lead to slightly suboptimal resource use of the species due to a mismatch between demand (restricted by a 50:50 sex ratio) and resource availability which can be more variable and dynamic. Second, consumer sexual dimorphism can promote coexistence when the more aggressively feeding sex (higher total attack rates) specializes on and suppresses what would otherwise be the competitively superior resource. This latter situation may be commonplace if sex-specific natural selection favours specialization, by the sex with higher resource requirements, on the most abundant resource. Finally, and related to the previous observation, the

effects of consumer sexual dimorphism are most pronounced when resources either differ in intrinsic per capita growth rates, or when consumer sexes differ in their total resource acquisition.

Although the assumption of complete female demographic dominance is common in theoretical models in ecology and evolutionary biology, there is little empirical support for this demographic extreme. Our simulation models instead focused on a harmonic mean birth function, which is biologically realistic in that it captures severely reduced birth rates when any one sex is rare, which is likely to be the case even if sperm limitation is unimportant at sex ratios near 1:1. Empirically evaluating alternative birth functions in real systems is difficult, requiring extreme variation in adult sex ratios rarely seen in many species. However, some empirical data exist and support the harmonic mean as the best-fitting description of sexual birth rates (Miller and Inouye 2011). Nonetheless, our analytical results (see Supplement A for details) indicate that our conclusions




FIGURE 5 Sexual dimorphism alters regions of coexistence of resource species. Panels show relative density of resource 1 ($R_1/[R_1 + R_2]$) from numerical simulations with starting conditions of $R_1 = R_2 = 1$ and $M = F = 1$. Panel a shows outcomes under sexual monomorphism ($\beta = 0$), unequal growth rates ($r_1 = 2, r_2 = 2.1$) across resources and no sex differences in total attack rate ($a_{M,\max} = a_{F,\max} = 1$). Panel b shows the outcome under the same parameter values but with strong consumer sexual dimorphism ($\beta = 0.7$). Panels c and d show the effect of sexual niche divergence when the sexes also differ in total attack rates (total resource acquisition), with c showing the outcomes under unequal total attack rates ($a_{M,\max} = 0.7, a_{F,\max} = 1.3$), sexual monomorphism in prey preference ($\beta = 0$) and differential resource growth rates ($r_1 = 1.6, r_2 = 1.8$), and d showing the outcome under the same parameter values but with moderate sexual dimorphism in prey preference ($\beta = 0.4$). Panels e and f show the effects of sex differences in mortality; in Panel e, $d_M = 0.018, d_f = 0.02$. In Panel f, $d_M = 0.02, d_f = 0.018$, while $\beta = 0.5$ in both panels. Note that in the absence of sex differences in mortality coexistence would be restricted to the bottom left quadrant represented by black dashed lines, which demark equal inter and intraspecific competition coefficients. White dashed box demarks coexistence determined by analytical invasion criteria into a two-species community under Equation 9

apply to any birth function that is first-order homogenous, that is $B(aM_{M_i}, aF_{F_i}) = aB(M_{M_i}, F_{F_i})$ for any positive constant $a > 0$. These functions include all functions typically considered candidate birth functions by demographers (Caswell and Weeks 1986, Miller and Inouye 2011). Our conclusions do depend on the assumption that male abundance makes some contribution to birth rate.

Although even moderate sex differences in attack rates can change the conditions under which resources coexist, the strongest effects are observed when the consumer population approaches complete sexual dimorphism in resource-specific attack rates. Cases of such extreme sex differences in diet preference exist (Temeles et al. 2000), although whether they occur with any regularity is unclear. Diet divergence between the sexes, and sexual dimorphisms in trophic morphology are common (Shine 1989), although overlap in male and female diets can be substantial even in the presence of ecological sex differences (e.g. Stamps et al. 1997, De Lisle and Rowe 2015a). Yet overlap in diet content may be expected even in the case of large differences in attack rates, as diet content is the product of both attack rate and resource abundance. Further, males and females may fail to diverge in diet preference even when optimum diet nutritional content differs across the sexes (Reddiex et al. 2013). These challenges to understanding sex-specific resource acquisition highlight the need for further empirical studies disentangling expressed diet preference, male and female nutritional optima and the evolution of ecological sexual dimorphism.

We have assumed a Fisherian sex ratio, where the primary sex ratio is maintained at a stable 1:1 ratio (Fisher 1930), although our model allows for deviations from a 1:1 operational sex ratio via sex-specific intrinsic mortality. In fact, our results show that even slight sex differences in mortality rates can have dramatic consequences for community assembly under both pure apparent and direct competition between resource species. Sex-specific mortality is commonplace in natural populations and can occur for a variety of reasons, such as sex-specific costs of reproduction, predation or sexual conflict. In our model, sex-specific mortality and variation in consumer mating system (see Supplemental material) had strong but predictable consequences for resource abundance, leading to increases in the abundance of the resource favoured by the sex with higher intrinsic mortality. Although we did not model resource-dependent consumer mortality, these results suggest that an expansion of our model to include resource-dependent consumer mortality would exacerbate the already striking community-wide effects of consumer sexual dimorphism.

Our analysis also assumes that the male and female trophic traits are constant. In nature, however, the degree and even direction of sexual dimorphism can vary dramatically among closely related populations (e.g. Reimchen et al. 2016). Dimorphism clearly evolves rapidly in response to spatially varying sexual and natural selection, and resource availability. Future extensions to our strictly ecological model could add in eco-evolutionary dynamics of sexual dimorphism.

Male and female densities were reduced with increasing ecological sexual dimorphism, because consumer births are limited by resource acquisition in both sexes in our model. In the extreme case, where

the consumer cannot persist on a single resource (e.g. under complete sexual dimorphism), coexistence of the consumer depends on stable coexistence of the resource species (Supplement A). Our simulations suggest robust persistence of extremely dimorphic consumers in two-resource communities, albeit at reduced density. Both of these results—reduced consumer density and dependence on the presence of both resource species in the community—suggest that sexually dimorphic populations may face a higher risk of extinction due to demographic stochasticity. However, it is difficult from our purely deterministic ecological model to fully interpret the consequences of sexual dimorphism on extinction probability, and our emphasis on a null model of sexually monomorphic generalists is useful for understanding the ecological effects of sexual dimorphism but perhaps tenuous for interpreting our results in the context of the evolution of ecological sexual dimorphism. Evolution of sexual dimorphism from a monomorphic ancestor that specializes on a single resource could lead to increased population mean fitness, by increasing the total resource pool available across both sexes (Rand 1952, Selander 1966, Slatkin 1984, Li and Kokko 2021); such a process represents a form of within-species, between-sex ecological character displacement, and may be particularly likely to occur when the sexes interact in small demes (Li and Kokko 2021). More generally, evolution of sexual dimorphism is expected to be critical to population mean fitness whenever optima differ for males and females (Lande 1980), and empirical data suggest sexual dimorphism can be associated with reduced extinction probability at both the macroevolutionary scale and in extant populations (De Lisle and Rowe 2015b). Nonetheless, our results suggest a full understanding of sexual dimorphism's role in population persistence could require integrating theory and data from population/community ecology and evolutionary genetics. This conclusion is complemented by recent theory (de Vries and Caswell 2019) indicating sexual dimorphism in demographic parameters can have important consequences for maintaining genetic diversity.

Our model generally predicts that ecological sexual dimorphism may in some cases promote, and in other cases reduce, diversity at lower trophic levels during community assembly. Testing this prediction with correlative data (reviewed in Tsuji and Fukami 2020) would be possible but challenging. An alternative and non-exclusive hypothesis, that dimorphic predators are more likely to establish in communities with diverse prey assemblages, would also generate patterns consistent with our results. Alternatively, experiments that manipulate the expressed dimorphism of predators and track community dynamics at lower trophic levels are tractable in some systems. Similar experiments have been performed (Fryxell et al. 2015, Start and De Lisle 2018), in which the effects of predator sex ratio manipulation on prey communities are assessed in mesocosm designs. Although the results of these experiments do suggest sex differences can have important community consequences, no studies (to our knowledge) have compared communities in which the magnitude of morphological sexual dimorphism is manipulated under a stable sex ratio. Such designs are possible when distributions of male and female phenotypes exhibit substantial variation and would represent an ideal empirical test of the theoretical results presented here.

Ecological sex differences are commonplace, although the details of their evolutionary drivers and ecological consequences are unclear. Emerging theory and data indicate ecological sex differences may have important consequences for the evolutionary genetics of adaptation (Zajitschek and Connallon 2017), the dynamics of diversification (Bolnick and Doebeli 2003, De Lisle and Rowe 2015b) and community assembly (Fryxell et al. 2015, Pincheira-Donoso et al. 2018, Start and De Lisle 2018). Our results add to this body of work, indicating that sexual dimorphism can have substantial effects on the structure, abundance and dynamics of ecological communities, including changing conditions for coexistence between competing resource species.

ACKNOWLEDGEMENTS

The authors thank Tim Connallon and Miguel Gomez for comments on an early draft of this manuscript, as well as Gonzalo Hernando for assistance in the early stages of the project. This work was supported by funds from the University of Connecticut to D.I.B., an establishment grant from the Swedish Research Council to S.P.D.L. (2019-03706) and a US National Science Foundation Grant DMS176803 to S.J.S.

CONFLICT OF INTEREST

The authors have no conflict of interest to declare.

AUTHORS' CONTRIBUTIONS

D.I.B., S.P.D.L. and S.J.S. designed the study; S.J.S. performed the mathematical analysis; S.P.D.L. performed the simulations and drafted the manuscript. All authors revised the manuscript.

DATA AVAILABILITY STATEMENT

No data are. R script is archived at <https://doi.org/10.5281/zenodo.6324965> (De Lisle, Schreiber, & Bolnick, 2022).

ORCID

Stephen P. De Lisle  <https://orcid.org/0000-0001-9587-8665>

Sebastian J. Schreiber  <https://orcid.org/0000-0002-5481-4822>

Daniel I. Bolnick  <https://orcid.org/0000-0003-3148-6296>

REFERENCES

- Andersson, M. B. (1994). *Sexual selection*. Princeton University Press.
- Belovsky, G. E. (1978). Diet optimization in a generalist herbivore: The moose. *Theoretical Population Biology*, 14, 105–134.
- Bolnick, D. I., Amarasekare, P., Araujo, M. S., Burger, R., Levine, J. M., Novak, M., Rudolf, V. H. W., Schreiber, S., Urban, M. C., & Vasseur, D. A. (2011). Why intraspecific trait variation matters in community ecology. *Trends in Ecology & Evolution*, 26, 183–192.
- Bolnick, D. I., & Doebeli, M. (2003). Sexual dimorphism and adaptive speciation: Two sides of the same ecological coin. *Evolution*, 57, 2433–2449.
- Bolnick, D. I., Svanbäck, R., Fordyce, J. A., Yang, L. H., Davis, J. M., Hulsey, C. D., & Forister, M. L. (2003). The ecology of individuals: Incidence and implications of individual specialization. *The American Naturalist*, 161, 1–28.
- Bonduriansky, R. (2007). The evolution of condition-dependent sexual dimorphism. *The American Naturalist*, 169, 9–19.
- Camus, M. F., Fowler, K., Piper, M. W. D., & Reuter, M. (2017). Sex and genotype effects on nutrient-dependent fitness landscapes in *Drosophila melanogaster*. *Proceedings of the Royal Society of London Series B*, 284, 20172237.
- Caswell, H., & Weeks, D. E. (1986). Two-sex models: Chaos, extinction, and other dynamic consequences of sex. *American Naturalist*, 128, 707–735.
- Cortez, M. H., & Patel, S. (2017). The effects of predator evolution and genetic variation on predator-prey population-level dynamics. *Bulletin of Mathematical Biology*, 79, 1510–1538.
- Darwin, C. (1871). *The descent of man and selection in relation to sex*. J. Murray.
- Day, T., & Rowe, L. (2002). Developmental thresholds and the evolution of reaction norms for age and size at life-history transitions. *The American Naturalist*, 159, 338–350.
- De Lisle, S. P. (2019). Understanding the evolution of ecological sex differences: Integrating character displacement and the Darwin-Bateman paradigm. *Evolution Letters*, 3, 444–447.
- De Lisle, S. P., & Rowe, L. (2015a). Ecological character displacement between the sexes. *The American Naturalist*, 186, 693–707.
- De Lisle, S. P., & Rowe, L. (2015b). Independent evolution of the sexes promotes amphibian diversification. *Proceedings of the Royal Society B: Biological Sciences*, 282, 20142213.
- De Lisle, S. P., Schreiber, S., & Bolnick, D. I. (2022). Data from: Complex community wide consequences of consumer sexual dimorphism. *Zenodo*, <https://doi.org/10.5281/zenodo.6324965>
- de Roos, A. M. (2020). Effects of life history and individual development on community dynamics: A review of counterintuitive consequences. *Ecological Research*, 35, 930–946.
- de Vries, C., & Caswell, H. (2019). Selection in two-sex stage structured populations: Genetics, demography, and polymorphism. *Theoretical Population Biology*, 130, 160–169.
- Des Roches, S., Post, D. M., Turley, N. E., Bailey, J. K., Hendry, A. P., Kinnison, M. T., Schweitzer, J. A., & Palkovacs, E. P. (2018). The ecological importance of intraspecific variation. *Nature Ecology and Evolution*, 2, 57–64.
- Fisher, R. (1930). *The genetical theory of natural selection*. Oxford University Press.
- Fryxell, D. C., Arnett, H. A., Apgar, T. M., Kinnison, M. T., & Palkovacs, E. P. (2015). Sex ratio variation shapes the ecological effects of a globally introduced freshwater fish. *Proceedings of the Royal Society of London Series B*, 282, 20151970.
- Fryxell, D. C., Weiler, D., Kinnison, M., & Palkovacs, E. P. (2019). Eco-evolutionary dynamics of sexual dimorphism. *Trends in Ecology & Evolution*, 34, 591–594.
- Garlapow, M. E., Huang, W., Yarboro, M. T., Peterson, K. R., & Mackay, T. F. C. (2015). Quantitative genetics of food intake in *Drosophila melanogaster*. *PLoS ONE*, 10, e0138129.
- Giery, S. T., & Layman, C. G. (2019). Ecological consequences of sexually selected traits: An eco-evolutionary perspective. *The Quarterly Review of Biology*, 94, 29–74.
- Gomez-Llano, M. A., Bensch, H. M., & Svensson, E. I. (2018). Sexual conflict and ecology: Species composition and male density interact to reduce male mating harassment and increase female survival. *Evolution*, 72, 906–915.
- Haldane, J. B. S. (1937). The effect of variation on fitness. *The American Naturalist*, 71, 337–349.
- Holt, R. D. (1977). Predation, apparent competition, and the structure of prey communities. *Theoretical Population Biology*, 12, 197–229.
- Holt, R. D., & Lawton, J. H. (1993). Apparent competition and enemy-free space in insect host-parasitoid communities. *The American Naturalist*, 142, 623–645.

- Hughes, A. R., Inouye, B. D., Johnson, M. T. J., Underwood, N., & Vellend, M. (2008). Ecological consequences of genetic diversity. *Ecology Letters*, *11*, 609–623.
- Jensen, K., McClure, C., Priest, N. K., & Hunt, J. (2015). Sex-specific effects of protein and carbohydrate intake on reproduction but not lifespan in *Drosophila melanogaster*. *Aging Cell*, *14*, 605–615.
- Kawatsu, K. (2018). Ecological effects of sex differ with trophic positions in a simple food web. *Ecology and Evolution*, *8*, 1239–1246.
- Lande, R. (1980). Sexual dimorphism, sexual selection, and adaptation in polygenic characters. *Evolution*, *34*, 292–305.
- Li, X.-Y., & Kokko, H. (2021). Sexual Dimorphism driven by intersexual resource competition: Why it is rare, and where to look for it? *Journal of Animal Ecology*, *90*, 1831–1843.
- Lindström, J., & Kokko, H. (1998). Sexual reproduction and population dynamics: the role of polygyny and demographic sex differences. *Proceedings of the Royal Society of London Series B*, *265*, 483–488.
- Maklakov, A., Simpson, S. J., Zajitchek, F., Hall, M. D., Dessmann, J., Clissold, F., Raubenheimer, D., Bonduriansky, R., & Brooks, R. C. (2008). Sex-specific effects of nutrient intake on reproduction and lifespan. *Current Biology*, *18*, 1062–1066.
- Miller, T. E. X., & Inouye, B. D. (2011). Confronting two-sex demographic models with data. *Ecology*, *92*, 2141–2151.
- Patel, S., & Schreiber, S. J. (2015). Evolutionarily driven shifts in communities with intraguild predation. *The American Naturalist*, *186*, E98–E110.
- Patel, S., & Schreiber, S. J. (2018). Robust permanence for ecological equations with internal and external feedbacks. *Journal of Mathematical Biology*, *77*, 79–105. <https://doi.org/10.1007/s00285-017-1187-5>
- Pincheira-Donoso, D., Tregenza, T., Butlin, R. K., & Hodgson, D. J. (2018). Sexes and species as rival units of niche saturation during community assembly. *Global Ecology and Biogeography*, *27*, 593–603.
- R Core Team. (2018). *R: A language and environment for statistical computing*. R Foundation for Statistical Computing.
- Rand, A. L. (1952). Secondary sexual characters and ecological competition. *Fieldiana-Zoology*, *34*, 65–70.
- Reddiex, A. J., Gosden, T. P., Bonduriansky, R., & Chenoweth, S. F. (2013). Sex-specific fitness consequences of nutrient intake and the evolvability of diet preferences. *The American Naturalist*, *182*, 91–102.
- Reimchen, T., Steeves, D., & Bergstrom, C. A. (2016). Sex matters for defence and trophic traits of threespine stickleback. *Evolutionary Ecology Research*, *17*, 459–485.
- Rowe, L., & Houle, D. (1996). The lek paradox and the capture of genetic variance by condition dependent traits. *Proceedings of the Royal Society of London Series B - Biological Sciences*, *263*, 1415–1421.
- Schreiber, S. J. (2000). Criteria for C^1 robust permanence. *Journal of Differential Equations*, *162*, 400–426.
- Schreiber, S. J. (2021). The P^* rule in the stochastic Holt-Lawton model of apparent competition. *Discrete and Continuous Dynamical Systems B*, *26*, 633–644.
- Schreiber, S. J., Burger, R., & Bolnick, D. I. (2011). The community effects of phenotypic and genetic variation within a predator population. *Ecology*, *92*, 1582–1593.
- Selander, R. K. (1966). Sexual dimorphism and differential niche utilization in birds. *The Condor*, *68*, 113–151.
- Shärer, L., Rowe, L., & Arnqvist, G. (2012). Anisogamy, chance and the evolution of the sex roles. *Trends in Ecology & Evolution*, *27*, 260–264.
- Shine, R. (1989). Ecological causes for the evolution of sexual dimorphism: A review of the evidence. *Quarterly Review of Biology*, *64*, 419–461.
- Slatkin, M. (1984). Ecological causes of sexual dimorphism. *Evolution*, *38*, 622–630.
- Soetgart, K., Petzoldt, T., & Setzer, R. W. (2010). Solving differential equations in R: Package deSolve. *Journal of Statistical Software*, *33*, 1–25.
- Stamps, J. A., Losos, J. B., & Andrews, R. M. (1997). A comparative study of population density and sexual size dimorphism in lizards. *The American Naturalist*, *149*, 64–90.
- Start, D., & De Lisle, S. P. (2018). Sexual dimorphism in a top predator drives aquatic prey community assembly. *Proceedings of the Royal Society of London Series B*, *2018*, 1717.
- Svensson, E. I. (2019). Eco-evolutionary dynamics of sexual selection and sexual conflict. *Functional Ecology*, *33*, 60–72.
- Temeles, E. J. (1985). Sexual size dimorphism of bird-eating hawks: The effect of prey vulnerability. *The American Naturalist*, *125*, 485–499.
- Temeles, E. J., Pan, I. L., Brennan, J. L., & Horwitt, J. N. (2000). Evidence for ecological causation of sexual dimorphism in a hummingbird. *Science*, *289*, 441–443.
- Trivers, R. L. (1972). Parental investment and sexual selection. In B. Campbell (Ed.), *Sexual selection and the descent of man* (pp. 136–179). Aladine.
- Tsuji, K., & Fukami, T. (2018). Community-wide consequences of sexual dimorphism: Evidence from nectar microbes in dioecious plants. *Ecology*, *99*, 2476–2484.
- Tsuji, K., & Fukami, T. (2020). Sexual dimorphism and species diversity: From clades to sites. *Trends in Ecology & Evolution*, *35*, 105–114.
- Tuszynski, J. 2019. *caTools: Tools: Moving window statistics, GIF, Base64, ROC AUC, etc.* R package version 1.17.1.2. Retrieved from <https://CRAN.R-project.org/package=caTools>
- Weimerskirch, H., Le Corre, M., Gadenne, H., Pinaud, D., Kato, A., Ropert-Coudert, Y., & Bost, C.-A. (2009). Relationship between reversed sexual dimorphism, breeding investment and foraging ecology in a pelagic seabird, the masked booby. *Oecologia*, *161*, 637–649.
- Wilbur, H. M., & Collins, J. P. (1973). Ecological aspects of amphibian metamorphosis. *Ecological aspects of amphibian metamorphosis Science*, *182*, 1305–1314.
- Zajitschek, F., & Connallon, T. (2017). Partitioning of resources: The evolutionary genetics of sexual conflict over resource acquisition and allocation. *Journal of Evolutionary Biology*, *30*, 826–838.

SUPPORTING INFORMATION

Additional supporting information may be found in the online version of the article at the publisher's website.

How to cite this article: De Lisle, S. P., Schreiber, S. J. & Bolnick, D. I. (2022). Complex community-wide consequences of consumer sexual dimorphism. *Journal of Animal Ecology*, *91*, 958–969. <https://doi.org/10.1111/1365-2656.13685>