

Unveiling new species of Phragmidiaceae (Basidiomycota, Pucciniales) on rosaceous plants from Guizhou, China

Qinfang Zhang^{1,2}, Qianzhen Wu^{1,2}, Peng Zhao³, Kamran Habib¹, Yao Wang², Dexiang Tang¹, Muhammad Aijaz Ahmad¹, Yulin Ren^{1,4}, Xiangchun Shen¹, Qingde Long^{1,2}, Lili Liu¹, Qirui Li^{1,2}

¹ State Key Laboratory of Discovery and Utilization of Functional Components in Traditional Chinese Medicine & School of Pharmaceutical Sciences, Guizhou Medical University, Gui'an New District, Guizhou 550004, China

² The High Efficacy Application of Natural Medicinal Resources Engineering Center of Guizhou Province (The Key Laboratory of Optimal Utilization of Natural Medicine Resources), School of Pharmaceutical Sciences, Guizhou Medical University, University Town, Gui'an New District, Guizhou 550004, China

³ State Key Laboratory of Mycology, Institute of Microbiology, Chinese Academy of Sciences (CAS), Beijing 100101, China

⁴ Immune Cells and Antibody Engineering Research Center of Guizhou Province, Guizhou Medical University, Guiyang 550004, China

Corresponding authors: Qirui Li (lqrnd2008@163.com); Lili Liu (lililiu550025@163.com)

Abstract

Rust fungi associated with *Rubus* were collected across diverse locations in Guizhou Province, and three new species – *Gerwasia amphidasys* on *Rubus amphidasys*, *Phragmidium coreanicola* on *Rubus coreanus*, and *Phragmidium parvifolius* on *Rubus parvifolius* are introduced. These novel species are described based on morphological characteristics and phylogenetic analysis of the ITS and LSU loci. Additionally, *Gerwasia rubi-setchuenensis* is introduced as a new host record on *Rubus buergeri*. The study includes comprehensive morpho-anatomical descriptions, detailed illustrations, and a phylogenetic tree, providing insights into the taxonomic placement and relationships of these novel taxa within their respective lineages.



Key words: ITS, LSU, Phragmidiaceae, phylogeny, rust disease, taxonomy

Academic editor:

Samantha C. Karunarathna

Received: 13 January 2025

Accepted: 2 March 2025

Published: 28 March 2025

Citation: Zhang Q, Wu Q, Zhao P, Habib K, Wang Y, Tang D, Ahmad MA, Ren Y, Shen X, Long Q, Liu L, Li Q (2025) Unveiling new species of Phragmidiaceae (Basidiomycota, Pucciniales) on rosaceous plants from Guizhou, China. MycoKeys 115: 309–326. <https://doi.org/10.3897/mycokeys.115.146604>

Copyright: © Qinfang Zhang et al.

This is an open access article distributed under terms of the Creative Commons Attribution License (Attribution 4.0 International – CC BY 4.0).

Introduction

Pucciniales (Basidiomycota, Pucciniomycetes) represents approximately 25% of basidiomycete fungi and constitutes one of the most prevalent fungal groups, parasitizing leaves, fruits, and branches of plants, thereby inhibiting normal growth and development of plants and impacting yield and quality (Aime and McTaggart 2021; Zhao et al. 2022a). To date, seven suborders (i.e., Araucariomycetinae, Melampsorineae, Mikronegeriineae, Raveneliineae, Rogerpetersoniineae, Skierkineae, and Urediniineae) and 18 families (i.e., Araucariomycetaceae, Coleosporiaceae, Crossosporaceae, Gymnosporangiaceae, Melampsoraceae, Milesinaceae, Ochropsoraceae, Phakopsoraceae, Phragmidiaceae, Pileolariaceae, Pucciniaceae, Pucciniastraceae, Raveneliaceae, Rogerpetersoniaceae, Skierkaceae, Sphaerophragmiaceae, Tranzscheliaceae, and Zaghouniaceae) have been reported (Aime et al. 2017; Aime and McTaggart 2021; Zhao et al. 2022b).

According to estimations, there are approximately 35,000 higher plant species in China, which are categorized into 454 families and 3,818 genera (Wang

et al. 2015). It is further estimated that between 1,700 and 8,800 species of rust fungi may exist within the country, posing a significant threat to the health and productivity of these plant species (Zhao et al. 2021). More than 8,400 rust taxa have been documented worldwide (Zhao et al. 2021; Sun et al. 2022). Based on their morphological characteristics and host associations, 1,200 species from 71 genera across 15 families have been documented in China. Among them, more than 70 species of *Phragmidium* have been identified. (Dai 1979; Zhuang et al. 1998, 2003, 2005, 2012, 2021; Ji et al. 2019, 2020; Zhao et al. 2021; Sun et al. 2022, 2024). Plants are susceptible to Coleosporiaceae, Melampsoraceae, Phragmidiaceae, and Pucciniaceae, which are the dominant families of rust taxa in China (Zhao et al. 2021; Zhao et al. 2022b; Sun et al. 2024).

The genus *Phragmidium* (Pucciniales: Phragmidiaceae) was established by Link (1816). Phragmidiaceae comprises a diverse array of species, encompassing 14 genera and an estimated 200 species, primarily targeting the economically vital family Rosaceae (examples of hosts in Rosaceae: *Rosa*, *Rubus*, *Potentilla*, *Sanguisorba*, *Duchesnea*, and *Acaena*) (Zhao et al. 2021; Sun et al. 2024). A total of 115 *Phragmidium* species have been described (Cummins 1931; Arthur 1934; Wahyuno et al. 2001; Cummins and Hiratsuka 2003; Zhuang et al. 2012; Yang et al. 2015; Ali et al. 2017; Zhao et al. 2021; Sun et al. 2022; Ya 2024). Among these, 38 species specifically target and parasitize members of the genus *Rosa* (Ya 2024), and at least 39 *Phragmidium* species have been reported in China (Wei 1988; Zhuang et al. 2003, 2012; Yang et al. 2015; Liu et al. 2018).

The susceptibility of Rosaceae plants to Phragmidiaceae infections underscores the importance of understanding the biology, ecology, and management strategies of these fungal pathogens (Dietel 1905; Yun et al. 2011; Sun et al. 2022). During the field investigation of rust fungi on medicinal plants in Guizhou Province, China, three potential new species were found belonging to the genera *Gerwasia* and *Phragmidium* of Phragmidiaceae, which infect the members of Rosaceae, as well as one species that infects a new host. We conducted a phylogenetic analysis using multi-locus (ITS and LSU) phylogeny and morphological characteristics to better understand their taxonomic position. Descriptions, illustrations, and phylogenetic analysis results of all the novel species and the new host record are provided.

Materials and methods

Sample collection and preservation

Rust-infected specimens were collected in Guizhou Province, China, from August to November of each year from 2021 to 2023. All host and habitat information of specimens was recorded (Rathnayaka et al. 2024). Photographs of the infected plants were taken using a camera (Canon G15, Corporation, Tokyo, Japan). The samples were kept in blotting papers and were brought to the laboratory for examination. The collected specimens were partly kept in a refrigerator at 4 °C for spare parts and partly pressed and air-dried to make pressed specimens (Wei and Wang 2011; Wu et al. 2023). All specimens were deposited at the herbarium of Guizhou Medical University (**GMB**) and Kunming Institute of Botany, Chinese Academy of Sciences (**KUN-HKAS**).

Morphological characterization

Macroscopic characteristics were observed under a stereomicroscope (Olympus SZ61), and photographs were taken with a digital camera (Canon 700D) fitted with a light microscope (Nikon Ni). The infected portions were examined, and photographs were taken as described by Wu et al. (2023). More than 30 measurements were noted for each type of teliospores, urediniospores, and paraphyses for each sample using the Tarosoft (R) Image Frame Work v. 0.9.0.7. The images were arranged using Adobe Photoshop CS6 (Adobe Systems, the USA).

DNA extraction and PCR amplification

The infected portion of the rust fungus was scraped using a sterilized scalpel. DNA extraction was carried out following the manufacturer's protocols for the Biomiga Fungal gDNA Kit. The DNA samples were kept at -20°C . The rust-specific primer pairs Rust2inv (Aime 2006) and ITS4rust (Pfunder et al. 2001) were used for the PCR amplification of the regions of internal transcribed spacer (ITS), whereas the universal primer pairs LR0R (Hopple and Vilgalys 1999) and LR6 (Vilgalys and Hester 1990) were used for the large subunit ribosomal (LSU). The composition of a 25 μL PCR mixture comprised the following: 9.5 μL of double-distilled water, 12.5 μL of PCR Master Mix, 1 μL of each primer, as well as 1 μL of template DNA. The qualified PCR products were verified through 1.5% agarose gel electrophoresis, stained with GoldenView, and subsequently submitted to Sangon Co. China for sequencing (Zhao et al. 2016).

Phylogenetic analyses

All sequences were obtained in ABI file format and deposited in the GenBank (Table 1). The consensus sequences were blasted in GenBank using the BLAST algorithm. The similar sequences were retrieved from the GenBank database. The molecular phylogeny was inferred from a combined dataset of ITS and LSU sequences. The reference sequences retrieved from open databases originated from recently published literature (Zhao et al. 2021; Wu et al. 2023; Sun et al. 2024). All the ambiguous nucleotides were trimmed using BioEdit software v.7.0.5.3 and TrimAL (Hall 1999). Sequences were aligned using the MAFFT v.7.110 online tool (Katoh and Standley 2013). The alignments are available in TreeBASE (www.treebase.org/treebase-web/home.html) under ID31754 for LSU and ITS rDNA sequences. The maximum likelihood (ML) analysis was implemented in RAxML v.8.2.12 using the GTRGAMMA substitution model with 1,000 bootstrap replicates (Stamatakis 2015). The phylogenetic analyses were also performed using Bayesian inference in MrBayes v.3.2.1 (Ronquist et al. 2015) online. The Markov chain Monte Carlo (MCMC) sampling in MrBayes v.3.2.2 (Ronquist et al. 2015) was used to determine the Bayesian posterior probabilities (BYPP). Six simultaneous Markov chains were run for 3,000,000 generations, and trees were sampled every 1,000th generation. All analyses were run on the CIPRES Science Gateway v.3.3 web portal (Miller et al. 2010; Huelsenbeck and Ronquist 2001). The phylogenetic tree was visualized by Fig-Tree v.1.4.3 (Rambaut 2012).

Table 1. Taxa information and corresponding GenBank accession numbers of the sequences used in the phylogenetic analyses.

Species	Specimen No.	Host	Province, Country	GenBank Accession No.		References
				ITS	LSU	
<i>Gerwasia chinensis</i>	HMAS249978 (ZP-R5)	<i>Rubus parkeri</i>	Yunnan, China	MK519039	MK518737	Zhao et al. 2021
<i>G. chinensis</i>	HMAS249980 (ZP-R295)	<i>Rubus parkeri</i>	Yunnan, China	MK519038	MK518540	Zhao et al. 2021
<i>G. pittieriana</i>	BPI 843556	<i>Rubus</i> sp.	USA	KY764065	KY764065	Zhao et al. 2021
<i>G. rubi</i>	(ZP-R345)	<i>Rubus setchuenensis</i>	Sichuan, China	–	MK518735	Zhao et al. 2021
<i>G. amphidasys</i>	GMB4047*	<i>Rubus amphidasys</i>	Guizhou, China	PQ472136	PQ456450	This study
<i>G. amphidasys</i>	GMB4076	<i>Rubus amphidasys</i>	Guizhou, China	PQ472137	PQ456451	This study
<i>G. rubi-setchuenensis</i>	GMB4052	<i>Rubus buergeri</i>	Guizhou, China	PQ472135	PQ456449	This study
<i>G. rubi-setchuenensis</i>	GMB4075	<i>Rubus buergeri</i>	Guizhou, China	PQ472134	PQ456448	This study
<i>G. rubi-setchuenensis</i>	HGUP21168*	<i>Rubus setchuenensis</i>	Guizhou, China	OR470045	OR528540	Sun et al. 2024
<i>G. rubus-playfairianus</i>	HMAS249840 (ZP-R 1374)	<i>Rubus playfairianus</i>	Guangxi, China	MK518976	MK518674	Sun et al. 2024
<i>Gymnosporangium asiaticum</i>	CUP-0016*	<i>Juniperus chinensis</i>	Japan	MN642593	MN642617	Zhao et al. 2020
<i>Gymnosporangium sabinae</i>	TNM F0030477	<i>Pyrus communis</i>	Bulgaria: Sofia	KY964764	KY964764	Shen et al. 2017
<i>Hamaspora acutissima</i>	BRIP:55606	<i>Rubus rolfei</i>	Philippines	–	KT199398	McTaggart et al. 2016
<i>H. sinica</i>	ZP-R1	<i>Rubus setchuenensis</i>	Guangdong, China	MK519049	MK518636	Zhao et al. 2020
<i>H. longissima</i>	BPI 871506	<i>Rubus rigidu</i>	South Africa	–	MW049262	Aime and McTaggart 2021
<i>H. rubi-lambertianuse</i>	HGUP21164*	<i>Rubus lambertianus</i>	Guizhou, China	OR470053	OR528547	Sun et al. 2024
<i>H. rubi-parkerii</i>	HGUP21159*	<i>Rubus parkeri</i>	Guizhou, China	OR470055	OR528543	Sun et al. 2024
<i>H. rubi-alceifolii</i>	GMB0109*	<i>Rubus alceifolius</i>	Guizhou, China	OQ067094	OQ067532	Wu et al. 2023
<i>H. rubi-alceifolii</i>	GMB0116	<i>Rubus alceaefolius</i>	Guizhou, China	OQ067095	OQ067533	Wu et al. 2023
<i>Kuehneola uredinis</i>	LD1029	<i>Rubus</i> sp.	New York, America	GU058013	GU058013	Dixon et al. 2018
<i>Phragmidium barnardii</i>	HGUP21035	<i>Rubus parvifolius</i>	Guizhou, China	OL684828	OL684839	Sun et al. 2022
<i>P. charyuensis</i>	BJFC: R02532*	<i>Rosa duplicata</i>	China	MH128374	NG_064492	Liu et al. 2018
<i>P. cibatum</i>	BJFC: R02528	<i>Rubus niceus</i>	China	MH128370	NG_064491	Liu et al. 2018
<i>P. cymosum</i>	GMB0115	<i>Rosa cymosa</i>	Guizhou, China	OQ067097	OQ067531	Wu et al. 2023
<i>P. cymosum</i>	GMB0108*	<i>Rosa cymosa</i>	Guizhou, China	OQ067096	OQ067530	Wu et al. 2023
<i>P. coreanicola</i>	GMB0101*	<i>Rubus coreanus</i>	Guizhou, China	PQ472133	PQ456447	This study
<i>P. coreanicola</i>	GMB4071	<i>Rubus coreanus</i>	Guizhou, China	PQ472132	PQ456448	This study
<i>P. griseum</i>	HMAS56906	<i>Rubus crataegifolius</i>	Beijing, China	MH128377	MG669115	Liu et al. 2018
<i>P. griseum</i>	BJFCR 03451	–	Beijing, China	MN264713	MN264731	Liu et al. 2020
<i>P. griseum</i>	BJFCR03449	<i>Rubus crataegifolius</i>	Beijing, China	MN264712	MN264730	Liu et al. 2020
<i>P. japonicum</i>	HMAS41585	<i>Rosa laecigata</i>	Fujian, China	MN264716	MN264734	Liu et al. 2020
<i>P. jiangxiense</i>	BJFCR 03453*	<i>Rosa laecigata</i>	Jiangxi, China	MN264715	MN264733	Liu et al. 2020
<i>P. kanasense</i>	ZP-R1382	<i>Phoenix acaulis</i>	Yunnan, China	MK518980	MK518678	Zhao et al. 2021
<i>P. kanasense</i>	ZP-R491	<i>Rosa fedtschenkoana</i>	Xinjiang, China	–	MK518748	Zhao et al. 2021
<i>P. leucoaecium</i>	BJFCR 02118*	<i>Rosa</i> sp.	Yunnan, China	MN264719	MN264737	Liu et al. 2020
<i>P. mexicanum</i>	E14_5_1	<i>Potentilla indica</i>	Slovenia, Forestry	LN795901	LN795901	Piškur and Jurc 2017
<i>P. octolocularis</i>	HMAS140416	<i>Rubus biflorus</i>	China	MH128376	MG669119	Liu et al. 2018
<i>P. paucilocularis</i>	ZP-R318	<i>Rubus corchorifolius</i>	Guangxi, China	MK518874	MK518542	Zhao et al. 2021
<i>P. potentillae</i>	HGUP21034	<i>Acaena novae-zelandiae</i>	Tasmania, Australia	OL684827	OL684838	Sun et al. 2022
<i>P. potentillae</i>	HMJAU8609	<i>Potentilla chinensis</i>	China	MK296538	MK296520	Ji et al. 2019
<i>P. potentillae</i>	BJFCR 00961	<i>Potentilla chinensis</i>	Beijing, China	MN264720	MN264738	Liu et al. 2020

Species	Specimen No.	Host	Province, Country	GenBank Accession No.		References
				ITS	LSU	
<i>P. potentillae</i>	TJ-1F	<i>Potentilla chinensis</i>	China	PP272995	PP266810	Liu et al. 2020
<i>P. potentillae</i>	GMB4048	<i>Potentilla chinensis</i>	Guizhou, China	PQ472142	PQ456456	This study
<i>P. potentillae</i>	GMB4072	<i>Potentilla chinensis</i> .	Guizhou, China	PQ472143	PQ456457	This study
<i>P. rosae-multiflorae</i>	HGUP21158	<i>Rosa multiflora</i>	Guizhou, China	OR470059	OR528548	Sun et al. 2024
<i>P. rosae-multiflorae</i>	BIFCR 03454	<i>Rosa multiflora</i>	Jiangxi, China	MN264721	MN264739	Liu et al. 2020
<i>P. rosae-multiflorae</i>	HMAS94924	<i>Rosa multiflora</i>	Zhejiang, China	–	KU059175	Sun et al. 2022
<i>P. rosae-multiflorae</i>	GMB4044	<i>Rosa multiflora</i>	Guizhou, China	PQ472144	PQ456459	This study
<i>P. rosae-multiflorae</i>	GMB4073	<i>Rosa multiflora</i>	Guizhou, China	PQ472145	PQ456458	This study
<i>P. rosae-roxburghii</i>	HGUP21025*	<i>Rosa roxburghii</i>	Guizhou, China	OL684818	OL684831	Liu et al. 2018
<i>P. rosae-roxburghii</i>	GMB0104	<i>Rosa xanthina</i>	Guizhou, China	OQ067092	–	Wu et al. 2023
<i>Prosae-kwangtungensis</i>	HGUP21154*	<i>Rosa kwangtungensis</i>	Guizhou, China	OR470067	–	Sun et al. 2024
<i>P. rosae-cymosaii</i>	HGUP21147	<i>Rosa cymosa</i>	Guizhou, China	OR470062	OR528551	Sun et al. 2024
<i>P. rubi-idaei</i>	HMUT100470	<i>Rubus saxatilis</i>	Chongqing, China	OQ613354	OQ606768	–
<i>P. rubi-oldhami</i>	KSNUH1322	<i>Rubus pungens</i>	South Korea	ON180674	ON170371	Kim et al. 2022
<i>Phragmidium</i> sp.	HMJAU8613	–	Changchun, China	MK398297	MK398296	Ji et al. 2019
<i>P. parvifolius</i>	GMB4054*	<i>Rubus parvifolius</i>	Guizhou, China	PQ472140	PQ456454	This study
<i>P. parvifolius</i>	GMB4070	<i>Rubus parvifolius</i>	Guizhou, China	PQ472141	PQ456455	This study
<i>P. tormentillae</i>	GMB00114	<i>Potentilla simulatrix</i>	Guizhou, China	OQ067093	–	Wu et al. 2023
<i>P. tormentillae</i>	BPI 843392	<i>Duchesnea</i> sp.	USA: Maryland	–	DQ354553	Yun et al. 2011
<i>P. violaceum</i>	KRM 0035511	<i>Rubus</i> sp.	Germany	ON063390	ON063390	Bradshaw et al. 2023
<i>P. violaceum</i>	ZP-R1384	<i>Duchesnea indica</i>	Yunnan, China	MK518982	MK518680	Zhao et al. 2021
<i>P. yangii</i>	BJFCR 00338	<i>Rosa lichiangensis</i>	Beijing, China	MN264725	MN264743	Yang et al. 2015
<i>P. zangdongi</i>	BJFC: R02447*	<i>Rosa tibetica</i>	Tibet, China	MH128372	NG064490	Liu et al. 2018
<i>P. zhouquensis</i>	BJFCR01516	<i>Rosa omeiensis</i>	Yunnan, China	MN264728	MN264746	Yang et al. 2015
<i>P. zhouquensis</i>	BJFCR01529	<i>Rosa omeiensis</i>	Yunnan, China	MN264729	MN264747	Liu et al. 2020
<i>Trachyspora alchemillae</i>	BPI 843828	<i>Alchemilla vulgari</i>	Switzerland	DQ354550	DQ354550	Aime 2006

Notes: –: no data available; *: type specimens or strains.

Results

Phylogeny

In this study, 14 samples from twelve host plant species were collected in Guizhou Province. Through morphological and molecular systematic studies, a total of 6 species were identified, including three new species, one new host species, and two known species.

For the final phylogenetic analyses, taxa were selected based on their morphological and phylogenetic affinities, largely following the approach of Zhao et al. (2021). Both the RAXML and BYPP analyses produced similar overall tree topologies with no significant differences. The alignment includes 67 species (comprising one family: Phragmidiaceae, six genera: *Phragmidium*, *Gerwasia*, *Hamaspora*, *Kuehneola*, *Trachyspora*, and the outgroup genus *Gymnosporangium*) and contains 1245 characters, including gaps (ITS: 398 bp, LSU: 847 bp). Among these, three novel species have been identified (*Gerwasia amphidasydis* sp. nov., *Phragmidium parvifolius* sp. nov., and *P. coreanicola* sp. nov.). Additionally, one species (*G. rubi-setchuenensis*) was reported for the first time on *Rubus buergeri*.




Figure 1. RAxML tree of the family Phragmidiaceae based on rDNA ITS and LSU sequences. ML bootstrap supports ($\geq 75\%$) and Bayesian posterior probability (≥ 0.90) are indicated as ML/BYPP. The tree is rooted to *G. sabinae* and *G. asiaticum* (Yang et al. 2015; Aime et al. 2018). All species newly studied are indicated in red, with novel species highlighted in bold red. Type materials were highlighted in bold.

Taxonomy

Family Phragmidiaceae Corda, *Icon. Fung. (Prague)* 1: 6. 1837

Genus *Gerwasia* Racib. *Bull. Acad. Sci. Lett. Cracovie, Cl. Sci. Math. Nat. Sér. B, Sci. Nat.* 3: 270. 1909

Gerwasia amphidasidis Q. F. Zhang, Q. Z. Wu & Q. R. Li, *sp. nov.*

MycoBank No: 854995

Fig. 2

Type. CHINA • Guizhou Province, Zunyi City, Kuangkuoshui Nature Reserve (28°12'40"N, 107°10'22"E), 2227 m a.s.l., on leaves of *Rubus amphidasys*. 1 November 2022, Q. Z. Wu and Q. F. Zhang (holotype GMB4047, isotype KUN-HKAS144247);

Etymology. The epithet refers to the host species, *Rubus amphidasys* Focke ex Diels., from which the holotype was collected.

Description. *Spermogonia*, *Aecia*, and *Telia* not found. *Uredinia* 0.2–0.8 mm diam. produced on the abaxial leaf surface, scattered to gregarious, hypophyllous, covered by peridium, small, rounded, light yellow, or orange-yellow. Urediniospores 29–41 × 22–29 µm (av. = 34 × 26 µm, n = 30), globose to subglobose or ovoid, golden, yellow-brown, wall 1.2–2.5 µm thick at sides, hyaline, prominent sparsely echinulate, markings elongated longitudinally, 1.3–3.1 µm in distance, pore obscure, germ pores inconspicuous. Pedicel broken; paraphyses not seen.

Additional material examined. CHINA • Guizhou Province, Zunyi City, Kuangkuoshui Nature Reserve, (28°12'38"N, 107°10'21"E) 2214 m a.s.l., on the leaves of *Rubus amphidasys* (Rosaceae). 1 November 2022, Q. Z. Wu and Q. F. Zhang (GMB4076).

Notes. *Gerwasia amphidasidis* was the first species of *Gerwasia* described on *Rubus amphidasys*. Our phylogenetic analyses showed that *G. amphidasidis* formed a separate branch (Fig. 1). Morphologically, *G. amphidasidis* and *G. rubi* exhibit similar spines. Moreover, the difference between *G. amphidasidis* and *G. rubi* is that the former has bigger urediniospores (29–41 × 22–29 µm vs. 22–33 × 16–26 µm) (Ito 1950; Hiratsuka et al. 1992). *Gerwasia amphidasidis* and *G. guanganensis* have similar *uredinia* and *urediniospores*; however, *G. guanganensis* has longer spine distances compared to *G. amphidasidis* (4.0–6.0 µm vs. 1.3–3.1 µm) (Zhao et al. 2021). *Gerwasia amphidasidis* is distinguishable from *G. rubi-setchuenensis* by having larger urediniospores (29–41 × 22–29 µm vs. 18–29 × 15–22 µm) and a thinner wall (1.2–2.5 µm vs. 2.1–3.2 µm) (Sun et al. 2024).

Additionally, the LSU sequences of *Gerwasia amphidasidis* also differ from that of *G. rubi* with 93.74% similarity and from *G. rubi-setchuenensis* with 90.56% similarity. The ITS sequence for *Gerwasia rubi* is not available in the NCBI database, whereas the ITS sequence similarity between *Gerwasia amphidasidis* and *G. rubi-setchuenensis* is 98.56%.




Figure 2. *Gerwasia amphidasydis* (Holotype GMB4047) **A–D** host and its habitat **E–F** uredinia under a stereomicroscope **G–K** urediniospores. Scale bars: 1 mm (**D, F**); 0.5 mm (**E**); 10 μ m (**G–K**).

Genus *Phragmidium* Link, Mag. Gesell. Naturf. Freunde, Berlin 7: 30. 1816

***Phragmidium coreanicola* Q. F. Zhang, Q. Z. Wu & Q. R. Li, sp. nov.**

MycoBank No: 855005

Fig. 3

Type. CHINA • Guizhou Province, Guiyang City, Campus of Guizhou Medical University (26°22'48"N, 106°37'30"E), 1911 m a.s.l., on leaves of *Rubus coreanus* (Rosaceae), 7 October 2021, Q. Z. Wu (holotype GMB0101, isotype KUN-HKAS144249).

Etymology. The epithet refers to the host species, *Rubus coreanus* Miq. var. *coreanus*, from which the holotype was collected.

Description. *Spermogonia* and *Aecia* not found. *Uredinia* 0.1–0.7 mm diam., produced on the abaxial leaf surface, scattered to gregarious, hypophyllous, yellow spots, scattered, irregular patches. Urediniospores 20–29 × 14–25 µm (av. = 24 × 21 µm, n = 30), globose to subglobose or broadly elliptical to ellipsoidal, wall 0.8–2.1 µm thick (av. = 1.4 µm, n = 30), inconspicuous or smooth at the base; inclusions orange-yellow or pale-yellow; germ pores 2–3, sub-equatorial. *Telia* 0.1–0.9 mm diam., hypophyllous, dark brown to black, clustered or scattered, bacilliform. Teliospores 107–167 × 25–35 µm (av. = 134 × 30 µm, n = 30), cylindrical, 5–7 cells, often 6, reddish-brown to opaque, rounded at the apex, rounded or somewhat attenuate at the base, not or slightly constricted at the septum, pedicels sub-hyaline, persistent, 47–96 × 12–19 µm (av. = 68 × 16 µm, n = 30), with a swollen base that gradually shows orange-yellow contents towards the lower end. Pedicel broken; paraphyses not seen.

Additional material examined. CHINA • Guizhou Province, Qingzhen City (26°34'58"N, 106°28'28"E), 1972 m a.s.l., on leaves of *Rubus coreanus* (Rosaceae), 7 October 2021, Q.Z. Wu (GMB4071).

Notes. *Phragmidium coreanicola* formed a separate branch in our phylogenetic analyses (Fig. 1). Morphologically, *P. coreanicola* differs from *P. griseum* in having slightly wider urediniospores (14–25 µm vs. 13–21 µm) and larger teliospores (107–167 × 25–35 µm vs. 50–125 × 18–28 µm) (Liu et al. 2018). Additionally, *P. coreanicola* is reported on *Rubus coreanus*, whereas *P. griseum* was found on *Rubus crataegifolius*. *Phragmidium coreanicola* differs from *P. cibatum*, which is reported on *Rubus niveus*, by having bigger urediniospores (20–29 × 14–25 µm vs. 17–20 × 18–19 µm) and larger teliospores (107–167 × 25–35 µm vs. 80–100 × 20–30 µm) (Wei 1988; Hiratsuka et al. 1992; Liu et al. 2018). *Phragmidium coreanicola* has the same host species as *P. rubi-coreani* in Guiyang City. However, *P. coreanicola* has larger teliospores (107–167 × 25–35 µm vs. 29–74 × 14–37 µm) (Sun et al. 2022). The morphological comparison between *P. coreanicola* and *P. pauciloculare* shows that the *uredinia* of *P. coreanicola* are larger than those of *P. pauciloculare* (0.1–0.7 mm diam vs. 0.2–0.3 mm diam), and teliospores of *P. coreanicola* are also larger than those of *P. pauciloculare* (107–167 × 25–35 µm vs. 35–111 × 18–27 µm) (Wei 1988).

Furthermore, the ITS sequences of *P. coreanicola* and *P. griseum* exhibit significant differences, with a similarity of 87.33%. However, the LSU sequences of *P. coreanicola* and *P. griseum* have little variation, sharing a similarity of 99.38%.




Figure 3. *Phragmidium coreanicola* (Holotype, GMB0101) **A–C** host and its habitat **D–E** uredinia and telia **F, G** urediniospores **H–K** teliospores. Scale bars: 0.5 mm (**D, E**); 10 μ m (**F–K**).

***Phragmidium parvifolius* Q. F. Zhang, Q. Z. Wu & Q. R. Li, sp. nov.**

MycoBank No: 855006

Fig. 4

Type. CHINA • Guizhou Province, Guiyang City, Huaxi District (26°43'27.3"N, 106°67'14.4"E), 1,114 m a.s.l., on leaves of *Rubus parvifolius* (Rosaceae), 3 November 2022, Q. Z. Wu and Q. F. Zhang (holotype GMB4054, isotype KUN-HKAS144250).

Etymology. The epithet refers to the host species, *Rubus parvifolius* L., from which the holotype was collected.

Description. *Spermogonia*, *Aecia* and *Telia* not found. *Uredinia* 0.3–0.8 mm diam., produced on the abaxial leaf surface, scattered to gregarious, hypophyllous, rounded to irregular, powdery, orange, pulverulent, at first covered by the epidermis, later, not surrounded by host epidermis; Urediniospores 18–32 × 12–24 µm (av. = 22 × 18 µm, n = 30), globose, oblong, orange, wall 1.1–1.7 µm thick (av. = 1.3 µm, n = 30) at sides, regularly echinulate with stout spines; germ pores 2–3, supra-equatorial. Paraphyses 49–83 × 10–19 µm (av. = 65 × 15 µm, n = 30), hyaline, curved.

Additional material examined. CHINA • Guizhou Province, Guiyang City, Huaxi District (26°43'59.7"N, 106°67'66.5"E), 1114 m a.s.l., on leaves of *Rubus parvifolius* (Rosaceae), 3 November 2022, Q. Z. Wu and Q. F. Zhang (GMB4070).

Notes. Phylogenetically, *P. parvifolius* formed a sister branch to *P. barnardii* Plowr. & G. Winter (HGU21035), which was also reported on *Rubus parvifolius* (Fig. 1). Morphologically, *P. parvifolius* can be easily differentiated from *P. barnardii* by its larger urediniospores (18–32 × 12–24 µm vs. 16–19 × 15–18 µm) and larger paraphyses (49–83 × 10–19 µm vs. 26–39 × 10–13 µm) (Winter 1886; McTaggart et al. 2016; Sun et al. 2022). In terms of urediniospore size, *P. parvifolius* is similar to *P. griseum* (Dietel) Syd. However, *P. parvifolius* differs from *P. griseum* by having relatively larger paraphyses (49–83 × 10–19 µm vs. 34–70 × 7–16 µm) and by its host, *Rubus parvifolius* vs. *Rubus crataegifolius* (Wei 1988; Hiratsuka et al. 1992; Liu et al. 2018; Sun et al. 2024). Additionally, *P. parvifolius* differs from *P. pauciloculare* by its larger urediniospores (18–32 µm vs. 13–20 µm) (Wei 1988; Hiratsuka et al. 1992). *Phragmidium parvifolius* and *P. kanas* have the same urediniospores and paraphyses, but *P. parvifolius* has no teliospores, whereas *P. kanas* has them (Zhao et al. 2021).

Furthermore, the morphological comparison between *P. parvifolius* and *P. coreanicola* (this study) shows that the urediniospores of *P. parvifolius* are larger than those of *P. coreanicola* (18–32 µm vs. 20–29 µm), and *P. parvifolius* has no teliospores, whereas *P. coreanicola* has them. The ITS and LSU sequence similarities of *P. parvifolius* with *P. coreanicola* are 97.57% and 99.22%.

***Gerwasia rubi-setchuenensis* J.E. Sun, Yong Wang bis & K.D. Hyde, (2024)**

Fig. 5

Host. *Rubus buergeri* Miq.

Description. *Spermogonia*, *aecia*, and *telia* unknown. *Uredinia* 0.4–1.0 mm diam., hypophyllous, pulverulent, golden, scattered, irregular, surrounded by host epidermis. *Urediniospores* 24–30 × 19–25 µm (av. = 27.2 × 22.3 µm, n = 30), subglobose or fusiform, inclusions golden or bright yellow; wall 1.4–2.9 µm thick (av. = 2.0 µm, n = 30), colorless, irregularly elongated verrucae.




Figure 4. *Phragmidium parvifolius* (Holotype, GMB4054) **A–D** host and its habitat **E, F** uredinia under a stereomicroscope **G, H** urediniospores and paraphyses **I–L** urediniospores **M** paraphyses. Scale bars: 1 mm (**D**); 0.5 mm (**E, F**); 10 μ m (**G–M**).

Materials examined. CHINA • Guizhou Province, Zunyi City, Xishui County (28°49'38"N, 106°41'23"E), 1223 m a.s.l., on the leaves of *Rubus buergeri* Miq. (Rosaceae), 3 November, 2022, Q. Z. Wu and Q. F. Zhang (GMB4052); CHINA • Guizhou Province, Zunyi City, Xishui County (28°33'43"N, 106°24'3"E), 1997 m a.s.l., on *Rubus buergeri* (Rosaceae), 3 November 2022, Q. Z. Wu and Q. F. Zhang (GMB4075).




Figure 5. *Gerwasia rubi-setchuenensis* (GMB4052) **A–D** host and its habitat **E–F** uredinia under the stereomicroscope **G–K** urediniospores. Scale bars: 0.5 mm (**E–F**); 10 µm (**G–L**).

Notes. In the phylogram (Fig. 1), our collections (GMB4052 and GMB4075) clustered with *G. rubi-setchuenensis* (HGUP21168). The morphological characteristics of our specimen are consistent with the original description of *G. rubi-setchuenensis*, and the DNA sequence aligns with that of *G. rubi-setchuenensis* HGUP21168 (ITS 100%; LSU 99.67%) (Sun et al. 2024). The only difference observed between the descriptions and figure of *G. rubi-setchuenensis* in Sun et al. (2024) is the size of the urediniospores. The urediniospores of *G. rubi-setchuenensis* (GMB4075) are slightly wider than those of *G. rubi-setchuenensis* (HGUP21168) (19–25 µm vs. 15–22 µm). This study identifies *Rubus buergeri* as a new host for this fungus.

Discussion

The exploration of rust fungi in China began in the mid-19th century, and to date, over 1200 rust taxa have been documented (Zhuang et al. 1998, 2005, 2021; Zhao et al. 2022a, b; Sun et al. 2024). Molecular techniques have significantly advanced fungal species identification, but accurately identifying rust fungi remains challenging, necessitating a comprehensive approach incorporating morphology, host specificity, and phylogenetic analyses (Sun et al. 2024). In China, over 70 species of *Phragmidium* have been described, although numerous species remain without molecular data. According to our literature review, approximately 22 species of *Phragmidium* have been reported in Guizhou (Cummins 1931; Zhuang et al. 2012; Aime et al. 2018; Aime and McTaggart 2021; Zhao et al. 2021; Sun et al. 2022, 2024). These studies emphasize the critical role of integrating morphological data, host specificity, and phylogenetic insights for a comprehensive understanding and accurate identification of rust fungi.

In our investigation, three new species of Phragmidiaceae belonging to the genera *Gerwasia* and *Phragmidium* are introduced based on phylogenetic analysis of the ITS and LSU regions and morphological features. *amphidasys* sp. nov., *Phragmidium coreanicola* sp. nov., and *P. parvifolius* sp. nov. infected *Rubus amphidasys*, *Rubus coreanus*, and *Rubus parvifolius*, respectively. The host of the *P. coreanicola* is the same as that of *P. rubi-coreani*; however, *P. coreanicola* has larger teliospores (107–167 × 25–35 µm vs. 29–74 × 14–37 µm), and *P. rubi-coreani* possesses aeciospores (Sun et al. 2022). In addition, *Rubus buergeri* was identified as a new host plant for *Gerwasia rubi-setchuenensis*. Previously, it was known only on *Rubus setchuenensis* (Sun et al. 2024). At the same time, we discovered some samples of *Phragmidium rosae-multiflorae* Dietel, and *Phragmidium potentillae* (Pers.) P. Karst. Here, we provide their sequences.

The hosts of *Phragmidium* discussed in this study are mainly from the genus *Rubus* within the Rosaceae family, yet *P. coreanicola* represents a new species on a previously reported host, while *Gerwasia rubi-setchuenensis* originates from a different host. This demonstrates the host specificity and species diversity of rust fungi (Wei 1988; Zhuang et al. 2012; Yang et al. 2015; Liu et al. 2018). Investigations into the interaction between plant hosts and pathogens suggest that the variety of pathogenic fungi may stem from host-switching or cooperative coevolutionary processes. These findings prompt inquiries concerning the linkage between plant hosts and *Phragmidium*, as well as the evolutionary dynamics at play (Zhao et al. 2016; McTaggart et al. 2015, 2016). Addressing these inquiries necessitates further research into *Phragmidium* in the future.

Additional information

Conflict of interest

The authors have declared that no competing interests exist.

Ethical statement

No ethical statement was reported.

Funding

This research was supported by the Guizhou Province Ordinary Colleges and Universities Youth Science and Technology Talent Growth Project (2021)154; the Guizhou Medical University High Level Talent Launch Fund Project (2023-058); and the National Natural Science Foundation of China (31960005, 32000009).

Author contributions

Conceptualization: Qinfang Zhang, Qirui Li, and Lili Liu. Collection and morphological examinations: Qinfang Zhang, Qianzhen Wu, Yulin Ren. Molecular sequencing and phylogenetic analyses: Qinfang Zhang, Kamran Habib, Lili Liu. Specimen identification: Peng Zhao, Qirui Li. Plant identification: Qingde Long. Original draft preparation: Qinfang Zhang, Qirui Li. Review and editing, supervision: Muhammad Aijaz Ahmad, Peng Zhao, Yao Wang, Dexiang Tang, Xiangchun Shen, Qirui Li. All authors have read and agreed to the published version of the manuscript.

Author ORCIDs

Qinfang Zhang  <https://orcid.org/0009-0003-9408-4988>

Kamran Habib  <https://orcid.org/0000-0003-2572-0306>

Yao Wang  <https://orcid.org/0000-0002-1262-6700>

Muhammad Aljaz Ahmad  <https://orcid.org/0000-0003-2798-5500>

Yulin Ren  <https://orcid.org/0009-0003-9063-425X>

Xiangchun Shen  <https://orcid.org/0000-0002-4333-9106>

Qirui Li  <https://orcid.org/0000-0001-8735-2890>

Data availability

All the specimens are deposited in the Herbaria of Guizhou Medical University (GMB) and Kunming Institute of Botany, Chinese Academy of Sciences (KUN-HKAS). Sequences have been deposited in the GenBank. The alignment file can be obtained from the corresponding author.

References

- Aime MC (2006) Toward resolving family-level relationships in rust fungi (Uredinales). *Mycoscience* 47(3): 112–122. <https://doi.org/10.1007/S10267-006-0281-0>
- Aime MC, Bell CD, Wilson AW (2018) Deconstructing the evolutionary complexity between rust fungi (Pucciniales) and their plant hosts. *Studies in Mycology* 89: 143–152. <https://doi.org/10.1016/j.simyco.2018.02.002>
- Aime MC, McTaggart AR (2021) A higher-rank classification for rust fungi, with notes on genera. *Fungal Systematics and Evolution* 7(1): 21–47. <https://doi.org/10.3114/fuse.2021.07.02>

- Aime MC, McTaggart AR, Mondo SJ, Duplessis S (2017) Phylogenetics and phylogenomics of rust fungi. *Advances in Genetics* 100: 267–307. <https://doi.org/10.1016/bs.adgen.2017.09.011>
- Ali B, Sohail Y, Mumtaz AS, Berndt R (2017) *Phragmidium punjabense*, a new species of rust fungus on *Rosa brunonii* in the outer Himalayan ranges of Murree, Pakistan. *Nova Hedwigia* 105(3–4): 385–396. https://doi.org/10.1127/nova_hedwigia/2017/0418
- Arthur JC (1934) *Manual of the rust in United States and Canada*, Hafner Publishing Company, New York, USA.
- Bradshaw MJ, Carey J, Liu M, Bartholomew HP, Jurick WM II, Hambleton S, Hendricks D, Schnittler M, Scholler M (2023) Genetic time traveling: Sequencing old herbarium specimens, including the oldest herbarium specimen sequenced from kingdom Fungi, reveals the population structure of an agriculturally significant rust. *The New Phytologist* 237(4): 1463–1473. <https://doi.org/10.1111/nph.18622>
- Cummins GB (1931) *Phragmidium* species of North America: Differential teliospore and aecial characters. *Mycologia* 23(6): 433–445. <https://doi.org/10.1080/00275514.1931.12017054>
- Cummins GB, Hiratsuka Y (2003) *Illustrated genera of rust fungi*. 3rd Edn. American Phytopathological Society Press: 222–223.
- Dai FL (1979) *China fungus confluence*. Science Press, Beijing, China 1527.
- Dietel P (1905) Über die arten der Gattung *Phragmidium*. *Hedwigia* 44: 112–132.
- Dixon LJ, Castlebury LA, Aime MC, Glynn NC, Comstock JC (2018) Phylogenetic relationships of sugarcane rust fungi. *Mycological Progress* 9(4): 459–468. <https://doi.org/10.1007/s11557-009-0649-6>
- Hall TA (1999) BioEdit: A user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucleic Acids Symposium Series* 41: 95–98.
- Hiratsuka N, Sato S, Katsuya K, Kakishima M, Hiratsuka Y, Kaneko S, Ono Y, Sato T, Harada Y, Hiratsuka T, Nakayama K (1992) The rust flora of Japan. *Tsukuba Shuppankai* 41: 127–151.
- Hopple Jr JS, Vilgalys R (1999) Phylogenetic relationships in the mushroom genus *Coprinus* and dark-spored allies based on sequence data from the nuclear gene coding for the large ribosomal subunit RNA: Divergent domains, outgroups, and monophyly. *Molecular Phylogenetics and Evolution* 13(1): 1–19. <https://doi.org/10.1006/mpev.1999.0634>
- Huelsenbeck JP, Ronquist F (2001) MRBAYES: Bayesian inference of phylogenetic trees. *Bioinformatics (Oxford, England)* 17(8): 754–755. <https://doi.org/10.1093/bioinformatics/17.8.754>
- Ito S (1950) Basidiomycetes, no. 3. Uredinales-Pucciniaceae, Uredinales Imperfecti. *Mycological flora of Japan* 2.
- Ji JX, Li Z, Li Y, Kakishima M (2019) Two new species of *Pucciniastrum* producing dimorphic sori and spores from northeast of China. *Mycological Progress* 18(4): 529–540. <https://doi.org/10.1007/s11557-018-1460-z>
- Ji JX, Li Z, Li Y, Kakishima M (2020) Life cycle and taxonomy of *Chrysomyxa succinea* in China and phylogenetic positions of *Caeoma* species on *Rhododendron*. *Forest Pathology* 50(2): e12585. <https://doi.org/10.1111/efp.12585>
- Katoh K, Standley DM (2013) MAFFT multiple sequence alignment software version 7: Improvements in performance and usability. *Molecular Biology and Evolution* 30(4): 772–780. <https://doi.org/10.1093/molbev/mst010>
- Kim BR, Sim JY, Lee SH, Ryu JY, Lee JS, Choi YJ, Kim HJ, Jin HY, Kim YJ (2022) First report of *Phragmidium rubi-oldhami* causing rust disease on *Rubus pungens* in Korea. *New Disease Reports* 46(1): e12113. <https://doi.org/10.1002/ndr2.12112>

- Link HF (1816) Observations in ordines plantarum naturales. Magazinder Gesellschaft Naturforschenden Freunde Berlin 7: 25–45.
- Liu Y, Cao B, Tao S, Tian C, Liang Y (2018) *Phragmidium* species parasitizing species of Rosaceae in Tibet, China, with descriptions of three new species. Mycological Progress 17(8): 967–988. <https://doi.org/10.1007/s11557-018-1406-5>
- Liu Y, Liang YM, Ono Y (2020) Taxonomic revision of species of *Kuehneola* and *Phragmidium* on *Rosa*, including two new species from China. Mycologia 112(4): 742–752. <https://doi.org/10.1080/00275514.2020.1753426>
- McTaggart AR, Doungsa-Ard C, Geering ADW, Aime MC, Shivas RG (2015) A co-evolutionary relationship exists between *Endoraecium* (Pucciniales) and its *Acacia* hosts in Australia. Persoonia 35(1): 50–62. <https://doi.org/10.3767/003158515X687588>
- McTaggart AR, Shivas RG, Nest MA, Roux J, Wingfield BD, Wingfield MJ (2016) Host jumps shaped the diversity of extant rust fungi (Pucciniales). The New Phytologist 209(3): 1149–1158. <https://doi.org/10.1111/nph.13686>
- Miller MA, Pfeiffer W, Schwartz T (2010) Creating the CIPRES Science Gateway for inference of large phylogenetic trees. 2010 Gateway Computing Environments Workshop (GCE), New Orleans, Louisiana, 14 Nov 2010. IEEE, New York, 1–8. <https://doi.org/10.1109/GCE.2010.5676129>
- Pfunder M, Schürch S, Roy BA (2001) Sequence variation and geographic distribution of pseudoflower-forming rust fungi (*Uromyces pisi* s. lat.) on *Euphorbia cyparissias*. Mycological Research 105(1): 57–66. <https://doi.org/10.1017/S0953756200003208>
- Piškur B, Jurc D (2017) New findings of rusts (Pucciniales) on trees and other plants in Slovenia. Proceedings of the Zbornik Predavanj in Referatov 13.
- Rambaut A (2012) FigTree: Tree figure drawing tool 2006–2012, version 1.4.0. Edinburgh: Institute of Evolutionary Biology, University of Edinburgh.
- Rathnayaka AR, Tennakoon DS, Jones GE, Wanasinghe DN, Bhat DJ, Priyashantha AH, Stephenson SL, Tibpromma S, Karunaratna SC (2024) Significance of precise documentation of hosts and geospatial data of fungal collections, with an emphasis on plant-associated fungi. New Zealand Journal of Botany 63: 462–489. <https://doi.org/10.1080/0028825X.2024.2381734>
- Ronquist F, Nieves-Aldrey JL, Buffington ML, Liu Z, Liljeblad J, Nylander JA (2015) Phylogeny, evolution and classification of gall wasps: The plot thickens. PLoS One 10(5): e0123301. <https://doi.org/10.1371/journal.pone.0123301>
- Shen YM, Chung WH, Huang TC, Rodeva R, Hung TH (2017) Unveiling *Gymnosporangium corniforme*, *G. unicorn*, and *G. nitakayamense* sp. nov. in Taiwan. Mycoscience 59(3): 218–228. <https://doi.org/10.1016/j.myc.2017.11.003>
- Stamatakis A (2015) Using RAxML to infer phylogenies. Current Protocols in Bioinformatics 51(1): 6–14. <https://doi.org/10.1002/0471250953.bi0614s51>
- Sun JE, Fu L, Norphanphoun C, Chen XJ, Yu LF, Hyde KD, McKenzie EHC, Wang Y, Yang ZF, Liu FQ (2024) Study on the diversity of rust pathogens from different hosts in Guizhou Province, China. Mycosphere: Journal of Fungal Biology 15(1): 473–653. <https://doi.org/10.5943/mycosphere/15/1/4>
- Sun JE, Zhang Q, Luo WM, Yang YQ, An HM, Wang Y (2022) Four new *Phragmidium* (Phragmidiaceae, Pucciniomycetes) species from Rosaceae plants in Guizhou Province of China. MycoKeys 93: 193–213. <https://doi.org/10.3897/mycokeys.93.90861>
- Vilgalys R, Hester M (1990) Rapid genetic identification and mapping of enzymatically amplified ribosomal DNA from several *Cryptococcus* species. Journal of Bacteriology 172(8): 4238–4246. <https://doi.org/10.1128/jb.172.8.4238-4246.1990>

- Wahyuno D, Kakishima M, Ono Y (2001) Morphological analyses of urediniospores and teliospores in seven *Phragmidium* species parasitic on ornamental roses. *Mycoscience* 42(6): 519–533. <https://doi.org/10.1007/BF02460950>
- Wang L, Jia Y, Zhang X, Qin H (2015) Overview of higher plant diversity in China. *Shengwu Duoyangxing* 23(2): 217–224. <https://doi.org/10.17520/biods.2015049>
- Wei GR, Wang XJ (2011) Purification and storage of *Puccinia striiformis* f. sp. *Tritici*. *Henan Nongye Kexue* 40: 90–92.
- Wei SX (1988) A taxonomic study of the genus *Phragmidium* of China. *Junwu Xuebao* 1: 179–210.
- Winter G (1886) *Fungi australienses*. *Revue Mycologique Toulouse* 8: 207–213.
- Wu QZ, He MH, Liu TZ, Hu HM, Liu LL, Zhao P, Li QR (2023) Rust fungi on medicinal plants in Guizhou province with descriptions of three new species. *Journal of Fungi (Basel, Switzerland)* 9(9): 953. <https://doi.org/10.3390/jof9090953>
- Ya TY (2024) Species of the genus *Phragmidium* (Pucciniales) as parasites of roses (*Rosa*, Rosaceae) in Ukraine. *Ukrains'kyi Botanichnyi Zhurnal. Ukrains'ke Botanichne Tovarystvo* 81(3): 214–228. <https://doi.org/10.15407/ukrbotj81.03.214>
- Yang T, Chang W, Cao B, Tian C, Zhao L, Liang Y (2015) Two new *Phragmidium* species identified on *Rosa* plants native to China. *Phytotaxa* 217(2): 182–190. <https://doi.org/10.11646/phytotaxa.217.2.8>
- Yun HY, Minnis AM, Kim YH, Castlebury LA, Aime MC (2011) The rust genus *Frommeëlla* revisited: A later synonym of *Phragmidium* after all. *Mycologia* 103(6): 1451–1463. <https://doi.org/10.3852/11-120>
- Zhao P, Qi XH, Crous PW, Duan W, Cai L (2020) *Gymnosporangium* species on *Malus*: species delineation, diversity and host alternation. *Persoonia-Molecular Phylogeny and Evolution of Fungi* 45(1): 68–100. <https://doi.org/10.3767/persoonia.2020.45.03>
- Zhao P, Li Y, Liu F, Liang J, Zhou X, Cai L (2022a) Applying early divergent characters in higher rank taxonomy of Melampsorineae (Basidiomycota, Pucciniales). *Mycology* 14(1): 11–36. <https://doi.org/10.1080/21501203.2022.2089262>
- Zhao P, Liu F, Huang JE, Zhou X, Duan WJ, Cai L (2022b) *Cronartium* rust (Pucciniales, Cronartiaceae): Species delineation, diversity and host alternation. *Mycosphere: Journal of Fungal Biology* 13(1): 672–723. <https://doi.org/10.5943/mycosphere/13/1/7>
- Zhao P, Liu F, Li YM, Cai L (2016) Inferring phylogeny and speciation of *Gymnosporangium* species, and their coevolution with host plants. *Scientific Reports* 6(1): 29339. <https://doi.org/10.1038/srep29339>
- Zhao P, Zhang ZF, Hu DM, Tsui KM, Qi XH, Phurbu D, Gafforov Y, Cai L (2021) Contribution to rust flora in China I, tremendous diversity from natural reserves and parks. *Fungal Diversity* 110(1): 1–58. <https://doi.org/10.1007/s13225-021-00482-w>
- Zhuang JY, Wei SX, Wang YC (1998) *Flora Fungorum Sinicorum*. Vol 10. Uredinales (I). Science Press, Beijing, China.
- Zhuang JY, Wei SX, Wang YC (2003) *Flora Fungorum Sinicorum*. Vol 19. Uredinales (II). Science Press, Beijing, China.
- Zhuang JY, Wei SX, Wang YC (2005) *Flora Fungorum Sinicorum*. Vol 25. Uredinales (III). Science Press, Beijing, China.
- Zhuang JY, Wei SX, Wang YC (2012) *Flora Fungorum Sinicorum*. Vol 41. Uredinales (IV). Science Press, Beijing.
- Zhuang JY, Wei SX, Wang YC (2021) *Flora Fungorum Sinicorum*. Vol 62. Uredinales (V). Science Press, Beijing, China.