

# Based on minimal clinically important difference values, a moderate dose of tanezumab may be a better option for treating hip or knee osteoarthritis: a meta-analysis of randomized controlled trials

Di Zhao\*, Ming-hui Luo\*, Jian-ke Pan, Ling-feng Zeng, Gui-hong Liang, Yan-hong Han, Jun Liu\*<sup>ID</sup> and Wei-yi Yang<sup>†</sup>

## Abstract

**Background:** Tanezumab is a nerve growth factor monoclonal antibody that may regulate pain in hip or knee osteoarthritis (OA). This meta-analysis was performed to evaluate the efficacy and safety of low and moderate doses of tanezumab in treating hip or knee OA.

**Methods:** PubMed, EMBASE, the Cochrane Library, and Web of Science were comprehensively searched for clinical trials published before 1 May 2021. Patients were assessed *via* efficacy and safety outcomes.

**Results:** Twelve randomized controlled trials including 6022 patients were identified. Both low and moderate doses of tanezumab significantly improved efficacy outcomes. However, only the point estimates (mean difference, MD) of moderate-dose tanezumab significantly exceeded the minimal clinically important differences (MCIDs). There were no significant differences in the incidence of treatment-related adverse events (AEs), withdrawals due to AEs, serious AEs, and total joint replacement between the tanezumab and placebo groups, whereas the incidence of AEs was higher in the tanezumab group (relative risk, RR = 1.10; 95% confidence interval, 95% CI = 1.04–1.17). The incidence of rapidly progressive OA was significantly higher in the combined low- and moderate-dose tanezumab groups than in the placebo group (RR = 5.01; 95% CI = 1.17–21.33). Furthermore, both low and moderate doses of tanezumab significantly increased the incidence of abnormal peripheral sensation (RR = 1.99, 95% CI = 1.21–3.28; RR = 2.64, 95% CI = 1.91–3.67, respectively). Compared with nonsteroidal anti-inflammatory drugs (NSAIDs) and opioids, tanezumab showed significantly improved efficacy outcomes ( $p < 0.05$ ). However, the point estimates (MD) of tanezumab were not greater than the MCID. Pooled analysis showed no significant differences between tanezumab and NSAIDs and opioids in safety outcomes ( $p > 0.05$ ).

**Conclusion:** Tanezumab is efficacious in patients with hip or knee OA. Tanezumab is relatively well tolerated and safe but increases the incidence of AEs and reversible abnormal peripheral sensation. Additional studies on the occurrence of rapidly progressive OA are needed. A moderate dose of tanezumab may maximize the benefits for hip or knee OA.

**Keywords:** anti-NGF antibody, meta-analysis, OA, osteoarthritis, randomized controlled trials, tanezumab

Received: 3 August 2021; revised manuscript accepted: 1 December 2021.

*Ther Adv Musculoskel Dis*

2022, Vol. 14: 1–15

DOI: 10.1177/  
1759720X211067639

© The Author(s), 2022.  
Article reuse guidelines:  
sagepub.com/journals-  
permissions

Correspondence to:

**Jun Liu**  
Bone and Joint Research  
Team of Degeneration  
and Injury, Guangdong  
Provincial Academy of  
Chinese Medical Sciences,  
Guangzhou 510120, China

Guangdong Second  
Traditional Chinese  
Medicine Hospital  
(Guangdong Province  
Engineering Technology  
Research Institute of  
Traditional Chinese  
Medicine), Guangzhou,  
China.

The Fifth Clinical Medical  
College of Guangzhou  
University of Chinese  
Medicine, Guangzhou,  
China

[liujun3040@126.com](mailto:liujun3040@126.com)

**Wei-yi Yang**  
Bone and Joint Research  
Team of Degeneration  
and Injury, Guangdong  
Provincial Academy of  
Chinese Medical Sciences,  
Guangzhou 510120, China

Department of  
Orthopedics, The Second  
Affiliated Hospital of  
Guangzhou University  
of Chinese Medicine,  
Guangzhou, China  
[czyangwy@163.com](mailto:czyangwy@163.com)

**Di Zhao**  
The Second Clinical School  
of Guangzhou University  
of Chinese Medicine,  
Guangzhou, China

Bone and Joint Research  
Team of Degeneration  
and Injury, Guangdong  
Provincial Academy of  
Chinese Medical Sciences,  
Guangzhou, China

**Ming-hui Luo**  
**Jian-ke Pan**  
**Ling-feng Zeng**  
**Gui-hong Liang**  
**Yan-hong Han**

Bone and Joint Research  
Team of Degeneration  
and Injury, Guangdong  
Provincial Academy of  
Chinese Medical Sciences,  
Guangzhou, China

Department of

Orthopedics, The Second Affiliated Hospital of Guangzhou University of Chinese Medicine, Guangzhou, China

\*Di Zhao and Ming-hui Luo contributed equally.

†Jun Liu and Wei-yi Yang contributed equally.

## Introduction

Osteoarthritis (OA) is a common disease worldwide. The pain and impairment in joint function caused by OA can lead to a decline in the quality of life of patients and even disability.<sup>1,2</sup> The effective management of pain due to OA is a major concern for clinicians. Although the current recommendations suggest a combination of drug and nondrug treatments,<sup>3,4</sup> many patients still suffer from inadequate pain relief. Moreover, currently used analgesics, including nonsteroidal anti-inflammatory drugs (NSAIDs) and opioids, are poorly tolerated by many patients, and there are concerns about the safety and efficacy of the long-term use of these drugs.<sup>5,6</sup> Therefore, drugs that are more effective in relieving pain than are existing treatments and are well tolerated by patients urgently need to be developed.

With the emergence of in-depth research on pain mechanisms, new potential targets, especially the regulation of nociceptive and neurological pathways, have become the focus of recent drug development. Nerve growth factor (NGF) is a neurotrophic factor. After tissue injury or inflammation, NGF is released and binds to tropomyosin receptor kinase (Trk) A, which can lead to central sensitization,<sup>7</sup> induce the expression of peripheral and central pain-related substances, and make adjacent pain-sensing neurons sensitive to inflammation, thereby mediating pain.<sup>8-10</sup> NGF is also considered a pathological product of OA or an important pathogenic factor involved in the process of OA pain. The content of NGF in the synovial fluid of OA patients is higher than that in the synovial fluid of normal persons,<sup>11</sup> and NGF is also expressed in the synovial membrane of OA patients.<sup>12</sup> Furthermore, NGF expression increases significantly in traumatic and inflammatory sites.<sup>13</sup> The inhibition of NGF binding to its receptor and the downregulation of NGF expression can significantly reduce pain,<sup>14</sup> relieve OA symptoms, and improve limb function,<sup>15,16</sup> thereby relieving OA pain.

Tanezumab is an IgG2 NGF monoclonal antibody that blocks NGF from interacting with its receptors, thereby blocking signals from nociceptive neurons responsible for pain.<sup>17,18</sup> In 2010, tanezumab was tested in phase II human trials. The monoclonal antibodies against NGF currently used in clinical research include not only tanezumab but also fulranumab and fasinumab. However, there are few studies on fulranumab and fasinumab, and most of them are still in phase

II trials, while tanezumab is the most studied monoclonal antibody to inhibit NGF. Studies on tanezumab in the treatment of hip or knee OA have been reported in meta-analyses, but these results were based on only statistical significant differences, and reporting only these results might be misleading in terms of clinical decision making. Thus, we carried out a specific analysis based on minimal clinically important difference (MCID) values. In addition, because there is a safety issue, the US Food and Drug Administration (FDA) has mandated that tanezumab and NSAIDs not be used in combination and called for more research on tanezumab at lower doses. Therefore, this study aimed to evaluate the efficacy and safety of lower doses of tanezumab monotherapy for OA of the hip or knee.

## Methods

The systematic review was designed and reported based on the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) recommendations.<sup>19</sup>

### *Search strategy and selection criteria*

PubMed, EMBASE, the Cochrane Library, and Web of Science were comprehensively searched for English studies using the following keywords: 'nerve growth factor antibody', 'NGF', 'anti-NGF', 'tanezumab', 'OA', or 'osteoarthritis'. Studies published from the inception of the database to 1 May 2021 were retrieved. The references of the articles included were also searched, and each article was manually cross-checked. Disagreements among the reviewers were resolved by consensus.

### *Study selection*

With respect to the inclusion criteria, the two researchers independently screened the titles, abstracts, and full texts to determine which studies to include in this review. Any disagreements were discussed and resolved by a third reviewer.

### *Eligibility criteria*

The inclusion criteria were as follows: (1) studies in which the participants were adult patients diagnosed with hip or knee OA; (2) studies in which the experimental group was treated with low or moderate doses of tanezumab monotherapy, and the control group was treated with a placebo or an

active comparator; (3) studies in which at least the following outcome measures were included – Western Ontario and McMaster Universities Osteoarthritis Index (WOMAC) pain score (scale of 0–10), WOMAC physical function score (scale of 0–10), patient’s global assessment (PGA; 5-point Likert-type scale), adverse events (AEs), treatment-related AEs, serious adverse events (SAEs), withdrawals due to AEs, rapidly progressive OA (RPOA), and total joint replacement and abnormal peripheral sensation; and (4) studies that had a randomized controlled trial (RCT) design.

The exclusion criteria were as follows: (1) studies in which the experimental group was treated with tanezumab combined with other active treatments, and (2) studies with incomplete data.

#### *Data extraction*

The data were extracted independently by two researchers, and the variables included first author’s name, year of publication, phase of study, patient age, patient sex, sample size, Kellgren–Lawrence (K-L) grade, outcome measures, follow-up time, and type and dose of intervention. For studies by the same author, we selected the study with the longer follow-up time and complete data for the variables of interest. Disagreements were resolved through discussion, and if necessary, a third investigator made the final decision.

#### *Risk of bias assessment*

The quality of the methodology included in the study was assessed with the Cochrane Collaboration risk of bias assessment tool,<sup>20</sup> which evaluates the following domains: random sequence generation, allocation concealment, blinding of the participants and outcome assessors, incomplete outcome data, selective reporting, and other bias. Each component was considered to have a low, unclear, or high risk of bias.

#### *Statistical analysis*

We used Review Manager (version 5.3; Cochrane Collaboration, Oxford, UK) for statistical analysis of the data and Stata (version 14.0; StataCorp, College Station, Texas, USA) to assess publication bias. The mean baseline-to-end point changes and their standard deviations (SDs) were

obtained. For the efficacy outcomes, including WOMAC pain, WOMAC physical function, and PGA scores, we calculated the mean differences (MDs) and 95% confidence intervals (CIs). For the safety outcomes, we calculated the relative risk (RR) values and 95% CIs. If only the standard error was reported, the SD was calculated based on the reported sample size and standard error. When the mean, standard error, or SD were not provided in an article, we extracted the values from charts or graphs, as needed. The Cochran *Q* test and *I*<sup>2</sup> statistic were used to determine the magnitude of heterogeneity among studies. Heterogeneity levels of 0–25% were considered low, levels of 26–49% were considered moderate, and levels  $\geq 50\%$  were considered high. If the level of heterogeneity was  $\geq 50\%$ , a random-effects model was used; otherwise, a fixed-effects model was used. Publication bias of the outcome measures was assessed with Egger’s regression tests if the number of studies was  $> 10$ . Subgroup analysis for tanezumab at different doses was performed;  $p < 0.05$  indicated a significant difference. To assess whether the difference was clinically meaningful, the differences in the results reported by patients were directly compared with the MCIDs reported. According to previous reports,<sup>21</sup> the MCID of the WOMAC pain score (scale of 0–10) is 0.97, and the MCID of the WOMAC physical function score (scale of 0–10) is 0.93. However, to our knowledge, there are no well-established MCID criteria for PGA (5-point Likert-type scale). Therefore, only statistically significant differences are reported. When outcomes are statistically significant, and the point estimates (MD) are greater than the MCID, the result is considered to reflect a clinical effect.


## **Results**

#### *Selection of the included studies*

Through the literature search, a total of 368 studies were retrieved. After the titles and abstracts were screened, 30 studies remained, and then, the full texts were read. According to the inclusion and exclusion criteria, 11 studies were finally selected. See Figure 1 for details.

#### *Characteristics of the included studies*

A total of 12 RCTs, including 6022 patients, were included in the review. One of the studies<sup>22</sup> involved two RCTs. According to the K-L grade, the degree of hip or knee OA ranged from



**Figure 1.** Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) study selection flow diagram.

moderate to severe. Two studies<sup>23,24</sup> were phase II trials, and nine studies<sup>22,25–32</sup> were phase III trials. The characteristics of the included clinical trials are summarized in Table 1. The doses of tanezumab in phase II trials were 10, 25, 50, 100, and 200  $\mu\text{g}/\text{kg}$ , while in phase III trials, the doses of tanezumab were 2.5, 5, and 10 mg. However, the dose–response analysis and pharmacokinetic/pharmacodynamic study of tanezumab showed that sex and the dose regimen of adjusting an individual’s body weight negligibly reduced the variability in systemic exposure compared with that predicted with a fixed-dose regimen.<sup>33</sup> Therefore, the data from the phase II and phase III trials were combined in this study. At present, according to the FDA regulations, high doses of tanezumab cannot be used for experimental research; thus, we studied only low and moderate doses of tanezumab. The combination of the 10 and 25  $\mu\text{g}/\text{kg}$  doses in the phase II trials and the 2.5 mg doses in the phase III trials was studied and considered the low-dose group; the 50  $\mu\text{g}/\text{kg}$

doses in the phase II trials and 5 mg doses in the phase III trials were combined and defined as the moderate-dose group. In all subsequent analyses, we conducted a subgroup analysis based on low and moderate doses.

There were 10 placebo-controlled studies and 4 active-controlled studies. The active-controlled agents included oxycodone, celecoxib, and naproxen. However, in each of the active-controlled RCTs, low doses of tanezumab were not reported, so we analyzed only moderate doses of tanezumab.

#### *Risk of bias assessment*


The quality of all of the studies was relatively high. Although all studies were randomized controlled studies, five studies<sup>22,26,28,29,32</sup> did not report the specific randomization methods used. Two studies<sup>22,26</sup> did not clearly indicate whether the participants and outcome assessors were

**Table 1.** Characteristics of the included studies.

Study	Interventions	No. of subjects (sex: female)	Mean age (years)	K-L grade (mainly affected)	Efficacy outcomes (endpoints)	Safety outcomes (endpoints)
Berenbaum <i>et al.</i> <sup>25</sup> (phase III)	Tanezumab 2.5 mg Tanezumab 5 mg Placebo	849 (69.1%)	64.9	II, III, IV	WOMAC pain, WOMAC physical function, PGA 24 weeks	Treatment-related AEs, withdrawals due to AEs, SAEs, RPOA, TJR, abnormal peripheral sensation 48 weeks
Schnitzer <i>et al.</i> <sup>30</sup> (phase III)	Tanezumab 2.5 mg Tanezumab 5 mg Placebo	696 (65.1%)	60.9	II, III, IV	WOMAC pain, WOMAC physical function, PGA 16 weeks	Treatment-related AEs, withdrawals due to AEs, SAEs, RPOA, TJR, abnormal peripheral sensation 40 weeks
Birbara <i>et al.</i> <sup>26</sup> (phase III)	Tanezumab 2.5 mg Tanezumab 5 mg Placebo	209 (62.7%)	60.9	II, III, IV	WOMAC pain, WOMAC physical function, PGA 16 weeks	Withdrawals due to AEs, SAEs, abnormal peripheral sensation 24 weeks
Ekman <i>et al.</i> (A) <sup>22</sup> (phase III)	Tanezumab 5 mg Placebo Naproxen 500 mg bid	619 (60.0%)	61.1	II, III, IV	WOMAC pain, WOMAC physical function, PGA 16 weeks	Withdrawals due to AEs, SAEs, abnormal peripheral sensation 24 weeks
Ekman <i>et al.</i> (A) <sup>22</sup> (phase III)	Tanezumab 5 mg Placebo Naproxen 500 mg bid	627 (64.8%)	60.0	II, III, IV	WOMAC pain, WOMAC physical function, PGA 16 weeks	Withdrawals due to AEs, SAEs, abnormal peripheral sensation 24 weeks
Brown <i>et al.</i> <sup>27</sup> (phase III)	Tanezumab 5 mg Placebo	145 (57.2%)	57.1	II, III, IV	WOMAC pain, WOMAC physical function, PGA 16 weeks	Withdrawals due to AEs, SAEs, abnormal peripheral sensation 24 weeks
Brown <i>et al.</i> <sup>29</sup> (phase III)	Tanezumab 2.5 mg Tanezumab 5 mg Placebo	464 (63.8%)	62.1	II, III, IV	WOMAC pain, WOMAC physical function, PGA 16 weeks	SAEs, TJR, abnormal peripheral sensation 32 weeks
Spierings <i>et al.</i> <sup>32</sup> (phase III)	Tanezumab 5 mg Placebo Oxycodone 10–40 mg q12h	460 (62.4%)	57.5	II, III, IV	WOMAC pain, WOMAC physical function, PGA 8 weeks	Treatment-related AEs, withdrawals due to AEs, SAEs, abnormal peripheral sensation 16 weeks
Brown <i>et al.</i> <sup>28</sup> (phase III)	Tanezumab 2.5 mg Tanezumab 5 mg Placebo	516 (60.9%)	61.7	II, III, IV	WOMAC pain, WOMAC physical function, PGA 16 weeks	Withdrawals due to AEs, SAEs, TJR, abnormal peripheral sensation 32 weeks
Nagashima <i>et al.</i> <sup>24</sup> (phase II)	Tanezumab 10 µg/kg Tanezumab 25 µg/kg Tanezumab 50 µg/kg Placebo	61 (65.6%)	59.2	II, III	WOMAC pain, WOMAC physical function 8 weeks	Treatment-related AEs, withdrawals due to AEs, SAEs, abnormal peripheral sensation 13–17 weeks
Lane <i>et al.</i> <sup>23</sup> (phase II)	Tanezumab 10 µg/kg Tanezumab 25 µg/kg Tanezumab 50 µg/kg Placebo	296 (60.1%)	59.2	II, III, IV	WOMAC pain, WOMAC physical function 16 weeks	Treatment-related AEs, SAEs, abnormal peripheral sensation 26 weeks
Schnitzer <i>et al.</i> <sup>31</sup> (phase III)	Tanezumab 5 mg Naproxen 500 mg bid or celecoxib 100 mg bid	1080 (72.2%)	61.6	II, III, IV	WOMAC pain, WOMAC physical function 16 weeks	Withdrawals due to AEs, SAEs, RPOA, TJR, abnormal peripheral sensation 64 weeks

AEs, adverse events; K-L grade, Kellgren–Lawrence grade for classifying OA severity, ranging from 0 (no OA) to IV (severe OA); PGA, patient's global assessment; RPOA, rapidly progressive osteoarthritis; SAEs, serious adverse events; TJR, total joint replacement; WOMAC, Western Ontario and McMaster Universities Osteoarthritis Index.





**Figure 2.** Forest plot of WOMAC pain scores detailing mean differences and 95% CIs comparing tanezumab with placebo. CIs, confidence intervals; IV, inverse variance; WOMAC, Western Ontario and McMaster Universities Osteoarthritis Index.


blinded, so the risk of bias for this domain was judged as unclear. Allocation concealment was implemented in all studies. The complete results for the risk of bias assessment are presented in Supplemental Appendix Figures A1 and A2. Egger’s test suggested that there was no significant publication bias for the WOMAC pain score ( $p=0.627$ ), WOMAC physical score ( $p=0.589$ ), AEs ( $p=0.435$ ), SAEs ( $p=0.666$ ), or abnormal peripheral sensation ( $p=0.672$ ).

*Placebo-controlled trials*

**Efficacy. WOMAC pain score:** A total of 11 RCTs (4305 patients) reported WOMAC pain scores. The results of the pooled analysis showed that tanezumab significantly improved WOMAC pain scores compared with placebo (MD = -0.98, 95% CI = -1.15 to -0.81,  $p < 0.00001$ ,  $I^2 = 0\%$ ). In the subgroup analysis, 7 studies were included in the low-dose group, and 11 studies were included in the moderate-dose group. The results of the pooled analysis for the low-dose group showed that tanezumab significantly relieved patients’ pain compared with placebo (MD = -0.90, 95%

CI = -1.19 to -0.61,  $p < 0.00001$ ,  $I^2 = 35\%$ ). Similarly, moderate doses of tanezumab also significantly relieved pain in patients compared with placebo (MD = -1.02, 95% CI = -1.24 to -0.81,  $p < 0.00001$ ,  $I^2 = 0\%$ ). There were no significant differences between the two treatment groups (combined low-dose group versus combined moderate-dose group;  $p = 0.50$ ,  $I^2 = 0\%$ ; Figure 2). In comparison with the MCID (0.97), the point estimates (MD) of the low-dose tanezumab group were not greater than the MCID, but the point estimates (MD) of the moderate-dose tanezumab group significantly exceeded the MCID.

**WOMAC physical score:** A total of 11 RCTs (4305 patients) reported WOMAC physical scores. The results of the pooled analysis showed that tanezumab significantly improved WOMAC physical scores compared with placebo (MD = -1.04, 95% CI = -1.20 to -0.87,  $p < 0.00001$ ,  $I^2 = 0\%$ ). In the subgroup analysis, 7 studies were included in the low-dose group, and 11 studies were included in the moderate-dose group. The pooled estimate of the WOMAC physical score




**Figure 3.** Forest plot of WOMAC physical scores detailing mean differences and 95% CIs comparing tanezumab with placebo. CIs, confidence intervals; IV, inverse variance; WOMAC, Western Ontario and McMaster Universities Osteoarthritis Index.

demonstrated significant differences in favor of low doses of tanezumab (MD = -0.93, 95% CI = -1.21 to -0.65,  $p < 0.00001$ ,  $I^2 = 29\%$ ). Similarly, the moderate-dose group had significantly better WOMAC physical scores than the placebo group (MD = -1.09, 95% CI = -1.30 to -0.89,  $p < 0.00001$ ,  $I^2 = 0\%$ ). There were no significant differences between the two treatment groups ( $p = 0.35$ ,  $I^2 = 0\%$ ; Figure 3). In comparison with the MCID (0.93), the point estimates (MD) of the low-dose tanezumab group were not greater than the MCID, but the point estimates (MD) of the moderate-dose tanezumab group significantly exceeded the MCID.

**PGA:** A total of nine RCTs (3951 patients) reported PGA results. The results of the pooled analysis showed that tanezumab significantly improved PGA compared with the placebo (MD = -0.29, 95% CI = -0.35 to -0.23,  $p < 0.00001$ ,  $I^2 = 0\%$ ). In the subgroup analysis, five studies were included in the low-dose group, and nine studies were included in the moderate-dose group. Pooled analysis showed that low doses of tanezumab resulted in greater

improvements in PGA than the placebo (MD = -0.24, 95% CI = -0.34 to -0.13,  $p < 0.00001$ ,  $I^2 = 0\%$ ; Figure 4). Similarly, the moderate-dose group had significantly better PGA results than the placebo group (MD = -0.31, 95% CI = -0.39 to -0.24,  $p < 0.00001$ ,  $I^2 = 0\%$ ). There were no significant differences between the two treatment groups ( $p = 0.24$ ,  $I^2 = 27\%$ ).

**Safety. AEs:** A total of 11 RCTs (4372 patients) reported AEs. The results of the pooled analysis showed that the incidence of AEs was significantly higher in the tanezumab group than in the placebo group (RR = 1.10, 95% CI = 1.04–1.17,  $p = 0.0007$ ,  $I^2 = 1\%$ ). In the subgroup analysis, 7 studies were included in the low-dose group, and 11 studies were included in the moderate-dose group. The pooled results for the low-dose group showed that the incidence of AEs was significantly higher in the tanezumab group than in the placebo group (RR = 1.10, 95% CI = 1.01–1.21,  $p = 0.03$ ,  $I^2 = 38\%$ ). Similarly, moderate doses of tanezumab also led to a higher risk of AEs than placebo (RR = 1.10, 95% CI = 1.03–1.19,  $p = 0.008$ ,  $I^2 = 0\%$ ; Table 2). The RR values of the



**Figure 4.** Forest plot of patient's global assessment detailing mean differences and 95% CIs comparing tanezumab with placebo. CIs, confidence intervals; IV, inverse variance.

two treatment groups (low-dose group *versus* moderate-dose group) were the same, without a significant difference ( $p=0.99$ ,  $I^2=0\%$ ).

**Treatment-related AEs:** A total of five RCTs (2204 patients) reported treatment-related AEs. The results of the pooled analysis showed no significant difference in the incidence of treatment-related AEs between the tanezumab and placebo groups (RR = 1.15, 95% CI = 0.92–1.43,  $p=0.22$ ,  $I^2=26\%$ ). In the subgroup analysis, four studies were included in the low-dose group, and five studies were included in the moderate-dose group. Pooled analysis showed no significant difference between the low-dose tanezumab and placebo groups in terms of treatment-related AEs (RR = 1.17, 95% CI = 0.83–1.63,  $p=0.37$ ,  $I^2=46\%$ ). Similarly, there were no differences between the moderate-dose group and placebo group in treatment-related AEs (RR = 1.13, 95% CI = 0.85–1.52;  $p=0.40$ ,  $I^2=24\%$ ; Table 2). The RR values of the two treatment groups (low-dose group *versus* moderate-dose group) were similar, without a significant difference ( $p=0.90$ ,  $I^2=0\%$ ).

**Withdrawals due to AEs:** A total of nine RCTs (3612 patients) reported withdrawals due to AEs. The results of the pooled analysis showed that

there was no significant difference in the incidence of withdrawals due to AEs between the tanezumab and placebo groups (RR = 1.19, 95% CI = 0.74–1.91,  $p=0.47$ ,  $I^2=0\%$ ). In the subgroup analysis, five studies were included in the low-dose group, and nine studies were included in the moderate-dose group. The combined results for the low-dose group showed no significant difference in the incidence of withdrawals due to AEs between the tanezumab and placebo groups (RR = 1.20, 95% CI = 0.42–3.37,  $p=0.74$ ,  $I^2=0\%$ ). Similarly, there was no significant difference in the incidence of withdrawals due to AEs between the moderate-dose tanezumab and placebo groups (RR = 1.19, 95% CI = 0.70–2.02,  $p=0.52$ ,  $I^2=0\%$ ; Table 2). The RR values of the two treatment groups were similar, without a significant difference ( $p=0.99$ ,  $I^2=0\%$ ).

**SAEs:** A total of 11 RCTs (4372 patients) reported SAEs. The results of the pooled analysis showed no significant difference in the incidence of SAEs between the tanezumab and placebo groups (RR = 1.21, 95% CI = 0.87–1.66,  $p=0.25$ ,  $I^2=0\%$ ). In the subgroup analysis, 7 studies were included in the low-dose group, and 11 studies were included in the moderate-dose group. Pooled analysis showed no significant difference



**Table 2.** The safety outcomes of placebo-controlled trials.

Outcome measures	Low dose				Moderate dose				Combined results
	Outcome (tanezumab/placebo)	RR	95% CI	<i>p</i> value	Outcome (tanezumab/placebo)	RR	95% CI	<i>p</i> value	RR (95% CI)
Adverse events	62.0%/56.2%	1.10	[1.01–1.21]	0.03	56.2%/49.2%	1.10	[1.04–1.17]	0.008	1.10 (1.04–1.17)
Treatment-related adverse events	15.8%/13.6%	1.17	[0.83–1.63]	0.37	15.0%/13.1%	1.13	[0.85–1.52]	0.40	1.15 (0.92–1.43)
Withdrawals due to adverse events	1.5%/1.3%	1.20	[0.42–3.37]	0.74	2.4%/2.3%	1.19	[0.70–2.02]	0.52	1.19 (0.74–1.91)
Serious adverse events	4.9%/3.6%	1.30	[0.77–2.18]	0.33	4.0%/3.1%	1.15	[0.76–1.74]	0.50	1.21 (0.87–1.66)
Rapidly progressive OA	1.8%/0%	5.02	[0.65–9.02]	0.12	1.7%/0%	4.99	[0.64–8.78]	0.12	5.01 (1.17–21.33)
Total joint replacement	3.9%/3.3%	1.18	[0.64–2.16]	0.60	4.7%/3.1%	1.49	[0.82–2.72]	0.19	1.33 (0.87–2.04)
Abnormal peripheral sensation	7.4%/3.6%	2.00	[1.21–3.29]	0.006	8.9%/3.6%	2.64	[1.90–3.66]	<0.00001	2.42 (1.84–3.18)


CI, confidence interval; OA, osteoarthritis; RR, relative risk.

between the low-dose tanezumab and placebo groups in SAEs (RR=1.30, 95% CI=0.77–2.18,  $p=0.33$ ,  $I^2=0\%$ ). Similarly, the combined results for the moderate-dose group showed no significant difference in the incidence of SAEs between the tanezumab and placebo groups (RR=1.15, 95% CI=0.87–1.66,  $p=0.50$ ,  $I^2=16\%$ ; Table 2). The RR values were similar for the two treatment groups ( $p=0.72$ ,  $I^2=0\%$ ).

**RPOA:** A total of two RCTs (1545 patients) reported RPOA results. The results of the pooled analysis showed that the incidence of RPOA was significantly higher in the tanezumab group than in the placebo group (RR=5.01, 95% CI=1.17–21.33,  $p=0.03$ ,  $I^2=0\%$ ). In the subgroup analysis, two studies were included in the low-dose group, and two studies were included in the moderate-dose group. The pooled estimate of RPOA demonstrated that there were no significant differences between the low-dose tanezumab and placebo groups (RR=5.02, 95% CI=0.65–39.02,  $p=0.12$ ,  $I^2=0\%$ ). Similarly, there were no significant differences between the moderate-dose group and placebo group in treatment-related AEs (RR=4.99, 95% CI=0.64–38.78,  $p=0.12$ ,  $I^2=0\%$ ). The RR values of the two treatment groups were similar ( $p=1.00$ ,  $I^2=0\%$ ; see Figure 5 and Table 2).

**Total joint replacement:** A total of four RCTs (2525 patients) reported total joint replacement results. The results of the pooled analysis showed that there was no significant difference in the incidence of total joint replacement between the tanezumab and placebo groups (RR=1.33, 95% CI=0.87–2.04,  $p=0.19$ ,  $I^2=0\%$ ). In the subgroup analysis, four studies were included in the low-dose group, and four studies were included in the moderate-dose group. The pooled analysis results showed no significant differences in the incidence of total joint replacement between the low-dose tanezumab and placebo groups (RR=1.18, 95% CI=0.64–2.16,  $p=0.60$ ,  $I^2=0\%$ ). Similarly, there was no significant difference in the incidence of total joint replacement between the moderate-dose tanezumab and placebo groups (RR=1.49, 95% CI=0.82–2.72,  $p=0.19$ ,  $I^2=21\%$ ; Table 2). There was no significant difference between the two treatment groups ( $p=0.59$ ,  $I^2=0\%$ ).

**Abnormal peripheral sensation:** A total of 11 RCTs (4372 patients) reported abnormal peripheral sensation results. The results of the pooled analysis showed that the incidence of abnormal peripheral sensation was significantly higher in the tanezumab group than in the placebo group (RR=2.42, 95% CI=1.84–3.18,  $p<0.00001$ ,



**Figure 5.** Forest plot of rapidly progressive OA detailing relative risk and 95% CIs comparing tanezumab with placebo. CIs, confidence intervals; M-H, Mantel–Haenszel test; OA, osteoarthritis.

$P^2=16\%$ ). In the subgroup analysis, 7 studies were included in the low-dose group, and 11 studies were included in the moderate-dose group. The pooled results for the low-dose group showed that the incidence of abnormal peripheral sensation was significantly higher in the tanezumab group than in the placebo group (RR=2.00, 95% CI=1.21–3.29,  $p=0.006$ ,  $I^2=0\%$ ). Similarly, moderate doses of tanezumab also led to a higher risk of abnormal peripheral sensation than placebo (RR=2.64, 95% CI=1.90–3.66,  $p<0.00001$ ,  $I^2=44\%$ ; Table 2). There was no significant difference between the two treatment groups ( $p=0.35$ ,  $I^2=0\%$ ).

#### Active-controlled trials

**Efficacy. WOMAC pain score:** A total of four RCTs (2203 patients) reported WOMAC pain scores. The results of the pooled analysis showed that moderate doses of tanezumab relieved pain significantly more than the active comparator did (MD=−0.62, 95% CI=−0.85 to −0.39,  $p<0.00001$ ,  $I^2=0\%$ ; Supplemental Appendix Figure A3). However, the point estimates (MD) of the moderate-dose tanezumab group were not greater than the MCID (0.97).

**WOMAC physical score:** A total of four RCTs (2219 patients) reported WOMAC physical

scores. Pooled analysis showed that moderate doses of tanezumab resulted in greater improvements in the WOMAC physical score than did the active comparator (MD=−0.70, 95% CI=−0.92 to −0.47,  $p<0.00001$ ,  $I^2=0\%$ ; Supplemental Appendix Figure A4). However, the point estimates (MD) of the moderate-dose tanezumab group were not greater than the MCID (0.93).

**PGA:** A total of four RCTs (2218 patients) reported PGA results. The pooled estimate of PGA demonstrated significant differences in favor of moderate doses of tanezumab (MD=−0.20, 95% CI=−0.36 to −0.05,  $p=0.008$ ,  $I^2=70\%$ ; Supplemental Appendix Figure A5).

**Safety. Withdrawals due to AEs:** A total of four RCTs (2233 patients) reported withdrawals due to AEs. Pooled analysis showed no significant difference between moderate doses of tanezumab and the active comparator in terms of withdrawals due to AEs (RR=0.54, 95% CI=0.20–1.40,  $p=0.20$ ,  $I^2=83\%$ ; Supplemental Appendix Figure A6).

**SAEs:** A total of four RCTs (2233 patients) reported SAE results. The results of the pooled analysis showed no significant differences in the incidence of SAEs between the moderate-dose group and

active-comparator group (RR=0.95, 95% CI=0.67–1.34,  $p=0.76$ ,  $I^2=3\%$ ; Supplemental Appendix Figure A7).

## Discussion

The efficacy of tanezumab in treating OA has been assessed in some systematic reviews. Schnitzer and Marks<sup>34</sup> reported that the anti-NGF antibody was effective in treating hip or knee OA, and found that high doses (10 mg, and 100 or 200  $\mu\text{g}/\text{kg}$ ) of tanezumab showed greater efficacy than a placebo but were accompanied by a higher incidence of adverse reactions. Chen *et al.*<sup>35</sup> conducted a meta-analysis of studies on tanezumab in the treatment of hip or knee OA, and found that low and high doses of tanezumab had similar effects in relieving pain and improving function. However, the use of tanezumab significantly increased the incidence of withdrawals due to AEs. Similarly, Kan *et al.*<sup>36</sup> also found in their meta-analysis that the use of tanezumab could significantly relieve pain and improved function but increased the incidence of withdrawals due to AEs. However, when analyzing the incidence of AEs, the authors mixed different doses (low to high) of tanezumab and mixed studies using tanezumab monotherapy and tanezumab + other active interventions. We conducted a subgroup analysis of only low- and moderate-dose tanezumab monotherapy and did not find that tanezumab increased the incidence of withdrawals due to AEs. Therefore, high-dose tanezumab and tanezumab + other active interventions are likely to be responsible for the increased incidence of withdrawals due to AEs. The FDA also found these issues; as a result, the FDA suggests limiting the use of these drugs. Therefore, the focus of this study was to evaluate the efficacy and safety of low- and moderate-dose tanezumab monotherapy in the treatment of hip or knee OA, as well as to define the optimal dose to maximize benefits and minimize risks.

The results of this meta-analysis showed that low and moderate doses of tanezumab significantly relieved pain and improved function, and there were no significant differences in the incidence of treatment-related AEs, withdrawals due to AEs, serious AEs, and total joint replacement between the tanezumab and placebo groups; however, tanezumab increased the risk of AEs and reversible peripheral sensory abnormalities. To more accurately study the clinical efficacy of tanezumab in the treatment of OA, we also introduced the

MCID value for detailed evaluations. The clinical results based on the MCID values showed that the point estimates (MD) of moderate-dose tanezumab *versus* a placebo exceeded the MCID of the WOMAC pain and WOMAC physical function scores. However, the point estimates (MD) of low-dose tanezumab *versus* placebo were not greater than the MCID values. Thus, the clinical effect of moderate-dose tanezumab is more significant than that of low-dose tanezumab. However, the incidence of adverse reactions was similar between the two groups with different doses, and the incidence of adverse reactions did not increase with increasing dose.

There were four RCTs comparing the efficacy of moderate-dose tanezumab therapy and active-comparator therapy (two RCTs used naproxen, one RCT used oxycodone, and one RCT used naproxen or celecoxib). The final results showed that the therapeutic effects of tanezumab were significantly larger than those of the active comparator, but the MDs were smaller than those of tanezumab *versus* the placebo, and the point estimates (MD) were not greater than the MCID values for WOMAC pain and WOMAC physical function scores. In terms of safety, there were no significant differences between tanezumab and the active comparator.

The remarkable efficacy of tanezumab in the treatment of hip or knee OA is very promising. For moderate-to-severe OA of the hip or knee for whom standard analgesics were not effective or could not be taken, the onset of efficacy of subcutaneous tanezumab was within the first week, and efficacy was maintained through the 24-week treatment period.<sup>37</sup> However, because of safety risks and an inadequate plan to manage them, the European Medicines Agency and the FDA adopted a negative opinion for the tanezumab marketing authorization application. Therefore, its safety is still the current focus of attention. We evaluated the safety of low and moderate doses of tanezumab in the treatment of hip or knee OA from several aspects. Overall, the incidence rates of treatment-related AEs, withdrawals due to AEs, and SAEs with low and moderate doses of tanezumab were low, and there were no significant differences between the tanezumab groups and placebo groups or active-comparator groups. Although the incidence of AEs was higher in the tanezumab group than in the placebo group, the RR values were relatively small, and the RR values of tanezumab at low doses were similar to those at moderate doses.

Moreover, in the assessment of individual adverse reactions, there was no significant difference in the incidence of total joint replacement between tanezumab and the placebo.

Two studies reported RPOA, and there was no significant difference between tanezumab and the placebo in the subgroup analysis of low or moderate doses. But when the doses were combined, the use of tanezumab significantly increased the incidence of RPOA. At present, the pathological mechanism of RPOA caused by NGF inhibitors is still unclear. Some scholars have reported that NGF can promote tissue repair, and the increased expression of NGF in chondrocytes of osteoarthritic articular cartilage may play an important role in preventing harmful changes by regulating the proliferation and differentiation of chondrocytes. Therefore, NGF inhibitors may adversely affect bone remodeling, resulting in an increased risk of RPOA.<sup>38,39</sup> Patients with subchondral insufficiency fractures before treatment are at a greater risk of developing RPOA.<sup>40</sup> In addition, NSAIDs may also inhibit subchondral microfracture repair, which may explain why the simultaneous use of NSAIDs may increase the risk of AEs associated with anti-NGF therapy in patients with joint failure.<sup>41</sup> However, it is worth noting that the two RCTs<sup>25,30</sup> that reported the incidence of RPOA excluded predetermined imaging evidence of specific bone or joint safety (such as RPOA, atrophic or malnourished OA, subchondral fracture, spontaneous knee joint necrosis, osteonecrosis or pathological fracture, and patients with long-term use of NSAIDs). However, the final results of the meta-analysis showed that there was still an increased RPOA risk. Another explanation is that the analgesic effect of anti-NGF drugs is obvious, which may increase the load of the degenerative part of the involved joint, thus promoting the occurrence of RPOA. However, joint failure also occurs in non-weight-bearing joints, including the shoulders, and in joints other than the index joints evaluated in clinical studies.<sup>8</sup> Some scholars also believe that in most cases, the destructive arthropathy that leads to rapidly progressive joint injury may be a neuropathic joint disease since NGF may play a role in the regulation of neurosensory neurons.<sup>42</sup> Although the available data cannot completely rule out this possibility, there is a lack of evidence of severe neuropathy and loss of protective sensation in weight-bearing joints.<sup>40,42</sup> As joint-related adverse reactions include clinical entities with different radiological characteristics, there may be a variety

of mechanisms contributing to the development of RPOA.<sup>43</sup> Therefore, more research is needed to explore the causes of RPOA.

In terms of abnormal peripheral sensation, our results showed that compared with a placebo, both low and moderate doses of tanezumab could significantly increase the incidence of abnormal peripheral sensation (including abnormal sensation, hypoesthesia, and hypersensitivity). However, the severity of this condition is usually rated as mild or moderate, and most of these cases are short-lived and do not have any permanent sequelae. Most commonly, this sensory disorder disappears within 1 to 2 weeks or 1 month after the first dose.<sup>23,28,44,45,46</sup> In general, most patients with a final neurological diagnosis of new or worsening peripheral neuropathy are diagnosed with some form of mononeuropathy (mainly carpal tunnel syndrome) or radiculopathy based on clinically significant signs or diagnostic tests. Very few patients are diagnosed with multiple neuropathies.<sup>28,29,32,44,45,47</sup> These results do not reflect the expected pattern for neurotoxic compounds, which typically cause prolonged durations of multiple neuropathies in affected patients. The association of tanezumab with mononeuropathy symptoms suggests that these manifestations may be the result of the inhibitory effect of NGF.<sup>31,48</sup> An RCT that evaluated the safety of tanezumab in the treatment of OA showed that when tanezumab was used to treat chronic pain in individuals without known peripheral neuropathies, it was not associated with structural neurotoxic effects on large motor or sensory nerves, autonomic nerves, or small sensory nerve fibers in the skin.<sup>27</sup> Therefore, neurosensory symptoms induced by anti-NGF therapy may be caused by reversible functional changes or homeostasis in peripheral nerve activity. A short-term follow-up study of tanezumab in the treatment of two types of peripheral neuropathic pain also showed that patients' neurological condition did not deteriorate with treatment.<sup>49</sup> In addition, no significant adverse reactions of tanezumab were found in other assessments, such as cardiovascular, liver, renal function, gastrointestinal, and other types of laboratory tests.<sup>22,31</sup> Therefore, tanezumab is generally safe. Cost-effectiveness analysis showed that the pain relief provided by anti-NGF therapy was significant enough that even rapid progressive OA, which was present in 10% of patients, did not offset the overall improvement in quality-adjusted life years achieved<sup>50</sup> and that the cost of anti-NGF therapy might be as low as \$400 per dose.<sup>50</sup> Of course, it



should be noted that these analyses are based entirely on models that used arbitrary values of the costs related to pain and are intended to inform policy makers rather than clinicians, as individuals have significantly different perspectives on risks and benefits.<sup>51</sup>

At present, the existing research on tanezumab in the treatment of hip or knee OA is relatively extensive; thus, the total sample size studied is relatively large, and the final results are precise. Moreover, the homogeneity of the whole study is relatively good. However, some shortcomings of this article are still noteworthy. First, our study is based on only English articles and does not include non-English studies, so the results may be affected by publication bias. However, Egger's test results indicate that there is no risk of publication bias. For the risk assessment of RPOA, only 2 studies were included, and more studies still need to be conducted in the future. Finally, there is a known potential risk, and all of the studies included were sponsored by pharmaceutical companies.

### Conclusion

Based on the current research evidence, lower doses of tanezumab monotherapy can significantly improve pain, function, and PGA in patients with hip or knee OA. Tanezumab is relatively well tolerated and safe, but increases the incidence of AEs and reversible abnormal peripheral sensation. The occurrence of RPOA is still worthy of attention, and more studies are needed to explore this aspect. In terms of dose selection, moderate doses of tanezumab have a more pronounced clinical effect than low doses of tanezumab, and the safety and tolerance were similar between the two doses, with no significant differences. Therefore, a moderate dose of tanezumab may maximize the benefits for hip or knee OA.

### Author contributions

**Di Zhao:** Data curation; Methodology; Writing – original draft; Writing – review & editing.

**Minghui Luo:** Data curation; Methodology; Writing – original draft; Writing – review & editing.

**Jian-ke Pan:** Data curation; Methodology; Writing – original draft; Writing – review & editing.

**Ling-feng Zeng:** Data curation; Methodology; Writing – review & editing.

**Gui-hong Liang:** Data curation; Writing – original draft.

**Yan-hong Han:** Data curation; Writing – review & editing.

**Jun Liu:** Data curation; Funding acquisition; Methodology; Project administration; Supervision; Validation; Writing – original draft; Writing – review & editing.

**Weiyi Yang:** Data curation; Methodology; Project administration; Supervision; Writing – original draft; Writing – review & editing.


### Conflict of interest statement

The authors declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

### Funding

The authors disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: This study was supported by the National Natural Science Foundation of China (No. 81873314, No. 81473698, No. 82004386, No. 82004383), the Science and Technology Research Project of Guangdong Provincial Hospital of Chinese Medicine (No. YN2019ML08, No. YN2015MS15), Science and Technology Program of Guangzhou (No. 202102010273), the Project of Guangdong Provincial Department of Finance (No. [2014]157, No. [2018]8), and the Science and Technology Planning Project of Guangdong Province (No. 2020A1414050050).

### ORCID iD

Jun Liu  <https://orcid.org/0000-0002-2440-4064>

### Supplemental material

Supplemental material for this article is available online.

### References

1. Glyn-Jones S, Palmer AJ, Agricola R, *et al.* Osteoarthritis. *Lancet* 2015; 386: 376–387.
2. Rosemann T, Wensing M, Joest K, *et al.* Problems and needs for improving primary care of osteoarthritis patients: the views of patients, general practitioners and practice nurses. *BMC Musculoskelet Disord* 2006; 7: 48.
3. Bannuru RR, Osani MC, Vaysbrot EE, *et al.* OARSI guidelines for the non-surgical management of knee, hip, and polyarticular osteoarthritis. *Osteoarthritis Cartilage* 2019; 27: 1578–1589.



4. Nelson AE, Allen KD, Golightly YM, *et al.* A systematic review of recommendations and guidelines for the management of osteoarthritis: the chronic osteoarthritis management initiative of the U.S. bone and joint initiative. *Semin Arthritis Rheum* 2014; 43: 701–712.
5. Kalso E, Aldington DJ and Moore RA. Drugs for neuropathic pain. *BMJ* 2013; 347: f7339.
6. Turk DC, Wilson HD and Cahana A. Treatment of chronic non-cancer pain. *Lancet* 2011; 377: 2226–2235.
7. Mantyh PW, Koltzenburg M, Mendell LM, *et al.* Antagonism of nerve growth factor-TrkA signaling and the relief of pain. *Anesthesiology* 2011; 115: 189–204.
8. Bannwarth B and Kostine M. Targeting nerve growth factor (NGF) for pain management: what does the future hold for NGF antagonists? *Drugs* 2014; 74: 619–626.
9. Cohen SP and Mao J. Neuropathic pain: mechanisms and their clinical implications. *BMJ* 2014; 348: f7656.
10. Eskander MA, Ruparel S, Green DP, *et al.* Persistent nociception triggered by nerve growth factor (NGF) is mediated by TRPV1 and oxidative mechanisms. *J Neurosci* 2015; 35: 8593–8603.
11. Aloe L, Tuveri MA, Carcassi U, *et al.* Nerve growth factor in the synovial fluid of patients with chronic arthritis. *Arthritis Rheum* 1992; 35: 351–355.
12. Shakoor N, Lee KJ, Fogg LF, *et al.* The relationship of vibratory perception to dynamic joint loading, radiographic severity, and pain in knee osteoarthritis. *Arthritis Rheum* 2012; 64: 181–186.
13. Hefti FF, Rosenthal A, Walicke PA, *et al.* Novel class of pain drugs based on antagonism of NGF. *Trends Pharmacol Sci* 2006; 27: 85–91.
14. Malfait AM and Schnitzer TJ. Towards a mechanism-based approach to pain management in osteoarthritis. *Nat Rev Rheumatol* 2013; 9: 654–664.
15. Ashraf S, Bouhana KS, Pheneger J, *et al.* Selective inhibition of tropomyosin-receptor-kinase A (TrkA) reduces pain and joint damage in two rat models of inflammatory arthritis. *Arthritis Res Ther* 2016; 18: 97.
16. Rahman W and Dickenson AH. Emerging targets and therapeutic approaches for the treatment of osteoarthritis pain. *Curr Opin Support Palliat Care* 2015; 9: 124–130.
17. Abdiche YN, Malashock DS and Pons J. Probing the binding mechanism and affinity of tanezumab, a recombinant humanized anti-NGF monoclonal antibody, using a repertoire of biosensors. *Protein Sci* 2008; 17: 1326–1335.
18. Patel MK, Kaye AD and Urman RD. Tanezumab: therapy targeting nerve growth factor in pain pathogenesis. *J Anaesthesiol Clin Pharmacol* 2018; 34: 111–116.
19. Liberati A, Altman DG, Tetzlaff J, *et al.* The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate healthcare interventions: explanation and elaboration. *BMJ* 2009; 339: b2700.
20. Higgins JP, Altman DG, Gotzsche PC, *et al.* The Cochrane Collaboration’s tool for assessing risk of bias in randomised trials. *BMJ* 2011; 343: d5928.
21. Ehrich EW, Davies GM, Watson DJ, *et al.* Minimal perceptible clinical improvement with the Western Ontario and McMaster Universities osteoarthritis index questionnaire and global assessments in patients with osteoarthritis. *J Rheumatol* 2000; 27: 2635–2641.
22. Ekman EF, Gimbel JS, Bello AE, *et al.* Efficacy and safety of intravenous tanezumab for the symptomatic treatment of osteoarthritis: 2 randomized controlled trials versus naproxen. *J Rheumatol* 2014; 41: 2249–2259.
23. Lane NE, Schnitzer TJ, Birbara CA, *et al.* Tanezumab for the treatment of pain from osteoarthritis of the knee. *N Engl J Med* 2010; 363: 1521–1531.
24. Nagashima H, Suzuki M, Araki S, *et al.* Preliminary assessment of the safety and efficacy of tanezumab in Japanese patients with moderate to severe osteoarthritis of the knee: a randomized, double-blind, dose-escalation, placebo-controlled study. *Osteoarthritis Cartilage* 2011; 19: 1405–1412.
25. Berenbaum F, Blanco FJ, Guermazi A, *et al.* Subcutaneous tanezumab for osteoarthritis of the hip or knee: efficacy and safety results from a 24-week randomised phase III study with a 24-week follow-up period. *Ann Rheum Dis* 2020; 79: 800–810.
26. Birbara C, Dabezies EJ, Burr AM, *et al.* Safety and efficacy of subcutaneous tanezumab in patients with knee or hip osteoarthritis. *J Pain Res* 2018; 11: 151–164.
27. Brown MT, Herrmann DN, Goldstein M, *et al.* Nerve safety of tanezumab, a nerve growth factor inhibitor for pain treatment. *J Neurol Sci* 2014; 345: 139–147.

28. Brown MT, Murphy FT, Radin DM, *et al.* Tanezumab reduces osteoarthritic knee pain: results of a randomized, double-blind, placebo-controlled phase III trial. *J Pain* 2012; 13: 790–798.
29. Brown MT, Murphy FT, Radin DM, *et al.* Tanezumab reduces osteoarthritic hip pain: results of a randomized, double-blind, placebo-controlled phase III trial. *Arthritis Rheum* 2013; 65: 1795–1803.
30. Schnitzer TJ, Easton R, Pang S, *et al.* Effect of tanezumab on joint pain, physical function, and patient global assessment of osteoarthritis among patients with osteoarthritis of the hip or knee: a randomized clinical trial. *JAMA* 2019; 322: 37–48.
31. Schnitzer TJ, Ekman EF, Spierings EL, *et al.* Efficacy and safety of tanezumab monotherapy or combined with non-steroidal anti-inflammatory drugs in the treatment of knee or hip osteoarthritis pain. *Ann Rheum Dis* 2015; 74: 1202–1211.
32. Spierings EL, Fidelholtz J, Wolfram G, *et al.* A phase III placebo- and oxycodone-controlled study of tanezumab in adults with osteoarthritis pain of the hip or knee. *Pain* 2013; 154: 1603–1612.
33. Jonsson EN, Xie R, Marshall SF, *et al.* Population pharmacokinetics of tanezumab in phase 3 clinical trials for osteoarthritis pain. *Br J Clin Pharmacol* 2016; 81: 688–699.
34. Schnitzer TJ and Marks JA. A systematic review of the efficacy and general safety of antibodies to NGF in the treatment of OA of the hip or knee. *Osteoarthritis Cartilage* 2015; 23(Suppl. 1): S8–S17.
35. Chen J, Li J, Li R, *et al.* Efficacy and safety of tanezumab on osteoarthritis knee and hip pains: a meta-analysis of randomized controlled trials. *Pain Med* 2017; 18: 374–385.
36. Kan SL, Li Y, Ning GZ, *et al.* Tanezumab for patients with osteoarthritis of the knee: a meta-analysis. *PLoS ONE* 2016; 11: e157105.
37. Berenbaum F, Langford R, Perrot S, *et al.* Subcutaneous tanezumab for osteoarthritis: is the early improvement in pain and function meaningful and sustained? *Eur J Pain* 2021; 25: 1525–1539.
38. Holmes D. Anti-NGF painkillers back on track? *Nat Rev Drug Discov* 2012; 11: 337–338.
39. Seidel MF, Herguijuela M, Forkert R, *et al.* Nerve growth factor in rheumatic diseases. *Semin Arthritis Rheum* 2010; 40: 109–126.
40. Hochberg MC, Tive LA, Abramson SB, *et al.* When is osteonecrosis not osteonecrosis? Adjudication of reported serious adverse joint events in the tanezumab clinical development program. *Arthritis Rheumatol* 2016; 68: 382–391.
41. Teichtahl AJ and Cicuttini FM. Editorial: pain relief in osteoarthritis: the potential for a perfect storm. *Arthritis Rheumatol* 2016; 68: 270–273.
42. Bannwarth B and Kostine M. Nerve growth factor antagonists: is the future of monoclonal antibodies becoming clearer? *Drugs* 2017; 77: 1377–1387.
43. Schmelz M, Mantyh P, Malfait AM, *et al.* Nerve growth factor antibody for the treatment of osteoarthritis pain and chronic low-back pain: mechanism of action in the context of efficacy and safety. *Pain* 2019; 160: 2210–2220.
44. Kivitz AJ, Gimbel JS, Bramson C, *et al.* Efficacy and safety of tanezumab versus naproxen in the treatment of chronic low back pain. *Pain* 2013; 154: 1009–1021.
45. Katz N, Borenstein DG, Birbara C, *et al.* Efficacy and safety of tanezumab in the treatment of chronic low back pain. *Pain* 2011; 152: 2248–2258.
46. Sanga P, Polverejan E, Wang S, *et al.* Efficacy, safety, and tolerability of fulranumab as an adjunctive therapy in patients with inadequately controlled, moderate-to-severe chronic low back pain: a randomized, double-blind, placebo-controlled, dose-ranging, dose-loading phase II study. *Clin Ther* 2016; 38: 1435–1450.
47. Balanescu AR, Feist E, Wolfram G, *et al.* Efficacy and safety of tanezumab added on to diclofenac sustained release in patients with knee or hip osteoarthritis: a double-blind, placebo-controlled, parallel-group, multicentre phase III randomised clinical trial. *Ann Rheum Dis* 2014; 73: 1665–1672.
48. Tive L, Bello AE, Radin D, *et al.* Pooled analysis of tanezumab efficacy and safety with subgroup analyses of phase III clinical trials in patients with osteoarthritis pain of the knee or hip. *J Pain Res* 2019; 12: 975–995.
49. Bramson C, Herrmann DN, Carey W, *et al.* Exploring the role of tanezumab as a novel treatment for the relief of neuropathic pain. *Pain Med* 2015; 16: 1163–1176.
50. Losina E, Michl G, Collins JE, *et al.* Model-based evaluation of cost-effectiveness of nerve growth factor inhibitors in knee osteoarthritis: impact of drug cost, toxicity, and means of administration. *Osteoarthritis Cartilage* 2016; 24: 776–785.
51. Miller RE, Block JA and Malfait AM. What is new in pain modification in osteoarthritis? *Rheumatology* 2018; 57(Suppl. 4): v99–v107.